Cellular and Molecular biology in oesophageal cancer

Alfred Lam, MBBS, MD, PhD, FRCPA

INTROUDCTION

Histological differences

Oesophageal cancers comprises cancers of different histological types with diverse cellular and molecular bases [1-2]. The major histological types of oesophageal cancers are either squamous cell carcinoma or adenocarcinoma. It is important to note that there are histological variants of squamous cell carcinoma and adenocarcinoma like basaloid squamous cell carcinoma, spindle cell carcinoma, mucoepidermoid carcinoma and adenosquamous carcinoma [3-5]. In addition, neuroendocrine neoplasms such as small cell carcinoma of the oesophagus accounted for approximately 1% of primary oesophageal carcinoma [6]. All these carcinomas have distinct clinicopathological features. Limited studies revealed that the cellular and molecular biology of these uncommon types of oesophageal carcinomas are different from oesophageal squamous cell carcinoma or adenocarcinoma [3, 7-8].

In practice, the current understanding of cellular and molecular biology of oesophageal cancers focuses on oesophageal squamous cell carcinoma and oesophageal adenocarcinoma. The difference in prevalence of these two major histological types in different geographic regions is likely as a result of the complex interactions of genetic and environmental factors. In general, oesophageal squamous cell carcinoma predominates in areas with high incidence of oesophageal cancer whereas oesophageal adenocarcinoma is more common in areas with low incidence of oesophageal cancer. Also, the genetic mechanisms of oesophageal squamous cell carcinoma are complex as multiple genetic factors have been proposed for carcinogenesis of the squamous cell

carcinoma [2]. On the other hand, most of the oesophageal adenocarcinomas are known to be related to the genetic changes of the progression of lesions related to acid reflux. The histological progression from reflux oesophagitis to Barrett' metaplasia to dysplasia to adenocarcinoma is well known.

Applications of molecular and cellular biology

Oesophageal cancer remains one of the leading causes of cancer death worldwide despite the improvements in surgical and adjuvant therapies. The only hope to further increase the quality of life of patients with oesophageal cancer is to understand and apply the knowledge of cellular and molecular biology in the clinical management of these cancers. Thus, the study of cellular and molecular biology in oesophageal cancer could potentially serve the following purposes: (1) to establish the presence or absence of an infectious co-factor; (2) to understand the genetic mechanisms; (3) to provide prognostic information; and (4) to predict response to medical therapies and new modalities of treatment. In performing researches, interpreting and applying the knowledge in this area, it is important to bear in mind the histological difference in oesophageal cancers. At current, most of the studies in this cancer are still on the understanding the genetic mechanisms rather than the applications of the knowledge.

ESTABLISH INFECTIOUS CO-FACTOR

Human papilloma virus

In oesophageal cancers, the main infectious co-factor under intensive studies was human papilloma virus (HPV). In squamous cell carcinomas from the upper aerodigestive tract, in particular in oral regions and oropharynx, identification of presence of human papilloma virus (HPV) in the carcinomas is of important values [9-11]. Patients with

cancers having HPV detected have better prognosis when compared to patients with cancers without HPV. The detection of HPV in squamous cell carcinomas from the oral and oropharynx also predicts better response to radiotherapy. Also, the detection of HPV in clinical setting is indirectly by the identification of increased expression of p16 protein by immunohistochemistry.

Oesophagus is distal to the oropharynx and histologically lined by stratified squamous epithelium as in the oral regions or oropharynx. Researches to investigate HPV in squamous cell carcinomas in oesophagus were noted for 30 years [12-13]. Thus, there is considerable data on the role of HPV infection in the development of oesophageal cancer. Majority of the studies were performed in oesophageal squamous cell carcinoma.

It is worth noting some data were obtained in oesophageal adenocarcinoma. Pooled - analysis from the literature revealed that HPV prevalence in oesophageal adenocarcinoma was 35.0% and HPV-16 prevalence was 11.4% [14]. Due to the limited number of studies on oesophageal adenocarcinoma, no detailed analysis of the impact was available. Nevertheless, it has postulated that progressive acid damage to the oesophagus increases the likelihood of mucosal breaks and allows the virus to enter the basal layer of the transformation zone. Recently, transcriptionally active HPV was noted be strongly associated with Barrett dysplasia and oesophageal adenocarcinoma, suggesting a potential role of HPV in oesophageal carcinogenesis.

In oesophageal squamous cell carcinoma, summarised HPV prevalence from both early and recent meta-analysis in oesophageal squamous cell carcinoma was 22% [14]. In general, HPV prevalence was found to be higher in the studies some countries from Asia and was much lower in studies reported in western countries such as in Europe and America [2]. Stratified analysis by localisation of cancer showed that oesophageal squamous cell carcinoma in was only slightly higher but not significantly higher than the

middle or lower portion of oesophagus [14-15].

With respect to HPV DNA detection in meta-analysis, the prevalence of oesophageal squamous cell carcinoma produced by type-specific primer PCR method (30.4%) was observed to be significantly higher than that by broad-spectrum primers (20.8%) [14]. P16 protein detection in by immunohistochemical method was not often studied. Nevertheless, the current data on using p16 detection in oesophageal squamous cell carcinoma did not reflect the HPV status in the cancer [16].

From pooled data, HPV-16 was the most frequently observed subtype with a summarised prevalence of 11.4% [2, 14]. The other six most frequent HPV individual types identified in oesophageal squamous cell carcinoma, in order of decreasing prevalence, were HPV-18 (2.9%), HPV-6 (2.1%), HPV-11(2.0%), HPV-52 (1.1%), HPV-33 (0.8%) and HPV-31 (0.6%).

Overall, human papillomavirus infection was observed to be associated with an increased risk of oesophageal squamous cell carcinoma. However, the association was not as strong as oral/oropharyngeal squamous cell carcinoma or cervical squamous cell carcinoma. In addition, it is unlikely that HPV detection and/or p16 expression will predict radiotherapy response as it does for upper aerodigestive tract squamous cell carcinoma. At this stage, routine evaluation of HPV status and/or p16 is not required in management of the cancer.

Epstein bar virus

The association of Epstein bar virus (EBV) has also been studied in oesophageal cancer. The results are variable with prevalence detected ranging from 0% to 35% [17-18]. The differences are likely to be related to variety of racial, geographical and detection methods used. It is worth noting the lymphocytes in the cancer stroma can harbour EBV and

thus detection of virus in oesophageal cancer by PCR based methods may shows false-positive results [18]. On the other hand, immunohistochemistry *in situ* hybridization may provide false negative result due to higher rate of RNA degradation. Most studies have shown that EBV-associated oesophageal cancer demonstrates similar morphologic findings to undifferentiated carcinoma of nasopharynx which is known to be associated with EBV. At the current status, more data is needed for the study of EBV in oesophageal cancer. It has no clinical application at the moment.

Bacteria

Helicobacter pylori is a Gram-negative spiral bacterium that is present in nearly half the world's population. It is the major cause of peptic ulcer disease and a recognized cause of gastric carcinoma. Some strains of *Helicobacter pylori* may protect patients from gastroesophageal reflux disease and oesophageal adenocarcinoma [17]. Overall, there is no consensus on the role of *Helicobacter pylori* in oesophageal/cardia adenocarcinoma, with substantial differences between Asian and Western studies. Other than this, many bacteria have been identified in the oesophagus by metagenomics [17, 19]. Metagenomics is the study of microbiota in their natural habitat using next-generation sequencing through a PCR-based analysis of bacterial 16S rRNA genes. Two distinct clusters: a predominantly Gram-positive cluster (type I) and a predominately Gram-negative cluster (type II) were found. Type II cluster may stimulate expression of different proteins and genes leading to reflux and trigger the process of adenocarcinoma.

UNDERSTANDING GENETIC MECHANISMS

Genetic profiles

The genetic profiles of oesophageal cancers have been studied for decades. The

roles of oncogenes, tumour suppressor genes, metastatic genes, apoptosis genes, proliferation related factors, epigenetic factors as well as proteins related to metastases have been studied in both oesophageal squamous cell carcinoma and oesophageal adenocarcinoma [2, 20]. It is worth noting that adenocarcinoma from the oesophagogastric junction share similar histological features and profiles. However, some of the genetic changes may be related to gastric cardia. In general, for both histological subtypes, p53 mutation is an important genetic change [21-22].

Starting from around 2000, comparative genomic hybridization (CGH) and expression array were used to identify the different in genetic profiles of oesophageal cancer when compared to non-cancer oesophageal tissue [23-26]. Chromosomal regions with amplification may harbor oncogenes, and chromosomal regions with deletion may harbor tumour-suppressor genes. CGH can identify the whole profile of cytogenetic changes in an individual cancer. By this approach, novel genes were identified in both oesophageal squamous and oesophageal adenocarcinoma [27-31]. These provide more information of the carcinogenesis of oesophageal cancers as well as finding gene candidates as prognostic markers and molecular targets for therapy.

The introduction of next-generation sequencing has brought forth the ability to generate vast quantities of genetic data at a low cost [32]. These recent technologies allow researchers to sequence DNA and RNA much more quickly and economically than the previously used Sanger sequencing, and as such have revolutionised the study of genomics and molecular biology. The first commercial available next-generation sequencer was available in 2007. In addition, the International Cancer Genome Consortium (ICGC) has been organized to launch and coordinate a large number of research projects that have the common aim of elucidating comprehensively the genomic changes present in many cancers [33]. The preliminary meeting was held in 2007 and the consortium launched to

the public notice in 2010. The primary goals of the ICGC are to generate comprehensive catalogues of genomic abnormalities (somatic mutations, abnormal expression of genes, epigenetic modifications).

The oesophageal squamous cell carcinoma genome study is being taken care of by researchers in China whereas the oesophageal adenocarcinoma genome study is by researchers in United Kingdom. It is likely the more comprehensive knowledge of the genetic events in the oesophageal carcinoma will be available soon. The whole genome sequencing data on oesophageal cancer started to appear in the literature in 2010 [34-37]. At current, a few studies were noted in literature on both oesophageal squamous cell carcinomas and adenocarcinomas. Data were seen in the website of the ICGC. A large volume of information is available for both histological subtypes of oesophageal cancer. They provide huge resources for future research directions for better management of patients with oesophageal carcinoma

In oesophageal squamous cell carcinoma, eight significantly mutated genes, of which six are well known tumour-associated genes (*TP53*, *RBI*, *CDKN2A*, *PIK3CA*, *NOTCH1*, *NFE2L2*), and two have not previously been described in oesophageal squamous cell carcinomas (*ADAM29* and *FAM135B*) [34]. Genomic analyses suggest that oesophageal squamous cell carcinoma and head/neck squamous cell carcinoma share some common pathogenic mechanisms.

In oesophageal adenocarcinoma, genes like *p53* and *SMAD4* were confirmed to be implicated in stage-specific manner and confined to high grade dysplasia and adenocarcinoma [35]. Many of the recurrently mutated genes in oesophageal adenocarcinoma were also mutated in non-dysplastic Barrett epithelium. Thus, mutations in oesophageal adenocarcinoma driver genes generally occur exceptionally early in disease development with profound implications for diagnostic and therapeutic strategies.

Also, new significantly mutated genes include chromatin-modifying factors and candidate contributors *SPG20*, *TLR4*, *ELMO1* and *DOCK2* were also identified.

MicroRNAs (miRNAs)

MicroRNAs (miRNAs) are a class of small, well-conserved, non-coding RNAs that regulate the translation of RNAs. Their biological and pathological roles and have been studies in numerous cancers [38-41]. Research works were also being done to see if the miRNA can be detected in serum of the patients with oesophageal cancer [42-43]. Studies showed that there are a number of candidate miRNAs that have so far been found to be up or down-regulated at various stages in the progression from Barrett oesophagus to oesophageal adenocarcinoma [44-47]. In oesophageal squamous cell carcinoma, many miRNAs were identified as diagnostic and prognostic markers [45-47]. It may be related to the fact that oesophageal squamous cell carcinoma has a more complex carcinogenesis than oesophageal adenocarcinoma. However, further work is required in order to be able to use miRNAs for clinical use.

Cancer stem cells

Cancer stem cells, as a subgroup of cancer cells, resemble critical properties of embryonic stem cells such as self-renewal and maintenance of stemness state. Only the cancer stem cells (CSCs) have tumour-initiating properties [48]. They play an important role in the resistance to adjuvant therapies for cancer. Also, epithelial-mesenchymal transition (EMT) may be involved in epithelial cell immortalization and enrichment of stemness. These immortal cells may regain their original properties via mesenchymal-epithelial transition (MET) and maintain epithelial stem cell properties [49]. In oesophageal squamous cell carcinoma, a few stem cell markers like CD44, ALDH1A1, SALL4 and SOX2 have been

identified [50]. The expressions of some of these markers were noted to be correlated to cancer staging. Cancer stem cells or tumour-initiating cells are present in oesophageal adenocarcinoma [51]. However, antibodies directed against novel surface antigens are needed to detect them.

Development of specific markers and signalling molecules to target the stem cells of oesophageal carcinomas and the validation of these stem cells might provide the basis for a revolutionary treatment approach for the elimination and /or differentiation of cancer stem cells in oesophageal cancer. Emerging therapeutic tools based on specific properties and function of CSCs may improve clinical output of the diseases. Therefore, innovative insight into CSC biology and CSC-targeted therapies will help to achieving effective management of oesophageal cancers.

PROGNOSTIC INFORMATION

Predication of progression

Aneuploidy (detected by FISH/flow cytometry), promoter hypermethylation (methylation profiles such as p16, APC, etc) and cyclin A protein expression has been shown to be correlated to the progression from Barrett oesophagus to oesophageal adenocarcinoma [52]. Despite these results, there is generally lack of large prospective study to validate the use of these markers in clinical practice. The likely candidate of clinical application is p53 protein over-expression as determined by immunohistochemistry. Overexpression of p53 protein was noted to be correlated with the neoplastic progression to oesophageal adenocarcinoma. It could be a useful adjunct to determine the grade of dysplasia in Barrette oesophagus. Also the results have been validated in some studies and the procedure used is simple.

Many data exists for oesophageal adenocarcinoma and has found that expression or

identification of cellular and molecular markers could predict survival of the patients with oesophageal adenocarcinoma [53, 54]. Some of the more commonly described markers are epidermal growth factor receptors (1 and 2); Transforming growth factor (TGF α and β 1); P53, Ki-67, cyclin dependent kinase inhibitor 1 (p21); B-cell lymphoma 2 (bcl-2); Cyclooxygenase-2 (COX-2); Nuclear factor- κ B (NF- κ B); vascular endothelial growth factor (VEGF); tissue inhibitor of metalloproteinase (TIMP) and microsatellite instability (MSI). At present, routine testing for all or any of these markers is not warranted as none of these markers have been validated adequately in prospective studies.

In oesophageal squamous cell carcinoma, many molecular and cellular markers have been proposed to be related to the prognosis of the patients with cancer [54-59]. Expressions of p21, p53, cyclin D1, Ki-67, E-cadherin and VEGF have been shown to provide some prognostic information. However, this method has not been widely used.

GUIDLLINES FOR MEDICAL THERAPIES

Predict response to medical therapies

Pre-operative chemo-radiation is adopted as a standard way of treatment of oesophageal cancers. In patients who underwent neoadjuvant chemo-radiation therapy, histological regression of the primary cancer indicated by percentage of residual viable cells is an important prognostic factor in addition to nodal status and gender [60].

It is thus important to have means to predict the response for chemo-radiation. It has been reported that grade of oesophageal squamous cell carcinoma can predict the response to pre-operative chemotherapy [61]. Many molecular makers have been studied [62-65]. p53 protein is expected to be a representative biomarker. The cell cycle markers- CDC25B and 14-3-3sigma have potential as response biomarkers independent of the p53 status. The DNA repair markers, p53R2 or ERCC1, VEGF and hedgehog

signalling pathway factor Gli-1 also have potential as predictive biomarkers. However, proper validations are needed. In oesophageal adenocarcinoma, expression of EGFR, vascular endothelial growth factor, nuclear factor- κ B and cDNA microarray are noted to acting as predictive factor for pre-operative chemo-radiation.

Predictors for targeted therapy

Targeted therapy is currently being investigated as an addition to these regimes and for the treatment of metastatic disease. The therapy usually involves targeting a specific gene mutation in the oesophageal cancer. Promising results were noted in melanoma, breast cancer and colorectal cancer [66-70]. Testing the cancer tissues for molecular markers is useful to predict the response of the patients to these targeted therapies.

Of the potential targets trialled to date in oesophageal cancer, epithelial growth factor receptor 2 - EGFR (HER1 and HER2) and vascular endothelial growth factor – VEGF surface receptor antagonists have shown the most promise [71-75]. Of these, Her 2 over expression has been most extensively studied. Trastuzumab-based therapy offered a significant survival advantage for patients with HER2 overexpressing locally advanced, recurrent or metastatic gastric and gastro- oesophageal junctional adenocarcinomas compared to conventional therapy alone. Currently the relevance in oesophageal adenocarcinomas and squamous carcinomas is unclear. It is likely that pathologists will be required to determine the HER 2 status in biopsy or resection material in gastroesophageal junction as well as in metastatic sites. HER 2 status could be assessed by immunohistochemistry and FISH testing. Other targeted therapies in oesophageal cancers are less well studied. For instance, over expression of EGFR -1 is found in 1/3 to 2/3 of oesophageal adenocarcinoma and squamous cell carcinoma. To date, clinical trials using the EGFR-1 antagonist, cetuximab, have revealed varied results. VEGF is over expressed in 30 – 60%

of oesophageal adenocarcinoma. VEGF antagonist bevacizumab has shown promising results when added to multimodality treatment. However, at this stage, there is not much data of the use of these therapies and predictive markers in oesophageal cancers.

RESEARCH SOURCES OF MOLECULAR AND CELLULAR STUDIES IN OESOPHAGEAL CANCERS

Tissues studies

Human cancer can be studied at tissue level when it is surgically removed from human body. These cancer tissues are without blood supply and degeneration will quickly occur. Cancer studies on these tissues can be performed in several ways. In clinical settings, the cancer tissues were fixed in formalin and embedded in paraffin. They were useful for different molecular studies. In fact, many of the research findings in oesophageal cancers were obtained by this method. This approach has the benefit of providing superior morphological features for studying the histological features as well as localization of biomarkers in the cellular level when compared with other methods (Figure 1).

In the recent years, tissue microarray (TMA) was increasing use in testing molecular markers in large number of samples by either immunohistochemistry or FISH (Figure 2). In the tissue microarray technique, a hollow needle is used to remove tissue cores as small as 0.6 mm in diameter from regions of interest in paraffin-embedded tissues. These tissue cores are then inserted in a recipient paraffin block in a precisely spaced, array pattern.

The drawback of working on paraffin-embedded tissues is formalin irreversibly cross-links proteins via the amino groups thus preserving the structural integrity of the cells so they can be stained with dyes used to analyse for abnormalities in the tissue that indicate cancer. The effect of these cross-linking fixatives on the nucleic acids and proteins will impair the molecular works. In recent decades, oesophageal cancer tissues were prospectively collected.

The cancer tissue snapped frozen the oesophageal cancer tissues in liquid nitrogen and stored the tissue in -80°C. The collection included a lot of clinical and scientific efforts and the tissues were superior in quality for molecular works. On the other hand, the morphological features are inferior to those obtained by paraffin embedded sections (Figure 3).

It is worth noting that research by these approaches could not provide functional dynamic studies of oesophageal cancers.

Cancer cell lines

For functional studies in oesophageal cancer, the works could be done on cancer cell lines derived from tissues obtained fresh after the operation. Cancer cell lines need proper medium to grow. Cancer cell lines often grow without attaching to a surface and they can proliferate to a very much higher density in a culture dish. The resulting transformed cancer cell lines, in reciprocal fashion, can often cause tumours if injected into a susceptible animal for making animal model. Cancer cell can be harvested from the animal and forming a more stable cell cancer cell line. In oesophageal cancers, some of the common used cell lines actually are secondary cell line. The cancer cell lines can allow functional studies to be done. They can be stored in liquid nitrogen for an indefinite period and retain their viability when thawed.

In oesophageal cancers, there are published cancer cell lines available for both adenocarcinoma and squamous carcinoma [76-79]. When compared to oesophageal squamous cell carcinoma, oesophageal adenocarcinoma is more uniform in characteristics as the risk factors and pathogenesis more established. Model research on oesophageal adenocarcinoma relies almost entirely on a relatively small set of established cancer cell lines. The high genomic similarities between the oesophageal cell lines and their original

cancers provide rationale for their use. Nonetheless, cancer cell lines nearly always differ in important ways from the original cancer from which they were derived.

Animal model

Animal model is important to study the effects of cancer in vivo and also production of cancer cell lines. In more clinically relevant applications, animal model is a must for developing therapeutic strategies. Cancer development is a complex process with accumulation of genetic alterations and their downstream effects as well as interactions of the microenvironment in different tissues. The cancer microenvironment and its interactions with the cancer are important in determining growth dynamics of different cancers.

Injection of cancer or cancer cells in the subcutaneous tissue of the skin of immunodeficiency mice is a common practice to produce a cancer model in animal (Figure 4). In oesophageal cancers, this approach cannot recapitulate the microenvironment of the oesophagus as well as the response to the targeting carcinogens. The current approach is to make an orthotopic (occurring at normal place) model for both oesophageal squamous cell carcinoma and oesophageal adenocarcinoma [80-81]. The orthotopic model provides the most optimum environment for cancer growth and drug testing. In the anatomical setting of oesophageal cancer, the site is very difficult to approach surgically. Different approaches have been applied but many of these had some shortcomings. The establishment of these orthotopic models need to involve radiological guidance (magnetic resonance imaging and fluorescence imaging) so the cancer and the metastases can be visualized in real time [82]. In addition, pathological examination is important to clarify the histological typing, microscopic location and the microenvironment of the cancer in the animal [83].

Figures Legends

Figure 1. Histological features of oesophageal cancer prepared by formalin fixation and paraffin embedded revealing a well-differentiated squamous cell carcinoma with good histological details

Figure 2. Tissue microarray of oesophageal squamous cell carcinoma showing multiple cores of cancer tissue in a block to allow testing of molecular markers simultaneously of many cases in one run of experiment.

Figure 3. Histological features of oesophageal cancer prepared by sectioning of frozen tissues showing inferior histological details when compared to Figure 1.

Figure 4. Tumour produced on an immunodeficiency mouse after injection of primary oesophageal squamous cell carcinoma in the subcutaneous

REFERENCES

1. Lam KY, Ma L. Pathology of esophageal cancers: local experience and current insights. *Chin Med J* 1997;110:459-64.
2. Lam AKY. Critical review: Molecular biology of esophageal squamous cell carcinoma. *Crit Rev Oncol Hematol* 2000;33:71-90.
3. Lam KY, Law S, Tung PH, Wong J. Esophageal basaloid squamous cell carcinoma: an unique clinicopathological entity with telomerase activity as a prognostic indicator. *J Pathol* 2001;195:435-42.
4. Lam KY, Law SYK, Loke SL, Fok M, Ma LT. Double sarcomatoid carcinomas of the esophagus. *Path Res Pract* 1996;192:604-9.
5. Lam KY, Dickens P, Loke SL, Fok M, Ma L, Wong J. Squamous cell carcinoma of the esophagus with mucin-secreting component (mucoepidermoid carcinoma and adenosquamous cell carcinoma): a clinicopathologic study and a review of literature. *Eur J Surg Oncol* 1994;20:25-31.
6. Law SYK, Fok M, Lam KY, Loke SL, Ma LT, Wong J. Small cell carcinoma of the esophagus. *Cancer* 1994;73:2894-99.
7. Chow V, Law S, Lam KY, Luk JM, Wong J. Telomerase activity in small cell esophageal carcinoma. *Dis Esophagus* 2001;14:139-42.
8. Lam KY, Law S, Tung PH, Wong J. Esophageal small cell carcinoma.: clinicopathologic parameters, p53 overexpression, proliferative marker, and their impact on pathogenesis. *Arch Pathol Lab Med* 2000;124:228-33.
9. Garbuglia AR. Human papillomavirus in head and neck cancer. *Cancers (Basel)*. 2014;6:1705-26.
10. Salazar CR, Anayannis N, Smith RV, Wang Y, Haigentz M Jr, Garg M, Schiff BA, Kawachi N, Elman J, Belbin TJ, Prystowsky MB, Burk RD, Schlecht NF. Combined p16 and human papillomavirus testing predicts head and neck cancer survival. *Int J Cancer*. 2014;135:2404-12.
11. Westra WH. Detection of human papillomavirus (HPV) in clinical samples: evolving methods and strategies for the accurate determination of HPV status of head and neck carcinomas. *Oral Oncol*. 2014;50:771-79.
12. He D, Zhang DK, Lam KY, Ma L, Ngan HYS, Liu SS, Tsao SW. Prevalence of HPV infection in esophageal squamous cell carcinoma in Chinese patients and its relationship to the p53 gene mutation. *Int J Cancer* 1997;72:959-64.
13. Lam KY, He D, Ma L, Zhang D, Ngan HYS, Wan TSK, Tsao SW. Presence of human papillomavirus in esophageal squamous cell carcinomas of Hong Kong Chinese and its relationship with p53 gene mutation. *Hum Pathol* 1997;28:657-63.
14. Li X, Gao C, Yang Y, Zhou F, Li M, Jin Q, Gao L. Systematic review with meta-analysis: the association between human papillomavirus infection and oesophageal cancer. *Aliment Pharmacol Ther*. 2014;39:270-81.
15. Löfdahl HE, Du J, Näsman A, Andersson E, Rubio CA, Lu Y, Ramqvist T, Dalianis T, Lagergren J, Dahlstrand H. Prevalence of human papillomavirus (HPV) in oesophageal squamous cell carcinoma in relation to anatomical site of the tumour. *PLoS One*. 2012;7:e46538.
16. Vaiphei K, Kochhar R, Bhardawaj S, Dutta U, Singh K. High prevalence of human papillomavirus in esophageal squamous cell carcinoma: a study in paired samples. *Dis Esophagus*. 2013;26:282-87.
17. Al-Haddad S, El-Zimaity H, Hafezi-Bakhtiari S, Rajendra S, Streutker CJ, Vajpeyi R, Wang B. Infection and esophageal cancer. *Ann N Y Acad Sci*. 2014;1325:187-96.

18. Lam KY, Srivastava G, Leung ML, Ma L. Absence of Epstein-Barr virus in esophageal squamous cell carcinoma: a study of 74 cases using in-situ hybridization. *J Clin Pathol Mol Pathol* 1995;48:M188-90.
19. Wang ZK, Yang YS. Upper gastrointestinal microbiota and digestive diseases. *World J Gastroenterol.* 2013;19:1541-50.
20. Gibson MK, Dhaliwal AS, Clemons NJ, Phillips WA, Dvorak K, Tong D, Law S, Pirchi ED, Räsänen J, Krasna MJ, Parikh K, Krishnadath KK, Chen Y, Griffiths L, Colleepriest BJ, Farrant JM, Tosh D, Das KM, Bajpai M. Barrett's esophagus: cancer and molecular biology. *Ann N Y Acad Sci.* 2013 Oct;1300:296-314.
21. Lam KY, Tsao SW, Zhang D, Law S, He D, Ma L, Wong J. Prevalence and predictive value of p53 mutation in patients with esophageal squamous cell carcinomas: a prospective clinicopathological study and survival analysis of 70 patients. *Int J Cancer* 1997;74:212-19.
22. Appelman HD, Matejic M, Parker MI, Riddell RH, Salemme M, Swanson PE, Villanacci V. Progression of esophageal dysplasia to cancer. *Ann N Y Acad Sci.* 2014;1325:96-107.
23. Pack SD, Karkera JD, Zhuang Z, Pak ED, Balan KV, Hwu P, Park WS, Pham T, Ault DO, Glaser M, Liotta L, Detera-Wadleigh SD, Wadleigh RG. Molecular cytogenetic fingerprinting of esophageal squamous cell carcinoma by comparative genomic hybridization reveals a consistent pattern of chromosomal alterations. *Genes Chromosomes Cancer.* 1999;25:160-68.
24. Walch AK, Zitzelsberger HF, Bruch J, Keller G, Angermeier D, Aubele MM, Mueller J, Stein H, Braselmann H, Siewert JR, Höfler H, Werner M. Chromosomal imbalances in Barrett's adenocarcinoma and the metaplasia-dysplasia-carcinoma sequence. *Am J Pathol.* 2000;156:555-66.
25. Kwong D, Lam A, Guan X, Law S, Tai A, Wong J, Sham J. Chromosomal aberrations in esophageal squamous cell carcinoma among Chinese: gain of 12p predicts poor prognosis after surgery. *Hum Pathol.* 2004;35:309-16.
26. Qin YR, Wang LD, Fan ZM, Kwong D, Guan XY. Comparative genomic hybridization analysis of genetic aberrations associated with development of esophageal squamous cell carcinoma in Henan, China. *World J Gastroenterol.* 2008;14:1828-35.
27. Tang JCO, Lam KY, Law S, Wong J, Srivastava G. Detection of genetic alterations in esophageal squamous cell carcinomas and adjacent normal epithelia by comparative DNA fingerprinting using inter-simple sequence repeat PCR. *Clin Cancer Res* 2001;7:1539-45.
28. Hu YC, Lam KY, Law S, Wong J, Srivastava G. Identification of differentially expressed in esophageal squamous cell carcinoma (ESCC) by cDNA expression array: overexpression of Fra-1, Neogenin, Id-1 and CDC25B genes in ESCC. *Clin Cancer Res* 2001;7:2213-21.
29. Fatima S, Chui CH, Tang WK, Hui KS, Au HW, Li WY, Wong MM, Cheung F, Tsao SW, Lam KY, Beh PS, Wong J, Law S, Srivastava G, Ho KP, Chan AS, Tang JC. Transforming capacity of two novel genes JS-1 and JS-2 located in chromosome 5p and their overexpression in human esophageal squamous cell carcinoma. *Int J Mol Med.* 2006;17:159-70.
30. Tang WK, Chui CH, Fatima S, Kok SH, Pak KC, Ou TM, Hui KS, Wong MM, Wong J, Law S, Tsao SW, Lam KY, Beh PS, Srivastava G, Chan AS, Ho KP, Tang JC. Oncogenic properties of a novel gene JK-1 located in chromosome 5p and its overexpression in human esophageal squamous cell carcinoma. *Int J Mol Med.* 2007;19:915-23.
31. Goh XY, Rees JR, Paterson AL, Chin SF, Marioni JC, Save V, O'Donovan M, Eijk PP, Alderson D, Ylstra B, Caldas C, Fitzgerald RC. Integrative analysis of array-comparative genomic hybridisation and matched gene expression profiling data reveals novel genes with prognostic significance in oesophageal adenocarcinoma. *Gut.* 2011;60:1317-26.
32. Xuan J, Yu Y, Qing T, Guo L, Shi L. Next-generation sequencing in the clinic: promises and challenges. *Cancer Lett.* 2013;340:284-95.

33. Zhang J, Baran J, Cros A, Guberman JM, Haider S, Hsu J, Liang Y, Rivkin E, Wang J, Whitty B, Wong-Erasmus M, Yao L, Kasprzyk A. International Cancer Genome Consortium Data Portal--a one-stop shop for cancer genomics data. Database (Oxford). 2011;2011:bar026.
34. Song Y, Li L, Ou Y, Gao Z, Li E, Li X, Zhang W, Wang J, Xu L, Zhou Y, Ma X, Liu L, Zhao Z, Huang X, Fan J, Dong L, Chen G, Ma L, Yang J, Chen L, He M, Li M, Zhuang X, Huang K, Qiu K, Yin G, Guo G, Feng Q, Chen P, Wu Z, Wu J, Ma L, Zhao J, Luo L, Fu M, Xu B, Chen B, Li Y, Tong T, Wang M, Liu Z, Lin D, Zhang X, Yang H, Wang J, Zhan Q. Identification of genomic alterations in oesophageal squamous cell cancer. *Nature*. 2014;509:91-5.
35. Dulak AM, Stojanov P, Peng S, Lawrence MS, Fox C, Stewart C, Bandla S, Imamura Y, Schumacher SE, Shefler E, McKenna A, Carter SL, Cibulskis K, Sivachenko A, Saksena G, Voet D, Ramos AH, Auclair D, Thompson K, Sougnez C, Onofrio RC, Guiducci C, Beroukheim R, Zhou Z, Lin L, Lin J, Reddy R, Chang A, Landrenau R, Pennathur A, Ogino S, Luketich JD, Golub TR, Gabriel SB, Lander ES, Beer DG, Godfrey TE, Getz G, Bass AJ. Exome and whole-genome sequencing of esophageal adenocarcinoma identifies recurrent driver events and mutational complexity. *Nat Genet*. 2013;45:478-86.
36. Nones K, Waddell N, Wayte N, Patch AM, Bailey P, Newell F, Holmes O, Fink JL, Quinn MC, Tang YH, Lampe G, Quek K, Loffler KA, Manning S, Idrisoglu S, Miller D, Xu Q, Waddell N, Wilson PJ, Bruxner TJ, Christ AN, Harliwong I, Nourse C, Nourbakhsh E, Anderson M, Kazakoff S, Leonard C, Wood S, Simpson PT, Reid LE, Krause L, Hussey DJ, Watson DI, Lord RV, Nancarrow D, Phillips WA, Gotley D, Smithers BM, Whiteman DC, Hayward NK, Campbell PJ, Pearson JV, Grimmond SM, Barbour AP. Genomic catastrophes frequently arise in esophageal adenocarcinoma and drive tumorigenesis. *Nat Commun*. 2014;5:5224.
37. Lin DC, Hao JJ, Nagata Y, Xu L, Shang L, Meng X, Sato Y, Okuno Y, Varela AM, Ding LW, Garg M, Liu LZ, Yang H, Yin D, Shi ZZ, Jiang YY, Gu WY, Gong T, Zhang Y, Xu X, Kalid O, Shacham S, Ogawa S, Wang MR, Koeffler HP. Genomic and molecular characterization of esophageal squamous cell carcinoma. *Nat Genet*. 2014;46:467-73.
38. Gopalan V, Pillai S, Ebrahimi F, Salajegheh A, Lam TC, Le TK, Langsford N, Ho YH, Smith RA, Lam AK. Regulation of microRNA-1288 in colorectal cancer: Altered expression and its clinicopathological significance. *Mol Carcinog*. 2014; 53: E36-E44.
39. Ebrahimi F, Gopalan V, Smith RA, Lam AK. miR-126 in human cancers: Clinical roles and current perspectives. *Exp Mol Pathol*. 2014; 96:98-107.
40. Maroof H, Salajegheh A, Smith RA, Lam AK. Role of microRNA-34 family in cancer with particular reference to cancer angiogenesis. *Exp Mol Pathol*. 2014;97:298-304.
41. Vosgha H, Salajegheh A, Smith RA, Lam AK. The Important Roles of miR-205 in normal physiology, cancers and as a potential therapeutic target. *Curr Cancer Drug Targets*. 2014; 14:621-37.
42. Yu H, Duan B, Jiang L, Lin M, Sheng H, Huang J, Gao H. Serum miR-200c and clinical outcome of patients with advanced esophageal squamous cancer receiving platinum-based chemotherapy. *Am J Transl Res*. 2013;6:71-7.
43. Wu C, Li M, Hu C, Duan H. Clinical significance of serum miR-223, miR-25 and miR-375 in patients with esophageal squamous cell carcinoma. *Mol Biol Rep*. 2014;41:1257-66.
44. Matsuzaki J, Suzuki H. MicroRNAs in Barrett's esophagus: future prospects. *Front Genet*. 2014;5:69.
45. Sakai NS, Samia-Aly E, Barbera M, Fitzgerald RC. A review of the current understanding and clinical utility of miRNAs in esophageal cancer. *Semin Cancer Biol*. 2013;23:512-21.
46. Gu J, Wang Y, Wu X. MicroRNA in the pathogenesis and prognosis of esophageal cancer. *Curr Pharm Des*. 2013;19:1292-300.

47. Amin M, Lam AK. Current perspectives of mi-RNA in oesophageal adenocarcinoma: Roles in predicting carcinogenesis, progression and values in clinical management. *Exp Mol Pathol.* 2015; 98: 411-418.
48. Islam F, Qiao B, Smith RA, Gopalan V, Lam AK. Cancer stem cell: Fundamental experimental pathological concepts and updates. *Exp Mol Pathol.* 2015; 98: 184-91.
49. Qiao B, Gopalan V, Chen Z, Smith RA, Tao Q, Lam AKY. Epithelial-mesenchymal transition and mesenchymal-epithelial transition are essential for the acquisition of stem cell properties in hTERT-immortalised oral epithelial cells. *Bio Cell.* 2012; 104: 476-89.
50. Islam F, Gopalan V, Smith RA, Lam AK. Translational potential of cancer stem cells: A review of the detection of cancer stem cells and their roles in cancer recurrence and cancer treatment. *Exp Cell Res.* 2015; 335:135-47.
51. Wang Z, Da Silva TG, Jin K, Han X, Ranganathan P, Zhu X, Sanchez-Mejias A, Bai F, Li B, Fei DL, Weaver K, Carpio RV, Moscovitz AE, Koshenkov VP, Sanchez L, Sparling L, Pei XH, Franceschi D, Ribeiro A, Robbins DJ, Livingstone AS, Capobianco AJ. Notch signaling drives stemness and tumorigenicity of esophageal adenocarcinoma. *Cancer Res.* 2014;74:6364-74.
52. di Pietro M, Alzoubaidi D, Fitzgerald RC. Barrett's esophagus and cancer risk: how research advances can impact clinical practice. *Gut Liver.* 2014;8:356-70.
53. Ong CAJ, Lao-Sirieix P and Fitzgerald RC. Biomarkers in Barrett's esophagus and esophageal adenocarcinoma: Predictors of progression and prognosis. *World J Gastroenterol* 2010; 16:5669-81.
54. Chen M, Huang J, Zhu Z, Zhang J, Li K. Systematic review and meta-analysis of tumor biomarkers in predicting prognosis in esophageal cancer. *BMC Cancer.* 2013;13:539.
55. Shang L, Liu HJ, Hao JJ, Jiang YY, Shi F, Zhang Y, Cai Y, Xu X, Jia XM, ZhanQM, Wang MR. A panel of overexpressed proteins for prognosis in esophageal squamous cell carcinoma. *PLoS One.* 2014 Oct 22;9:e111045.
56. Chung Y, Lam AK, Luk JM, Law S, Chan KW, Lee PY, Wong J. Altered E-cadherin expression and p120 catenin localization in esophageal squamous cell carcinoma. *Ann Surg Oncol.* 2007;14:3260-7.
57. Chan D, Tsoi MY, Liu CD, Chan SH, Law SY, Chan KW, Chan YP, Gopalan V, Lam AK, Tang JC. Oncogene GAEC1 regulates CAPN10 expression which predicts survival in esophageal squamous cell carcinoma. *World J Gastroenterol.* 2013; 19:2772-80.
58. Lam KY, Law S, Lo T, Tung HM, Wong J. The clinicopathological significance of p21 and p53 expression in esophageal squamous cell carcinoma: an analysis of 153 patients. *Am J Gastroenterol;* 1999; 94: 2060-68.
59. Lam KY, Law SYK, So MKP, Fok M, Ma LT, Wong J. Prognostic implication of proliferative markers MIB-1 and PC 10 in esophageal squamous cell carcinoma. *Cancer* 1996;77:7-13.
60. Tong DK, Law S, Kwong DL, Chan KW, Lam AK, Wong KH. Histological regression of squamous esophageal carcinoma assessed by percentage of residual viable cells after neoadjuvant chemoradiation is an important prognostic factor. *Ann Surg Oncol.* 2010;17: 2184-92.
61. Lam KY, Law S, Ma LT, Ong SK, Wong J. Pre-operative chemotherapy for squamous cell carcinoma of the esophagus: do histological assessment and p53 overexpression predict chemo-responsiveness? *Eur J Cancer* 1997;33:1221-25.
62. Okumura H, Uchikado Y, Setoyama T, Matsumoto M, Owaki T, Ishigami S, Natsugoe S. Biomarkers for predicting the response of esophageal squamous cell carcinoma to neoadjuvant chemoradiation therapy. *Surg Today.* 2014;44:421-8.

63. Zhang SS, Huang QY, Yang H, Xie X, Luo KJ, Wen J, Cai XL, Yang F, Hu Y, Fu JH. Correlation of p53 status with the response to chemotherapy-based treatment in esophageal cancer: a meta-analysis. *Ann Surg Oncol*. 2013;20:2419-27.
64. Bain GH, Petty RD. Predicting response to treatment in gastroesophageal junction adenocarcinomas: combining clinical, imaging, and molecular biomarkers. *Oncologist*. 2010;15:270-84.
65. Imdahl A, Jenkner J, Ihling C, Rückauer K, Farthmann EH. Is MIB-1 proliferation index a predictor for response to neoadjuvant therapy in patients with esophageal cancer? *Am J Surg*. 2000;179:514-20.
66. Ung L, Lam AK, Morris DL, Chua TC. Tissue-based biomarkers predicting outcomes in metastatic colorectal cancer: a review. *Clin Transl Oncol*. 2014; 16: 425-435.
67. Pakneshan S, Salajegheh A, Smith RA, Lam AK. Clinicopathological relevance of BRAF mutations in human cancer. *Pathology*. 2013;45:346-56.
68. Rahman MA, Salajegheh A, Smith RA, Lam AK. BRAF inhibitors: From the laboratory to clinical trials. *Crit Rev Oncol Hematol*. 2014;90:220-32.
69. Rahman MA, Salajegheh A, Smith RA, Lam AK. BRAF Inhibitor therapy for melanoma, thyroid and colorectal cancers: development of resistance and future prospects. *Curr Cancer Drug Targets*. 2014;14:128-43.
70. O'Sullivan CC, Connolly RM. Pertuzumab and its accelerated approval: evolving treatment paradigms and new challenges in the management of HER2-positive breast cancer. *Oncology (Williston Park)*. 2014;28:186-94.
71. Wiedmann MW, Mössner J. New and emerging combination therapies for esophageal cancer. *Cancer Manag Res*. 2013;5:133-46.
72. Kordes S, Cats A, Meijer SL, van Laarhoven HW. Targeted therapy for advanced esophagogastric adenocarcinoma. *Crit Rev Oncol Hematol*. 2014;90:68-76.
73. Xu WW, Li B, Lam AK, Tsao SW, Law SY, Chan KW, Yuan QJ, Cheung AL. Targeting VEGFR1- and VEGFR2-expressing non-tumor cells is essential for esophageal cancer therapy. *Oncotarget*. 2015; 6:1790-805.
74. Nakajima M, Kato H. Treatment options for esophageal squamous cell carcinoma. *Expert Opin Pharmacother*. 2013;14:1345-54.
75. Boland PM, Burtneess B. Esophageal carcinoma: are modern targeted therapies shaking the rock? *Curr Opin Oncol*. 2013 ;25:417-24.
76. Tang JCO, Wan TSK, Wong N, Pang E, Lam KY, Law SYK, Chow LMC, Ma ESK, Chan LC, Wong J, Srivastava G. Establishment and characterization of a new xenograft-derived human esophageal squamous cell line SLMT-1 of Chinese origin. *Cancer Genet Cytogenet* 2001;124:36-41.
77. Hu YC, Lam KY, Wan TSK, Fang WG, Ma ESK, Chan LC, Srivastava G. Establishment and characterization of HKESC-1, a new cancer cell line from human esophageal squamous cell carcinoma. *Cancer Genet Cytogenet* 2000;118:112-20.
78. Hu YC, Lam KY, Law SYK, Wan TSK, Ma ESK, Kwong YL, Chan LC, Wong J, Srivastava G. Establishment, characterization, karyotyping, and comparative genomic hybridization analysis of HKESC-2 and HKESC-3: two newly established human esophageal squamous cell lines. *Cancer Genet Cytogenet* 2002;135:120-27.
79. Boonstra JJ, Tilanus HW, Dinjens WN. Translational research on esophageal adenocarcinoma: from cell line to clinic. *Dis Esophagus*. 2013 doi:10.1111/dote.12095.
80. Gros SJ. Orthotopic models of esophageal carcinoma and their use in drug discovery. *Curr Protoc Pharmacol*. 2011 Sep; Chapter 14:Unit14.20. doi:10.1002/0471141755.ph1420s54.
81. Furihata T, Sakai T, Kawamata H, Omotehara F, Shinagawa Y, Imura J, Ueda Y, Kubota K, Fujimori T. A new in vivo model for studying invasion and metastasis of esophageal squamous cell carcinoma. *Int J Oncol*. 2001;19:903-7.

- 82.Hori T, Yamashita Y, Ohira M, Matsumura Y, Muguruma K, Hirakawa K. A novel orthotopic implantation model of human esophageal carcinoma in nude rats: CD44H mediates cancer cell invasion in vitro and in vivo. *Int J Cancer*. 2001;92:489-96.
- 83.Ip JC, Ko JM, Yu VZ, Chan KW, Lam AK, Law S, Tong DK, Lung ML. A versatile Orthotopic nude mouse model for study of esophageal squamous cell carcinoma. *Biomed Res Int*. 2015;2015:910715.