Opportunistic predation by small fish on epibiota of jetty pilings in urban waterways

S. MOREAU^{1,3}, C. PÉRON¹, K. A. PITT^{1*}, R. M. CONNOLLY¹, S. Y. LEE¹ & T. MEZIANE²

¹ Australian Rivers Institute – Coasts and Estuaries, and The Griffith School of Environment, Gold Coast Campus, Griffith University, QLD 4222 Australia. ² UMR 5178, Biologie des Organismes Marins et Ecosystèmes, Département Milieux et Peuplements Aquatiques, MNHN, CP 53, 61 rue Buffon, 75231, Paris CEDEX 05, France. ³ Institut des sciences de la mer (ISMER) Université du Québec à Rimouski (UQAR) 310 allée des Ursulines Rimouski, Qc G5L 3A1, Canada.* Corresponding author

(Received 26 January 2007, Accepted 13 September 2007)

We first determined if small fish associated with the pilings of jetties in urban waterways prey on the epibiota attached to the pilings. Epibiota was sampled on nine small jetties in the tidal, urban canals of southeast Queensland, Australia. Epibiota was dominated by barnacles, filamentous and foliose algae and ranged in thickness from 0.4 – 1.1 cm. The two species of fish that associated most closely with jetty pilings, *Pandaka lidwilli* (Gobiidae) and *Monodactylus argenteus* (Monodactylidae), were sampled twice during the day and twice during the night for analysis of stomach contents. During the day the diet of *P. lidwilli* was dominated by amphipods (~70%, by weight of organic content), with copepods, bivalves and bryozoans each contributing < 10%. At night, amphipods contributed less (~45%) and copepods more (~35%). The diet of *M. argenteus* was dominated by filamentous algae (55%) and amphipods (20%) during the day and filamentous algae (70%) and barnacle cirri (23%) at night. Epibiota, therefore, made a substantial contribution to the diet of the fish but was not the sole source of food for either species. As jetties were the only structures that supported epibiota in the area, fish were likely to source their epibiota from the pilings of the jetties. We therefore tested whether fish depended on the epibiota using a manipulative Before After Control Impact (BACI) study. Three jetties were assigned randomly to each of three treatments: 1) epibiota removed from pilings, 2) epibiota cut and damaged (a procedural control) and, 3) epibiota left undisturbed. Abundances of *P. lidwilli* and *M. argenteus* around jetty pilings remained similar across all treatments from before to after the removal of epibiota. These results indicate that although fish consumed epibiota on the jetties, they did not depend on the epibiota of the jetties for food.

INTRODUCTION

Coastal urbanisation has resulted in increasing numbers of artificial structures being deployed in the sea. Artificial structures, such as jetties and pontoons, support distinct epibiotic assemblages (Connell & Glasby, 1999; Glasby, 1999) and are rapidly colonized by fish (Molles, 1978; Beets, 1989; Ibrahim *et al.*, 1996; Rilov & Benayahu, 1998). In artificial waterways (e.g. canal estates) artificial structures may provide a significant, and sometimes the only, source of hard habitat available to fish. Understanding the mechanism that attracts fish to jetties is the first step towards maximising their value as fish habitat in such systems.

Positive correlations have been observed between the amount of epibiota on artificial structures and abundances of fish (e.g. Rooker *et al.*, 1997) but only two studies (Coleman & Connell, 2001 and Clynick *et al.*, 2007) have used manipulative experiments to demonstrate a causal relationship between the presence of epibiota and some species of fish. While epibiota is clearly important in attracting some fish to artificial structures it is still unclear why the epibiota is important. Fish may associate with structures because the complexity of the epibiota provides a source of shelter (Beets, 1989; Steele, 1999; Coleman & Connell, 2001). Other fish feed on the attached (Keough, 1984; Connell & Anderson, 1999; Connell, 2001a) or motile (Caine 1987; Deudero & Morales-Nin 2000; Relini *et al.*, 2002; Fabi *et al.*, 2006) epibiota associated

with the structures and may, therefore, use the epibiota on artificial structures as a source of food. Studies that have observed predation by fish on the epibiota of artificial structures, however, have not used manipulative experiments to determine the importance of epibiota in maintaining the relationship between fish and artificial structures.

The addition of structure to coastal waters typically increases the biomass of fish at a site (Bohnsack, 1989). This has led to a substantial research effort attempting to determine whether structure acts to increase overall fish biomass (i.e. production) or concentrate fish spatially from a larger area (i.e. attraction). The two factors are not easily separated (Pickering & Whitmarsh, 1997), and it is not our intention to test their importance here. We can, however, provide insights into the mechanisms underlying the links between structure and fish densities, an important step towards separating production and attraction (Brickhill *et al.*, 2005).

Canal estates are increasingly being constructed in coastal regions throughout the world to provide waterfront residential housing. Canal estates provide unique habitats for fish and other biota (Lindall *et al.*, 1975; Baird *et al.*, 1981; Connolly, 2003). In southeast Queensland, hundreds of kilometres of artificial waterways have been constructed since the 1970s (Connolly, 2003) and approximately one third of the waterfront houses have small jetties or pontoons for mooring boats. These artificial structures provide the major source of hard substratum in the canals, support epibiota and attract large numbers of fish.

The aims of this study were to 1) characterise the epibiota on the pilings of jetties in the canals; 2) to use stomach content analysis to test the hypothesis that fish that associate closely with jetties feed on epibiota; and 3) to use a manipulative experiment to test the hypothesis that abundances of fish that prey on epibiota would decrease around jetty pilings after epibiota was removed.

MATERIALS AND METHODS

STUDY SITE AND JETTIES

The study was undertaken in two adjoining artificial lakes (Wonderland and Rumrunner) that form part of the extensive system of canals associated with the Nerang River (27°56' S, 153°26' E), southeast Queensland, Australia (see Morton, (1989) for general location description). The lakes are located ~ 15 km from the Nerang River's entrance to the ocean, have a tidal range of ~ 1 m and had a salinity of ~ 17 psu during the period of the study (Hughes & Mondon, unpubl. data). Twelve jetties were sampled. Nine jetties (five in Lake Rumrunner, four in Lake Wonderland) were used to quantify the distribution, abundance and thickness of epibiota and to assess the effects of removing epibiota on abundances of fish. Fish for stomach content analyses were sampled from three additional jetties (two in Lake Rumrunner, one in Lake Wonderland) to avoid removing fish from jetties that were to be used in the manipulative experiment. Jetties ranged in size from 11 to 24 m² (area of deck) and had between 5 and 13 pilings. All jetties occurred on sandy substrata. Narrow sandy beaches lined the lakes and the canal walls were not inundated with water, even at a high tide. In Lake Wonderland, a small rocky berm lined the edges of some parts of the lake. The berm was, however, only inundated at high tide and supported little epibiota. The jetties, therefore, were the dominant hard substrata in the area and the major source of epibiota. The jetties were separated by > 30 m and were considered to be independent sampling units because although small fish may be capable of swimming more than 30 m, the species targeted have never been caught away from jetty pilings, despite extensive sampling in this area (M. Brickhill, unpubl. data). It was assumed, therefore, that the fish rarely swam distances > 30 m over open sand. The jetties were constructed of three different types of materials. Five had pilings made of PVC, six of fibreglass, and one had concrete pilings. Pilings ranged in circumference from 56 to 79 cm. The jetties that were constructed from different materials and had different circumferences were spatially interspersed.

PERCENTAGE COVER & THICKNESS OF EPIBIOTA

Percentage covers of epibiota were measured using a 25 x 25 cm quadrat with 49 intersection points (Connell, 2001b; Bacchiocchi & Airolidi, 2003). Quadrats were placed 25 cm below the high water line,

which was defined as the highest point at which epibiota occurred (Coleman & Connell, 2001). The primary and secondary covers of epibiota were recorded under each intersection point (Bacchiocchi & Airolidi, 2003). Primary cover was defined as growth directly attached to the piling, and secondary cover as growth attached to the primary cover. Three pilings were randomly selected on each jetty and one quadrat was placed at each cardinal direction on each piling (i.e. $n = 4$). Only sessile organisms were recorded. Rare species that were observed in quadrats but did not occur under an intersection point were given a nominal value of 0.5% (Holloway & Connell, 2002; Bacchiocchi & Airolidi, 2003).

Percentage covers of dominant taxa were analysed using 2-way ANOVAs. The factors were Jetty (random) and Piling (random, nested within Jetty). Prior to analyses, the assumption of homoscedasticity was tested using Cochran's test. When ANOVAs detected significant differences, post-hoc Student-Newman-Keuls (SNK) tests were used to determine where the differences occurred.

The thickness of epibiota on each jetty piling was estimated by comparing the circumferences of each piling below and above the high water mark (i.e. where no epibiota grew) (*sensu* Coleman & Connell, 2001). The average circumference below the high water mark was calculated from measurements at three heights distributed evenly between the high water mark and the substratum. The average thickness of epibiota on each piling was estimated by subtracting the radius of the bare piling from the below-water radius that included epibiota.

ANALYSIS OF FISH DIETS

Two species of fish, *Pandaka lidwilli* (McCulloch) (Gobiidae) and *Monodactylus argenteus* L. (Monodactylidae), occurred abundantly around the jetty pilings. Since these two species only occur around jetty pilings within these lakes (M. Brickhill; unpubl. data) and occurred in reasonable numbers, they were selected to test our hypotheses. *P. lidwilli* and *M. argenteus* were collected for stomach content analysis at low tide using a hand net. Fish were collected from the pilings of three jetties that were independent of those used in the epibiota removal experiment. Fish were collected twice during the day (14 and 28 March) and twice during the night (23 and 28 March) and were frozen immediately upon capture. Stomach contents were quantified separately for animal prey and algae.

Prey were identified by taxon or functional group, counted, and measured using a graticule on a dissecting microscope. Frequency of occurrence (FOC) of each type of prey was calculated as the percentage of fish with that prey in their stomachs (Berg 1979). Given that different types of prey vary in size, the dietary importance of prey was quantified using ash-free dry weight (AFDW). We estimated AFDW of each prey taxon in each individual fish by summing the AFDW of individual prey (Hollingsworth & Connolly, 2006), using Edgar's (1990) algorithms relating prey size and AFDW.

Dry weights for filamentous and foliose algae were estimated by suspending the algae in 1 mL of water in a haemocytometer and then counting under the microscope the proportion of 25 grid points on the haemocytometer under which algae occurred. Prior to this, we had used algae from fish not included in the present study to calculate a relationship between grid point scores and algal weights (measured directly after drying, which for algae is equivalent to AFDW since there is negligible inorganic matter). This relationship was used to convert point estimates to weights.

EXPERIMENTAL REMOVAL OF EPIBIOTA

A BACI (Before-After-Control-Impact) experimental design was used to test the hypothesis that abundances of *P. lidwilli* and *M. argenteus* around the jetty pilings would decrease after epibiota was removed. The same nine jetties used for the epibiota surveys were used for this experiment. The experiment consisted of three treatments: a Removal treatment, a Control and a Procedural Control. Each treatment was applied to three randomly chosen jetties. For the Removal treatment, all epibiota was removed from all the pilings of each jetty using a paint scraper. Care was taken to collect all debris from around the base of the pilings, since the accumulation of damaged epibiota could have artificially attracted fish. Epibiotic assemblages were left undisturbed for the Control jetties. The Procedural Control aimed to control for the disturbance created during removal of the epibiota (e.g. the release of attractive or repellent chemicals) and

consisted of cutting and damaging the epibiota on all pilings on the jetties within this treatment. Fish were sampled twice before (10/11 and 18/19 March) and twice after (5/6 and 13/14 April) the removal of the epibiota. The first “after” sampling occurred five days after the removal of the epibiota to allow the effects of the initial disturbance to dissipate. Based on previous studies (Coleman & Connell, 2001; Clynick *et al.*, 2007) five days was considered sufficient time for an effect to be detected, should there be one. Fish were sampled using a 1 m deep net with a mesh size of 1 mm. Fish were sampled from around three pilings that were randomly selected from the pilings that were covered with water to a depth of 1 m at the time of sampling. The bottom of the net was first tied around the base of the piling and the two edges of the net were drawn together and fastened using Velcro, to encircle the piling. The net was narrower at the base so that when it was raised it formed a cone. The top of the net was quickly raised to the surface of the water to trap the fish around the piling. The epibiota was gently tickled by hand to ensure that fish sheltering within the epibiota were dislodged and collected. All fish caught in the net were identified, counted and released at the piling from which they had been captured. The net was efficient at capturing fish that associate closely with pilings (e.g. *P. lidwilli*) but was not efficient at capturing schooling species that swam rapidly among pilings (e.g. Ambassidae). *M. argenteus* can move rapidly and was sometimes observed to dart between pilings during the study. Numbers of *M. argenteus* around each piling, therefore, were counted visually by divers before the net was deployed. The numbers caught in the net were comparable to those assessed visually, indicating that the net was efficient at capturing this species. Jetties were sampled within 2 h either side of the low tide. Since the pilings of different jetties varied in circumference, abundances of fish were standardized to numbers per m² of piling.

Abundances of *P. lidwilli* and *M. argenteus* were analysed using ANOVAs. Although three pilings were selected randomly at each jetty, at each sampling time, due to the small number of pilings available to be sampled on each jetty, some pilings were sampled multiple times. Due to concerns regarding independence of data, separate two-way ANOVAs were done for each of the four times sampled. The factors were Treatment (fixed, orthogonal) and Jetty (random, nested in Treatment)

RESULTS

DISTRIBUTION OF EPIBIOTA

Barnacles (*Balanus variegatus* and *Balanus trigonus*) and filamentous and foliose algae were the most common epibiota on the jetties. The primary cover was dominated by barnacles with an average cover of $73 \pm 4\%$ (SE). Bare space occupied $20 \pm 4\%$ of primary cover. On average, $54 \pm 14\%$ of the epibiota had a secondary cover growing on it. The secondary cover was dominated by foliose algae with an average cover of $45 \pm 8\%$. The hairy mussel, *Trichomya hirsuta* and bryozoans occurred rarely on pilings. Percentage covers of barnacles and bare space did not vary among jetties but did vary among pilings within some jetties (Table I). Percentage covers of foliose algae did vary among jetties but post-hoc tests revealed that the difference only occurred between the jetties with the largest and smallest covers. The average thickness of epibiota ranged from 0.4 to 1.1 cm.

FISH DIETS

The stomachs of most (78%) *P. lidwilli* specimens contained food. Of these, amphipods and copepods occurred in ~ 70% and bivalves and lophophores of bryozoans were found in ~ 20% (Fig. 1a). The different types of prey occurred in similar frequencies during the day and night. Amphipods comprised > 70% of the AFDW in the stomachs of *P. lidwilli* during the day, with copepods, bivalves and bryozoans each contributing < 10% (Fig. 1b). At night, amphipods contributed less (~ 45%) and copepods more (~ 35%).

Food was found in the stomachs of all *M. argenteus* during the day and in 69% at night. Filamentous algae occurred in ~ 80% of stomachs during both the day and night but, although barnacle cirri and foliose algae occurred in > 40% during the night, they occurred in < 15% of stomachs during the day (Fig. 2a). Amphipods occurred in 70% of stomachs during the day but only 20% at night, and copepods, bivalves, and the lophophores of bryozoans occurred infrequently (in < 20% of the stomachs) and only during the day. By weight, *M. argenteus* stomach contents were dominated by filamentous algae (55%) and amphipods

(20%) during the day (Fig. 2b). Filamentous algae dominated by weight at night (70%) but barnacle cirri were also relatively abundant (23%).

TABLE I. Results of ANOVA of percentage cover of dominant taxa on pilings. All Cochran's tests were non-significant

Source of variation	<i>P</i> values		
	Barnacles (I)	Bare space (I)	Foliose algae (II)
Jetty	>0.05	>0.05	<0.01
Piling	<0.01	<0.01	<0.01

I, primary cover; II, secondary cover.

FIG. 1. (a) Frequency of occurrence (F_{OC}) and (b) mean $\pm$ S.E. relative abundances of ash-free dry mass (M_{AFD}) of different taxa in the stomachs of *Pandaka lidwilli* during the day (■, $n = 50$) and night (□, $n = 30$).

IMPACT OF REMOVAL OF EPIBIOTA ON ABUNDANCES OF FISH

Abundances of *P. lidwilli* did not vary among treatments at any of the four times sampled (Table II, Fig. 3a). Abundances also did not vary among jetties within treatments during the two 'before' or first 'after'

sampling times, but did vary among jetties within the removal and control treatments during the last sampling time. The abundances of *M. argenteus* did not vary among any of the factors (Table III) although there was a slight trend for abundances of *M. argenteus* to decrease across all treatments after the removal of epibiota (Fig. 3b).

FIG. 2. (a) Frequency of occurrence (F_{OC}) and (b) mean $\pm$ S.E. relative abundances of ash-free dry mass (M_{AFD}) of different taxa in the stomachs of *Monodactylus argenteus* during the day (■; $n = 14$) and night (□; $n = 9$). Uncommon taxa were omitted.

DISCUSSION

Epibiota was ingested by both *P. lidwilli* and *M. argenteus* and as the jetties were the only major source of hard substratum supporting epibiota in the area, it is likely that both species of fish grazed directly on the epibiota attached to the pilings of the jetties. *M. argenteus* appeared to ingest more epibiota than *P. lidwilli*, with filamentous algae contributing $> 60\%$ of the organic content overall and barnacle cirri contributing $> 20\%$ at night. *P. lidwilli* grazed on bivalves and bryozoans but these made up a relatively small proportion of the organic content in the stomachs. Instead, amphipods and copepods were the dominant prey captured. Amphipods were very abundant among the epibiota on the pilings (C. Péron, pers. obs.) and other studies have shown that amphipods associated with artificial structures are preyed upon by fish (Caine, 1987). The

jetties, therefore, may have provided a readily available source of amphipods for the fish, but the possibility that amphipods were captured from other habitats cannot be excluded.

TABLE II. Results of one-way ANOVA of abundance of *Pandaka lidwilli* and *Mono-dactylus argenteus* around pilings at four times during the study. All Cochran's tests were non-significant

Source of variation	P-values			
	10–11 March	18–19 March	5–6 April	13–14 April
<i>P. lidwilli</i>				
Treatment	>0.05	>0.05	>0.05	>0.05
Jetty (treatment)	>0.05	>0.05	>0.05	<0.01
<i>M. argenteus</i>				
Treatment	>0.05	>0.05	>0.05	>0.05
Jetty (treatment)	>0.05	>0.05	>0.05	>0.05

Despite a strong apparent trophic link between *P. lidwilli* and *M. argenteus* and the jetty pilings, removal of epibiota had no significant influence on the abundances of fish. If epibiota was an important source of food for fish associated with jetties, then we would have expected fish to either leave jetties from which epibiota was removed or to have suffered greater mortality on these jetties due to lack of food. Instead, numbers of fish did not change around the jetties where epibiota was removed or around the control and procedural control jetties. The lack of change around the procedural control jetties indicated that the disturbance created by removal of epibiota had no influence on the numbers of fish and the lack of change around control jetties suggested that no large-scale processes (such as recruitment or immigration) occurred concurrently with the experiment, which may have confounded interpretation of the results. The duration of the experiment should have been sufficient to detect changes in abundances of fish due to emigration, since in previous studies, emigration has been detected within 5 days (Coleman and Connell 2001; Clynick *et al.* 2007). Starvation would probably have acted more slowly to reduce abundances of fish around jetties from which epibiota was removed and it is possible that the effects of starvation were not detected within the timeframe of the experiment. Small fish, however, tend to be less able to withstand periods of starvation (e.g. Bystroem *et al.* 2006). Given that the average size of *P. lidwilli* caught was 12.5 $\pm$ 0.13 mm, it is probably unlikely that this species could endure starvation for periods of >2 weeks. Starved fish are also more susceptible to predation (Jonas and Wahl 1998) and are more likely to take risks when feeding, which further increases mortality (Biro *et al.* 2005). Consequently we are confident that, had *P. lidwilli* been dependent on the epibiota for food, a decrease in abundance of this species would have been apparent following removal of epibiota, within the timeframe of the experiment.

Although epibiota was ingested by both *P. lidwilli* and *M. argenteus*, epibiota was not the sole source of food for either species. Indeed, copepods comprised, on average, >20% of the organic content found in the stomachs of *P. lidwilli* and also occurred in the stomachs of *M. argenteus*. Both species of fish, therefore, may be considered generalist predators that preyed opportunistically on the epibiota but were not entirely dependent on it as a source of food. Other studies have similarly found that fish that associate with artificial structures feed in habitats adjacent to the artificial structure as well as on the epibiota of the structure itself (Lindquist *et al.*, 1994; Relini *et al.*, 2002). Following removal of epibiota both *P. lidwilli* and *M. argenteus* may have preyed more intensively on copepods or amphipods that may have been available in adjacent habitats. Removal of the epibiota, however, clearly did not significantly reduce the abundance or accessibility of food for the species of fish studied. Predation on epibiota, therefore, can be excluded as being the reason why these fish associate with jetties.

An alternative model to explain the association of fish and epibiota is that the epibiota provides refuge for fish (Coleman & Connell, 2001). The epibiota on the pilings in this study consisted mainly of small barnacles (diameter $\sim$ 2 cm) and was less structurally complex and much thinner than that observed in other studies (maximum of $\sim$ 3 cm thick cf. $\sim$ 10 cm Coleman pers. comm.). The interstices between adjacent

FIG. 3. Mean $\pm$ S.E. abundances of (a) *Pandaka lidwilli* and (b) *Monodactylus argenteus* on pilings of jetties in the epibiota removal (■), procedural control (□) and control (■) treatments before (10 and 18 March 2006) and after (5 and 13 April 2006) the removal of epibiota.

barnacles may have still provided suitable-sized shelters for *P. lidwilli*. The speckled brown colouration of *P. lidwilli* would have also camouflaged it well against the epibiota. This species was observed to remain within 1 – 2 cm of the epibiota and it was considered likely, therefore, that it may use the epibiota as a refuge. In contrast, *M. argenteus* is a longer and deeper-bodied species, and although the individuals caught in this study were juveniles, they were nevertheless larger than *P. lidwilli* (max TL 4.2 cm, average 2.5 cm). Individual *M. argenteus* remained several centimetres away from the pilings and often darted between adjacent pilings and it is less likely that this species used the epibiota as a source of shelter. Despite the different sizes and behavioural characteristics, both species failed to respond to removal of epibiota.

Predators can remove up to 60% of small fish over periods of 96 hours when access to refugia is limited (Holbrook and Schmitt 2002). As numbers of fish in the present study did not vary over a period of 2 weeks following removal of epibiota, we conclude that epibiota is not a critical source of shelter for either species.

Although Clynick *et al.* (2007) found that several species of fish responded to removal of epibiota, small cryptic blennies, which appeared to associate very closely with the epibiota, did not respond to removal of epibiota. Clynick *et al.* (2007) suggested that difficulties in using visual censuses to sample small cryptic species may have contributed to their inability to detect a difference in abundances of blennies. We are confident, however, that the net used to sample fish in the present study was efficient at sampling the small fish that associated closely with the pilings and that the lack of a response by the fish was a real effect and unrelated to sampling efficiency.

The lack of change in abundances of fish following the removal of epibiota indicates that the fish associate with the pilings themselves, rather than the epibiota growing on them. Jetties without epibiota may still provide a source of refuge. In particular, the shade created by the jetties and pilings may reduce the visibility of fish to both aerial and aquatic predators (Helfman, 1981; Hair *et al.*, 1994). The pilings themselves may also provide a physical barrier behind which fish may seek refuge from predators, in the same way coarse woody debris provide a refuge from predation (Everett & Ruiz, 1993). Indeed, when captured within the conical net, both *P. lidwilli* and *M. argenteus* rapidly retreated to the opposite side of the piling to the person doing the sampling. The presence of pilings is likely, therefore, to reduce the risk of predation for these two species, but further experiments are required to test these models.

Artificial structures provide a significant habitat for fish in artificial waterways. Given that increasing numbers of artificial structures are likely to be deployed, the design of these structures should be optimised so that they provide the best quality fish habitat. The first step to achieving this is to identify what attracts fish to these structures. We conclude that although *P. lidwilli* and *M. argenteus* prey opportunistically on the epibiota associated with jetty pilings, the association of small fish and jetties does not depend on the presence of epibiota. Alternative models involving factors such as the shade and refuge provided by jetties need to be tested.

Acknowledgements

We thank M. Arthur, J. Warnken, M. Brickhill and the numerous volunteers who assisted in the field. We also thank two anonymous reviewers for valuable comments.

References

- Bacchiocchi, F., Airolidi, L. (2003). Distribution and dynamics of epibiota on hard structures for coastal protection. *Estuarine, Coastal and Shelf Science* **56**, 1157-1166.
- Baird, D., Marais, J. F. K. & Wooldridge, T. (1981). The influence of a marina canal system on the ecology of the Kromme Estuary, St. Francis Bay. *South African Journal of Zoology* **16**, 21-34.
- Beets, J. (1989). Experimental evaluation of fish recruitment to combination of Fish Aggregating Devices and benthic artificial reefs. *Bulletin of Marine Science* **44**, 973-983.
- Biro, P. A., Post, J. R. & Abrahams, M. V. (2005). Ontogeny of energy allocation reveals selective pressure promoting risk-taking behaviour in young fish cohorts. *Proceedings of the Royal Society of London, Series B* **272**, 1443-1448.
- Bohnsack, J. A. (1989). Are high densities of fishes at artificial reefs the result of habitat limitation or behavioral preference? *Bulletin of Marine Science* **44**, 631-645.
- Brickhill, M. J., Lee, S. Y. & Connolly, R. M. (2005). Fish associated with artificial reefs: attributing changes to attraction or production using novel approaches. *Journal of Fish Biology* **67**, 53-71.
- Brystrom, P., Andersson, J., Kiessling, A. & Eriksson, L-O. (2006). Size and temperature dependent foraging capacities and metabolism: consequences for winter starvation mortality in fish. *Oikos* **115**, 43-52.
- Caine, E. A. (1987). Potential effect of floating dock communities on a South Carolina estuary. *Journal of Experimental Marine Biology and Ecology* **108**, 83-91.

- Clynick, B. G., Chapman, M. G. & Underwood, A. J. (2007). Effects of epibiota on assemblages of fish associated with urban structures. *Marine Ecology Progress Series* **332**, 201-210.
- Coleman, M. A. & Connell, S. D. (2001). Weak effects of epibiota on the abundances of fishes associated with pier pilings in Sydney Harbour. *Environmental Biology of Fishes* **61**, 231-239.
- Connell, S. D. (2001a). Predatory fishes do not always affect the early development of epibiotic assemblages. *Journal of Experimental Marine Biology and Ecology* **260**, 1-12.
- Connell, S. D. (2001b). Urban structures as marine habitats: an experimental comparison of the composition and abundance of subtidal epibiota among pilings, pontoons and rocky reefs. *Marine Environmental Research* **52**, 115-125.
- Connell, S. D. & Anderson, M. J. (1999). Predation by fish on assemblages of intertidal epibiota: effects of predator size and patch size. *Journal of Experimental Marine Biology and Ecology* **241**, 15-29.
- Connell, S. D. & Glasby, T. M. (1999). Do urban structures influence local abundance and diversity of subtidal epibiota? A case study from Sydney Harbour, Australia. *Marine Environmental Research* **47**, 373-387.
- Connolly, R. M. (2003). Differences in trophodynamics of commercially important fish between artificial waterways and natural coastal wetlands. *Estuarine, Coastal and Shelf Science* **58**, 929-936.
- Deudero, S. & Morales-Nin, B. (2000). Occurrence of *Polyprion americanus* under floating objects in western Mediterranean oceanic waters, inference from stomach contents analysis. *Journal of the Marine Biological Association of the United Kingdom* **80**, 751-752.
- Edgar, G. J. (1990). The use of size structure of benthic macrofaunal communities to estimate faunal biomass and secondary production. *Journal of Experimental Marine Biology and Ecology* **137**, 195-214.
- Everett, R. A. & Ruiz, G. M. (1993). Coarse woody debris as a refuge from predation in aquatic communities. *Oecologia* **93**, 475-486.
- Fabi, G., Manoukian, S. & Spagnolo, A. (2006). Feeding behaviour of three common fishes at an artificial reef in the northern Adriatic Sea. *Bulletin of Marine Science* **78**, 39-56.
- Glasby, T. M. (1999). Differences between subtidal epibiota on pier pilings and rocky reefs at marinas in Sydney, Australia. *Estuarine, Coastal and Shelf Science* **48**, 281-290.
- Hair, C. A., Bell, J. D. & Kingsford, M. J. (1994). Effects of position in the water column, vertical movement and shade on settlement of fish to artificial habitats. *Bulletin of Marine Science* **55**, 434-444.
- Helfman, G. S. (1981). The advantage to fishes of hovering in shade. *Copeia* **1981**, 392-400.
- Holbrook, S. J. & Schmitt, R. J. (2002) Competition for shelter space causes density-dependent predation mortality in damselfishes. *Ecology* **83**, 2855-2868.
- Hollingsworth, A. & Connolly, R. M. (2006). Feeding by fish visiting inundated subtropical saltmarsh. *Journal of Experimental Marine Biology and Ecology* **336**, 88-98.
- Holloway, M. G. & Connell, S. D. (2002). Why do floating structures create novel habitats for subtidal epibiota? *Marine Ecology Progress Series* **235**, 43-52.
- Ibrahim, S., Ambak, M. A., Shamsudin, L. & Samsudin, M. Z. (1996). Importance of fish aggregating devices (FADs) as substrates for food organisms of fish. *Fisheries Research* **27**, 265-273.
- Jonas, A. L. & Wahl, D. H. (1998). Relative importance of direct and indirect effects of starvation for young walleyes. *Transactions of the American Fisheries Society* **127**, 192-205.
- Lindall, W. N. J., Fable, W. A. J. & Collins, L. A. (1975). Additional studies of the fishes, macroinvertebrates, and hydrological conditions of upland canals in Tampa Bay, Florida. *Fishery Bulletin* **73**, 81-85.
- Lindquist, D. G., Cahoon, L. P., Clavijo, I. E., Posey, M. H., Bolden, S. K., Pike, L. A., Burk, S. W. & Cardullo, P. A. (1994). Reef fish stomach contents and prey abundance on reef and sand substrata associated with adjacent artificial and natural reefs in Onslow Bay, North Carolina. *Bulletin of Marine Science* **55**, 308-318.
- Molles, M. C. (1978). Fish species diversity on model and natural patch reefs: experimental insular biogeography. *Ecological Monographs* **48**, 289-305.
- Morton, R. M. (1989). Hydrology and fish fauna of canal developments in an intensively modified Australian estuary. *Estuarine Coastal and Shelf Science* **28**, 43-58.
- Pickering, H. & Whitmarsh, D. (1997). Artificial reefs and fisheries exploitation: a review of the 'attraction versus production' debate, the influence of design and its significance for policy. *Fisheries Research* **31**, 39-59.

- Relini, G., Relini, M., Torchia, G. & De Angelis, G. (2002). Trophic relationships between fishes and an artificial reef. *ICES Journal of Marine Science* **59**, S36-S42.
- Rilov, G. & Benayahu, Y. (1998). Vertical artificial structures as an alternative habitat for coral reef fishes in disturbed environments. *Marine Environmental Research* **45**, 431-451.
- Rooker, J. R., Dokken, Q. R., Pattengill, C. V. & Holt, G. J. (1997). Fish assemblages on artificial and natural reefs in Flower Garden Banks National Marine Sanctuary. *Coral Reefs* **16**, 83-92.
- Steele, M. A. (1999). Effects of shelter and predators on reef fishes. *Journal of Experimental Marine Biology and Ecology* **233**, 65-79.
- Underwood, A. J. (1997). *Experiments in Ecology*. Cambridge University Press, United Kingdom