Mucosal immunization against respiratory bacterial pathogens

A Ruth Foxwell[†], Jennelle M Kyd and Allan W Cripps

Bacterial respiratory diseases remain a major cause of morbidity and mortality throughout the world. The young and the elderly are particularly susceptible to the pathogens that cause these diseases. Therapeutic approaches remain dependent upon antibiotics contributing to the persistent increases in antibiotic resistance. The main causes of respiratory disease discussed in this review are *Mycobacterium tuberculosis*, *Corynebacterium diphtheriae*, *Bordetella pertussis*, *Streptococcus pneumoniae*, nontypeable *Haemophilus influenzae*, *Moraxella catarrhalis* and *Pseudomonas aeruginosa*. All these organisms initiate disease at the mucosal surface of the respiratory tract and thus the efficacy of the host's response to infection needs to be optimal at this site. Vaccines available for diseases caused by many of these pathogens have limitations in accessibility or efficacy, highlighting the need for improvements in approaches and products. The most significant challenges in both therapy and prevention of disease induced by bacteria in the respiratory tract remain the development of noninjectable vaccines and delivery systems/immunization regimens that improve mucosal immunity.

Expert Rev. Vaccines 2(4), 551–560 (2003)

Respiratory disease due to bacteria remains a major cause of morbidity and mortality in both developed and developing communities. The major groups affected by bacterial respiratory illness are: the young, with a high incidence of otitis media (OM) and lower incidence of pneumonia and whooping cough; all age groups, with tuberculosis or with conditions that compromise the immune system; and the elderly, with a susceptibility for chronic bronchitis, chronic obstructive pulmonary disease (COPD) and pneumonia. Bacterial pathogens that colonize or invade through mucosal surfaces are a major cause of disease in all these groups. Prevention of disease should therefore be targeted at prevention of invasion or disturbance of these mucosal membranes.

This review will briefly summarize the epidemiology, including burden of disease, of the bacterial respiratory pathogens causing greatest human and economic impact; comment on optimal and available therapy and its 'limitations'; provide a synopsis of the 'state of the art'

of mucosal vaccines against bacterial respiratory pathogens; and present key issues for future research and clinical strategies.

Respiratory disease & bacterial pathogens Epidemiology & burden of disease

Respiratory infections account for 50 million deaths worldwide annually [1], with 3 million of these deaths being caused by a single microbial agent, *Mycobacterium tuberculosis* [2]. Children of less than 5 years of age in developing countries are at greatest risk of mortality from respiratory tract infections, with 2 million deaths in this cohort in 2002 [3]. Over the last decade mortality from respiratory tract infections has tended to decrease due to broader access to antibiotics. However, with increasing antibiotic resistance, for example, as seen with *Streptococcus pneumoniae* [4,5] and both resistance and disease resurgence, as seen with *M. tuberculosis* [6], this trend is already reversing. Vaccine-preventable illnesses that have high priority with the World Health Organization include tuberculosis, diphtheria and whooping cough.

Respiratory disease & bacterial pathogens
Disease prevention
Vaccine development for protection
Vaccine delivery to target
mucosal surfaces
Expert opinion
Key issues
Conclusions
References
Abbreviations

†Correspondence:
Senior Lecturer in Allergy and
Food Research Centre in Medical
and Health Sciences, University of
Canberra, Canberra ACT 2601,
Australia.
Tel: +61 2 6129 2000
Fax: +61 2 620 1240
E-mail: a.cripps@canberra.edu.au

KEYWORDS:
allergens; delivery systems;
immune mechanism; mucosal
infection; otitis media; respiratory
infection

There is a significant human and economic burden resulting from bacterial respiratory disease morbidity. While the use of antibiotics has reduced the mortality associated with bacterial diseases, such as pneumonia, it has had little overall effect on the morbidity [7]. Other lung-based bacterial-associated conditions, such as chronic bronchitis and COPD, have been estimated to cost up to US\$3000 per clinical case per year, almost twice that reported for asthma [8]. The sleeping giant of lung morbidity is tuberculosis with an estimated 2 billion individuals infected worldwide. The majority of these individuals do not exhibit overt signs of disease but are at increased risk of reactivating tuberculosis should their immune system become compromised, as it does with age or through contracting other infectious agents, such as HIV [2].

Respiratory tract infections also include infections of the middle ear cavity. OM is the most common reason for a child under 3 years to visit a general practitioner in the USA and most developed countries. In the USA, it is estimated that more than 20 million episodes of acute OM occur annually [9], resulting in a cost of \$4–5 billion per year [10]. Developing communities, as well as certain ethnic groups, such as the Australian Aborigines, have higher rates of OM at younger ages and longer term morbidity [11]. In developed countries, extensive usage of antibiotics in the treatment for OM remains a controversial issue amongst specialists in the field. Complications are increasing due to drug resistance and sequelae, such as mastoiditis, meningitis and chronic suppurative OM [12].

Bacterial respiratory pathogens

Seven organisms are responsible for the majority of bacterial respiratory infections, *S. pneumoniae*, nontypeable *Haemophilus influenzae* (NTHi), *Moraxella catarrhalis*, *Pseudomonas aeruginosa*, *M. tuberculosis*, *Corynebacterium diphtheriae* and *Bordetella pertussis*. *S. pneumoniae*, NTHi and *M. catarrhalis* are often found as part of the normal microbiota of the upper airways and as such, can be considered 'commensal pathogens'.

S. pneumoniae, a commensal of the upper respiratory tract, is carried in 95% of healthy children less than 3 years of age and 40% of adults [13]. Infection often follows acquisition of a new serotype and results in a variety of mucosal infections, such as OM and pneumonia and/or leads to invasive cases of septicemia and meningitis. The morbidity associated with this micro-organism is high and it remains a leading cause of mortality in young children [14]. NTHi and, to a lesser extent, *M. catarrhalis*, follow similar patterns of colonization and disease to that seen by *S. pneumoniae* [15,16]. The organisms that commonly colonize the nasopharynx and their links to the disease state are well reviewed in a recent publication by García-Rodríguez and Martínez [17]. The impact that an effective vaccine can have is demonstrated by the marked decrease in pneumonia and other debilitating infections associated with *H. influenzae* type b since the vaccine was introduced in 1988 [18]. Thus, it is a priority to develop effective and preferably multivalent vaccines for these pathogens.

M. tuberculosis is a complex pathogen, capable of establishing persistent infection in the host despite the development of an appropriate cell-mediated immune response [19]. Recent evidence would suggest that secretory immunoglobulin (Ig)A (sIgA) is also part of the effector mechanisms in disease prevention, however, more studies need to be performed to establish its role as a protective agent against primary disease or disease reactivation [20]. The bacterium has been estimated to infect one in three people worldwide and although many people survive the acute phase, 2–23% will develop disease symptoms in later life with approximately 90 million expected to die of tuberculosis within the next 30 years [2]. Targeted research to develop better preventative therapies to this microorganism is still required.

C. diphtheriae, like *H. influenzae* type b, was a cause of serious disease prior to an effective vaccine. Death as a result of infection has been reported to be as high as 10% [21]. Vaccine efficacy has been high and Europe was considered almost 'diphtheria free' until a lack of vaccine compliance resulted in an outbreak in the newly formed Independent States of the former Soviet Union between 1990 and 1999, resulting in more than 150,000 cases and 5000 deaths [4,21]. The vaccine, active against a toxin produced by the bacteria, does not affect the bacteria itself and the molecular characterization of isolates from the European outbreak identified the strain as being part of the endemic reservoir that existed in Russia prior to the epidemic [22]. The general community has become complacent over time and there is a clear need to continually remind the public, through active media programs, of the need to maintain vaccination against bacteria such as *C. diphtheriae*.

B. pertussis, the cause of whooping cough, remains circulating in the population despite the introduction of community-wide vaccination programs. Officially reported statistics of incidence and complications in whooping cough are grossly underestimated. Of the 330/100,000 cases identified in a study in Birmingham, England, only 4/100,000 were officially recorded in the statutory notifications for England and Wales [23]. In areas of good vaccine compliance, such as in England and France, there has been an age shift in organism colonization and a subsequent disease shift from the young to older age groups [24]. Hence, this highlights the issue of duration of immunological memory for some vaccines. Whooping cough can be a severe illness which can lead to brain damage and death and, therefore, it has been suggested that immunization coverage be extended from infants to include adolescent and adult populations [24]. The reduction in side effects following the introduction of the acellular pertussis vaccine will increase acceptability in all population age groups.

Disease prevention

S. pneumoniae, NTHi, *M. catarrhalis* and *C. diphtheriae* are constantly circulating between individuals in the community and colonize the mucosal epithelium of the upper respiratory tract. In the majority of individuals the acquisition of a new bacterial strain is harmless – a neutral phenomenon to the

Table 1. Vaccine delivery technology and application for mucosal delivery of respiratory bacterial antigens.

	Technology	Advantage	Antigens against infection	Mucosal application
Adjuvants				
Endotoxins	Endotoxin of <i>V. cholerae</i> used either as CT or the subunit CTB [42] and heat-labile toxin of <i>E. coli</i> [42]	CT more potent mucosal adjuvant either conjugated or mixed. CTB not as effective orally	Extensive number of studies for many microbial and dietary antigens	Yes as well as transcutaneously for mucosal protection [73]
Monophosphoryl lipid A (MPL)	Chemically modified LPS from <i>Salmonella minnesota</i> R595	Potent adjuvant + low toxicity [74]	TT, HbsAg, FluShield® [45], [75] Pneumococcal-CRM197 [44]	Yes
Oligodeoxynucleotides containing CpG	Immunostimulatory sequences, synthetic	Potent adjuvant of Th1 immune response when delivered mucosally	<i>Haemophilus influenzae</i> type b-CRM197 [47]	Yes
Lipid core peptide (LCP)	Synthetic	Antigen + carrier + adjuvant as single molecule	<i>Chlamydia trachomatis</i> [76] Group A Streptococci [49]	Not yet reported
Bacterial systems				
<i>Salmonella</i> strains	Attenuated live carrier expressing foreign antigens	Targets antigen to mucosal lymphoid tissues and can deliver improved antigen load [52]	Many antigens in addition to model antigens and includes <i>M. tuberculosis</i> and pneumococcal antigens as well as DNA with eukaryotic expression systems [52,77-80]	Yes
<i>Mycobacterium</i> strains	Attenuated strain and now recombinant technology incorporated	Licensed vaccine, although current BCG vaccine is questionable for effective protection against TB	Against tuberculosis plus carrier of foreign antigens and immunomodulators [60,61]	Yes and was used as an oral vaccine until the 1940s
Lactic acid bacteria	Expression of antigens in <i>Lactococcus</i> and <i>Lactobacillus</i> [81]	Lactic acid are well tolerated as food microbes and have been used as probiotics	TT, pneumococcal polysaccharide, <i>H. influenzae</i> antigen [82-85]	
Bacterial ghosts	Recombinantly engineered nonliving bacteria	Potential for the bacterial envelope to be both carrier and vaccine	Mostly as homologous vaccine, but some recent advances as a carrier as well [53-55,86,87]	
<i>Bacillus subtilis</i> spores	Recombinant spores with chimeric antigen expression	Recombinant system, organism used as probiotic	TT delivered orally and nasally [88]	
Microparticles				
ISCOMs	Saponin mixture Quil A, cholesterol and phospholipids	Potent adjuvant and available in veterinary vaccines	Influenza virus, hepatitis A, diphtheria and tetanus toxoids, but mostly model antigens for mucosal applications [63]	Yes but not extensively and new formulations being investigated [64,89,90]
Liposomes	These are particles with a bilayer structure with different phospholipid-based compositions [63]	Potency depends on formulation, antigen and route of immunization as does type of response, Th1 and/or Th2	Range of antigens including DNA and influenza [66,67,91-93]	Yes, especially for oral delivery
Nanoparticles	Cationic cross-linked polysaccharides enclosed in lipidic bilayer	Designed for nasal delivery	DNA and bacterial antigens [69-71]	Yes, nasal
Plants				
Edible plants	Recombinant technology in potato, tomato, cereals, alfalfa, legumes, banana	Simplicity of delivery in a food, but technology still needs further development	Diarrheal antigens, Hepatitis B antigens in human trials [72]	Yes

BCG: Bacillus Calmette-Guérin; CT: Cholera toxin; CTB: Cholera toxin B subunit; *E. coli*: *Escherichia Coli*; HbsAg: Hepatitis B surface antigen; ISCOM: Immunostimulatory complex; LPS: Lipopolysaccharide; TB: Tuberculosis; Th: T helper cell; TT: Tetanus toxoid; *V. Cholerae*: *Vibrio Cholerae*.

host. Infections occur when the host-commensal balance is disturbed for some reason or in individuals more susceptible due to age, genotype, current immune status and other similar factors. Hence, these bacteria are part of the normal microflora, so the question that must be asked is whether a vaccine that creates sterile immunity would open a niche for new pathogens to colonize the nasopharynx. Is it even necessary that vaccines against 'commensal pathogens' induce sterile immunity? A number of studies have suggested that it is high bacterial load that predisposes to disease, so the ideal vaccine may be one that controls bacterial load sufficiently such that disease is prevented. Clearly a better understanding of immune responses that enable control of the micro-organisms at the mucosal surface as well as an understanding of the microbial ecology of the upper airways is essential to optimize therapeutic initiatives.

P. aeruginosa is ubiquitous within the environment and tends to rely on an underlying disease-state or structural damage to tissue before the host becomes susceptible to infection. As such, it is a significant cause of acute and chronic disease in specific population groups. It is these groups that need to be targeted for more effective prophylactic and therapeutic approaches, rather than the general population.

In the case of *C. diphtheriae*, it is the immunity induced by vaccination against the toxin that prevents disease and not immunity preventing bacterial colonization and, as stated above, vaccine programs must be maintained. *M. tuberculosis* and *B. pertussis* are specific human pathogens that are not normal commensals and thus prevention of initial colonization on the mucosal epithelium is essential.

Ideal outcome for protection

The question as to whether elimination, from the host, of the bacteria that are responsible for the different respiratory diseases is the optimal outcome for therapy or if control of a potential disease state is more desirable still needs to be addressed.

In the case of absolute pathogens, such as *M. tuberculosis* and *B. pertussis*, bacterial elimination from the host is essential. This is also probably the most desirable outcome for *C. diphtheriae*, however, the current toxoid vaccine is safe and effective and, despite research into changes in the delivery mode (TABLE 1), it is unlikely that the injectable vaccine will be changed at this point in time. In the case of the normal respiratory commensals, such as *S. pneumoniae*, NTHi and *M. catarrhalis*, the possibility of bacterial replacement with an organism that will be more harmful to the host has created debate.

New conjugate pneumococcal vaccines have become available and the results of several studies on populations in which the vaccines have been trialed have been examined. Trials in the American Indian population looked at the efficacy of the parenterally administered heptavalent formulation conjugated to a nontoxic diphtheria variant (Prevnam[®], Wyeth). The study found that the overall pneumococcal load in the respiratory tract remained unchanged postvaccination despite a significant

reduction in the vaccine-type pneumococcal carriage rate, thus indicating the vaccine was associated with replacement carriage in the nasopharynx [25]. Furthermore, clinical trials in Finland evaluating the effect of pneumococcal vaccines on OM demonstrated that the number of episodes of OM caused by nonvaccine serotypes of pneumococcus increased by 33% despite a demonstration of protection against serotypes included in the vaccine [26]. A similar replacement of pathogenic bacteria was not observed in the respiratory tract following mucosal vaccination with a whole-cell NTHi preparation in Papua New Guinea [27], yet most individuals make strain-specific immune responses postinfection with NTHi.

Control of absolute bacterial numbers and age of colonization are important in the predisposition to infection in individuals with high levels of bacterial colonization. Studies in developed and developing communities have established that the risk for development of OM increases when either the number of potential pathogenic bacteria, *S. pneumoniae*, NTHi, *M. catarrhalis*, colonizing the nasopharynx is increased or the age of the patient is less than 3 months when primary colonization occurs [28–31].

Antibiotics have had mixed success in the treatment of respiratory infections. Acute OM is the most common reason for prescribing antibiotics in young children [9]. However, lack of diagnostic specificity for bacterial causes; self-resolution of infection, in some cases in short time periods; and increased bacterial resistance, have resulted in mixed recommendations and outcomes in the use of antibiotic treatment for OM [12]. Effective immune responses at the mucosal site that control colonization and are upregulated to effectively prevent infection would therefore be a preferable therapeutic outcome. This could only be achieved by the development of new vaccines for most of these respiratory pathogens.

Factors likely to affect choice of therapy

Mucosal immunization

Since the late 18th Century, of the 25–30 vaccines available for human use, very few have been administered nonparenterally [32], despite one of the most successful disease eradication programs being achieved using an oral vaccine. The control of polio was achieved in many countries in the world through oral vaccination programs. However, once eradicated from a population, the continued use of the vaccine has proven questionable due to the live attenuated virus posing a risk of disease. Since the mucosal portals of the body are the entry for many infectious diseases, vaccines that increase the integrity of the mucosal barriers through specific immune mechanisms should have a significant impact on disease incidence. It is clear from both animal and human studies that when antigen is presented directly to immune-induction sites, such as the gut-, nasal- and bronchial-associated lymphoid tissues (GALT, NALT and BALT, respectively), protective responses are induced within the respiratory tract. Animal experiments have provided a great deal of information on the mechanisms of mucosal immune regulation such that it is now possible to

predict and certainly determine, whether a given antigen formulation will result in tolerance or polarize the mucosal response towards T-helper (Th)1 or Th2 [33]. Determination of pathogen-specific protective responses required in the respiratory tract will enable relatively well-targeted vaccine development programs.

The GALT and NALT are the major inductive sites of the mucosal immune system, although other minor networks have been described [34,35]. For the development of effective human mucosal vaccines against pathogenic respiratory bacteria, an oral or other mucosally delivered formulation would be most desirable for a number of reasons. First, oral administration is most likely to pose the least number of complications, although there are advances that are simplifying airway delivery. Second, no sophisticated administration procedures are required. Third, a number of mucosal vaccine formulations are stable without refrigeration, eliminating the 'cold chain' requirement of many vaccines. Fourth, mucosal administration is not invasive and, as such, would most likely have better subject compliance.

While these are very positive advantages for mucosal vaccination, one issue that remains unresolved is that at this point in time many mucosal vaccines do not induce adequate long-term immunity. Three essential criteria have been determined for a successful mucosal immunization strategy [36]:

- The effective delivery of antigen to the mucosal immune induction site
- The enhancement of mucosal immune responses by the use of mucosal immunomodulators (such as bacterial enterotoxins, cytokines, adjuvants and delivery systems)
- Choosing a regime and route of immunization that will induce protective responses and immunological memory at the desired mucosal site and preferably systemically as well

Antigen delivery

Delivering antigens by means other than injection remains one of the most significant challenges facing new generation vaccines and a high priority area in vaccine development [4]. Therefore, why is it that delivering vaccines via the mucosal route has posed such a challenge? A number of factors contribute to the problem. These include:

- The desire for nonreplicating agents due to perceived risks with live or attenuated microorganisms
- The need to overcome mucosal tolerance [37]
- The capacity to deliver the required antigen doses to the inductive sites of mucosal tissues [38]
- The choice of inductive site since this can influence the resulting acquired immune response [39,40]
- The choice of adjuvant and/or carrier system
- Whether these will preserve antigen conformation [41].

This is in addition to knowledge of the nature of both the protective antigens and immune responses that are required (not discussed in this review but is a key issue). It is the complexity of these issues that highlights the fact that there

appears to be no single answer to effective mucosal vaccination approaches, although there are many innovative and promising developments within the field of mucosal vaccine delivery systems (TABLE 1).

A number of adjuvants have been investigated for potential use with mucosal delivery. The most commonly used in animal studies of mucosal immunization have been the protein enterotoxins, cholera toxin (CT) and *E. coli* heat-labile toxin (LT). The use of these enterotoxins as native or modified adjuvant subunits have recently been reviewed [42]. One of the potential limitations with these enterotoxins is that the native adjuvant activity is related to the enterotoxicity and so some of the current challenges are to develop modified CT and LT products that retain adjuvant activity but not toxicity [43]. Both these adjuvants have shown promising results as mucosal adjuvants for oral and nasal delivery of a variety of antigens, inducing both mucosal and serum antibody responses. Therefore, forms of these adjuvants safe for human use would be quite advantageous.

Monophosphoryl lipid A (MPL) is an adjuvant that has been used in humans and has potential for mucosal delivery formulation. MPL in the pneumococcal-CRM₁₉₇ vaccine was safe when administered parenterally to healthy toddlers. The results showed a dose-dependent effect on both Th1 and Th2 responses to the carrier protein but did not enhance serum IgG to any of the polysaccharide antigens [44]. Nasally formulated hepatitis B surface antigen (HbsAg), tetanus toxoid (TT) and FluShield[®] influenza vaccine in MPL in animal studies resulted in enhanced mucosal and systemic immunity [45]. The most significant difference between the mucosal and parenteral delivery in these studies was the production of specific IgA in mucosal secretions following intra nasal immunization. MPL-based adjuvants have also been successfully used to boost immune responses in mouse studies, resulting in substantial protection from aerosol challenge with *M. tuberculosis* [46]. This result is particularly exciting due to the poor efficacy of the BCG in countries with endemic *M. tuberculosis* and could be used to orally boost responses using a subunit vaccine.

Another agent that has recently shown promise as a mucosal adjuvant is the synthetic oligodeoxynucleotides containing CpG stimulatory sequences. This adjuvant has been used with the *H. influenzae* type b polysaccharide (PRP) vaccine and when delivered mucosally, resulted in both systemic and mucosal antibody, particularly mucosal IgA [47] (as observed for the vaccines in the Baldridge study [45]). To date, application using CpG DNA sequences are predominantly in animal studies. Very limited data are available on any human clinical trials, however, it is reported that the preliminary data from some early stage human trials are encouraging. Hence, this may be a very promising technology with both mucosal and parenteral adjuvant application [48]. Other adjuvants are regularly reported with model antigens but recently, a lipid core peptide (LCP) adjuvant has been used with group A streptococci antigens [49] that currently appears to have potential as a mucosal adjuvant suitable for human application.

Many studies have investigated bacterial systems for delivery of antigens. These have ranged from whole-killed-cell vaccines [50] to more complex engineered micro-organisms [51]. Some of the advantages of bacterial delivery systems are that they contain their own adjuvants, they naturally form and maintain their microsphere structure and once the recombinant technologies have been incorporated, they can be efficient to produce. Some of the limitations are potential toxic effects, poor or ineffective immune responses to the vaccine antigens, possible pathogenic issues with live micro-organisms and issues with the use of antibiotic resistance genes in their preparation. As such, while some of our existing vaccines are based on whole-cell systems and many of the bacterial systems indicated in TABLE 1 show definite promise, a broader acceptance for human use needs to be demonstrated. *Salmonella* species have been extensively used as a system for investigating live bacterial carrier systems and paved the way for investigations into a range of such carriers [52]. The range of human clinical trials using these systems has not been extensive and in those that have been done, the results would suggest that there is still a need for more development of these systems (reviewed in [52]).

Studies with nonliving bacterial systems in mice using *Vibrio cholerae* ghosts expressing a major outer membrane protein from *Chlamydia trachomatis* [53] and *E. coli* ghosts expressing an *H. influenzae* antigen in recombinant S-layers [54], showed induction of immune responses to the respective antigens but demonstrated a need for further development of the system. The bacterial ghost technology is limited to Gram-negative bacteria and has been effective against the pig respiratory pathogen, *Actinobacillus pleuropneumoniae* [55]. Such nonliving bacterial systems are most likely quite safe as oral vaccines but whether they will meet the requirements of drug regulators remains an issue.

Mycobacterium sp. has been evaluated for mucosal delivery as both a vaccine and carrier of other antigens or immunomodulators. Oral and nasal delivery of formulated mycobacteria and BCG vaccines have been shown to be protective in mouse studies [56–59], demonstrating the feasibility of mucosal delivery as a safe and practical method for controlling tuberculosis. In addition, recombinant *M. bovis* BCG-based vaccines have been investigated for both immunopotentiality of the BCG vaccine and as delivery of heterologous antigens (reviewed in [60,61]). BCG is not the only currently available whole cell-based vaccine for bacterial respiratory infections to be investigated for mucosal delivery. The acellular pertussis vaccine delivered intranasally has been investigated in a mouse model for induction of both mucosal and systemic antibody responses in mice [62]. In this study, intranasal immunization induced titers of IgG in the serum and IgA in the lungs that were protective against bacterial challenge and which were reported to be more effective against infection than when mice were immunized subcutaneously.

Microparticles, such as ISCOMs and liposomes, have been investigated quite extensively [63]. ISCOMs are a

licensed carrier for some veterinary vaccines. As a mucosal carrier, it has been used in both oral and intranasal immunization studies and appears to produce a mixed Th1- and Th2-type immune response [64]. It has also been used to investigate delivery of a *M. tuberculosis* antigen [65]. Micro-particle technologies have been more extensively studied for viral respiratory pathogens but there have been few studies reported investigating the potential for delivery of bacterial antigens. Oral delivery of a liposome formulation containing outer membrane proteins from NTHi enhanced nasal clearance in a mouse model [66], whereas pneumococcal polysaccharide delivered in a liposome formulation induced a relatively poor antipolysaccharide response [67]. The development of new formulations of liposomes and other micro-particles continues as a means of targeting antigen to the mucosa-associated lymphoid tissues [68] and they certainly have potential for safe use in humans.

Research efforts are also focusing on even smaller particles for antigen delivery. Nanoparticles have been developed specifically for nasal immunization. These are cationic cross-linked polysaccharides enclosed in lipidic bilayer and comprise of different formulations. Among the antigens tested, nanoparticles have been used to deliver DNA [69], group C meningococcal antigens [70] and antineoplastic proteoglycans from *M. vaccae* [71]. The results of these studies demonstrate potential for this technology to be applied for delivery of other bacterial antigens.

In concluding, a section on mucosal delivery would not be complete without mentioning edible vaccines. These have recently been reviewed by Mason and colleagues [72]. Potato, tomato, banana, legumes, cereals and alfalfa are edible species that have been transformed with varying degrees of success. Various issues associated with both the production of and immunization with plants as vaccine delivery vehicles remain to be resolved. One possibility is that they may be effective in boosting immune responses that were primed by a parenteral vaccine, as has been found for a clinical trial of the HBsAg expressed in potato [Y. THANAVALA, ROSWELL PARK CANCER INSTITUTE, NY, USA. PERS. COMM.].

Despite the excellent research efforts, antigen delivery systems remain one of the key issues in the development of preventative therapies for respiratory bacterial infections. A recent issue of the journal *Vaccine* (vol. 21, issue 9–10, 2003) was specifically dedicated to immunomodulating agents and provides a more extensive review of this field. In this section we have highlighted a range of developments in the mucosal delivery of vaccines and cited examples where there have been studies associated with the respiratory bacteria discussed in this review. What is apparent is that there remains a need for more extensive evaluation of a number of the more promising technologies to determine their suitability for inducing protective immune responses in the respiratory tract against a broader range of bacterial pathogens. There is some excellent basic research with many of the technologies discussed but for some there are obstacles

affecting the progress to clinical evaluation. The lack of reliable outcome measures from clinical trials is an issue confronting development of all vaccines and delivery technologies. Efficacy of human vaccines is still frequently equated to antibody titers, yet preclinical evidence often indicates the importance of cellular immune responses. Other issues that delay the progress include the costs of clinical trials, a significant factor affected by commercial decisions and the processes of meeting the requirements of drug regulators. While it is necessary to ensure appropriate safety, sometimes the conditions applied by regulatory agencies to mucosal vaccines are more relevant to pharmaceuticals and parenteral vaccines than to these formulations.

Expert opinion

Mucosal adjuvants and new delivery technologies have demonstrated potential in the fight against diseases such as tuberculosis. The problems of poor efficacy of the current parenterally delivered vaccine may be an issue of the past if further trials on the mucosally delivered booster antigens have positive outcomes. However, while technology for enhancing immune responses against bacterial organisms and their subunits has advanced markedly in recent years, the move to clinical trials to confirm animal studies in humans remains an impediment. One of the major problems in the use of animal models and predicting results in humans are the effects the vaccines will have on the normal microbes colonizing the respiratory tract. As focussed on in the review, one of the major risks in vaccination is the replacement of bacterial colonizers with potentially more pathogenic microbes (as already seen with *S. pneumoniae*). Reliable data on the balance of colonization versus disease can only be confirmed once tested in human trials. Hence, efforts should continue on the development of new vaccines and the evaluation of those currently under trial. It is most likely that if replacement colonization were to occur, this would take some years. During this time, if an efficacious vaccine technology was established against the microbe, rapid adjustments could be made to formulations. In this scenario, regulatory authorities will be challenged to provide appropriate approval pathways.

This review has therefore looked at the epidemiology and burden of disease presented by bacterial pathogens in the respiratory tract. It has examined the questions of potential pathogen eradication from the respiratory tract versus prevention of disease and provided a brief synopsis on the state of the art in the delivery of vaccine antigens at the mucosal surface. Key issues for the future are addressed and an opinion on where we will be in 5 years is offered.

Five-year view

In 5 years it is feasible that some of the current developments in mucosal delivery systems will be in clinical trials. The availability of genomes for most of these bacteria means that a new range of antigens will have been evaluated as vaccine candidates. A new generation of pneumococcal vaccines will have been developed to overcome the efficacy and serotype coverage problems associated with the current conjugate vaccine. It is desirable that there will be acceptable delivery systems available for routine use in humans and that we will have knowledge from human vaccine studies that enables the identification of the type of immune mechanisms required for pathogen-specific protection.

Key issues

- Establishment of these respiratory pathogens as a priority in the development of effective multivalent vaccines.
- Knowledge of the immune responses that control the bacterial colonizers of the respiratory mucosal surfaces is essential to optimizing therapeutic initiatives.
- Identification of the measures of pathogen-specific protective responses required at the mucosal surfaces must be resolved to ensure vaccine efficacy.
- Determination of the requirement for a vaccine against a 'commensal pathogen' to induce sterile immunity against a microbe.
- Investigation on whether a vaccine that creates sterile immunity to a commensal would open a niche for new pathogens to colonize the nasopharynx.
- Adoption of appropriate and cost-effective pathways by regulatory authorities to enable rapid changes to vaccine formulation in cases where safety has been demonstrated and there is an established technology track record.
- Dissemination of information that ensures public awareness of the importance of maintaining vaccine compliance for pertussis and diphtheria vaccines.
- The design of therapeutic and prophylactic approaches that prevent resurgence of disease causing pathogens as has occurred with tuberculosis
- Although not discussed in this review, identification of key antigens for inclusion in vaccines remains an issue for a number of these bacteria
- Continued development and more timely advancement to clinical evaluation of effective multivalent vaccines as a more effective means of ensuring vaccine coverage against a range of pathogens.
- The development of mucosal delivery systems that are inexpensive, effectively target antigen to the immune induction sites and are safe for human use.

References

Papers of special note have been highlighted as:

• of interest

•• of considerable interest

- 1 Nicolau D. Clinical and economic implications of antimicrobial resistance for the management of community-acquired respiratory tract infections. *J. Antimicrob. Chemother.* 50, 61–70 (2002).
- Extensive analysis and review of antimicrobial resistance and what this means for the patient and health care system.
- 2 Zahrt TC. Molecular mechanisms regulating persistent *Mycobacterium tuberculosis* infection. *Microbes Infect.* 5, 159–167 (2003).
- 3 Center for Disease Control, http://www.cdc.gov/ncidod/dbmd/diseaseinfo/pneumchilddevcount_t.htm, accessed April, 2003
- 4 Center for Disease Control. Geographic variation in penicillin resistance in *Streptococcus pneumoniae* - selected sites, United States, 1997. *MMWR* 4830, 651–661 (1999).
- 5 Schrag S, Beall B, Dowell S. Limiting the spread of resistant pneumococci: biological and epidemiologic evidence for the effectiveness of alternative interventions. *Clin. Microbiol. Rev.* 13, 588–601 (2000).
- 6 Snell N. Examining unmet needs in infectious disease. *Drug Discov. Today* 8, 22–30 (2003).
- 7 Puli V, Clarke SC. Clinical aspects of pneumonia. *Br. J. Biomed. Sci.* 59, 170–172 (2002).
- 8 Miravittles M, Murio C, Guerrero T, Gisbert R. Costs of chronic bronchitis and COPD: a 1-year follow-up study. *Chest* 123, 784–791 (2003).
- 9 Dagan R, Klugman KP, Craig WA, Baquero F. Evidence to support the rationale that bacterial eradication in respiratory tract infection is an important aim of antimicrobial therapy. *J. Antimicrob. Chemother.* 47, 129–140 (2001).
- 10 Gates GA. Cost-effectiveness considerations in otitis media treatment. *Otolaryngol Head Neck Surg.* 114, 525–530 (1996).
- 11 Boswell J. Presentation of early otitis media in 'top-end' Aboriginal infants. *Aus. New Zeal. J. Pub. Health* 21, 100–103 (1997).
- 12 Dagan R, Leibovitz E. Bacterial eradication in the treatment of otitis media. *Lancet Infect. Dis.* 2, 593–604 (2002).
- 13 Obaro S, Adegbola R. The pneumococcus: carriage, disease and conjugate vaccines. *J. Med. Microbiol.* 51, 98–104 (2002).
- 14 Poland GA. The burden of pneumococcal disease: the role of conjugate vaccines. *Vaccine* 17, 1674–1679 (1999).
- 15 Verduin CM, Hol C, Fler A, van Dijk H, van Belkum A. *Monaxella catarrhalis*: from emerging to established pathogen. *Clin. Microbiol. Rev.* 15, 125–144 (2002).
- 16 Foxwell AR, Kyd JM, Cripps AW. Nontypeable *Haemophilus influenzae*: pathogenesis and prevention. *Microbiol. Molec. Biol. Rev.* 62, 294–308 (1998).
- 17 Garcia-Rodriguez JA, Fresnadillo Martinez MJ. Dynamics of nasopharyngeal colonisation by potential respiratory pathogens. *J. Antimicrob. Chemother.* 50(Suppl. C), 59–74 (2002).
- Excellent review on factors contributing to increased risk of carriage of potential respiratory pathogens and clinical infection.
- 18 Rosenstein N, Perkins B. Update on *Haemophilus influenzae* serotype B and meningococcal vaccines. *Pediatr. Clin. N. Am.* 47, 337–352 (2000).
- 19 Flynn JL, Chan J. Tuberculosis: latency and reactivation. *Infect. Immun.* 69, 4195–4201 (2001).
- 20 Reljic R, Williams A, Marsh P *et al.* IgA antibody modulation of tuberculosis infection *in vivo* and *in vitro*. *Mucosal Immunol. Update* 10, 2693 (2002).
- 21 Centre for Disease Control. Vaccine Fact Sheet. *National Vaccine Program Office*. http://www.cdc.gov/od/nvpo/fs_table1_doc3.htm (2003).
- 22 Skogen V, Cherkasova V, Maksimova N *et al.* Molecular characterization of *Corynebacterium diphtheriae* isolates, Russia 1957–1987. *Emerg Infect Dis* 8, 516–518 (2002).
- 23 Campins-Marti M, Cheng HK, Forsyth K *et al.* Recommendations are needed for adolescent and adult pertussis immunisation: rationale and strategies for consideration. *Vaccine* 20, 641–646 (2001).
- 24 von Konig C, Halperin S, Riffelmann M, Guiso N. Pertussis of adults and infants. *Lancet Infect. Dis.* 2, 744–750 (2002).
- 25 Pelton SI, Dagan R, Gaines BM *et al.* Pneumococcal conjugate vaccines: proceedings from an Interactive Symposium at the 41st Interscience Conference on Antimicrobial Agents and Chemotherapy. *Vaccine* 21, 1562–1571 (2003).
- 26 Eskola J, Kilpi T, Palmu A *et al.* Efficacy of a pneumococcal conjugate vaccine against acute otitis media. *New Engl. J. Med.* 344, 403–409 (2001).
- Demonstration of vaccine efficacy and strain replacement.
- 27 Lehmann D, Coakley KJ, Coakley CA *et al.* Reduction in the incidence of acute bronchitis by an oral *Haemophilus influenzae* vaccine in patients with chronic bronchitis in the highlands of Papua New Guinea. *Am. Rev. Respir. Dis.* 144, 324–330 (1991).
- 28 Aniansson G, Alm B, Andersson B *et al.* A prospective cohort study on breast-feeding and otitis media in Swedish infants. *Pediatr. Infect. Dis. J.* 13, 83–88 (1994).
- 29 Harabuchi Y, Faden H, Yamanaka N *et al.* Nasopharyngeal colonization with nontypeable *Haemophilus influenzae* and recurrent otitis media. *J. Infect. Dis.* 170, 862–866 (1994).
- 30 Faden H, Duffy L, Wasielewski R *et al.* Relationships between nasopharyngeal colonization and the development of otitis media in children. *J. Infect. Dis.* 175, 1440–1445 (1997).
- 31 Leach AJ, Boswell JB, Asche V, Nienhuys TG, Mathews JD. Bacterial colonisation of the nasopharynx predicts very early onset and persistence of otitis media in Australian Aboriginal infants. *Pediatr. Infect. Dis. J.* 13, 983–989 (1994).
- 32 Ogra PL, Faden H, Welliver RC. Vaccination strategies for mucosal immune responses. *Clin. Microbiol. Rev.* 14, 430–445 (2001).
- Good review on mucosal immune vaccination strategies
- 33 Anderson AO. Peripheral and Mucosal Immunity: Critical issues for oral vaccine design. In: *Peripheral and Mucosal Immunity after Oral vaccination*, <http://www.geocities.com/CapeCanaveral/Hangar/1962/crossregulation.html> 1–8. (2001).
- 34 Eriksson K, Holmgren J. Recent advances in mucosal vaccines and adjuvants. *Curr. Opin. Immunol.* 14, 666–672 (2002).
- 35 Russell MW, Martin MH, Wu H-Y *et al.* Strategies of immunization against mucosal infections. *Vaccine* 19, S122–127 (2001).
- 36 Cripps AW, Kyd JM, Foxwell AR. Vaccines and mucosal immunization. *Vaccine* 19, 2513–2515 (2001).
- 37 Mowat AM, Weiner HL. Oral tolerance: physiological basis and clinical applications. In: *Mucosal Immunology*. Ogra PL, Mestecky J, Lamm ME, Strober W, McGhee JR, Bienenstock J (Eds). Academic Press, NY, USA, 587–582 (1999).
- 38 Neutra MR, Pringault E, Kraehenbuhl JP. Antigen sampling across epithelial barriers and induction of mucosal immune responses. *Ann. Rev. Immunol.* 14, 275–300 (1996).

- 39 Murphy TF, Kyd JM, John A, Kirkham C, Cripps AW. Enhancement of pulmonary clearance of *Monaxella* (*Branhamella*) *catarrhalis* following immunization with outer membrane protein CD in a mouse model. *J. Infect. Dis.* 178, 1667–1675 (1998).
- 40 Mestecky J, Michalek SM, Moldoveanu Z, Russell MW. Routes of immunization and antigen delivery systems for optimal mucosal immune responses in humans. *Behring Inst. Mitt.* 98, 33–43 (1997).
- 41 Mahon BP. The ration design of vaccine adjuvants for mucosal and neonatal immunization. *Curr. Med. Chem.* 8, 1057–1075 (2001).
- 42 Elson CE, Dertzbaugh MT. Mucosal adjuvants. In: *Mucosal Immunology*. Ogra PL *et al.* (Eds). Academic Press, NY, USA, 817–838, (1999).
- 43 Pizzo M, Giuliani MM, Fontana MR *et al.* Mucosal vaccines: non toxic derivatives of LT and CT as mucosal adjuvants. *Vaccine* 19, 2534–2542 (2001).
- 44 Vernacchio L, Bernstein H, Pelton S *et al.* Effect of monophosphoryl lipid A (MPL) on T-helper cells when administered as an adjuvant with pneumococcal-CRM₁₉₇ conjugate vaccine in healthy toddlers. *Vaccine* 20, 3658–3667 (2002).
- 45 Baldridge JR, Yorgensen Y, Ward JR, Ulrich JT. Monophosphoryl lipid A enhances mucosal and systemic immunity to vaccine antigens following intranasal administration. *Vaccine* 18, 2416–2425 (2000).
- 46 Doherty TM, Olsen AW, van Pinxteren I, Andersen P. Oral vaccination with subunit vaccines protects animals against aerosol infection with *Mycobacterium tuberculosis*. *Infect. Immun.* 70, 3111–3121 (2002).
- **Demonstration of mucosal vaccination as a boosting mechanism.**
- 47 Mariotti S, Teloni R, von Hunolstein C *et al.* Immunogenicity of anti-*Haemophilus influenzae* type b CRM₁₉₇ conjugate following mucosal vaccination with oligonucleotide containing immunostimulatory sequences as adjuvant. *Vaccine* 20, 2229–2239 (2002).
- 48 Agrawal S, Kandimalla ER. Medicinal chemistry and therapeutic potential of CpG DNA. *Trends Mol. Med.* 8, 114–121 (2002).
- **Novel adjuvant.**
- 49 Olive C, Batzloff M, Horváth A *et al.* Potential of lipid cord peptide technology as a novel self-adjuvanting vaccine delivery system for multiple different synthetic peptide immunogens. *Infect. Immun.* 71, 2373–2383 (2003).
- 50 Kyd JM, Cripps AW. Killed whole bacterial cells, a mucosal delivery system for the induction of immunity in the respiratory tract and middle ear: An overview. *Vaccine* 17, 1775–1781 (1999).
- 51 Szostak MP, Hensel A, Eko FO *et al.* Bacterial ghosts: non living candidate vaccines. *J. Biotechnol.* 44, 161–170 (1996).
- 52 Pasetti MF, Levine MM. Animal models paving the way for clinical trials of attenuated *Salmonella enterica* serovar Typhi live oral vaccines and live vectors. *Vaccine* 21, 401–418 (2003).
- **Demonstration of *Salmonella* in human clinical trials.**
- 53 Eko FO, Lubitz W, McMillan L *et al.* Recombinant *Vibrio cholerae* ghosts as a delivery vehicle for vaccinating against *Chlamydia trachomatis*. *Vaccine* 21, 1694–1703 (2003).
- 54 Riedmann EM, Kyd JM, Smith AM *et al.* Construction of recombinant S-layer proteins (rSbsA) and their expression in bacterial ghosts – a delivery system for the nontypeable *Haemophilus influenzae* antigen Omp26. *FEMS Immunol. Med. Microbiol.* 37, 185–192 (2003).
- 55 Huter V, Hensel A, Brand E, Lubitz W. Improved protection against lung colonization by *Actinobacillus pleuropneumoniae* ghosts: characterization of a genetically inactivated vaccine. *J. Biotechnol.* 83, 161–172 (2000).
- 56 Aldwell FE, Tucker IG, de Lisle GW, Buddle BM. Oral delivery of *Mycobacterium bovis* BCG in a lipid formulation induces resistance to pulmonary tuberculosis in mice. *Infect. Immun.* 71, 101–108 (2003).
- 57 Manabe YC, Scott CP, Bishai WR. Naturally attenuated, orally administered *Mycobacterium microti* as a tuberculosis vaccine is better than subcutaneous *Mycobacterium bovis* BCG. *Infect. Immun.* 70, 1566–1570 (2002).
- 58 Falero-Diaz G, Challacombe S, Banerjee D *et al.* Intranasal vaccination of mice against infection with *Mycobacterium tuberculosis*. *Vaccine* 18, 3223–3229 (2000).
- 59 Biet F, Kremer L, Wolowczuk I, Delacré M, Lochet C. Immune response induces by recombinant *Mycobacterium bovis* BCG producing the cholera toxin B subunit. *Infect. Immun.* 71, 2933–2937 (2003).
- 60 Dietrich G, Viret J-F, Hess J. *Mycobacterium bovis* BCG-based vaccines against tuberculosis: novel developments. *Vaccine* 21, 667–670 (2003).
- **Good review on *Mycobacterium* vaccines.**
- 61 Dietrich G, Viret J-F, Hess J. Novel vaccination strategies based on recombinant *Mycobacterium bovis* BCG. *Int. J. Med. Microbiol.* 292, 441–451 (2003).
- 62 Isaka M, Yasuda Y, Taniguchi T *et al.* Mucosal and systemic antibody responses against an acellular pertussis vaccine in mice after intranasal co-administration with recombinant cholera toxin B subunit as an adjuvant. *Vaccine* 21, 1165–1173 (2003).
- 63 Kersten GFA, Crommelin DJA. Liposomes and ISCOMs. *Vaccine* 21, 915–920 (2003).
- 64 Smith RE, Donachie AM, Mowat AM. Immune stimulating complexes as mucosal vaccines. *Immunol. Cell Biol.* 76, 263–269 (1998).
- 65 da Fonseca DP, Frerichs J, Singh M, Snippe H, Verheul AFM. Induction of antibody and T-cell responses by immunization with ISCOMS containing the 38-kilodalton protein of *Mycobacterium tuberculosis*. *Vaccine* 19, 122–131 (2000).
- 66 Suzuki M, Kurono Y, Kodama H, Shigemori H, Mogi G. Enhancement of nasal clearance of nontypeable *Haemophilus influenzae* by oral immunization with outer membrane proteins. *Acta Otolaryngol.* 118, 864–869 (1998).
- 67 VanCott JL, Kobayashi H, Yamamoto M *et al.* Induction of pneumococcal polysaccharide-specific mucosal immune responses by oral immunization. *Vaccine* 14, 392–398 (1996).
- 68 Zho F, Neutra M. Antigen delivery to mucos-associated lymphoid tissues using liposomes as a carrier. *Biosci. Rep.* 22, 355–369 (2002).
- 69 Cui Z, Mumper RJ. Intranasal administration of plasmid DNA-coated nanoparticles results in enhanced immune responses. *J. Pharm. Pharmacol.* 54, 1195–1203 (2002).
- 70 Baudner BC, Bolland O, Giuliani MM *et al.* Enhancement of protective efficacy following intranasal immunization with vaccine plus a nontoxic LTK63 mutant delivered with nanoparticles. *Infect. Immun.* 70, 4785–4790 (2002).
- 71 Tian XX, Groves MJ. Formulation and biological activity of antineoplastic proteoglycans derived from *Mycobacterium vaccae* in chitosan nanoparticles. *J. Pharm. Pharmacol.* 51, 151–157 (1999).
- 72 Mason HS, Warzecha H, Mor T, Arntzen CJ. Edible plant vaccines: applications for prophylactic and therapeutic molecular medicine. *Trends Molec. Med.* 8, 324–329 (2002).

- 73 Glenn GM, Scharon-Kersten T, Vassell R *et al.* Cutting edge: transcutaneous immunization with cholera toxin protects mice against lethal mucosal toxin challenge. *J. Immunol.* 161, 3211–3214 (1998).
- 74 Ulrich JT, Myers KR. Monophosphoryl lipid A as an adjuvant. In: *Vaccine Design: The Subunit and Adjuvant Approach*. Powell MF, Newman MJ (Eds). Plenum Press, NY, USA 465–524 (1995).
- 75 Childers NK, Miller KL, Tong G *et al.* Adjuvant activity of monophosphoryl lipid A for nasal and oral immunization with soluble or liposome-associated antigen. *Infect. Immun.* 68, 5509–5516 (2000).
- 76 Zong G, Toth L, Reid R, Brunham RC. Immunogenicity evaluation of a lipidic amino acid based synthetic peptide vaccine for *Chlamydia trachomatis*. *J. Immunol.* 151, 3728–3736 (1993).
- 77 Nayak AR, Ting SA, Tart RC *et al.* A live recombinant avirulent oral *Salmonella* vaccine expressing pneumococcal surface protein A induces protective responses against *Streptococcus pneumoniae*. *Infect. Immun.* 66, 3744–3751 (1998).
- 78 Hess J, Kaufmann SH. Live antigen carriers as tools for improved anti-tuberculosis vaccines. *FEMS Immunol. Med. Microbiol.* 23, 165–173 (1999).
- 79 Huang Y, Hajishengallis G, Michalek SM. Induction of protective immunity against *Streptococcus mutans* colonization after mucosal immunization with attenuated *Salmonella enterica* serovar typhimurium expressing an *S. mutans* adhesin under the control of *in vivo*-inducible *nirB* promoter. *Infect. Immun.* 69, 2154–2161 (2001).
- 80 Kang HY, Srinivasan J, Curtiss R, III. Immune responses to recombinant pneumococcal PspA antigen delivered by live attenuated *Salmonella enterica* serovar Typhimurium vaccine. *Infect. Immun.* 70, 1739–1749 (2002).
- 81 Wells JM, Robinson K, Chamberlain LM, Schofield KM, Page RWFL. Lactic acid bacteria as vaccine delivery vehicles. *Lactic Acid Bacteria: Curr. Adv. Metabol. Genetics Applications* 70, 317–330 (1996).
- 82 Kyd J, Cripps A. Identifying vaccine antigens and assessing delivery systems for the prevention of bacterial infections. *J. Biotechnol.* 83, 85–90 (2000).
- 83 Gilbert C, Robinson K, Le Page RWF, Wells JM. Heterologous expression of an immunogenic pneumococcal type 3 capsular polysaccharide in *Lactococcus lactis*. *Infect. Immun.* 68, 3251–3260 (2000).
- 84 Norton P, Wells J, Brown H, Macpherson A, Page RL. Protection against tetanus toxin in mice nasally immunized with recombinant *Lactococcus lactis* expressing tetanus toxin fragment C. *Vaccine* 15, 616–619 (1997).
- 85 Grangette C, Muller-Alouf H, Coudercourt D *et al.* Mucosal immune responses and protection against tetanus toxin after intranasal immunization with recombinant *Lactobacillus plantarum*. *Infect. Immun.* 69, 1547–1553 (2001).
- 86 Eko FO, Mayr UB, Attridge SR, Lubitz W. Characterization and immunogenicity of *Vibrio cholerae* ghosts expressing toxin-coregulated pili. *J. Biotechnol.* 83, 115–123 (2000).
- 87 Szostak MP, Mader H, Truppe M *et al.* Bacterial ghosts as multifunctional vaccine particles. *Behring Inst. Mitt.* 98, 191–196 (1997).
- 88 Duc LH, Hong HA, Fairweather N, Ricca E, Cutting SM. Bacterial spores as vaccine vehicles. *Infect. Immun.* 71, 2810–2818 (2003).
- 89 Mowat AM, Donachie AM, Jagewall S *et al.* CTA1-DD-immune stimulating complexes: a novel, rationally designed combined mucosal vaccine adjuvant effective with nanogram doses of antigen. *J. Immunol.* 167, 3398–3405 (2001).
- 90 Furrie E, Smith RE, Turner MW, Strobel S, Mowat AM. Induction of local innate immune responses and modulation of antigen uptake as mechanisms underlying the mucosal adjuvant properties of immune stimulating complexes (ISCOMS). *Vaccine* 20, 2254–2262 (2002).
- 91 Klavinskis LS, Gao L, Barnfield C, Lehner T, Parker S. Mucosal immunization with DNA-liposome complexes. *Vaccine* 15, 818–820 (1997).
- 92 Hyde SC, Southern KW, Gileadi U *et al.* Repeat administration of DNA/liposome to the nasal epithelium of patients with cystic fibrosis. *Gene Ther* 7, 1156–1165 (2000).
- 93 Joseph A, Louria-Hayon I, Plis-Finarov A *et al.* Liposomal immunostimulatory DNA sequence (ISS-ODN): an efficient parenteral and mucosal adjuvant for influenza and hepatitis B vaccines. *Vaccine* 20, 3342–3354 (2002).

Affiliations

- A Ruth Foxwell, PhD, Senior Lecturer in Microbiology, Gadi Research Centre for Medical and Health Sciences, University of Canberra, Canberra ACT 2601, Australia
Tel.: +61 2 6201 2089
Fax: +61 2 6201 2461
ruth.foxwell@canberra.edu.au
- Jennelle M Kyd, PhD, Professor of Forensic Sciences, Gadi Research Centre for Medical and Health Sciences, University of Canberra, Canberra ACT 2601, Australia.
jennelle.kyd@canberra.edu.au
Tel.: +61 2 6201 2160, Fax: +61 2 6201 2461
- Allan W Cripps, PhD, Professor of Health, Gadi Research Centre for Medical and Health Sciences, University of Canberra, Canberra ACT 2601, Australia.
Tel.: +61 7 5552 8595
Fax: +61 7 5552 8777
allan.cripps@griffith.edu.au