**Monitoring by telemetry reveals differences in movement and survival
following hatchery or wild-rearing of an endangered fish**

B. C. Ebner^{a, b*} and J. D. Thiem^{a, c}

^aParks, Conservation and Lands, Department of Territory and Municipal Services, ACT
Government, GPO Box 158, Canberra ACT 2601, Australia.

^bPresent address: Australian Rivers Institute, Griffith University, Nathan, QLD 4111,
Australia

^cPresent address: Institute for Applied Ecology, University of Canberra, Bruce, ACT
2601, Australia

* Author to whom correspondence should be addressed. Tel.: +61 7 4092 5768; email:
b.ebner@griffith.edu.au

Abstract

Species reintroduction is a management strategy used to conserve endemic fish biodiversity. We investigated stocking on-grown endangered trout cod (*Maccullochella macquariensis*) in the Murrumbidgee River, Australia. We also tested the hypothesis that post-juvenile dispersal underpins the long term scarcity of adults recorded at fingerling stocking locations. Radio-tracking was used to quantify dispersal of stocked sub-adults (two-year old, hatchery fish, n=27) compared with fish originally stocked as fingerlings (unknown age, wild fish, n=31), but we encountered poor survivorship of the former group (survivorship = 9% and 95%, respectively, at 13 months post-release). The hatchery group exhibited both limited dispersal and large scale dispersal (up to 55 km) downstream from the release site. Wild fish exhibited limited net dispersal, occupying home-ranges within a 13-km reach, and occasionally undertook large scale excursions (10–70 km). We conclude that a) re-establishment of cod populations based on release of on-grown fish is not straightforward, and b) that adults of this species have an ability to disperse away from stocking sites. The study demonstrates the benefit of using radio-tracking to monitor the movement and survivorship of stocked threatened fish and indicates a need to consider the effects of hatchery-rearing, when conducting fish reintroductions.

Key words: radio-tracking; dispersal; survivorship; reintroduction; hatchery; *Maccullochella*

49

50 **Introduction**

51

52 Species reintroduction programs (Wallace 2000), including those for threatened fishes
53 (Brown and Day 2002) should be based on a detailed and transparent platform of
54 experimental research (Fischer and Lindenmayer 2000). Translocation of wild fish and
55 stocking of hatchery-reared fish are common approaches to re-establishing populations
56 of threatened species (Minckley 1995) and enhancing fisheries (Molony *et al.* 2003).
57 Comparison of wild and hatchery-reared fish in the hatchery (Metcalf *et al.* 2003;
58 Salvenes and Braithwaite 2005) and the wild (Thorstad *et al.* 1998; Dieperink *et al.*
59 2001; Bettinger and Bettoli 2002) has provided useful insights into shortcomings of
60 stocking hatchery-reared fish, especially salmonids in the Northern Hemisphere.

61 Behavioural changes due to domestication often reduce the survival of stocked fish,
62 resulting in highly variable success among fish stocking programs (Dieperink *et al.*,
63 2001; Bettinger and Bettoli 2002).

64

65 Familiarity of wild fish with the river environment is probably a major advantage
66 relative to newly released hatchery fish (Brown and Day 2002). For instance the
67 movement and dispersal of hatchery and wild-reared fish is likely to differ if wild fish
68 are capable of homing to a territory or home-range that they have occupied previously,
69 whereas, hatchery-reared fish are naïve to the new environment. In this regard, studies
70 comparing search behaviour, homing and home-range occupation (e.g. Crook 2004a, b)
71 of fish offer promise for informing species reintroductions programs. Individuals can
72 disperse immediately following release or later undertake home-range shifts (Crook

2004a), leading to emigration and insufficient numbers of adults at the release site to establish a self-sustaining population (Armstrong and Seddon 2008).

In Australia relative to the Northern Hemisphere there has not been a comparable research focus on fish stocking in freshwaters, despite widespread stocking of freshwater fishes including species of the family Percichthyidae (Lintermans *et al.* 2005; Gillanders *et al.* 2006; Lintermans 2006). In temperate mainland Australia, the family Percichthyidae is the most identifiable group of endemic freshwater fishes from the public perspective. This is because the family includes large-bodied species of freshwater cod in the genus *Maccullochella* and the perches of *Macquaria* (Harris and Rowland 1996). These fishes are targeted by recreational anglers and in a number of cases represent conservation listed species (Morris *et al.* 2001; Lintermans and Phillips 2004).

Trout cod *Maccullochella macquariensis* (Cuvier) is a nationally endangered freshwater fish of the family Percichthyidae, endemic to rivers in the southeast of the Murray-Darling Basin, Australia (Ingram and Douglas 1995). The species is conservation-listed as endangered in the Australian Capital Territory (ACT) and New South Wales (NSW) and critically endangered in Victoria (Morris *et al.* 2001; DSE 2003; ACT Government 2007). Efforts to recover the species have involved the largest freshwater conservation-stocking program in the country (Lintermans *et al.* 2005; Lintermans 2006). Since 1986, hatchery-produced fingerlings have been stocked into areas where the natural population was locally extinct in NSW, Victoria and the ACT, in an effort to re-establish populations (Douglas *et al.* 1994; Gilligan 2005). Large rivers have been the

focus of stocking efforts as they were assumed to be the primary habitats of *M. macquariensis*. There has generally been minimal or sporadic monitoring at reintroduction sites, and knowledge of successful survivorship has been mainly from angler reports. Survival and growth of fingerlings through to the age of 2–3 years (juveniles) has been documented at a number of sites (Faragher *et al.* 1993; Lintermans 1995; Brown 1998; Brown *et al.* 1998; Douglas and Brown 2000; Gilligan 2005; NSW DPI 2005). The number of 3–5 year old individuals (sub-adult or young adults, Harris and Rowland 1996) captured in monitoring programs is much lower. Reports of large adult individuals have been uncommon. Possible explanations for the scarcity of adults in re-stocked populations are: 1) larger fish (adults) are subject to high mortality including angling pressure and/or natural mortality, 2) larger fish remain at the release site but are undetected by monitoring, and/or 3) juveniles remain at the release site until they reach sub-adulthood at which stage they disperse, becoming difficult to detect.

Generally, fish populations suffer greatest natural mortality in early life history phases (Jones 1991) and this presumably applies to *M. macquariensis* (Todd *et al.* 2004). It is unlikely that adult fish are subject to high mortality, though, an understanding of mortality in this species is a major knowledge gap in recovery of the species (Simon Nicol, Department of Sustainability and Environment Victoria, pers. comm.). It is also unlikely that larger fish remain at the site and are not detected by monitoring. Electro-fishing has been a commonly employed monitoring technique for *M. macquariensis* and other freshwater cod (Faragher *et al.* 1993; Growns *et al.* 2004; Ebner *et al.* 2008) and has proven successful in detecting a wide range of size classes at a subset of sites (Ebner *et al.* 2006).

The main hypothesis that we tested was that post-juvenile dispersal is the reason for the scarcity of sub-adult and adult *M. macquariensis* recorded at stocking locations. Despite indications that juveniles can be gregarious (Douglas *et al.* 1994), it is possible that as individuals grow and mature they become territorial, leading to dispersal from the release sites, since under hatchery conditions individuals become particularly aggressive towards one another from a very early age (Brett Ingram, Department of Primary Industries Victoria, pers. comm.). Therefore, to improve re-establishment of populations of *M. macquariensis* it is important to determine the movement patterns of sub-adults following stocking in rivers. Hatchery-reared *M. macquariensis* released into upland streams (Ebner *et al.* 2005) and hatchery-reared salmonids released in the Northern Hemisphere have been reported to disperse predominantly in a downstream direction (e.g. Bettinger and Bettoli 2002). In the context of *M. macquariensis* limited dispersal has been observed previously in a lowland river (Koehn *et al.* 2008) and percichthyids are known to home following release (Crook 2004a, b). We tested the hypothesis that hatchery and wild-reared *M. macquariensis* would disperse differently. Additionally, stocking of on-grown fish is also examined as an alternative reintroduction strategy to releasing fingerlings. We also aimed to determine the habitat use of this species to inform river restoration programs in the Murrumbidgee River.

Methods

Site description

The study was done in a lowland reach of the Murrumbidgee River near Narrandera (146°36'E; 34°46'S), between Berembed and Gogeldrie weirs (105 river km) in southern NSW, Australia (Fig. 1). This is one of twelve *M. macquariensis* stocking sites in the Murrumbidgee catchment, with 85 000 fingerlings (c. 30–50 mm TL) stocked at this location (Narrandera) between 1996 and 2000 (Gilligan 2005). Good numbers of wild fish captured by subsequent surveys indicate the reach to be the most successful stocking site for the species in NSW (Gowns *et al.* 2004; Gilligan 2005). In this reach, water is diverted at Berembed, Yanco and Gogeldrie weirs to meet irrigation requirements between spring and autumn. In-stream flow remains a function of rainfall during winter (Ebsary 1992). The geomorphology includes a transition from lower confined floodplains to open floodplains (Young *et al.* 2001). River red-gum *Eucalyptus camaldulensis* is common along much of the riparian edge of the river and channel widths are in the order of 70 m (Gowns *et al.* 2004). River depths of 3 to 5 m are commonly encountered on outside bends. The dominant in-stream habitat comprises structural woody habitat (SWH) consisting of fallen trees or branches, particularly of river red-gum.

Source of fish

Two experimental release groups were used for this study. The first (hatchery) group, comprised 29 *M. macquariensis* [two years of age, mean $\pm$ SE total length (TL) 394.00 $\pm$ 6.71 mm, range 310–429 mm; weight 1159.97 $\pm$ 53.16 g, range 513–1567 g] sourced from Snobs Creek Research Station (Department of Primary Industries) in Victoria and transported to the Narrandera Fisheries Centre in NSW. These individuals were bred

from wild broodstock (Murray River natural population). Initially post-larvae were reared in a fertilised earthen pond under semi-natural conditions. After harvesting the pond, fingerlings were transferred to hatchery facilities where they were weaned onto an artificial extruded diet then on-grown in 500-L circular fibreglass tanks that were part of an intensive recirculating aquaculture system. Methods used for fry rearing, weaning and on-growing were similar to those employed for Murray cod [*M. peelii peelii* (Mitchell)], which are described in more detail by Ingram (2004). The second (wild) group, comprised 32 *M. macquariensis* (age unknown, mean $\pm$ SE TL 451.90 ± 10.58 mm, range 370–635 mm; weight 1258.19 ± 116.50 g, range 599–3704 g) originally stocked as fingerlings and subsequently collected by boat electro-fishing from the Murrumbidgee River at Narrandera prior to surgery (Ebner *et al.* 2006). These individuals were collected between 10.3 km downstream and 4.7 km upstream of their later release site, and a GPS record was obtained at each capture location (Ebner *et al.* 2006). Fulton's condition index (Ricker 1975) was applied to both groups. A two-sample t-test identified that hatchery *M. macquariensis* were characterised by a higher condition index than the wild group ($T = 18.8$; d.f. = 48.6; $p < 0.05$) due to large fat deposits in the peritoneal cavity.

Surgery and release

Radio-tags [F1830, 35, 40 and 50, 11–25 g, Advanced Telemetry Systems (ATS), Isanti, USA, two-stage radio-transmitters, 150–152 MHz, pulse coded, duty cycle of 5 s on and 7 s off] were surgically implanted into hatchery and wild fish from 12–16 September and 16–24 September 2003, respectively. Radio-tag to body weight ratios ranged from 1.1–2.1% and 0.6–2.1% for hatchery and wild fish, respectively. The

surgery method was similar to that described by Ebner *et al.* (2007), with the exception of incisions being 2–3 cm in the current study. For external identification individuals were also tagged with a dart tag in the dorsal musculature between the second and third dorsal spines. Individuals initially recovered in a darkened enclosure holding 200-L of aerated water at Practical Salinity of 5 (based on the Practical Salinity Scale of 1978). Upon regaining swimming ability individuals were transferred to large circular concrete enclosures that held between 500 and 1000-L at Practical Salinity of 5. Hatchery and wild individuals were held in separate enclosures to prevent aggressive interactions. One wild individual died immediately post-surgery and autopsy revealed internal bleeding from a severed artery within the peritoneal cavity.

On 25 September 2003, 21 hatchery individuals implanted with radio-tags were released into the Murrumbidgee River, five river kilometres upstream of Narrandera (Fig. 1). On 26 September, 29 wild individuals implanted with radio-tags were released at the same location. Surgery was repeated on a further 10 individuals (two wild and eight hatchery individuals) that showed signs of loose sutures and/or open incisions, with three hatchery individuals rejecting their first radio-tags. On 1 October, two hatchery individuals were euthanased since they had developed severe infection around the incision and showed no sign of healing externally (despite swimming and behaving similarly to other healthy individuals). The remaining six hatchery and two wild individuals had healed incisions and were released on 1 October.

Radio-telemetry

A two-person crew, aboard a 3-m aluminium punt powered by an 8-Hp, two-stroke outboard motor, undertook manual radio-tracking approximately fortnightly for the first 3 months and then monthly up until 13 months post-release. Individuals were located during daylight hours using a scanning receiver (Australis 26k, Titley Electronics, Ballina, Australia or R4100, ATS) and a three-element Yagi antenna (Titley Electronics or ATS). Locations of individuals were recorded by taking three waypoints with a hand-held GPS [Garmin GPSII Plus or Garmin GPS 76 Marine Navigator; Figure of merit (F.O.M.) ≤ 5.0 at about 95% of locations, F.O.M. ≤ 7.0 at 100% of locations].

Record was made of the habitat occupied by each individual, including positioning within either a straight, the inside bend or the outside bend of a river channel. Water depth (± 0.1 m) was recorded from a depth sounder (Eagle™ Strata 128, Catoosa, USA) mounted to the stern of the boat. Distance from the bank was estimated by both crew members and averaged. Structural habitat that each individual was contacting was characterised as structural woody habitat, clay bank, macrophytes, or an absence of structure referred to as open-water. Habitat was determined visually and from depth sounder images obtained on repeated passes. Water temperature was collected at half-hourly intervals five river kilometres downstream from the tagged fish release site using a Hydrolab MS5 set at 0.5 m below the water surface (Sue Vink, CSIRO Land and Water, unpublished data) and averaged to provide a daily record. Temperature stratification was unlikely to occur at this site due to the predominance of summer irrigation flows. River discharge was recorded at daily intervals (NSW DNR 2004).

From 13 November 2003 to 14 November 2004, remote radio-telemetry data loggers (DCCII Model D5041, ATS connected to an ATS R4100 receiver, powered by 12 V power supply) were deployed at Berembed and Yanco weirs, 75 km apart (see Fig. 1). These loggers scanned 60 frequencies sequentially (58 implanted radio-transmitters, and two reference transmitters that were retained in the vehicle used to conduct field exercises) every 30 min. The approximate detection range of these loggers was 1 km.

Data analysis

The average of three replicate spatial points recorded at the location of an individual was used to minimise GPS error (except where one of these replicates was clearly an outlier i.e. >10 m from either of the other replicate points in which case the two similar replicates were averaged). Large-scale movement of individuals was quantified from records of the remote telemetry logger stations combined with those arising from manual radio-tracking. Movement distances were calculated using ArcView 3.2™ based on maps digitised at the 1:25 000 scale. For large-scale movement and dispersal, each spatial location was shifted onto the river mid-line. A polyline was generated based on the sequential locations of each individual using the Animal Movement Extension in ArcView (Hooge and Eichenlaub 1997). This was used to construct a time series of a) the proximity of individuals to the release point, and b) distance moved between consecutive radio-tracking events. A post-hoc power analysis (performed in Statistica version 7) revealed that sample sizes were adequate for detecting differences in dispersal of the wild and hatchery groups.

Dispersal between the two release-groups was compared using two-sample t-tests of log-transformed data at one and three months post-release. Homing was categorised as return to a location within one home-range, in either an upstream or downstream direction, from where an individual was originally captured (relevant to wild fish only), based on a maximum home-range size of 272 m for a subset of this group (Thiem *et al.* 2008). To determine if homing success was a function of translocation distance from capture sites a two-sample t-test was applied to square-root-transformed data. To determine if translocation direction (upstream or downstream from the release site) affected homing success a chi-square analysis was used. Total range was calculated as the distance between the furthest upstream and downstream location attained by each individual in the study. This did not include the capture locations of wild individuals but did incorporate records from the remote telemetry loggers. Home-range was estimated from monthly locations of individuals beginning 8 weeks after release, as at this stage >90% of wild individuals had established fidelity to a site, and so any potential biases from the use of a single release site are removed. We tested the hypothesis that following initial dispersal and/or homing *M. macquariensis* undertook comparatively limited movement, based on a two-sample t-test to compare total range with home-range of wild individuals. Site fidelity was classified as repeated monthly locations (i.e. a minimum of two locations) <272 m upstream or downstream from a previous location. In general, movements were either contained within this length of river or vastly exceeded this distance. Linear home-range estimates were taken as the minimum direct distance between upstream and downstream extremities, where site fidelity was clearly established (inaccuracy of the base-map prevented calculation of linear home-range along the river channel midline). Minimum convex polygons (MCP) were used to

estimate home-range area (Hooge and Eichenlaub 1997), where a minimum of 10 monthly fixes was obtained unless otherwise stated.

All statistical analyses were conducted in Statistix for Windows (version 8.1). T-tests were conducted following F-tests for homogeneity of variance. Degrees of freedom in t-tests are calculated using Satterwaite's approximation (Snedecor and Cochran 1980) and hence are expressed as a decimal number. Wilk-Shapiro normality tests were conducted, with data transformed where necessary to achieve approximate normal distribution (Tabachnick and Fidell 1989). The significance level for hypothesis tests was $p=0.05$.

Results

Dispersal

Dispersal from the release site was characterised by a different scale of movement between the two *M. macquariensis* groups (Fig. 2) and was significantly greater for the hatchery than the wild group at both one ($T= 2.45$; d.f.= 40.8; $p<0.05$) and three months ($T= 2.14$; d.f.= 41.8; $p<0.05$) post-release. Individuals from the wild group remained within 5 km of the release site (upstream or downstream) for the majority of the study with the exception of a maximum of four individuals on any fortnightly or monthly radio-tracking exercise (Fig. 2). Typically these exceptions involved movements up to 15 km upstream or downstream from the release site, although one individual was recorded 40 km upstream of the release site over a four and a half month period (Fig. 2). Conversely, the hatchery group exhibited pronounced modal classes 0–5 km and 5–10

km downstream of the release site, plus large-scale downstream movement of individuals within one month of being released (Fig. 2). Two individuals that had passed through Yanco Weir and were 40–50 km downstream of the release site represented the largest of these downstream movements. At 7–8 weeks post-release the location of individuals from the hatchery group was similar to the previous fortnightly fix and generally remained stable until the completion of the study (Fig. 2).

Additionally, a hatchery individual that was undetected at 4 weeks post-release was located 10–15 km upstream of the release site 7 weeks post-release where it remained for the entire study (Fig. 2).

Up until 5–6 weeks after release both the hatchery and wild groups remained close to the release site, with the hatchery group also dispersing downstream whilst the wild group moved in either direction (Fig. 3a and b). After this time, there was minimal movement by hatchery individuals (Fig. 3b). Similarly, there was minimal movement by the majority of wild individuals, however, a subset of individuals displayed pronounced upstream and/or downstream movement (Fig. 3a). These movements were not synchronised, occurring in November and December 2003, and April, May and September 2004 based on monthly radio-tracking exercises (Fig. 3a). These movements showed no discernable relationship with changes in discharge, temperature (Fig. 3c) or expected spawning period (Koehn and O'Connor 1990; Koehn and Harrington 2006).

In addition to movements detected by monthly and fortnightly radio-tracking, the downstream (Yanco Weir) telemetry logger (set-up after two hatchery individuals had passed through the weir (Fig. 2)) detected the presence of five individuals (one hatchery

and four wild *M. macquariensis*) encountering Yanco Weir. Interestingly all five individuals did not stay within detection range of the logger for more than a single logger cycle (approximately 30 mins). The hatchery individual was the first record on the logger (November 2003), travelling approximately 11 km downstream from its normal home-range and then returning to its previous location. This individual displayed the fastest recorded swimming speed, covering 10.9 km in a maximum time of four hours. In all four instances of wild individuals encountering Yanco Weir, individuals vacated an established home-range to undertake downstream movements of between 27 and 33 km, followed by return journeys to the vacated home-range. One of these movements occurred in December 2003, two in February 2004 and another in April 2004. All journeys were circular, with fish subsequently located at the previously occupied SWH. Minimum swimming speeds for these individuals varied from 0.04 to 0.32 km h⁻¹.

Confirmed survival of individuals from the two groups, wild and hatchery, differed greatly throughout the course of this study (Fig. 4). In the case of the wild group, 29 individuals were alive six months after release (Fig. 4a). One individual had a failed radio-transmitter (heard failing during radio-tracking) and one individual was suspected (and later confirmed) as a mortality or radio-tag rejection. Twelve months after release 19 individuals were alive and being radio-tracked, with 11 radio-transmitter failures and no further radio-tag rejections or mortality (Fig. 4a). A light aircraft was used to search for a number of these failed transmitters and verified they were not operational in the study area or surrounding areas. One individual was reported as an angler capture and

subsequent release. This fish was radio-tracked following capture for three months until its radio-transmitter failed.

For the hatchery group, survival of individuals declined rapidly, with only 14 individuals alive after one-month (Fig. 4b). This number decreased to six individuals two months after release, with three individuals alive after 12 months. Three radio-tags were recovered from shallow water during low-flow events in the river at three, six and nine months after release. Further, a large number of radio-fixes for hatchery individuals were taken in repeated locations for the duration of the study from ‘suspect’ locations in either relatively shallow water, or open water devoid of suitable structure, or associated with access points along the river. Three of these radio-tags were verified as radio-tag rejections or mortality through 24-h radio-tracking (i.e. no movement detected), as was one radio-tag from the wild group. In these cases it could not be determined if rejection of radio-tags or mortality had occurred. In contrast, the majority of wild individuals regularly moved short but detectable distances. As a consequence, further analyses were solely conducted on individuals from the wild group.

Homing in wild fish

Sixteen of 31 wild individuals homed to their original capture location in this study. The majority of individuals originally captured within 3 km either side of the release site (91%) homed to their capture location at some stage during the study. In contrast individuals captured between 3 and 5 km either side of the release site only homed on 40% of occasions. At capture distances greater than 5 km from the release site fish did not home back to the original capture site. Homing success was a function of an

individual having been captured in close proximity to the release site ($T = -3.66$; d.f. = 29, $p < 0.005$). The direction (upstream or downstream of release site) did not effect homing ability ($\chi^2 = 2.00$; $p = 0.1575$).

Typically, homing to a previous capture location was rapid. Ten of sixteen individuals had homed by the second radio-track at 12 days post-release. Five of these individuals homed in less than seven days to locations within 2 km of the upstream side of the release site. The time required to home was greater for the remaining six individuals, taking between 21 and 67 days.

Individuals exhibiting homing behaviour typically remained at their home-site once they had returned, with the proportion of subsequent radio-locations at the home-site ranging between 31 and 100%. However, on one occasion an individual took 12 days to home, stayed in this location for approximately 8 weeks and then relocated to a different home-site.

Total range and home-range of individuals

The total range [mean $\pm$ standard error (SE)] of wild *M. macquariensis* was 8840 ± 2151 m for the entire study. This was significantly different ($T = 4.16$; d.f. = 43.8; $p < 0.005$) when compared to total range following establishment of home-ranges for 90% of the group (6660 ± 2202 m). This finding is consistent with localised home-range movements following initial dispersal and/or homing.

Twenty-nine of the 31 wild *M. macquariensis* either returned to their pre-capture home-range or established new home-ranges after initial dispersal movements. One individual was not recorded establishing a home-range as its radio-transmitter failed early in the project. A second individual was excluded from home-range analysis as it was verified as a radio-tag rejection or mortality following 24 hr radio-tracking. Based on the remaining 29 individuals, the mean linear home-range was 78 ± 13 m and ranged between 8 and 270 m. In the case of three individuals that had undertaken home-range shifts, more than one linear home-range was estimated where at least two radio-tracking fixes were obtained. Each of these estimates was of comparable size to individuals that did not undertake home-range shifts. The home-range area of 19 individuals (with a minimum of 10, monthly radio-tracks) based on minimum convex polygons was between 53 and 4073 m², with a mean of 1070 ± 302 m². Home-range overlap occurred for a small number of individuals immediately upstream of the release site including two individuals which inhabited the same log upon establishment of home-ranges, for the duration of the study.

A number of different types of movement over the 13-month study period were observed following the establishment of home-ranges for 29 wild *M. macquariensis*. These were grouped into three categories of movement that are discussed in turn. Examples of each are provided in Fig. 5. Hatchery fish are not incorporated into this discussion or the subsequent section on habitat use due to a lack of spatial data resulting from poor survivorship.

(1) *Sedentary (occupation of a home-range)*

Following home-range establishment, through either homing or dispersal, 18 of 29 *M. macquariensis* exhibited restricted movements for the entire study (Fig. 5a). For example, nine of these individuals exhibited homing behaviour to their capture location post-dispersal, established tight home-ranges that were seemingly not vacated for the remainder of the study (or until transmitters failed in some cases). This occurred for homing in upstream or downstream directions. The remaining nine individuals also exhibited strong site fidelity and only localised movements were observed following dispersal, however, these individuals set up homes in different locations to their capture location, in both upstream and downstream directions from the release site. It should be noted that one of these individuals either took much longer to establish a clear home-range or had a larger home-range than was calculated as it did not exhibit fidelity to a restricted area for consecutive periods of time until six months after release. Collectively this group of 18 individuals comprised 10 females, two males and six individuals of unknown sex (Table 1). Length and weight (mean $\pm$ SE) of individuals was 443 ± 10 mm and 1147 ± 95 g, respectively (Table 1).

(2) Home-range shifts

Five individuals exhibited home-range shift in this study (Fig. 5b). This occurred when an established home-range (based on consecutive fixes at repeated locations) was vacated and a new home-range was established. Two female and three unknown sex *M. macquariensis* exhibited home-range shift behaviour and mean $\pm$ SE length and weight for this group was 407 ± 11 mm and 838 ± 71 g, respectively (Table 1). In the case of four of these individuals, single home-range shifts were recorded, varying in distance from 863 m to 2.4 km (see example W12, Fig. 5b). The fifth individual in this group

(W6) exhibited four home-range shifts, at least one of which was to a previous home-range (Fig. 5b). These shifts ranged in distance from 1 to 36 km, with the last recorded movement of 36 km to within five metres of a previous location five months earlier (W6, Fig. 5b).

(3) *Circular journeys*

Six individuals exhibited movements not associated with home-range movements or shifts (Fig. 5c). Four of these individuals undertook single large-scale return movements and were recorded on the Yanco Weir logger. These movements were characterised by an initial downstream movement of 27 to 33 km, a single record on the downstream logger and a subsequent return of the same distance to a home-range (see W26, Fig. 5c). Another individual (W31, Fig. 5c) undertook four large-scale movements following the establishment of a home-range. These movements ranged between 2.3 and 20 km and always culminated in a return to a discernible home-range. It should be noted that this individual was detected for 9.5 of the 13 months in a distinct home-range and exhibited fidelity to this location. The remaining individual in this group was located within its home-range on all except for two consecutive radio-tracking occasions. The scale of movement undertaken during these two months is unknown. The six *M. macquariensis* in this group were one female, two males and three individuals of unknown sex (Table 1). Length and weight (mean $\pm$ SE) for this group was 494 ± 20 mm and 1633 ± 223 g, respectively (Table 1).

Habitat-use

Wild *M. macquariensis* displayed a preference for outside bend habitats (53.3% of locations) within their home-range. The locations of home-range movements were less frequently encountered on straight sections of river channel (23.7%), inside bends (17.3%) and mid-channel (5.7%). *M. macquariensis* also displayed distinct patterns of habitat use within home-ranges. Structural woody habitat was the predominant habitat type used comprising 83% of fixes. Open water (13%), rock (2%), indeterminate habitat (2%) and undercut banks (<1%) were infrequently used and *M. macquariensis* were not found to use emergent macrophytes.

Discussion

Direction and magnitude of dispersal

The current study represents the first attempt to simultaneously investigate movement of a hatchery and wild *Maccullochella* species. Dispersal patterns of the hatchery and the wild group differed. Specifically the wild group exhibited limited dispersal, returning to the reach, 13 km in length, from which they had originally been collected. In comparison, the hatchery group remained near the release site or dispersed in a downstream direction. Similar findings arose from a study of rainbow trout, *Oncorhynchus mykiss* (Walbaum) in the Clinch River, USA (Bettinger and Bettoli 2002). Fish were released immediately below a dam in that study, whereas, there was opportunity for upstream dispersal in the current study.

The current study demonstrates that *M. macquariensis* is capable of large-scale dispersal. Previous reports identified that this species operates at smaller scales of less than 10 km (Koehn 1997; Ebner *et al.* 2005; Ebner *et al.* 2007; Nicol *et al.* 2007; Koehn *et al.* 2008; Thiem *et al.* 2008). Our finding may well be a result of the relatively large sample sizes that we used in addition to the fact that a number of previous studies have concentrated on aspects of home-range behaviour and habitat use at small spatial or temporal scales (e.g. Nicol *et al.* 2007, Thiem *et al.* 2008). Collectively, these radio-tracking studies of *M. macquariensis* indicate that hatchery-reared individuals are likely to disperse downstream and that individuals stocked as fingerlings or that naturally occur are capable of fidelity to a river reach. The current study indicates that habitat restoration (e.g. Nicol *et al.* 2004) may be maximised by targeting reaches downstream of existing populations or stocking sites for *M. macquariensis*. However, the downstream movement and dispersal observed in the current study poses a paradox for the management of this endangered species. It is widely accepted by river and fisheries managers that weirs pose a major ecological threat to riverine ecosystems and specifically migratory fishes within the Murray-Darling Basin (Harris and Mallen-Cooper 1994; Lintermans and Phillips 2004). Conversely, weirs may prove useful initially for concentrating *M. macquariensis* within small reaches in order to establish minimum viable populations (Gilpin and Soulé 1986; Caughley 1994; Ebner *et al.* 2005; Armstrong and Seddon 2008). This approach to stocking would facilitate concentrated research, monitoring and compliance efforts (Ebner *et al.* 2005) and is especially pertinent in view of the low annual hatchery production of this species (Gilligan 2005) relative to that predicted as necessary for establishing wild populations (Todd *et al.* 2004).

Homing

Half of the wild group returned to their original capture location. It is considered that this represents an ability to home. Further, the rapid homing of a substantial number of these individuals, including 30% of the wild group within a fortnight indicates effective searching and/or navigational capability of this species. It also represents the first evidence of experimental displacement and subsequent homing in *M. macquariensis* or any of the *Maccullochella*. Large-scale (10–100s km) return migrations have also been recorded by radio-tracking *M. peelii peelii* (Koehn 1997), *M. peelii mariensis* (Rowland) (Simpson and Mapleston 2002) and *M. ikei* (Rowland) (Gavin Butler, Southern Cross University, unpublished data), following release at capture locations. Collectively these findings indicate that all extant species of *Maccullochella* are capable of homing.

Homing success was a function of an individual having been captured in close proximity to the release site in this study. Similarly, more than half of a sample of *Macquaria ambigua* (Richardson) were found to home following displacements of about 2 km (Crook 2004a; n=10) though not following displacement of ~25 km (Crook 2004b; n=15). Crook (2004a) suggested that a number of non-homing individuals encountered high quality habitat not long after release and probably elected not to home. If in the future threatened species of *Maccullochella* were to be translocated, it would be informative to have some understanding of the maximum scale at which they home and whether or not unsuccessful homing behaviour will affect the success of translocation.

Individuals were found to have overlapping home-ranges (also see Thiem *et al.* 2008) and in some cases individuals exhibited long-term (1 year) co-occupation of SWH in the current study. There is little comparable information of intraspecific interaction in percichthyids. Butler (2001) reported that *M. ikei* individuals foraged in close proximity to one another at the head of a pool in the Nymboida River. Collectively, these observations challenge the idea that Australian freshwater cod are wholly territorial and solitary. Understanding intraspecific interactions that occur in the home-range occupation phase is a necessary step toward re-establishing self-sustaining populations of *Maccullochella*.

Home-range establishment

Much of the movement recorded in this study is comparable with that of Crook (2004a, 2004b) and is largely in agreement with the revised home-range shift model proposed in the second of these publications. Specifically, three main behaviours, home-range occupation, return movements and home-range shifts, were observed.

However, we also observed behaviour that indicating fish have a detailed spatial knowledge of the riverscape. We are referring to an individual that undertook a return home-range shift in the current study (Fig. 6; also see W6 in Fig. 5). Specifically, this is an individual returning to occupy a past home-range, though not its most recent home-range. The behaviour typically occurs in a subset of home-range shift cases (Fig. 6). Return home-range shifts have been recorded in studies of other percichthyids including *M. peelii peelii* (Koehn 1997), *M. p. mariensis* (Simpson and Mapleston 2002) and *M. ambigua* (Crook 2004b).

569

570 The cause of the return movements on a scale of tens of kilometres in the current study
571 is unknown. Crook (2004b) considered explorations to be searches for more profitable
572 habitat. Large-scale adult movement has often been ascribed to reproductive activity in
573 percichthyids (Koehn 1997; O'Connor *et al.* 2005). There is also an indication that high
574 flow may be an important factor or a contributing stimulus for percichthyid movement
575 (Koehn 1997; O'Connor *et al.* 2005). The return movements observed in the current
576 study were asynchronous among individuals and occurred outside the known spawning
577 period of the species. In this regard our findings are similar to that for *M. p. mariensis*
578 (Simpson and Mapleston 2002).

579

580 Only large individuals performed return movements in this study. Focussing on a range
581 of size classes and in particular large individuals (e.g. >2 kg) may prove useful in future
582 investigations of the movement of this species. Clearly, there is a need for better
583 understanding of the ecological mechanisms underpinning these return movements
584 (Ebner *et al.* 2006). Presumably this will be achieved by complementing the radio-
585 tracking approach with other techniques including confirmation of reproductive status,
586 manipulation of the dominance hierarchy (if indeed a hierarchy exists) or investigating
587 resource availability.

588

589 Within their home-range wild fish displayed a preference for SWH and outside bends
590 based on 13 months of monthly radio-tracking 29 individuals in a large river reach
591 (>10 km). The association with SWH is comparable with findings from electro-fishing
592 1 km of the Murrumbidgee River (Gowns *et al.* 2004) and radio-tracking studies based

on small samples sizes (Koehn 1997, n=4; Nicol *et al.* 2007, n=15; Thiem *et al.* 2008, n=10). Nicol *et al.* (2007) demonstrated that the location of SWH within the channel is also of importance to this species. Specifically, Nicol *et al.* (2007) found that association with SWH was greatest in narrow sections and more relaxed in wider sections of the Murray River, and predicted that these results would apply to this species in other river systems. The current study provides a level of validation for this prediction in that *M. macquariensis* were frequently found associated with woody debris in the main channel on outside banks of the Murrumbidgee River.

Possible rejection of radio-tags

It is possible that hatchery fish rejected radio-tags following release into the Murrumbidgee River in the current study. This is indicated by a small number of cases of rejection or signs of unsatisfactory recovery from surgery (e.g. incomplete closing of the incision, tearing of tissue at suture entry points) that occurred within days of implantation prior to release and contrasts with rapid healing of the incision and complete retention of radio-tags by wild fish (although two individual wild fish showed minor signs of reduced incision closure). The surgical technique was initially validated on hatchery *M. macquariensis* (n=9) that showed complete recovery in an aquaria trial (Ebner *et al.* 2005). The unsatisfactory post-surgery recovery of about one third of hatchery individuals initially in the current study was probably a function of their notably rotund abdomens relative to the wild fish. However it is unlikely that radio-tag rejection was widespread following release of the hatchery group since two-thirds of the sample were healing well following initial surgery and the remainder showed signs of healing following re-surgery (with the exception of the two individuals that were

euthanased). Radio-tagging of excessively fat hatchery fish should be avoided in future studies. It would also be useful to minimize incision length and radio-tag size, and provide an increased holding period for observation post-surgery.

Survivorship

Survivorship of wild and hatchery fish at 13 months post-release was 95% and 9%, respectively. Poor survivorship of hatchery fish has also been recorded for *M. macquariensis* in releases in the upper Murrumbidgee River catchment (Ebner *et al.* 2005; Ebner *et al.* 2007). Ebner *et al.* (2007) attributed some of the observed mortality to predation by cormorants and the common water rat. While collectively, studies of releases of on-grown *M. macquariensis* have revealed poor survivorship (Ebner *et al.* 2005; Ebner *et al.* 2007, this study) the mechanisms for mortality remain largely unexplained. Comparable studies have centred on salmonids (Thorstad *et al.*, 1998; Dieperink *et al.*, 2001; Bettinger and Bettoli 2002). Of these, higher survival of wild than hatchery fish is reported by both Dieperink *et al.* (2001) and Bettinger and Bettoli (2002). Predation mediated by morphological and behavioural differences was suggested as the explanation for the outcome in those two studies. Similarly, differences in the body shape and dispersal of hatchery and wild groups (Fig. 2) represent possible explanations for higher survival of wild than hatchery groups in the current study. In contrast, Thorstad *et al.* (1998) recorded 77% and 9% survival of hatchery and wild salmon, respectively. It was argued that greater energy stores of hatchery fish resulted in their higher survival over winter (Thorstad *et al.* 1998). Better condition of hatchery fish relative to wild fish did not obviously confer a benefit to the former group in the current

study. More likely, the familiarity of wild fish with the river environment was critical to their success (Brown and Day 2002).

The causes of mortality were unclear in the current study. A number of factors including low visibility, SWH and moderately deep water prevented retrieval of radio-tags or location of fish remains in most instances. In releases of hatchery *M. macquariensis* in upland rivers, the location of radio-tags provided a useful indicator of avian and mammal predation or scavenging (Ebner *et al.* 2007). For instance, radio-tags were found under the roosts of cormorant flocks. Similar reports relate to salmonids overseas (e.g. Dieperink *et al.* 2001). Based on the position of radio-tags in the current study, (primarily associated with SWH), a distinction could not be made between radio-tag rejection and predator effects, let alone among broad predator groups (i.e. avian, riparian mammal, fish, anglers) that might have been involved. This highlights a need for improved methods for detecting the timing and cause of mortality of *M. macquariensis* (e.g. Dieperink *et al.* 2001) especially following releases in turbid lowland rivers.

The current study demonstrates the benefit of using radio-tracking to monitor the movement and survivorship of stocked threatened fish to determine if there is fidelity to the release site. Our study also highlights the need to consider the effects of hatchery-rearing, when conducting fish reintroductions. Similar views have been expressed following poor survivorship or dispersal of other hatchery-reared fishes (e.g. Dieperink *et al.* 2001; Bettinger and Bettoli 2002).

Post-juvenile dispersal

The current study has not clarified if post-juvenile *M. macquariensis* dispersal is the explanation for the apparent disappearance of *M. macquariensis* from fingerling-stocking sites. The complete mortality of the hatchery group partly compromised our study by reducing the total sample size and the number of radio-tagged sub-adults. However, radio-tracking of the wild group revealed that limited emigration occurred in 13 months in the Murrumbidgee River at Narrandera, despite adults exhibiting the capability to move on a scale of tens of kilometres. It may be possible that the wild group comprised individuals that were unlikely to disperse over large distances, since they were collected near their original stocking site (*cf.* the Restricted Movement Paradigm, Gowan *et al.* 1994).

Observation of five individuals encountering and not passing Yanco Weir in 13 months may also be important considering the possibility that this behaviour may be more common in the population when scaled to the life-span of *M. macquariensis*. It highlights a fish passage issue at that weir, and more generally highlights the potential for maintaining a threshold population size. Dispersal away from a release site can lead to insufficient adults to maintain a self-sustaining population, an issue expressed in the small population paradigm (Caughley 1994; Armstrong and Seddon 2008).

To test if post-juvenile *M. macquariensis* dispersal is the explanation for the apparent disappearance of *M. macquariensis* from fingerling-stocking sites, hatchery fish should not be used as a surrogate for examining the dispersal of wild fish in the future. However, the current study has shown the benefit of using radio-tracking to monitor

individual movement and estimate survivorship from a sample of stocked threatened fish. Therefore, future efforts to conserve *M. macquariensis* could use this approach to monitor the dispersal and survivorship of large juveniles and sub-adults at sites where reintroductions have been less effective than is the case in the Murrumbidgee River at Narrandera (*cf.* Gilligan 2005). This study demonstrates the likely benefit of using radio-tracking as the basis for monitoring the fate of widespread stockings of large bodied percichthyid species in Australia.

Acknowledgements

The idea for this comparison of wild and hatchery trout cod came from discussion with I. Wooden. We are also grateful to the project Steering Committee. V. Carracher, D. Gilligan, S. Godschalx, M. Jekabsons, L. Rava, M. Stimson, and I. Wooden assisted with wild fish collection. Staff at the Snobs Creek Research Station, including P. Boyd, B. Ingram and P. Ryder reared fish and provided support. S. Godschalx assisted in surgery. L. Baumgartner, B. Broadhurst, S. Godschalx, M. Jekabsons, L. Johnston, M. Lintermans, P. Maclean, M. McLellan, M. Stimson and S. Thurstan provided field assistance. S. Thurstan also provided advice that influenced the project. State Water staff including L. Beck, M. Deaton, M. McNamara and E. Taylor mounted the telemetry aerials at Yanco Weir. M. Dunford provided invaluable GIS support, M. Evans and E. Harrison provided statistical advice and S Vink provided water quality data. Input from D. Gilligan, K. Frawley, B. Ingram, M. Lintermans and K. Ryan improved versions of this manuscript. Two anonymous reviewers also improved this manuscript. This research was funded by the Fisheries Research and Development Corporation,

Environment ACT and NSW Department of Primary Industries with additional contributions from Victorian Department of Primary Industries, Cooperative Research Centre for Freshwater Ecology, Murray-Darling Basin Commission, the Arthur Rylah Institute for Environmental Research, RecFish Australia, NSW State Water and the Department of Infrastructure, Planning and Natural Resources. This study was performed under NSW Fisheries Animal Care and Ethics Committee authorisation 03/07 in Wiradjuri Country.

References

- ACT Government. (2007). Ribbons of life: ACT aquatic species and riparian zone conservation strategy. Environment and Recreation, Canberra.
- Armstrong, D. P., and Seddon, P. J. (2008). Directions in reintroduction biology. *Trends in Ecology & Evolution* **23**, 20–25. doi:10.1016/j.tree.2007.10.003
- Bettinger, J. M., and Bettoli, P. W. (2002). Fate, dispersal and persistence of recently stocked and resident Rainbow Trout in a Tennessee tailwater. *North American Journal of Fisheries Management* **22**, 425–432. doi:10.1577/1548-8675(2002)022
- Brown, P. (1998). Goulburn River trout cod stock assessment, November 1997: growth and mortality estimates. Marine and Freshwater Resources Institute, Victoria.
- Brown, A. M., Nicol, S. J., and Koehn, J. D. (1998). Recovery plan for the trout cod *Maccullochella macquariensis* 1998–2005. Aquatic Ecosse Pty Ltd and Department of

734 Natural Resources and Environment, Melbourne.

735 Brown, C., and Day, R. (2002). The future of stock enhancements: lessons for hatchery
 736 practice from conservation biology. *Fish and Fisheries* **3**, 79–94. doi: 10.1046/j.1467-
 737 2979.2002.00077.x

738 Butler, G. (2001). Age, growth and telemetric tracking of the eastern freshwater cod,
 739 *Maccullochella ikei* (Pisces: Percichthyidae) within the Mann–Nymboida River System,
 740 NSW. BSc (Hons) Thesis, Southern Cross University, Australia.

741 Caughley, G. (1994). Directions in conservation biology. *The Journal of Animal*
 742 *Ecology* **63**, 215–244.

743 Crook, D. A. (2004a). Is the home range concept compatible with the movements of two
 744 species of lowland river fish? *Journal of Animal Ecology* **73**, 353–366. doi:
 745 10.1111/j.0021-8790.2004.00802.x

746 Crook, D. A. (2004b). Movements associated with home-range establishment by two
 747 species of lowland river fish. *Canadian Journal of Fisheries and Aquatic Sciences* **61**,
 748 2183–2193. doi:10.1139/f04-151

749 Dieperink, C., Pedersen, S., and Pedersen, M. I. (2001). Estuarine predation on
 750 radiotagged wild and domesticated sea trout (*Salmo trutta* L.) smolts. *Ecology of*
 751 *Freshwater Fish* **10**, 177–183. doi: 10.1034/j.1600-0633.2001.100307.x

752 Douglas, J. W., and Brown, P. (2000). Notes on successful spawning and recruitment of
 753 a stocked population of the endangered Australian freshwater fish, trout cod,
 754 *Maccullochella macquariensis* (Cuvier) (Percichthyidae). *Proceedings of the Linnaean*

755 *Society of New South Wales* **122**, 143–147.

756 Douglas, J. W., Gooley, G. J., and Ingram, B. I. (1994). Trout Cod, *Maccullochella*
757 *macquariensis* (Cuvier) (Pisces: Percichthyidae) resource handbook and research and
758 recovery plan. Department of Conservation and Natural Resources, Melbourne.

759 DSE (Victorian Department of Sustainability and Environment). (2003). Advisory list
760 of threatened vertebrate fauna in Victoria. Victorian Department of Sustainability and
761 Environment, Melbourne.

762 Ebner, B., Johnston, L., and Lintermans, M. (2005). Re-introduction of trout cod into the
763 Cotter River catchment. Final Report to the Natural Heritage Trust, Environment ACT,
764 Canberra.

765 Ebner, B.C., Thiem, J.D., Gilligan, D.M., Lintermans, M., Wooden, I.J., and Linke, S.
766 (2008). Estimating species richness and catch per unit effort from boat electro-fishing in
767 a lowland river in temperate Australia . *Austral Ecology* In press.

768 Ebner, B., Thiem, J., Lintermans, M., and Gilligan, D. (2006). An ecological approach
769 to re-establishing Australian freshwater cod populations: an application to trout cod in
770 the Murrumbidgee catchment. Parks, Conservation and Lands, Canberra.

771 Ebner, B. C., Thiem, J. D., and Lintermans, M. (2007). Fate of 2 year-old, hatchery-
772 reared trout cod *Maccullochella macquariensis* (Percichthyidae) stocked into two
773 upland rivers. *Journal of Fish Biology* **71**, 182–199. doi: 10.1111/j.1095-
774 8649.2007.01481.x

775 Ebsary, R. (1992). Regulation of the Murrumbidgee River. In ‘The Murrumbidgee: Past

776 and Present'. (Eds J. Roberts and R. Oliver.) pp. 49–59. (CSIRO Division of Water
777 Resources: Griffith, Australia.)

778 Faragher, R. A., Brown, P., and Harris, J. H. (1993). Population surveys of the
779 endangered fish species trout cod (*Maccullochella macquariensis*) and eastern cod (*M.*
780 *ikei*). NSW Fisheries, Cronulla.

781 Fischer, J., and Lindenmayer, D. B. (2000). An assessment of the published results of
782 animal relocations. *Biological Conservation* **96**, 1–11. doi:10.1016/S0006-
783 3207(00)00048-3

784 Gillanders, B., Elsdon, T., and Munro, A. (2006). Impacts of native fish stocking on fish
785 within the Murray–Darling Basin. Murray–Darling Basin Commission, Canberra.

786 Gilligan, D. (2005). Fish communities of the Murrumbidgee catchment: status and
787 trends. NSW Department of Primary Industries, Cronulla.

788 Gilpin, M. E., and Soulé, M. E. (1986). Minimum viable populations: processes of
789 species extinction. In 'Conservation Biology: The Science of Scarcity and Diversity'.
790 (Ed M. E. Soulé) pp. 19–34. (Sinauer Associates: Massachusetts.)

791 Gowan, C., Young, M. K., Fausch, K. D., and Riley, S. C. (1994). Restricted movement
792 in resident stream salmonids: a paradigm lost? *Canadian Journal of Fisheries and*
793 *Aquatic Sciences* **51**, 2626–2637.

794 Gowns, I., Wooden, I., and Schiller, C. (2004). Use of instream wood habitat by Trout
795 Cod *Maccullochella macquariensis* (Cuvier) in the Murrumbidgee River. *Pacific*
796 *Conservation Biology* **10**, 261–265.

797 Harris, J. H., and Mallen-Cooper, M. (1994). Fish-passage development in the
 798 rehabilitation of fisheries in mainland south-eastern Australia. In ‘Rehabilitation of
 799 Freshwater Fisheries’. (Ed I. G. Cowx.) pp. 185–193. (Fishing News Books UK,
 800 University of Hull.)

801 Harris, J. H., and Rowland, S. J. (1996). Family Percichthyidae: Australian freshwater
 802 cods and basses. In ‘Freshwater Fishes of South-Eastern Australia’. 2nd edn. (Ed R. M.
 803 McDowall.) pp. 150–163. (Reed Books: Australia.)

804 Hooge, P. N., and Eichenlaub, B. (1997). ‘Animal Movement Extension to Arcview
 805 1.1’. (US Geological Survey, Anchorage.)

806 Ingram, B. A., and Douglas, J. W. (1995). Threatened fishes of the world:
 807 *Maccullochella macquariensis* (Cuvier, 1829) (Percichthyidae). *Environmental Biology*
 808 *of Fishes* **43**, 38. doi:10.1007/BF00001815

809 Ingram, B. A. (2004). Murray cod fingerling production and growout. In ‘Development
 810 of Intensive Commercial Aquaculture Production Technology for Murray Cod’. (Eds
 811 B.A. Ingram and S. S. DeSilva.) pp. 25–65. (Primary Industries Research: Victoria.)

812 Jones, G. P. (1991). Post recruitment processes in the ecology of coral reef fish
 813 populations: a multifactorial perspective. In ‘The Ecology of Fishes on Coral Reefs’.
 814 (Ed P. F. Sale.) pp. 294–398. (Academic Press: California.)

815 Koehn, J. D., and O’Connor, W. G. (1990). Biological information for management of
 816 native freshwater fish in Victoria. Department of Conservation and Environment,
 817 Victoria.

818 Koehn, J. (1997). Habitats and movements of freshwater fish in the Murray–Darling
819 Basin. In ‘Proceedings of the 1995 Riverine Environment Research Forum’. (Eds R. J.
820 Banens and R. Lehane.) pp. 27–32. (Murray–Darling Basin Commission: Canberra).

821 Koehn, J. D., and Harrington, D. J. (2006). Environmental conditions and timing for the
822 spawning of Murray cod (*Maccullochella peelii peelii*) and the endangered trout cod
823 (*M. macquariensis*) in southeastern Australian rivers. *River Research and Applications*
824 **22**, 327–342. doi: 10.1002/rra.897

825 Koehn, J. D., Nicol, S. J., McKenzie, J. A., Lieschke, J. A., Lyon, J. P., and Pomorin, K.
826 (2008). Spatial ecology of an endangered native Australian Percichthyid fish, the trout
827 cod *Maccullochella macquariensis*. *Endangered Species Research* **4**, 219–225.
828 doi:10.3354/esr00075

829 Lintermans, M. (1995). The status of trout cod *Maccullochella macquariensis* in
830 Bendora Reservoir, ACT: results of 1992/93 and 1995 monitoring. ACT Parks and
831 Conservation Service, Canberra.

832 Lintermans, M. (2006). A review of stock enhancement activities for threatened fish
833 species in Australia. In ‘Research, Development and Extension Priorities for Stock
834 Enhancement, Fish Stocking and Fish Recovery’. (Ed B. Sawnok.) pp. 30–43. (Recfish
835 Australia: Brisbane.)

836 Lintermans, M., and Phillips, B. (2004). Downstream movement of fish in the Murray–
837 Darling Basin: Statement, recommendations and supporting papers. Murray–Darling
838 Basin Commission, Canberra.

839 Lintermans, M., Rowland, S. J., Koehn, J., Butler, G., Simpson, R., and Wooden, I.

840 (2005). The status, threats and management of freshwater cod species in Australia. In
 841 'Management of Murray Cod in the Murray–Darling Basin: Statement,
 842 Recommendations and Supporting Papers'. (Eds M. Lintermans and B. Phillips.) pp.
 843 15–29. (Murray–Darling Basin Commission: Canberra.)

844 Metcalfe, N. B., Valdimarsson, S. K., and Morgan, I. J. (2003). The relative roles of
 845 domestication, rearing environment, prior residence and body size in deciding territorial
 846 contests between hatchery and wild juvenile salmon. *Journal of Applied Ecology* **40**,
 847 535–544. doi: 10.1046/j.1365-2664.2003.00815.x

848 Minckley, W. L. (1995). Translocation as a tool for conserving imperiled fishes:
 849 Experiences in western United States. *Biological Conservation* **72**, 297–309.
 850 doi:10.1016/0006-3207(94)00091-4

851 Molony, B. W., Lenanton, R., Jackson, G., and Norriss, J. (2003). Stock enhancement
 852 as a fisheries management tool. *Reviews in Fish Biology and Fisheries* **13**, 409–432.
 853 doi:10.1007/s11160-005-1886-7

854 Morris, S. A., Pollard, D. A., Gehrke, P. C., and Pogonoski, J. J. (2001). Threatened and
 855 potentially threatened freshwater fishes of coastal New South Wales and the Murray–
 856 Darling Basin. NSW Fisheries, Sydney.

857 Nicol, S. J., Lieschke, J., Lyon, J., and Koehn, J. D. (2004). Observations on the
 858 distribution and abundance of carp and native fish, and their responses to a habitat
 859 restoration trial in the Murray River, Australia. *New Zealand Journal of Marine and*
 860 *Freshwater Research* **38**, 541–551.

861 Nicol, S. J., Barker, R. J., Koehn, J. D., and Burgman, M. A. (2007). Structural habitat

862 selection by the critically endangered trout cod, *Maccullochella macquariensis*, Cuvier.
 863 *Biological Conservation* **138**, 30–37. doi:10.1016/j.biocon.2007.03.022
 864 NSW DNR. (2004). New South Wales Department of Natural Resources supported
 865 website <<http://waterinfo.dlwc.nsw.gov.au/>> [accessed 18 October 2005].
 866 NSW DPI (New South Wales Department of Primary Industries). (2005). Trout Cod
 867 (*Maccullochella macquariensis*) draft recovery plan. New South Wales Department of
 868 Primary Industries, Port Stephens, Australia.
 869 O'Connor, J. P., O'Mahony, D. J., and O'Mahony, J. M. (2005). Movements of
 870 *Macquaria ambigua*, in the Murray River, south-eastern Australia. *Journal of Fish*
 871 *Biology* **66**, 392–403. doi:10.1111/j.0022-1112.2005.00604.x
 872 Ricker, W. E. (1975). Computation and interpretation of biological statistics of fish
 873 populations. *Bulletin of the Fisheries Research Board of Canada* **191**, 1–382.
 874 Salvanes, A. G. V., and Braithwaite, V. A. (2005). Exposure to variable spatial
 875 information in the early rearing environment generates asymmetries in social
 876 interactions in cod (*Gadus morhua*). *Behavioral Ecology and Sociobiology* **59**, 250–
 877 257. doi:10.1007/s00265-005-0031-x
 878 Simpson, R. R., and Mapleston, A. J. (2002). Movements and habitat use by the
 879 endangered Australian freshwater Mary River cod, *Maccullochella peelii mariensis*.
 880 *Environmental Biology of Fishes* **65**, 401–410. doi:10.1023/A:1021129021451
 881 Snedecor, G.W., and Cochran, W. G. (1980). 'Statistical Methods'. (Iowa State
 882 University Press: Ames, Iowa.)

883 Tabachnick, B. G., and Fidell, L. S. (1989). ‘Using Multivariate Statistics’. (Harper and
884 Row Publishers: New York.)

885 Thiem, J. D., Ebner, B. C., and Broadhurst, B. T. (2008). Diel activity of the endangered
886 trout cod (*Maccullochella macquariensis*) in the Murrumbidgee River. *Proceedings of*
887 *the Linnean Society of New South Wales* **129**, 167–173.

888 Thorstad, E. B., Heggberget, T. G., and Økland, F. (1998). Migratory behaviour of adult
889 wild and escaped farmed Atlantic salmon, *Salmo salar* L., before, during and after
890 spawning in a Norwegian river. *Aquaculture Research* **29**, 419–428.
891 doi:10.1111/j.1365-2109.1998.tb01149.x

892 Todd, C. R., Nicol, S. J., and Koehn, J. D. (2004). Density-dependence uncertainty in
893 population models for the conservation management of trout cod, *Maccullochella*
894 *macquariensis*. *Ecological Modelling* **171**, 359–380.
895 doi:10.1016/j.ecolmodel.2003.06.002

896 Wallace, M. P. (2000). Retaining natural behaviour in captivity for re-introduction
897 programmes. In ‘Behavioural Conservation’. (Eds L. M. Gosling and W. J. Sutherland.)
898 pp. 300–314. (Cambridge University Press: Cambridge.)

899 Young, W. J., Roberts, J., and Hillman, T. J. (2001). The rivers of the Basin and how
900 they work. In ‘Rivers as Ecological Systems: The Murray–Darling Basin’. (Ed W. J.
901 Young.) pp. 3–43. (Murray Darling Basin Commission: Canberra.)

902

903

904

Table 1. Summary statistics for 29 wild *M. macquariensis* exhibiting different types of movements following the establishment of a home-range. TL and weight are expressed as mean $\pm$ SE.

	Sedentary	Shifts	Circular journeys
Number	18	5	6
Sex	10♀, 2♂, 6?	2♀, 3?	1♀, 2♂, 3?
TL (mm)	443 $\pm$ 10	407 $\pm$ 11	494 $\pm$ 20
Weight (g)	1147 $\pm$ 95	838 $\pm$ 71	1633 $\pm$ 223

Fig. 1. Location of the study reach along the Murrumbidgee River within New South Wales, Australia.

Fig. 2. Dispersal of hatchery (■) and wild (■) *M. macquariensis* groups in the Murrumbidgee River. Time after release: (a) 1 month, (b) 3 months, (c) 6 months, (d) 9 months, and (e) 12 months. Distance moved is expressed as upstream and downstream direction in 5 km categories.

Fig. 3. Minimum distances moved between radio-tracking intervals (as detected by manual radio-tracking and remote telemetry) by all (a) wild and (b) hatchery *M. macquariensis* and, (c) the water discharge (—) and temperature (--) in the Murrumbidgee River at Narrandera. Positive and negative values indicate a direction of movement upstream and downstream, respectively.

Fig. 4. Fate of radio-tagged (a) wild *M. macquariensis* and, (b) hatchery *M. macquariensis* released into the Murrumbidgee River near Narrandera.

Fig. 5. Typical examples of different types of *Maccullochella macquariensis* movements, representing: (a) sedentary home-range behaviour for the entire study, following dispersal, (b) home-range shifts that occurred once or multiple times, (c) large-scale movements outside of home-ranges that did not result in a home-range shift. The first point in each figure (zero) represents the release location (all fish were released at the same location).

Fig. 6. Categorisation of *M. macquariensis* (n=29) movement over the 13-month study.

The number of individuals exhibiting specific emigration home-range behaviour is shown. The occurrence of each behaviour is scored only once per individual