Metadata of the chapter that will be visualized online

Chapter Title	Introduction to Angiogenesis in Normal Physiology. Disease and Malignancy	
Copyright Year	2016	
Copyright Holder	Springer International Publishing Switzerland	
Corresponding Author	Family Name	Salajegheh
	Particle	
	Given Name	Ali
	Suffix	
	Division	School of Medicine, Menzies Health Institute Queensland
	Organization/University	Griffith University
	City	Gold Coast
	State	QLD
	Country	Australia
	Email	a.salajegheh@griffith.edu.au
Abstract	<p>Vasculogenesis is the de novo formation of blood vessels by mesodermal progenitors undergoing differentiation to endothelial cells. New vasculature formation from pre-existing vasculature occurs through the physiological remodelling process known as angiogenesis. Angiogenesis is implicated in the proliferation and growth of both physiologically normal and neoplastic tissues, through the establishment of vascular supply, essential for delivering growth requirements such as oxygen and nutrients. Many inhibitory and promoter genes regulate this process, however the role of specific genes in the shift from normal angiogenesis to tumour initiation is complex and thus poorly understood. This book investigates the process of angiogenesis in malignancies. The functional role of key regulatory factors will be examined in the context of normal healthy condition. Then the association of these factors to disease and malignancy, cancer proliferation and progression will be discussed. New insights into the role of angiogenesis and the therapeutic inhibition of its regulators will be investigated due to the great potential for exploitation in the development of a novel treatment for cancer.</p>	
Keywords (separated by “ - ”)	Angiogenesis - Normal physiology - Disease - Malignancy	

Chapter 11

Introduction to Angiogenesis in Normal2

Physiology. Disease and Malignancy3

Abstract Vasculogenesis is the de novo formation of blood vessels by mesodermal progenitors undergoing differentiation to endothelial cells. New vasculature formation from pre-existing vasculature occurs through the physiological remodelling process known as angiogenesis. Angiogenesis is implicated in the proliferation and growth of both physiologically normal and neoplastic tissues, through the establishment of vascular supply, essential for delivering growth requirements such as oxygen and nutrients. Many inhibitory and promoter genes regulate this process, however the role of specific genes in the shift from normal angiogenesis to tumour initiation is complex and thus poorly understood. This book investigates the process of angiogenesis in malignancies. The functional role of key regulatory factors will be examined in the context of normal healthy condition. Then the association of these factors to disease and malignancy, cancer proliferation and progression will be discussed. New insights into the role of angiogenesis and the therapeutic inhibition of its regulators will be investigated due to the great potential for exploitation in the development of a novel treatment for cancer.

Keywords Angiogenesis • Normal physiology • Disease • Malignancy19

1.1 Introduction to Angiogenesis20

There are two mechanisms for the formation of blood vessels in the body; angiogenesis and vasculogenesis. Both are essential in formation of the vascular network in our bodies.

Vasculogenesis is the formation of blood vessels from angioblasts. It is a dynamic process that involves cell–cell and cell–extracellular matrix interactions directed by growth factors and morphogens (Carmeliet 2005). This process includes differentiation of mesodermal stem cells into angioblasts, and then the differentiation of angioblasts into endothelial cells under the command of growth factors. It occurs inside the embryo and it initiates the first vascular system of our body.

Angiogenesis is the term used to describe the process of growth or formation of new blood vessels from pre-existing blood vessels (Folkman and Shing 1992). This process begins in utero and prolongs throughout life. Angiogenesis is an essential

process in normal physiology in the body for it has major roles in healing, reproduction, growth of tissues and also diseases (Felmeden et al. 2003). The role of angiogenesis and its contribution to disease has been a topic of discussion for the last few decades (Folkman 1996). Metabolically active tissues in the body are always in close proximity to bloody capillaries for essential processes such as exchange of nutrients and metabolites must occur (Adair and Montani 2010). It is responsible for both remodelling and expansion of the existing network formed by vasculogenesis. Angiogenesis exists as a very important process in the body due to the fact that all tissues require oxygen. Oxygen is delivered through capillaries which are formed through angiogenesis. The control of angiogenesis in the body is dependent on a specific tissue's metabolic needs. The reason angiogenesis has gained much publicity in the world of health is due to its influence on both many diseases and cancer. Malignant growth is dependent on the formation of these new blood vessels, that supply nutrition to the tumour site, and provide a way for certain cells and survival factors to enter and leave circulation (Folkman 2006; Seaman et al. 2007). Chemical factors such as angiogenic growth factors, angiogenesis inhibitors and other genetic factors can cause abnormalities in the process of angiogenesis (Seaman et al. 2007). This lack of control results in either too much or too less angiogenesis occurring (Albini et al. 2012). The stimulation of angiogenesis is now considered therapeutic approach in terms of treating peripheral arterial disease, ischemic heart disease, and in wound care, however, in cases of ophthalmic conditions, cancer, rheumatoid arthritis, and other diseases, limitation of angiogenesis seems a better option (Folkman and Shing 1992).

In general, angiogenesis occurs in a series of steps that are initiated by a certain cell not having enough oxygen. This cell will release proangiogenic growth factors which attract inflammatory and endothelial cells that then proliferate and grow. Endothelial cells in the target location then secrete metalloproteases which digest the blood vessel walls allowing these cells to go towards the angiogenic stimulus. The process is completed through anastomosis which joins the capillaries coming from the arterioles and venules, establishing proper blood flow in the body (Carmeliet 2005). For angiogenesis to occur in the body, its positive regulation through growth factors must overcome the factors that inhibit angiogenesis. Angiogenesis is a complex process composed of many steps and a combination of factors such as cells, soluble factors and extracellular matrix (ECM) components (Liekens et al. 2001). Embryonic mesenchymal cells differentiate into endothelial cells to develop into blood vessels (Felmeden et al. 2003). There are growth factors that have been described to be associated with the positive regulation of angiogenesis; these include vascular endothelial growth factor (VEGF), transforming growth factors (TGF-beta), fibroblast growth factors (FGF), epidermal growth factor (EGF). On the other hand, there are also growth factors that have been shown to have little effect on proliferation of endothelial cells in angiogenesis (Carmeliet 2005). There are two types of angiogenesis that occur in adults: these are sprouting angiogenesis and intussusceptive angiogenesis.

Sprouting angiogenesis essentially refers to branching of existing blood vessels via enzymatic degradation of capillary basement membrane, proliferation and migration of endothelial cells towards angiogenic stimulus through the extracellular

matrix, formation of a tube with lumen, fusion of vessels to form network and finally stabilization of vessels by recruitment of pericytes (Adair and Montani 2010). Angiogenic stimuli induce the endothelial cells of existing vessels to proliferate and increase in permeability. Next, the extracellular matrix of the proximal tissue gets degraded to give space for the new blood vessel. The endothelial cells continue sprouting and proliferating into the provisional extracellular space, and form the lumen of the developing capillary. In the final stages of angiogenesis stabilization occurs, where a mature vascular basement membrane covers and pericytes surround the new vessel to protect and support it (Hong et al. 2014; Senger and Davis 2011).

In terms of sprouting angiogenesis, the first growth factor was discovered in the 1980s which was fibroblast growth factor. FGF triggers all the main steps in the angiogenesis process and is produced by many cells, among which are macrophages and tumour cells. Despite the fact that FGF is not released on a regular basis, after its secretion into the extracellular matrix the angiogenesis cascade is initiated. Various research points to the fact that vascular endothelial growth factor (VEGF) is a very important part of the initiation of angiogenesis as well. Essentially, it activates the proliferation of endothelial cells and is also a powerful initiator of vasodilation in blood vessels. Furthermore, it is a stimulator of metalloproteinases and plasminogens (Hillen and Griffioen 2007).

To go into more depth on the process, we can use the role of VEGF as an example. VEGF provides an angiogenic stimulus that helps the endothelial tip cell sprout through the extra-cellular matrix thus forming the foundation for a new capillary to develop. The tip cells are aided by processes on them called filopodia which release proteolytic enzymes that establish a path through the ECM. The sprouting of the tip cells with their filopodia is also heavily dependent on the VEGF gradient as the filopodia are covered in many VEGF receptors. Following the tip cells, are endothelial stalk cells which cause the elongation of the sprout. These stalk cells are what forms the capillary lumen. The uniting of two tip cells at a centre of VEGF release forms one continuous lumen through which blood can perfuse. The VEGF stimulus dies down after the tissues in that area are sufficiently oxygenated (Carmeliet 2005).

Intussusceptive angiogenesis, in comparison, is a much newer discovery which was first observed in the lungs (Burri and Djonov 2002). It is a much faster and efficient form of angiogenesis as there is no endothelial cell proliferation and it is merely the splitting of a blood vessel into two by reorganization of the existing endothelial cells so that the cells extend into the lumen (Adair and Montani 2010; Carmeliet 2005). This ultimately causes an increase in the overall complexity of the network and occurs throughout life like sprouting angiogenesis.

1.2 Angiogenesis in Normal Physiology

During embryonic development the primary capillary plexus is developed by vasculogenesis. Vasculogenesis differs from angiogenesis in that the formation of new vasculature does not happen from pre-existing vessels but rather from endothelial

cell precursors, called angioblasts. The primary capillary plexus is then remodelled by the process of angiogenesis (Papetti and Herman 2002). The growth of newly developed cells is angiogenesis-dependent; all cells require nutrition, oxygen and removal of wastes to be able to function (Folkman 1996). The growth of the vascular network that is the cardiovascular system is one of the earliest events of angiogenesis occurring during embryonic development (DeWitt 2005). After birth the blood vessels remain quiescent however it may also arise during adulthood such as seen in reproduction. During the menstrual cycle the development of the endometrium occurs resulting in highly-developed vascularity. The endometrium eventually sheds causing breakages in the surrounding blood vessels, which then undergo angiogenesis in order to repair the endometrial tissue (Smith 2001; Folkman 2006). Another instance where angiogenesis may occur later in life, and is of quite significance is in the regeneration of damaged tissues during the wound repair and healing process. During the process of wound healing, sprouting angiogenesis takes place resulting in endothelial projections that enter the blood clot over the wound (Tonnesen et al. 2000). The organised vascularity assists in the deposition of a collagen matrix within the granulation tissue and as it matures into a scar, the dense network of blood vessels eventually decreases (Tonnesen et al. 2000). Angiogenesis should be also enhanced in numerous nonrelated diseases such as in ischemic tissues or cardiac failure where reperfusion is required to improve the disease conditions; such event is also referred to as compensatory angiogenesis (Carmeliet 2003).

It is evident from the aforementioned roles in normal physiology that angiogenesis requires regulation in order to switch the process on and off when required. Certain naturally occurring inhibitors of this process prevent the formation of neovasculature under pathologic conditions (Folkman and Shing 1992). It can therefore be concluded that a dynamic relationship must exist between growth factors and inhibitors, and if disrupted could lead to disease.

1.3 Angiogenesis in Disease

Several chronic inflammatory diseases are, in fact, causative agents of excessive angiogenesis as part of the pathology (Carmeliet 2003). In such conditions, the balance between stimulant and inhibitory angiogenic factors tilt towards the stimulant outcome leading to 'angiogenic switch,' resulting in profound angiogenesis (Carmeliet 2005). These diseases include atherosclerosis, rheumatoid arthritis, diabetes, psoriasis, endometriosis and cancer. To treat such disorders, therapeutic inhibition of angiogenesis can be an option of improvement (Carmeliet 2003).

Multiple studies revealed that the immune system is critically impacted by angiogenesis. Studies have shown that during angiogenesis exposure of endothelial cells to angiogenic factors such as FGF-2 and VEGF down-regulates adhesion molecule expression on the endothelial cells. In addition, endothelial adhesion molecules controlled by cytokines such as TNF is severely hampered by angiogenic stimulus which interferes with activation and adhesion of leukocytes to endothelial tissues at

the site of neovascularization (Carmeliet 2003). This was an important finding as it suggested that those chronic inflammatory disorders which cause angiogenesis were capable of escaping immune surveillance of the body. Such chronic inflammatory diseases are also capable of causing angiogenesis by introducing angiogenesis stimulatory factors.

In cases of rheumatoid arthritis, the rheumatoid synovial endothelium is enriched with VEGF and FGF-2 which are angiogenesis-promoting cytokines (Carmeliet 2003). Clinical trials in attempt to treat rheumatoid arthritis deals with prescribing TNF- α -blocking antibodies which in turn revealed a significant decrease in serum VEGF levels, decreasing the synovial vascularity (Carmeliet 2003). Growth factors like PlGF which plays an angiogenic role in disease without activating quiescent vessels in healthy tissues become an attractive target in the development of a safe anti-angiogenic treatment (Luttun et al. 2004).

In addition, many leukocyte subtypes were found to produce multiple angiogenic factors such as VEGF, PDGF and FGF and various interleukins and proteinases. For example, neutrophils and natural killer cells are implicated in cyclical uterine angiogenesis and in abnormal angiogenesis in endometriosis. This revealed that different leukocytes have the ability to promote many angiogenic processes. Nevertheless leukocytes also generate angiogenesis inhibitors. Hence their overall role in initiating or terminating angiogenesis depends on the temporal and spatial balance of the released modulators. Mast cells are also found to release angiogenic factors when they encounter allergens and pathogens in the skin and mucosa (Mori et al. 2013).

In conditions such as ischemic heart disease or preeclampsia, the angiogenic switch is insufficient which leads to endothelial cell dysfunction, vessel malformation or regression and revascularisation, therefore in such conditions promoting revascularization or therapeutic angiogenesis can be beneficial (Carmeliet 2005). Currently the angiogenic factors such as VEGF and FGF are the genes of interest to be used as a treatment for disorders such as ischaemic tissues.

1.4 Angiogenesis in Malignancy

Cancer is a broad term collectively used to describe a number of diseases in which abnormal cells divide uncontrollably and have the ability to invade other tissues. Cancerous cells are able to spread to other parts of the body via lymph and blood systems. In the normal state, our body's plethora of cell types is constantly undergoing growth, division and eventually replacement when they die or become damaged. The origin of cancer can be attributed to a genetic mutation within cell DNA that affects the normal cell growth and division. Normal cell death does not occur and instead unwarranted new cell formation and growth occurs leading to the development of a mass of tissue known as a tumour. It is important to note that not all tumours are necessarily cancerous. Benign tumours are non-cancerous, do not spread to other parts of the body and can be removed without reoccurrence. Malignant tumours however are much more dangerous and can metastasise to other

parts of the body (Nishida et al. 2006). Metastasis is the term used to describe cancer spread and the potential of cancer cells to leave the original site, penetrate into blood vessels and lymphatics and move through blood vessels to other organs (Seaman et al. 2007). Years of cancer research and meticulous studies have clearly shown that the angiogenesis phenomenon plays a pivotal role in the spread of cancer and is a fundamental concept involved in cancer metastasis; specifically the transition of benign tumours to those that become malignant. It was of previous thought prior to the 1960s that blood supply to tumours was actually carried out by vasodilation of pre-existing blood vessels. The new theories however support the angiogenesis phenomenon as the key blood supply for growth of tumours.

Angiogenesis, tumour growth and metastasis are also very closely related. In terms of malignancies, the extra blood vessels sustain the growth of tumours and in fact malignant tissue development, expansion and its spread are all angiogenesis-dependent (Folkman and Shing 1992). For a lethal tumour mass to develop, it is not only the proliferation of the malignant cells that is required but also the formation of new blood vessels (Folkman 2006). In a malignancy, endothelial cells of the neo-vasculature grow in a disorderly manner presenting with many structural shortcomings. Their role as a functional barrier is lost, and they cannot withstand the increased interstitial pressure, which then alters the blood flow and flux of leukocytes reaching the tumour site (Folkman 1996). Thus, angiogenesis not only sustains the tumour cells by meeting their increasing metabolic needs. They also, resulting from their incomplete structural build (i.e. a defective basement membrane and lack of normal perivascular connective tissue), provide these cells with many advantages; most importantly an easy penetration of metastatic tumour cells into circulation. Tumour cells can simply fall into the lumen and spread to distant sites and organs through the blood flow (Papetti and Herman 2002). Certain genetic changes promote the growth of malignant cells. So-called oncogenes – genetic material that carry the potential to induce cancer – give rise to uncontrolled proliferation and hindered apoptosis of the tumour cells (Papetti and Herman 2002). For every one endothelial cell there are approximately a hundred tumour cells, surrounding the vessels in a cylindrical fashion (Folkman 1995, 1996).

Cancer cells are able to release molecules that in fact act as signals to activate angiogenesis. These include many different many proteins and many other molecules. Among these, the two most angiogenic proteins are Vascular Endothelial Growth Factor (VEGF) and Basic Fibroblast Growth Factor (bFGF). Oncogenes induce angiogenesis primarily through an increased expression of the vascular endothelial growth factors (VEGF) in those cells (Folkman 1996, 2006). The tumour cells themselves are able to synthesize both VEGF and bFGF before they are secreted into surrounding tissue. From here, they bind to receptors (specific proteins) that are located on the outer surface of endothelial cells. Subsequent binding of protein to appropriate receptors initiates a signal transmission from a series of relay proteins into the endothelial cell nucleus (Munoz-Chapuli et al. 2004). This is a stimulus for a gene group to then make products required for endothelial cell growth.

Activated endothelial cells secrete a matrix called metalloproteinases (MMPs) into the surrounding tissue. MMPs are a distinct class of degradative enzymes and

are responsible for breakdown of extracellular matrix that is primarily consisted of proteins and polysaccharides. This degradation is required for the proliferation and migration of endothelial cells. Over time, the endothelial cells begin to divide as they proliferate and organize into hollow tubes that progressively advance into a mature network of blood vessels (Lamalice et al. 2007).

Given the fact that angiogenesis is a key step in the development and spread of cancer, blocking angiogenesis seems a plausible solution to stop or slow down tumour growth. Numerous studies have and are currently being undertaken to find synthetic and natural angiogenesis inhibitors. Angiogenesis inhibitors obstruct upon the binding multistep binding process of the signalling molecules such as VEGF and bFGF.

One of the most well-known of the angiogenesis inhibitors is a drug called *Bevacizumab* or *Avastin* (Samant and Shevde 2011; Tanne 2011). *Bevacizumab* is a monoclonal antibody – a laboratory synthesised molecule that is made in such a way that they mimic the antibodies that the human body naturally produces as part of the immune system. The antibody binds to VEGF and blocks signalling of the molecule, leading to suppressed formation of new blood vessel growth (angiogenesis). Reduced nutrient supply to the tumour can slow or stop its growth. The U.S. Food and Drug Administration (FDA) approves of *Bevacizumab* to be used alone for glioblastoma (tumours that originate from connective tissue in brain) and when in combination with other drugs for treatment of non-small cell lung cancers, metastatic colorectal cancer as well as metastatic renal cell cancer (Samant and Shevde 2011; Tanne 2011). Other FDA approved antiangiogenic drugs are *Sorafenib* (for hepatocellular carcinoma and kidney cancer), *Sunitinib* (kidney cancer and neuroendocrine tumours), *Pazopanib* (kidney cancer) and *Everolimus* (kidney cancer and neuroendocrine tumours) (Samant and Shevde 2011; Tanne 2011). Alternative theories also suggest that *Bevacizumab* induces more stability within the tumour blood vessels (usually leaky) thus allowing for more effective penetration of chemotherapy into cancer cells. Other drugs such as *Endostatins* inhibits angiogenesis directly rather than through the signalling pathway. This drug cause apoptosis of endothelial cells or destruction of proliferating endothelial cells (Samant and Shevde 2011). As outlined previously, breakdown of the extracellular matrix is essential for endothelial cell proliferation and migration. Therefore drugs that can target the MMP's within the matrix can also act at angiogenesis inhibitors.

It is important to distinguish between angiogenesis inhibitors and conventional anticancer drugs. As the name suggests, angiogenesis inhibitors inhibit the growth of blood vessels but does not necessarily kill tumours. Instead, they have a high chance of preventing tumours from growing so it is suggested that that the therapy must be spanned over a long period to have any major effect.

In some cases, the inhibitors will be more effective in combination with other remedies such as chemotherapies. Like other medications there are reports of common side effects for these inhibitors such as high blood pressure, fatigue, rash, dry and itchy skin, hand-foot syndrome (tender, thickened areas of skin), diarrhoea, wound healing problems/cuts reopening and low blood counts (O'Reilly et al. 1997).

In conclusion, angiogenesis plays a key role in the development of cancer. Upon this observation, a suggested strategy against tumour growth is to not only target the tumour cells alone but also the endothelial cells of the newly formed blood vessels – a technique that has so far shown to be effective in the battle against cancer (Folkman 2006). This book attempts to discuss and summarise the role of important regulators of angiogenesis in regards to normal physiology, disease and malignancy.

References

- Adair TH, Montani J-P (2010) Overview of angiogenesis. In: Rafael S (ed) Angiogenesis. Morgan & Claypool Life Sciences, San Rafael
- Albini A, Tosetti F, Li VW, Noonan DM, Li WW (2012) Cancer prevention by targeting angiogenesis. *Nat Rev Clin Oncol* 9(9):498–509. doi:[10.1038/nrclinonc.2012.120](https://doi.org/10.1038/nrclinonc.2012.120)
- Burri PH, Djonov V (2002) Intussusceptive angiogenesis – the alternative to capillary sprouting. *Mol Aspects Med* 23(6):1–27. doi:[10.1016/s0098-2997\(02\)00096-1](https://doi.org/10.1016/s0098-2997(02)00096-1)
- Carmeliet P (2003) Angiogenesis in health and disease. *Nat Med* 9(6):653–660. doi:[10.1038/nm0603-653](https://doi.org/10.1038/nm0603-653) nm0603-653 [pii]
- Carmeliet P (2005) Angiogenesis in life, disease and medicine. *Nature* 438(7070):932–936. doi:[10.1038/nature04478](https://doi.org/10.1038/nature04478)
- DeWitt N (2005) Angiogenesis. *Nature* 438(7070):931–931
- Felmeden DC, Blann AD, Lip GY (2003) Angiogenesis: basic pathophysiology and implications for disease. *Eur Heart J* 24(7):586–603
- Folkman J (1995) Angiogenesis in cancer, vascular, rheumatoid and other disease. *Nat Med* 1(1):27–31
- Folkman J (1996) New perspectives in clinical oncology from angiogenesis research. *Eur J Cancer* 32a(14):2534–2539
- Folkman J (2006) Angiogenesis. *Annu Rev Med* 57:1–18. doi:[10.1146/annurev.med.57.121304.131306](https://doi.org/10.1146/annurev.med.57.121304.131306)
- Folkman J, Shing Y (1992) Angiogenesis. *J Biol Chem* 267(16):10931–10934
- Hillen F, Griffioen AW (2007) Tumour vascularization: sprouting angiogenesis and beyond. *Cancer Metastasis Rev* 26(3–4):489–502. doi:[10.1007/s10555-007-9094-7](https://doi.org/10.1007/s10555-007-9094-7)
- Hong H, Chen F, Zhang Y, Cai W (2014) New radiotracers for imaging of vascular targets in angiogenesis-related diseases. *Adv Drug Deliv Rev* 76:2–20. doi:[10.1016/j.addr.2014.07.011](https://doi.org/10.1016/j.addr.2014.07.011)
- Lamalice L, Le Boeuf F, Huot J (2007) Endothelial cell migration during angiogenesis. *Circ Res* 100(6):782–794. doi:[10.1161/01.res.0000259593.07661.1e](https://doi.org/10.1161/01.res.0000259593.07661.1e)
- Liekens S, De Clercq E, Neyts J (2001) Angiogenesis: regulators and clinical applications. *Biochem Pharmacol* 61(3):253–270
- Luttun A, Autiero M, Tjwa M, Carmeliet P (2004) Genetic dissection of tumor angiogenesis: are PlGF and VEGFR-1 novel anti-cancer targets? *Biochim Biophys Acta* 1654(1):79–94. doi:[10.1016/j.bbcan.2003.09.002](https://doi.org/10.1016/j.bbcan.2003.09.002)
- Mori S, Tran V, Nishikawa K, Kaneda T, Hamada Y, Kawaguchi N, Fujita M, Takada YK, Matsuura N, Zhao M, Takada Y (2013) A dominant-negative FGF1 mutant (the R50E mutant) suppresses tumorigenesis and angiogenesis. *PLoS One* 8(2):e57927. doi:[10.1371/journal.pone.0057927](https://doi.org/10.1371/journal.pone.0057927)
- Munoz-Chapuli R, Quesada AR, Angel Medina M (2004) Angiogenesis and signal transduction in endothelial cells. *CMLS Cell Mol Life Sci* 61(17):2224–2243. doi:[10.1007/s00018-004-4070-7](https://doi.org/10.1007/s00018-004-4070-7)
- Nishida N, Yano H, Nishida T, Kamura T, Kojiro M (2006) Angiogenesis in cancer. *Vasc Health Risk Manag* 2(3):213–219. doi:[10.2147/vhrm.2006.2.3.213](https://doi.org/10.2147/vhrm.2006.2.3.213)
- O'Reilly MS, Boehm T, Shing Y, Fukai N, Vasios G, Lane WS, Flynn E, Birkhead JR, Olsen BR, Folkman J (1997) Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell* 88(2):277–285. doi:[10.1016/s0092-8674\(00\)81848-6](https://doi.org/10.1016/s0092-8674(00)81848-6)

- Papetti M, Herman IM (2002) Mechanisms of normal and tumor-derived angiogenesis. *Am J Physiol Cell Physiol* 282(5):C947–C970. doi:[10.1152/ajpcell.00389.2001](https://doi.org/10.1152/ajpcell.00389.2001)
- Samant RS, Shevde LA (2011) Recent advances in anti-angiogenic therapy of cancer. *Oncotarget* 2(3):122–134
- Seaman S, Stevens J, Yang MY, Logsdon D, Graff-Cherry C, St Croix B (2007) Genes that distinguish physiological and pathological angiogenesis. *Cancer Cell* 11(6):539–554. doi:[10.1016/j.ccr.2007.04.017](https://doi.org/10.1016/j.ccr.2007.04.017)
- Senger DR, Davis GE (2011) Angiogenesis. *Cold Spring Harb Perspect Biol* 3(8):a005090. doi:[10.1101/cshperspect.a005090](https://doi.org/10.1101/cshperspect.a005090)
- Smith SK (2001) Angiogenesis and reproduction. *BJOG* 108(8):777–783
- Tanne JH (2011) FDA cancels approval for bevacizumab in advanced breast cancer. *BMJ* 343:d7684. doi:[10.1136/bmj.d7684](https://doi.org/10.1136/bmj.d7684)
- Tonnesen MG, Feng X, Clark RA (2000) Angiogenesis in wound healing. *J Invest Dermatol Symp Proc Soc Invest Dermatol Inc Eur Soc Dermatol Res* 5(1):40–46. doi:[10.1046/j.1087-0024.2000.00014.x](https://doi.org/10.1046/j.1087-0024.2000.00014.x)