Manuscript title:

Clinical audits to improve critical care: Part 2: Analyse, benchmark and feedback

Authors:

Gillian Ray-Barruel

RN; BSN; GC ICU; BA (Honours); PhD

g.ray-barruel@griffith.edu.au

Postdoctoral Research Fellow: National Centre of Research Excellence in Nursing; Alliance for Vascular Access Teaching and Research (AVATAR) Group; Menzies Health Institute Queensland, Griffith University, Australia.

Honorary Research Fellow: Centre for Clinical Nursing; Royal Brisbane and Women's Hospital, Australia.

Visiting Scholar: Nursing Practice Development Unit, Princess Alexandra Hospital, Australia

Amanda J Ullman

RN; GC Paediatric ICU; MAppSci; PhD

Senior Lecturer: School of Nursing and Midwifery; Alliance for Vascular Access Teaching and Research (AVATAR) Group; Menzies Health Institute Queensland, Griffith University, Australia.

Honorary Research Fellow: Centre for Clinical Nursing; Royal Brisbane and Women's Hospital, Australia.

Honorary Research Fellow: Paediatric Critical Care Research Group; Lady Cilento Children's Hospital, Brisbane, Australia.

Claire M Rickard

RN; PhD

Professor of Nursing: National Centre of Research Excellence in Nursing, Alliance for Vascular Access Teaching and Research (AVATAR) Group, Menzies Health Institute Queensland, Griffith University, Australia

Honorary Research Fellow: Royal Brisbane and Women's Hospital, Australia

Honorary Research Fellow: Princess Alexandra Hospital, Woolloongabba, Australia

Honorary Research Fellow: The Prince Charles Hospital, Australia

Marie Cooke

RN; PhD

Professor of Nursing: National Centre of Research Excellence in Nursing, Alliance for Vascular Access Teaching and Research (AVATAR) Group, Menzies Health Institute Queensland, Griffith University, Nathan, QLD, Australia

Contributions of authors:

Conception and design of the study: all authors

Acquisition of data, or analysis and interpretation of data: all authors

Drafting the article and revising it critically for important intellectual content: all authors

Final approval of the version to be submitted: all authors

Acknowledgements:

All authors have approved the final article and acknowledge that all those entitled to authorship are listed as authors.

Title: Clinical audits to improve critical care: Part 2: Analyse, benchmark and feedback

Abstract

Clinical audits are an essential part of the cycle designed to ensure that patients receive the best quality of care. By measuring the care delivered against established best practice standards, it becomes possible to identify shortcomings and to plan targeted strategies and processes for continuous improvement. The success of a clinical audit depends upon defined goals, motivation of stakeholders, appropriate tools and resources, and clear communication.

In part 1 of this series, an overview of the structures and processes needed to prepare and collect data for clinical audits in the critical care setting was provided. In part 2, we discuss how to analyse the collected audit data, benchmark findings with internal and external data sets, and feedback audit results to critical care clinicians to promote evidence-based practice and improve patient outcomes.

Key words: Quality improvement; clinical audit; critical care; evidence-based practice; feedback

Total word count: 2523

1 Introduction

2 The importance of clinical audits in measuring performance and tracking progress is well recognised
3 by critical care leaders. Clinical audits are a quality improvement process, systematically undertaken
4 to improve clinical practice and subsequent patient outcomes. By providing objective and quantifiable
5 data, clinical audits enable clinicians to compare current performance with explicit, defined criteria,
6 and identify areas for improvement.¹ Clinical audits can be used to monitor and track both clinical
7 practice and service delivery changes, and they provide a useful and objective tool to motivate
8 healthcare staff to engage in the process of continuous quality improvement.

10 Undertaking a clinical audit in the critical care unit need not be a daunting task. With careful
11 preparation and an informed approach, clinical audits can provide a useful and valuable tool for
12 critical care clinicians. Preparing and committing to a clear audit strategy creates transparency in the
13 process, ensures validity and reliability of the data, and builds confidence in the findings. The success
14 of an audit will depend upon clearly defined goals, motivation of key opinion leaders and
15 stakeholders, appropriate tools and resources (time, staff, equipment), and clear communication.

17 This is Part Two of a two-paper series regarding clinical audits in critical care. In the previous article,
18 the structures and processes needed to prepare and collect data for clinical audits in the critical care
19 setting were addressed. This article features an overview of the skills and resources needed to analyse,
20 benchmark and feedback audit data. Together, these articles provide a step-by-step guide, and the
21 remaining five steps are outlined here. These are:

- 22 7. Identify appropriate techniques for analysing audit data
- 23 8. Identify internal and external audit data for benchmarking
- 24 9. Feedback to clinical area and management
- 25 10. Plan targeted strategies and processes for continuous improvement
- 26 11. Repeat the audit cycle: re-identify areas for improvement

Step Seven: Identify appropriate techniques for analysing audit data

Once the data collection is complete, the data must be collated and analysed. The primary goal of data analysis is to understand the key audit findings, so these can be presented to the clinical staff and other stakeholders, and action plans can be developed, if needed. Before any data analysis can begin, it is important to check the data are “clean”, that is, no typographic errors have been made during data entry. This does not mean that all the audit data entry needs to be double-checked (although this is ideal), but attention is at least needed to confirm/correct any missing data, and check any extreme results (high and low), as well as any impossible results (e.g., age = 187 years old). Ideally, the person checking the data should have a thorough understanding of the data collection tool and an excellent eye for detail, as well as knowledge of the clinical practice or service being audited.

An in-depth knowledge of statistics is not necessary when analysing audit data. For the purposes of clinical audit, it is perfectly acceptable in many cases to present the data using simple percentages and absolute numbers, incorporating appropriate numerators and denominators whenever possible.² Percentages represent a calculation of the number of times an event occurred based on the total number of people, whereas rates represent the probability of a certain event. For instance, to determine the percentage of pressure injuries in the critical care unit, a simple snapshot audit could identify the number of pressure injuries detected (numerator) in the total sample of patients (denominator) and multiply by 100. To calculate the rate of pressure injuries for your unit, it would be more accurate to keep track of the total number of pressure injuries per 1,000 patient days. While the simple percentage provides a neat starting point to identify if the unit has a problem with pressure injuries, the rate provides a better picture of the extent of the problem over time.

The primary purpose of the clinical audit is to improve practice, and an essential component of this is building staff engagement in the continuous quality improvement process, therefore, simple calculations and descriptive statistics³ that can be easily presented and understood by busy clinicians are preferred. Software programs such as Microsoft Excel® are satisfactory for this level of analysis. If more detailed enquiry is desired, statistical software packages can be used, but this is certainly not

essential. If capacity exists, inferential statistics, such as Chi-square, T-tests, and Mann-Whitney, can be used to determine meaningful differences between samples. More information regarding the appropriate use of descriptive and inferential statistics can be found in the series of statistics articles published by Australian Critical Care.³⁻⁹

Step Eight: Identify internal and external audit data for benchmarking

The presentation of results using internal and external benchmarking is effective in provoking discussion surrounding the results and strategies for improvement.^{10, 11} Benchmark criteria indicate a desired level of care in the critical care area.¹² Therefore, clinical practice guidelines should be used to inform the audit criteria, as discussed in Step Three. These criteria may be in the form of rates or percentages (e.g., percentage of patients receiving enteral nutrition within 48 hours of ICU admission), but need to provide a realistic and attainable goal for the local unit.

Benchmarking can be either internal or external, and decisions about benchmarking criteria should be made with the key stakeholders to ensure the benchmarks chosen are clinically relevant and appropriate. Internal benchmarking can be accomplished via repeated audits over time in the same institution, using the same audit tools. For the majority of data types, results may be presented using graphical illustrations such as Statistical Process Control (SPC) Charts. Figure 1 illustrates how tools such as SPC Charts can be used to present data over time for repeated measures, with built-in thresholds (upper and lower control lines) to highlight significant variations in practice. These significant variations emphasise results outside of the normal fluctuations in the statistical 'norm' of care and reflect three standard deviations above (upper control line) or below (lower control line) the mean (central line).¹³ These statistical norms are most valuable when external benchmarks do not exist. Resources are readily available to support the development of these charts which use programmes including Microsoft Excel® (see Table 1: NHS Institute for Innovation and Improvement and iSixSigma®).

<Insert Figure 1>

1
2 External benchmarking provides another form of goal setting for improvement, but should be chosen
3 in consultation with stakeholders to ensure their relevance to the institution. External benchmarks can
4 be identified within a range of resources including international institutions (e.g., World Health
5 Organization), national institutions (e.g., Australian Commission on Safety and Quality in
6 Healthcare), discipline-specific clinical registries (e.g., Australian and New Zealand Intensive Care
7 Society registry), or published audits using similar methods of assessment. Some examples of
8 common indicators to benchmark ICU performance include: the rate of central-line associated
9 bloodstream infection (CLABSI) (expressed as number of CLABSI occurrences per 1,000 line days);
10 the Standard Mortality Ratio (actual deaths divided by the predicted number of deaths at each ICU);
11 the rate of ICU readmissions, declined admissions, or after-hours discharge.¹⁴ An example of external
12 international benchmarking is the International Nutrition Survey, in which 150 ICUs participate
13 annually.¹⁵

15 **Step Nine: Feedback to clinical area and management**

16 Providing feedback about the audit results is one of the most important but frequently undervalued
17 aspects of the audit cycle.^{16, 17} The goal of feedback is to raise awareness and challenge beliefs about
18 current practice and clinical outcomes, with the assumption that people will be motivated to change
19 when presented with suboptimal results and a clear action plan.¹⁶ Developing appropriate feedback
20 strategies should be the result of in-depth, collaborative discussion between the original project
21 stakeholders. Audit feedback is most effective when delivered by a well-respected supervisor or
22 colleague, rather than an external party.^{16, 18} This person should ideally be passionate about improving
23 practice and have a good rapport with the critical care staff. Importantly, feedback needs to be timely,
24 individualised and non-punitive in order to be effective in improving performance.^{19, 20} A lag time of
25 months between collecting the data and providing feedback to staff can lead to disinterest in the
26 findings.²¹

1 Current audit results can be displayed beside previous audit results or the results of another area in the
2 same hospital. Alternatively, results can be displayed alongside benchmarking criteria, such as clinical
3 practice guidelines or external audit data, to enable comparison of the current audit findings with the
4 expected level of performance. How to best deliver and display audit feedback depends on the
5 intended audience, but a range of strategies is likely to be more effective.¹⁶ Possible strategies could
6 include interactive workshops, in-service sessions, laminated posters, display boards, email bulletins,
7 newsletters, and screensavers, among others. For maximal impact, visual and verbal presentations
8 should be delivered in a clear, comprehensive format, using tables, charts and graphs (for example,
9 presentation software, such as Microsoft PowerPoint®, is useful). The prominent display of audit
10 results in a populated area of the unit, such as the office, hallway or tearoom can encourage staff to
11 read and engage with the audit findings. During in-service sessions and workshops, staff can be
12 prompted to discuss the findings and their relevance to their own unit practice, as well as their
13 adherence or deviance from best practice guidelines, and they should be encouraged to be creative in
14 coming up with possible strategies for improvement. Encouraging patients and visitors to take an
15 interest in the unit's quest for improvement by displaying the audit results in the waiting room can
16 also be a proactive strategy for consumer engagement and participation.²²

17

18 Targeted feedback strategies should incorporate relevance to the local unit, individual hospital and
19 wider audience. As identified in Step Two ('Engaging of stakeholders'), discussed in part 1 of this
20 article series, key stakeholders play an essential role in preparing the audit, developing the audit tools,
21 data collection, and disseminating the findings of the audit. Staff will be more motivated to embrace
22 the audit process and adopt change if they can see the relevance of the audit findings to the local
23 situation.^{18, 20} Not all staff will experience the same degree of motivation, so it can be helpful to
24 engage key stakeholders to identify and work with those staff members who are more driven to
25 improve practice and encourage them to champion the desired change.¹⁸ Remember that if there have
26 been many recent changes in practices or processes within the unit, implementing further changes
27 could be difficult, therefore it is also important to consider the best timing for undertaking audits and
28 implementing new practices.

Step Ten: Plan targeted strategies and processes for continuous improvement

The successful implementation of improvements requires investment by all stakeholders involved in the audit process.¹⁹ If gaps between audited practice and clinical practice guidelines or high-quality evidence exist, this provides a good starting point to develop the practice improvement strategies. Consider the development of an action plan with timeframes to record these recommendations. This plan should include clear targets and the name of each person who agreed to take responsibility for each element.¹² It is preferable to focus on one or two areas needing improvement, rather than attempt to change too many things at once. Be warned, however, systematic reviews have demonstrated that practice change via rapid dissemination of educational materials²³ or clinical practice guidelines alone is rarely effective at improving patient outcomes.¹²

Improvements should be implemented using a combination of initiatives, tailored to the local critical care audience and resources.^{24, 25} The stakeholder group, including interdisciplinary clinicians, patient and family representatives, safety and quality experts, and institutional management, needs to plan appropriate improvement interventions and prospectively identify factors that will encourage or discourage their application (i.e., the determinants of practice).²⁶ Understanding these determinants (e.g., a lack of knowledge, lack of staff and time) and using this information to devise appropriate strategies will improve their effectiveness.²⁶ Tailored practice improvement strategies can include targeted educational bedside visits, reminders (manual or computerised), checklists, interactive educational meetings, and multiple educational resources.^{24, 25, 27} Areas highlighted by the audit findings that require improvement may also be the focus for developing research and on-going, targeted quality improvement activities so as to build and maintain a focus on care improvements in the unit.^{16, 17} It may also be possible to connect with critical care special interest groups, patient safety groups or consumer advisory panels, to determine what activities and improvements other clinical areas have instituted.

1 Audit results that demonstrate notable improvements should be acknowledged and staff commended
2 for their efforts. Audit drivers and change champions can be encouraged to present the findings at unit
3 meetings, hospital quality and safety meetings, journal clubs, the intranet and local conferences.
4 Sharing experiences with audit processes and subsequent practice improvements in critical care
5 newsletters and journals is strongly recommended. Not only is this an excellent way to develop
6 research skills, but it also can open up exciting avenues for further networking and benchmarking
7 with colleagues from other critical care units.

9 **Step Eleven: Repeat the audit cycle: re-identify areas for improvement**

10 After a period of implementing improvements, it is important to evaluate them at least once, and
11 preferably at regular intervals, to see if they were successful. Re-auditing also monitors whether any
12 improvements were maintained after the initial post-audit enthusiasm. If the goals were not reached, a
13 continued program of strategies to achieve change must continue, and the auditing process repeated
14 until it can be demonstrated that the desired practice has been successfully implemented and sustained
15 over time. Once it has been decided that a practice has improved and is being sustained, a new focus
16 for improvement can be addressed.

18 Performance indicators can be used to drive the continuous quality improvement cycle. Progress
19 reports should be communicated and copies made readily available to staff and managers. A regular
20 auditing time period should be mutually agreed upon by the stakeholder group, providing enough time
21 to implement improvements while also being sustainable for the critical care area. The goal is always
22 to improve clinical practice and patient outcomes, so the planned frequency of audits will depend on
23 the practice being audited. While there is no one size fits all solution for deciding how often to audit,
24 as a general guide, the frequency of auditing should be based upon your unit's needs and capacity,
25 compliance regulations, and risk assessment. For example, ongoing auditing and reporting of hospital
26 patient falls data is required by the Australian Council on Healthcare Standards, whereas monitoring
27 insulin infusion management following the introduction of a new guideline²⁸ would likely be a time-
28 limited unit audit.

473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531

1
2
3
4
5
6
7
8
9
10
11
12
13
14

Conclusion

Clinical audits can be effective tools to promote best practice, reducing errors in the critical care setting,²⁹ but their success relies on several characteristics. Most importantly, it is essential to focus and define the clinical audit goals from the beginning. Next, identify and engage with key opinion leaders and local stakeholders throughout the audit process. Spend time developing the appropriate method and tool for undertaking the clinical audit. Remember, benchmarking and providing goals for improvement will strengthen the audit, but make sure it is the right benchmark for the clinical area. Finally, communicate the results of the audit, including a plan for improvement. Keep going! Auditing is a cyclical process and should be repeated regularly to ensure improvements are effective.

Acknowledgements

No funding received.

References

1. Copeland G. A practical handbook for clinical audit. In: Team NCGS, editor.: NHS; 2005.
2. Verma R. Data quality and clinical audit. *Anaesthesia and intensive care medicine*. 2009;**10**(8):400-2.
3. Fisher MJ, Marshall AP. Understanding descriptive statistics. *Aust Crit Care*. 2009;**22**(2):93-7.
4. Burmeister E, Aitken LM. Sample size: How many is enough? *Aust Crit Care*. 2012;**25**(4):271-4.
5. Fethney J. Statistical and clinical significance, and how to use confidence intervals to help interpret both. *Aust Crit Care*. 2010;**23**(2):93-7.
6. Fisher MJ, Marshall AP, Mitchell M. Testing differences in proportions. *Aust Crit Care*. 2011;**24**(2):133-8.
7. Pereira SMC, Leslie G. Hypothesis testing. *Aust Crit Care*. 2009;**22**(4):187-91.
8. Prematunga RK. Correlational analysis. *Aust Crit Care*. 2012;**25**(3):195-9.
9. Rickard CM. Statistics for clinical nursing practice: An introduction. *Aust Crit Care*. 2008;**21**(4):216-9.
10. Ettorchi-Tardy A, Levif M, Michel P. Benchmarking: a method for continuous quality improvement in health. *Healthc Policy*. 2012;**7**(4):e101-e19.
11. Kiefe CI, Allison JJ, Williams OD, Person SD, Weaver MT, Weissman NW. Improving quality improvement using achievable benchmarks for physician feedback: a randomized controlled trial. *JAMA*. 2001;**285**(22):2871-9.
12. Benjamin A. Audit: how to do it in practice. *Br Med J*. 2008;**336**(7655):1241-5.
13. Noyez L. Control charts, Cusum techniques and funnel plots. A review of methods for monitoring performance in healthcare. *Interact Cardiovasc Thorac Surg*. 2009;**9**(3):494-9.
14. Australian and New Zealand Intensive Care Society. ANZICS Centre for Outcome and Resource Evaluation Annual Report 2014-2015. Melbourne: ANZICS; 2016.
15. Clinical Evaluation Research Unit (CERU). Critical Care Nutrition. Ontario. 2017 [Available from: <http://criticalcarenutrition.com/ins>].
16. Ivers N, Jamtvedt G, Flottorp S, Young JM, Odgaard-Jensen J, French SD, et al. Audit and feedback: effects on professional practice and healthcare outcomes. *Cochrane Database Syst Rev*. 2012;**6**:Cd000259.
17. Gardner B, Whittington C, McAteer J, Eccles MP, Michie S. Using theory to synthesise evidence from behaviour change interventions: the example of audit and feedback. *Soc Sci Med*. 2010;**70**(10):1618-25.
18. Christina V, Baldwin K, Biron A, Emed J, Lepage K. Factors influencing the effectiveness of audit and feedback: nurses' perceptions. *J Nurs Manag*. 2016;**24**(8):1080-7.
19. Hysong SJ, Best RG, Pugh JA. Audit and feedback and clinical practice guideline adherence: making feedback actionable. *Implement Sci*. 2006;**1**:9.
20. Grainger A. Clinical audit: shining a light on good practice. (cover story). *Nurs Manag*. 2010;**17**(4):30-3.
21. Colquhoun H, Michie S, Sales A, Ivers N, Grimshaw JM, Carroll K, et al. Reporting and design elements of audit and feedback interventions: a secondary review. *BMJ Qual Saf*. 2016.
22. Australian Commission on Safety and Quality in Health Care. National Safety and Quality Health Service Standards. In: ACSQHC, editor. Sydney: Commonwealth of Australia; 2012.
23. Giguere A, Legare F, Grimshaw J, Turcotte S, Fiander M, Grudniewicz A, et al. Printed educational materials: effects on professional practice and healthcare outcomes. *Cochrane Database Syst Rev*. 2012;**10**:Cd004398.
24. Foy R, Eccles M, Jamtvedt G, Young J, Grimshaw J, Baker R. What do we know about how to do audit and feedback? Pitfalls in applying evidence from a systematic review. *BMC Health Serv Res*. 2005;**5**(1):50.
25. Johnson MJ, May CR. Promoting professional behaviour change in healthcare: what interventions work, and why? A theory-led overview of systematic reviews. *BMJ Open*. 2015;**5**(9):e008592.

591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649

- 1 26. Baker R, Camosso-Stefinovic J, Gillies C, Shaw EJ, Cheater F, Flottorp S, et al. Tailored
2 interventions to address determinants of practice. *Cochrane Database Syst Rev*.
3 2015;4:Cd005470.
- 4 27. O'Brien MA, Rogers S, Jamtvedt G, Oxman AD, Odgaard-Jensen J, Kristoffersen DT, et al.
5 Educational outreach visits: effects on professional practice and health care outcomes.
6 *Cochrane Database Syst Rev*. 2007(4):Cd000409.
- 7 28. Breeding J, Welch S, Buscher H, Nair P, Frost C, Newman S, et al. A retrospective audit of
8 insulin infusion management involving a locally developed dynamic insulin infusion guideline
9 in a tertiary ICU. *Australian Critical Care*.28(3):149-59.
- 10 29. Sinuff T, Cahill NE, Dhaliwal R, Wang M, Day AG, Heyland DK. The value of audit and feedback
11 reports in improving nutrition therapy in the intensive care unit: a multicenter observational
12 study. *JPEN J Parenter Enteral Nutr*. 2010;34(6):660-8.

13

Figure 1. Example of a Statistical Process Chart demonstrating unplanned ICU readmissions per quarter