**Comparative Studies of Mammalian Acid Lipases: Evidence for a New
Gene Family in Mouse and Rat (*Lipo*)**

Roger S Holmes^{1,4}, Laura A Cox^{1,2} and John L VandeBerg^{1,2}

¹Department of Genetics and ²Southwest National Primate Research Center, Southwest Foundation for Biomedical Research, San Antonio, TX, USA, and ³School of Biomolecular and Physical Sciences, Griffith University, Nathan, QLD, Australia

⁴Corresponding Author:

Roger S Holmes, DSc.

Department of Genetics

Southwest National Primate Research Center

Southwest Foundation for Biomedical Research

San Antonio, TX, USA 78227

Email: rholmes@sfbrgenetics.org

Phone: 210-258-9687

Fax: 210-670-3344

Keywords: Mammals; amino acid sequence; acid lipases; evolution; gene duplication.

Running Head: Mammalian acid lipases: comparative genomics and proteomics

Summary

At least six families of mammalian acid lipases (E.C. 3.1.1.-) catalyse the hydrolysis of triglycerides in the body, designated as LIPA (lysosomal), LIPF (gastric), LIPJ (testis) and LIPK, LIPM and LIPN (epidermal), which belong to the AB hydrolase superfamily. In this study, *in silico* methods were used to predict the amino acid sequences, secondary and tertiary structures, and gene locations for acid lipase genes and encoded proteins using data from several mammalian genome projects. Mammalian acid lipase genes were located within a gene cluster for each of the 8 mammalian genomes examined, including human (*Homo sapiens*), chimpanzee (*Pons troglodytes*), rhesus monkey (*Macacca mulatta*), mouse (*Mus musculus*), rat (*Rattus norvegicus*), cow (*Bos taurus*), horse (*Equus caballus*) and dog (*Canis familiaris*), with each containing 9 coding exons. Human and mouse acid lipases shared 44-87% sequence identity and exhibited sequence alignments and identities for key amino acid residues and conservation of predicted secondary and tertiary structures with those previously reported for human gastric lipase (LIPF) (Roussel *et al.*, 1999). Evidence for a new family of acid lipase genes is reported for mouse and rat genomes, designated as *Lipo*. Mouse acid lipase genes are subject to differential mRNA tissue expression, with *Lipa* showing wide tissue expression, while others have a more restricted tissue expression in the digestive tract (*Lipf*), salivary gland (*Lipo*) and epidermal tissues (*Lipk*, *Lipm* and *Lipn*). Phylogenetic analyses of the mammalian acid lipase gene families suggested that these genes are products of gene duplication events prior to eutherian mammalian evolution and derived from an ancestral vertebrate *LIPA* gene, which is present in the frog, *Xenopus tropicalis*.

Introduction

At least six mammalian acid lipase genes have been reported, including *LIPA*, encoding lysosomal acid lipase/cholesteryl ester hydrolase (E.C.3.1.1.13) (Anderson & Sando, 1991; Anderson *et al.*, 1994; Ameis *et al.*, 1994); *LIPF*, encoding a gastric lipase (E.C.3.1.1.3) (Bodmer *et al.*, 1987; Lohse *et al.*, 1997) ; *LIPJ*, expressed in testis (Thierry-Mieg & Thierry-Mieg, 2006) and three other genes (*LIPK*, *LIPM* and *LIPN*), which are expressed in epidermal cells of the body (Toulza *et al.*, 2007) and form part of an acid-lipase gene complex on human chromosome 10 (Deloukas *et al.*, 2004). Acid lipases have the capability to withstand acid conditions and lack any significant homology (<20%) with previously described neutral lipases (Bodmer *et al.*, 1987), including endothelial lipase (LIPE) (Hirata *et al.*, 1999; Jaye *et al.*, 1999), hepatic lipase (LIPC) (Martin *et al.*, 1998), lipoprotein lipase (LIPL) (Wion *et al.*, 1987) and pancreatic lipase (LIPP) (Lowe *et al.*, 1989), which perform specialized roles in lipid metabolism in various tissues and cells of the body.

LIPA catalyses the deacylation of triacylglycerols and cholesteryl esters of lysosomal low density lipoproteins (LDLs), an essential intracellular lipid catabolic process (Goldstein *et al.*, 1975; Wang *et al.*, 2008). Two major genetic diseases, a severe infantile-onset Wolman disease (Patrick & Lake, 1969; Hoeg *et al.*, 1984) and a milder late-onset cholesteryl ester storage disease (CESD) (Assmann *et al.*, 1973), are caused by mutations of the *LIPA* gene. *LIPF* is involved with the metabolism of dietary triglycerides under acidic conditions, being synthesized by gastric chief cells in the fundic mucosa of the stomach and responsible for 30% of triglyceride digestion in humans (Bodmer *et al.*, 1987). Structures for other acid lipase genes have been determined, including *LIPJ*, *LIPK*, *LIPM* and *LIPN*, and derived from whole genome sequences for human chromosome 10 (Deloukas *et al.*, 2004; Toulza *et al.*, 2007) and mouse chromosome 19 (The MGC Project Team, 2004; Carninci *et al.*, 2005), which contain acid lipase gene clusters in each case. Human *LIPK*, *LIPM* and *LIPN* genes are specifically expressed in epidermal cells and may play a role in differentiated keratinocyte cells in the body (Toulza *et al.*, 2007). Mammalian acid lipase genes usually contain 9 coding exons of DNA encoding enzyme sequences which undergo exon shuffling generating several acid lipase isoproteins (Thierry-Mieg and Thierry-Mieg, 2006). Predictive three-dimensional structural analyses of human *LIPA* have been undertaken using the human gastric lipase as a model, and key residues and sequences have been identified (Roussel *et al.*, 1999).

This paper reports the predicted gene structures and amino acid sequences for several mammalian acid lipase genes and proteins, including human (*Homo sapiens*), chimpanzee (*Pons troglodytes*), rhesus monkey

(*Macacca mulatta*), mouse (*Mus musculus*), rat (*Rattus norvegicus*), cow (*Bos taurus*), horse (*Equus caballus*) and dog (*Canis familiaris*). Predicted secondary and tertiary structures for mammalian acid lipases are also described, as well as the structural, phylogenetic and evolutionary relationships of these genes and enzymes with other mammalian lipase gene families. In addition, evidence for a new family of acid lipase genes is reported for mouse and rat genomes, designated as *Lipo*.

Methods

***In silico* mammalian acid lipase gene and protein identification.**

BLAST (Basic Local Alignment Search Tool) studies were undertaken using web tools from the National Center for Biotechnology Information (NCBI) (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) (Altschul *et al.*, 1997). Protein BLAST analyses used mammalian acid lipase amino acid sequences previously described (Table 1). Non-redundant protein sequence databases for several mammalian and vertebrate genomes were examined using the blastp algorithm, including human (*Homo sapiens*) (International Human Genome Sequencing Consortium, 2001); chimpanzee (*Pan troglodytes*) (Chimpanzee Sequencing & Analysis Consortium, 2005); orangutan (*Pongo abelii*) (Orangutan Genome Project, 2007); rhesus monkey (*Mucaca mulatta*) (Gibbs *et al.*, 2007), horse (*Equus caballus*) (Horse Genome Project, 2008), cow (*Bos Taurus*) (Bovine Genome Project, 2008); mouse (*Mus musculus*) (Mouse Genome Sequencing Consortium, 2002); rat (*Rattus norvegicus*) (Rat Genome Sequencing Consortium, 2004); guinea pig (*Cavia porcellus*) (MGC Project Team, 2004); (dog (*Canis familiaris*) Dog Genome Project, 2005); and frog (*Xenopus tropicalis*) (*Xenopus tropicalis* Genome Project, 2005). This procedure produced multiple BLAST ‘hits’ for each of the protein databases which were individually examined and retained in FASTA format, and a record kept of the sequences for predicted mRNAs and encoded acid lipase-like proteins . These records were derived from annotated genomic sequences using the gene prediction method: GNOMON and predicted sequences with high similarity scores generated.

BLAT analyses were subsequently undertaken for each of the predicted acid amino acid sequences using the UC Santa Cruz genome browser [<http://genome.ucsc.edu/cgi-bin/hgBlat>] (Kent *et al.* 2003) with the default settings to obtain the predicted locations for each of the mammalian acid lipase genes, including predicted exon boundary locations and gene sizes (see Table 1). Structures for mouse acid lipase isoforms were obtained using the AceView website (<http://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/index.html?human>) to examine predicted gene and protein structures to interrogate this database of mouse mRNA sequences for mouse *Lipa*, *Lipf*, *Lipk*, *Lipm*, *Lipn* and *Lipo1* genes(Thierry-Mieg and Thierry-Mieg, 2006).

Predicted Structures and Properties of Mouse Acid Lipases.

Predicted secondary and tertiary structures for mouse acid lipases were obtained using the PSIPRED v2.5 web site tools provided by Brunel University [<http://bioinf.cs.ucl.ac.uk/psipred/psiform.html>] (McGuffin *et al.* 2000) and the SWISS MODEL web tools [<http://swissmodel.expasy.org>], respectively (Guex & Peitsch 1997; Kopp & Schwede 2004). The reported tertiary structure for human gastric lipase (LIPF) (Roussel *et al.*, 1999) served as the reference for the predicted mouse acid lipase tertiary structures, with a modeling range of residues 22-395 (LIPA); 20-395 (LIPF); 27-394 (LIPK); 40-409 (LIPM); 28-398 (LIPN); and 24-392 (LIPO1). Theoretical isoelectric points and molecular weights for mammalian acid lipases were obtained using Expasy web tools (http://au.expasy.org/tools/pi_tool.html). SignalP 3.0 web tools were used to predict the presence and location of signal peptide cleavage sites (<http://www.cbs.dtu.dk/services/SignalP/>) for each of the predicted mammalian acid lipase sequences (Emanuelsson *et al.* 2007). The NetNGlyc 1.0 Server was used to predict potential N-glycosylation sites for human, mouse and rat acid lipases (<http://www.cbs.dtu.dk/services/NetNGlyc/>). Predictions of subcellular locations for mammalian acid lipases were conducted using PSORT 11 (<http://psort.ims.u-tokyo.ac.jp/form2.html>) (Horton & Nakai, 1997).

Mouse Acid Lipase Gene Expression

The mouse genome browser (<http://genome.ucsc.edu>) (NCBI37/mm9 2007 assembly) (Kent *et al.* 2003) was used to examine GNF Expression Atlas 2 data using various expression chips for mouse acid lipase genes *Lipa*, *Lipf*, *Lipk*, *Lipm*, *Lipn* and *Lipo1* (using GenBank ID AI747699) (<http://biogps.gnf.org>). Mouse chip expression 'heat maps' were examined for comparative gene expression levels among mouse tissues showing high (red); intermediate (black); and low (green) expression levels.

Phylogenetic Studies and Sequence Divergence

Alignments of acid lipase protein sequences and percentages of sequence identities were assembled using BioEdit v.5.0.1 and the default settings (Hall, 1999). Alignment ambiguous regions, including the amino and carboxyl termini, were excluded prior to phylogenetic analysis (BioEdit v.5.0.1) yielding alignments of 365 residues for comparisons of mammalian LIPA, LIPF, LIPJ, LIPK, LIPM, LIPN and LIPO sequences with the frog (*Xenopus tropicalis*) LIPA sequence (Table 1). Evolutionary distances were calculated using the Kimura option (Kimura, 1983) in TREECON (Van de Peer & de Wachter, 1994). Phylogenetic trees were constructed from evolutionary distances using the neighbor-joining method (Saitou & Nei, 1987). Tree topology was

reexamined by the boot-strap method (100 bootstraps were applied) of resampling and only values that were highly significant (≥ 90) are shown (Felsenstein, 1985).

Results and Discussion

Alignments of Mouse Acid Lipase Amino Acid Sequences.

Amino acid sequence alignments for five previously reported mouse acid lipases [LIPA, LIPF, LIPK, LIPM and LIPN (Carninci *et al.*, 2005)] are shown in Figure 1 together with predicted sequences for four new acid lipases (designated LIPO1, LIPO2, LIPO3 and LIPO4). The relative values of sequence identities (41-60%) and comparisons of amino acid sequence alignments for the mouse LIPA, LIPF, LIPK, LIPM, LIPN and LIPO1 sequences strongly suggest that these proteins are products of distinct but related gene families, whereas the predicted mouse LIPO1, LIPO2, LIPO3 and LIPO4 sequences exhibited higher levels of identities (76-96%), indicating that these are members of the same family, designated as LIPO (or *Lipo* for the gene family) (Table 2).

Amino acid sequences for these nine mouse acid lipase proteins contained between 395 (LIPF) and 422 (LIPM) residues with the latter exhibiting extended N- and C-termini (Figure 1). The results of three dimensional structural studies for human LIPF were used to identify key residues which are likely to contribute to the catalytic and structural features for these enzymes (sequence numbers refer to mouse LIPA) (Roussel *et al.*, 1999). These included the catalytic triad for the active site (Ser172; Asp343; His372); the active site motif (Gly-Xaa-Ser-Yaa-Gly) (residues 172-176); residues Leu89 and Gln175 (replaced with 175Glu for chicken LIPA) which stabilize the 'oxyanion' transition state during catalysis; and cysteine residues which form a disulfide bond (Cys248/Cys257 [37] to support the enzyme's structure.

The hydrophobic N-terminus signal peptide function (residues 1-18 for mouse LIPA) has been retained for all of the mouse acid lipase sequences examined, although these vary in length from 18 for LIPA (residues 1-18) to 33 (residues 1-33 for LIPM) (Figure 1). The mannose-6-phosphate containing N-glycosylation site (residues 161-3: Asn-Lys-Thr for mouse LIPA) (Sleat *et al.*, 2006) was not present for other mouse acid lipase sequences, with the exception of mouse LIPN, which supports the reported microlocalization of LIPA within lysosomes (Goldstein *et al.*, 1975). A basic amino acid 'patch' at the mouse LIPA C-terminus (residues 394Lys-395Lys) is present only within the LIPA sequence, which may interact with lysosomal UDP-N-acetylglucosamine phosphotransferase, causing phosphorylation of specific LIPA residues, which are proposed to target this enzyme for lysosomal location (Baranski *et al.*, 1990). Two other high probability N-glycosylation sites predicted for mouse LIPA (Asn36-Val37-Ser38; and Asn273-274Met-275Ser) were also observed for all other human, mouse and rat acid lipase sequences

examined. Other high probability N-glycosylation sites are described in Table 3 and Figure 1, including site 2 for human LIPA (72Asn-73His-74Ser) and mouse LIPO1 and LIPO3 sequences; site 3 for mouse LIPA (99Asn-100Ser-101Ser), mouse LIPF (98Asn-99Asn-100Ser), human LIPJ (68Asn-69Asn-70Ser), mouse LIPM (113Asn-114Asn-115Ser) and mouse LIPN (102Asn-103Gly-104Ser); site 5 for human LIPF (185Asn-186Pro-187Ser) and mouse LIPO3 (183Asn-184Gln-185Ser); site 8 for human LIPA (321Asn-322Gln-323Ser), human LIPJ (288Asn-289Gln-290Ser), human LIPN (320Asn-321Gln-322Ser) as well as mouse and rat LIPO sequences (316Asn-317Gln-318Ser for mouse LIPO1); site 9 for human LIPF (327Asn-328Val-329Thr) and LIPK (327 Asn-328Ile-329Thr); site 10 for mouse LIPO sequences (357Asn-358Leu-359Thr); and site 11 for human LIPN (413Asn-414Leu-415Ser). Four N-glycosylation sites have been previously identified for human LIPJ by three dimensional studies (Roussel et al., 1999) which may contribute to the stability and activity of this enzyme in acid environments. Individual differences were observed for the theoretical isoelectric points (pI) of the human, mouse and rat acid lipases examined, with higher values (pI values > 8) predicted for human LIPK, mouse LIPK and LIPM and rat LIPK, LIPM and LIPN, as compared with the other acid lipases examined, which exhibited lower predicted pI values (Table 1).

Predicted Secondary and Tertiary Structures for Mammalian Acid Lipases.

Analyses of predicted secondary structures for mammalian acid lipase sequences were compared with the previously reported secondary structure for human LIPF, or human gastric lipase, and the predicted structure for human LIPA (Roussel et al., 1999) (Figure 1). Similar α -helix β -sheet structures were observed for all of the mammalian acid lipases examined, particularly near key residues or functional domains, including the α -helix within the N-terminal signal peptides, the β -sheet and α -helix structures surrounding the active site Ser172 (for mouse LIPA) and the α -helix enclosing the lysosomal targeting signal residues (Asn-Lys-Thr residues 159-161 for mouse LIPA). The pattern of secondary structures were very similar to those reported for human LIPF and predicted for human LIPA and are numbered according to Roussel and coworkers (1999). These have been previously described as globular enzymes which are α/β hydrolase-like, contain a core domain between residues 26-200 and 326-396 (see Figure 1 for mouse LIPA), and a central β -sheet composed of 8 strands, designated as β 1 – β 8, and 6 α -helices, designated as α 1, α A, α B1/B2, α C1/C2, α E and α F, with 3 helices on each side of the central β -sheet. In addition, a ‘Cap’ domain is described for human LIPF and LIPA with 8 helices (designated as α e1- α e8) within residues 203-329 for human LIPA (Roussel *et al.*, 1999). This domain may serve as a ‘lid’ for the active site Ser174, restricting access to the aqueous environment but enabling cholesteryl ester and other substrate entry when the ‘lid’ opens. All of these secondary structures have been retained for the mammalian acid lipases examined however these are based on predictions and may not reflect fully structures *in vivo*.

The predicted tertiary structures for mouse LIPF, LIPK, LIPA, LIPM, LIPN and LIPO1 were sufficiently similar to the previously reported human LIPF (gastric acid lipase) and the predicted human LIPA structures (Roussel *et al.*, 1999) (Figure 2) to enable predictions of these tertiary structures which were based on incomplete sequences for these enzymes (residues 22-393 for mouse LIPA). The predictions observed suggest that the secondary and tertiary structures for human LIPF and LIPA resemble those for each of the six mouse acid lipase proteins examined, reflecting conservation of the major structural features for these enzymes.

Predicted Gene Locations and Exonic Structures for Mammalian Acid Lipase Genes.

Table 1 summarizes the predicted locations for mammalian acid lipase genes based upon BLAT interrogations of several mammalian genomes using the reported sequences for human and mouse acid lipases, LIPA (Anderson *et al.*, 1994; Ameis *et al.*, 1994), LIPF (Bodmer *et al.*, 1987; Lohse *et al.*, 1997), LIPJ, LIPK, LIPM and LIPN (Deloukas *et al.*, 2004; Toulza *et al.*, 2007), and the predicted sequences for rat acid lipases (see Table 1 for sources) and the UC Santa Cruz Genome Browser (Kent *et al.* 2003). The mammalian acid lipase genes were located in a gene cluster in each case, although the gene order underwent changes for different species, including an addition of one (rat) or four (mouse) acid lipase genes, designated as LIPO (*Lipo* for the gene family). Supplementary Table 1 also provides data for other mammalian acid lipases genes, including those predicted for chimpanzee, orangutan, horse, cow, guinea pig and dog genomes. These BLAT interrogations of mammalian genomes with the corresponding acid lipase sequences suggested that the gene cluster was syntenic for chromosomes 10 (human, chimp and orangutan), 9 (rhesus monkey), 19 (mouse), 1 (rat and horse) and 26 (cow and dog). Figure 1 summarizes the predicted exonic start sites for human, mouse and rat acid lipase genes with each having 9 coding exons, in identical or similar positions to those reported for the human acid lipase genes (Deloukis *et al.*, 2004).

Comparative Mouse Acid Lipase Gene Expression and Transcripts.

Figure 3 illustrates the comparative predicted structures of mRNA mouse acid lipase gene transcripts (<http://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/index.html?mouse>) for the major transcript isoform in each case (Theirry-Mieg & Thierry-Mieg, 2006). There were 9 introns present for the mRNA transcripts for all mouse acid lipase genes with the exception of *Lipo1*, which contained 10 introns, and 9 coding exons. Mouse *Lipa* and *Lipo1* transcripts were encoded on the minus strand whereas other mouse acid lipases were encoded on the positive strand (Table 1; Figure 3). With the exception of mouse *Lipn* transcripts, mouse acid lipase transcripts contained extended 3'-noncoding sequences.

Figure 4 presents ‘heat maps’ showing comparative gene expression for various mouse tissues obtained from GNF Expression Atlas Data using the U74a (*Lipa*), GNF1N (*Lipf*, *Lipk*, *Lipn* and *Lipo*) and U74b (*Lipm*) mouse chips (<http://genome.ucsc.edu>; <http://biogps.gnf.org>). These data supported a broad tissue expression for mouse *Lipa* (Du et al., 1996); mouse *Lipf* expression in tissues associated with digestion, including pancreas, stomach and salivary gland (Bodmer et al., 1987; and *Lipk*, *Lipm* and *Lipn* expression at higher levels particularly in epidermal tissues, but also in tongue, trachea, liver and kidney (*Lipk*), trachea, bone marrow and eye (*Lipm*) and in liver and kidney (*Lipn*). In contrast, *Lipo* expression (data available only for the *Lipo1* gene) showed higher levels only in the salivary gland. Mouse acid lipase gene expression levels were compared with the expression for average mouse gene (see Table 1) (Theiry-Mieg & Thierry-Mieg, 2006) (<http://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/index.html?mouse>). Average or higher expression levels were observed for mouse *Lipa*, *Lipf* and *Lipo1* acid lipase genes, while *Lipj*, *Lipk*, *Lipm* and *Lipn* showing lower than average expression in the mouse. Similar results were observed for human acid lipase genes with *LIPA* and *LIPF* having much higher levels of expression than the average human gene and the other acid lipase genes, *LIPJ*, *LIPK*, *LIPM* and *LIPN*.

Sequence Identities and Phylogeny of Mammalian Acid Lipases

Table 2 summarizes the percentages of identity for human and mouse acid lipase family sequences (and the rat *LIPO* sequence) which are $\geq 74\%$ identical in comparison with the 44-62% identities observed comparing sequence identities between acid lipase families. This supports a proposal for at least 7 mammalian acid lipase gene families, namely *LIPA*, *LIPF*, *LIPJ*, *LIPK*, *LIPM*, *LIPN* and *LIPO* (designated as *Lipo* for mouse and rat genes for consistency with other rodent acid lipase genes).

Phylogenetic trees (Figure 5) were constructed from alignments of mammalian acid lipase-like amino acid sequences with the frog (*Xenopus tropicalis*) *LIPA* sequence. The dendrogram showed clustering of the sequences into 7 mammalian acid lipase gene family groups. This is consistent with these gene families being present throughout mammalian evolution and of an origin of more than ~100 million years ago, which corresponds to the time of appearance for the eutherian mammalian common ancestor (Woodburne *et al.*, 2003; Donoghue & Benton, 2007). Figure 5 also shows the number of times a clade (sequences common to a node or branch) occurred in the bootstrap analyses with replicate values of 90 or more (which are highly significant) for the 100 replicates undertaken in each case. Of particular interest are the nodes demonstrating highly significant separations for each of the mammalian acid lipase gene family sequences (*LIPA*, *LIPF*, *LIPJ*, *LIPK*, *LIPM*, *LIPN* and *LIPO*) sequences during mammalian evolution, which supports the separate family status for each of

these genes. There were however species differences in the distribution of these mammalian gene families, with *LIPJ* apparently absent in rodents (mouse, rat and guinea pig), while the *Lipo* gene family was found only in mouse and rat genomes among the mammalian species studied (Table 1; Figure 5). The highly significant clustering of the mammalian *LIPA* clade with the single frog acid lipase sequence reported (designated as frog *LIPA*) (Table 1) suggests that *LIPA* may have served as a primordial gene for subsequent gene duplication events generating the 7 families of mammalian acid lipases. The sequence and timing however for these proposed acid lipase gene duplication events remain to be determined.

Conclusions

The results of this present study support previous studies for at least 6 mammalian acid lipase genes and encoded acid lipases, namely *LIPA* (encoding lysosomal lipase), *LIPF* (encoding gastric lipase), *LIPJ* (encoding a human testis lipase), and *LIPK*, *LIPM* and *LIPN* (encoding epidermal lipases). This report also reports evidence for a new acid lipase gene family in mouse and rat (designated as *Lipo*), for which the mouse genome contains 4 *Lipo*-like genes, designated as *Lipo1*, *Lipo2*, *Lipo3* and *Lipo4*, whereas the rat genome contains a single *Lipo* gene. All of these mammalian acid lipase sequences share key conserved sequences and predicted secondary and tertiary structures that have been reported for human *LIPJ* and *LIPA* (Roussel *et al.*, 1999), including active site and catalytic transition state supporting residues, as well as disulfide bond forming cysteine residues. A specific N-glycosylation site involved in the localization of mammalian *LIPA* within lysosomes was also conserved within mammalian *LIPA* sequences. Comparative gene expression data showed that human and mouse *LIPA* and *LIPF* genes are expressed at higher levels than those for the average gene (as defined by Theirry-Mieg & Thierry-Mieg, 2006; <http://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/index.html?mouse>) . This is consistent with key metabolic roles for these enzymes in lysosomal cholesterol ester and triglyceride metabolism and in gastric triglyceride metabolism, respectively. A high level of expression for mouse *Lipo1* was also observed in the salivary gland, which may indicate a supporting role for this acid lipase in triglyceride hydrolysis, either during mastication of food or the subsequent digestion in the stomach . Phylogeny studies using several mammalian acid lipases (human, chimp, orangutan, mouse, rat, guinea pig, horse, cow and dog) indicated that these genes have apparently appeared prior to the eutherian common ancestor more than 100 million years ago, and may have evolved from one or more vertebrate acid lipase gene common ancestors, which include the vertebrate *LIPA* gene.

Acknowledgements:

This project was supported by NIH Grants P01 HL028972 and P51 RR013986. In addition, this investigation was conducted in facilities constructed with support from Research Facilities Improvement Program Grant Numbers 1 C06 RR13556, 1 C06 RR15456, 1 C06 RR017515. We gratefully acknowledge the expert assistance of Dr Bharet Patel of Griffith University, Brisbane Australia with the phylogenetic analyses. We also acknowledge the expert advice of Ms Lois Maltais (Mouse Genome Nomenclature) of the Jackson Laboratory and Dr Stan Lauderkind (Rat Genome Database) of the Medical College of Wisconsin regarding the nomenclature for the new acid lipase gene (*Lipo*) described in this paper.

REFERENCES

- Altschul, F., Vyas, V., Cornfield, A., Goodin, S., Ravikumar, T.S., Rubin, E.H., Gupta, E., 1990. Basic local alignment search tool. *J. Mol. Biol.* 215, 403-410.
- Ameis, D., Merkel, M., Eckerskorn, C., Greten, H., 1994. Purification, characterization and molecular cloning of human hepatic lysosomal acid lipase, *Eur. J. Biochem.*, 219, 905–914.
- Anderson, R.A., Sando, G.N., 1991. Cloning and expression of cDNA encoding human lysosomal acid lipase/cholesteryl ester hydrolase: similarities to gastric and lingual lipases, *J. Biol. Chem.*, 266, 22479-22484.
- Anderson, R.A., Byrum, R.S., Coates, P.M., Sando, G.N., Mutations at the lysosomal acid cholesteryl ester hydrolase gene locus in Wolman disease, 1994. *Proc. Natl. Acad. Sci. USA* 91, 2718-2722.
- Assmann, G., Krauss, R.M., Frederickson, D.S., Levy, R.I., 1973. Positional specificity of triglyceride lipases in post-heparin plasma 1973. *J. Biol. Chem.* 240, 7184-7190.
- Baranski, T.J., Cantor, A.B., Kornfield, S., 1991. Mapping and molecular modelling of a recognition domain for lysosomal targeting. *J. Biol. Chem.* 266, 23365-23372.
- Bodmer, M.W., Angal, S., Yarranton, G.T., Harris, T.J., Lyons, A., King, D.J., Pieroni, G., Riviere, C., Verger, R., Lowe, P.A., 1987. Molecular cloning of a human gastric lipase and expression of the enzyme in yeast. *Biochim. Biophys. Acta* 909, 237-244.
- Bovine Genome Project., 2008. <http://hgsc.bcm.tmc.edu/projects/bovine>
- Carninci P., Kasukawa T., Katayama S., Gough J., Frith M.C., Maeda N., Oyama R., Ravasi T., Lenhard B., Wells C., Kodzius R., Shimokawa K., Bajic V.B., Brenner S.E., Batalov S., Forrest A.R., Zavolan M., Davis M.J. Hayashizaki Y., 2005. The transcriptional landscape of the mammalian genome. *Science* 309,1559-1563.
- Chenna, R., Sugawara, H., Koike, T, Lopez, R, Gibson, T.J., Higgins, D.J., Thompson, J.D., 2003. Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.* 31, 3497-3500.
- Chimpanzee Sequencing and Analysis Consortium., 2005. Initial sequence of the chimpanzee genome and comparison with the human genome. *Nature* 437, 69-87.
- Deloukas, P., Earthrowl, M.E., Grafham, D.V., Rubenfield, M., French, L., Steward, C.A., Sims, S.K., Jones, M.C., Searle, S., Scott, C., Howe, K., Hunt, S.E., Andrews, T.D., Gilbert, J.G., Swarbreck, D., Ashurst, J.L., Taylor, A., Battles, J., Bird, C.P., Ainscough, R., Almeida, J.P., Ashwell, R.I., Ambrose, K.D., Babbage, A.K., Bagguley CL, Bailey J, Banerjee R, Bates K, Beasley H, Bray-Allen S, Brown AJ, Brown JY, Burford DC, Burrill, W., Burton, J., Cahill, P., Camire, D., Carter, N.P., Chapman, J.C., Clark, S.Y., Clarke, G., Clee, C.M., Clegg, S., Corby, N., Coulson, A., Dhami, P., Dutta, I., Dunn, M., Faulkner, L., Frankish, A., Frankland, J.A., Garner, P., Garnett, J., Gribble, S., Griffiths, C., Grocock, R., Gustafson, E., Hammond, S., Harley, J.L., Hart, E., Heath, P.D., Ho, T.P., Hopkins, B., Horne, J., Howden, .PJ., Huckle, E., Hynds, C., Johnson, C., Johnson, D., Kana, A., Kay, M., Kimberley, A.M., Kershaw, J.K., Kokkinaki, M., Laird, G.K., Lawlor, S., Lee, H.M/

Leongamornlert, D.A., Laird, G., Lloyd, C., Lloyd, D.M., Loveland, J., Lovell, J., McLaren, S., McLay, K.E., McMurray, A., Mashreghi-Mohammadi, M., Matthews, L., Milne, S., Nickerson, T., Nguyen, M., Overton-Larty, E., Palmer, S.A., Pearce, A.V., Peck, A.I., Pelan, S., Phillimore, B., Porter, K., Rice, C.M., Rogosin, A., Ross, M.T., Sarafidou, T., Sehra, H.K., Shownkeen, R., Skuce, C.D., Smith, M., Standring, L., Sycamore, N., Tester, J., Thorpe, A., Torcasso, W., Tracey, A., Tromans, A., Tsolas, J., Wall, M., Walsh, J., Wang, H., Weinstock, K., West, A.P., Willey, D.L., Whitehead, S.L., Wilming, L., Wray, P.W., Young, L., Chen, Y., Lovering, R.C., Moschonas, N.K., Siebert, R., Fechtel, K., Bentley, D., Durbin, R., Hubbard, T., Doucette-Stamm, L., Beck, S., Smith, D.R., Rogers, J. 2004. The DNA sequence and comparative analysis of human chromosome 10, *Nature*, 429, 375-381.

Dog Genome Project, 2005. <http://www.broadinstitute.org/mammals/dog>.

Donoghue, P.C.J., Benton, M.J., 2007. Rocks and clocks: calibrating the tree of life using fossils and molecules. *Trends Genet.* 22, 424-431.

Du, H., Witte, D.P., Grablowski, G.A., 1996. Tissue and cellular specific expression of murine lysosomal acid lipase mRNA and protein, *J. Lipid Res.*, 37, 937-949.

Emmanuelsson, O., Brunak, S., von Heijne, G., Nielson, H., 2007. Locating proteins in the cell using TargetP, SignalP and related tools. *Nature Protocols* 2, 953-971.

Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783-791.

Gibbs, R.A., Rogers, J., Katze, M.G., Bumgarner, R., Weinstock, G.M., Mardis, E.R., Remington, K.A., Strausberg, R.L., 2007. Evolutionary and biomedical insights from the rhesus macaque genome. *Science* 316, 222-34.

Goldstein, J.L., Dana, S.E., Faust, J.R., Beaudet, A.L., Brown, M.S., 1975. Role of lysosomal acid lipase in the metabolism of plasma low density lipoprotein. Observations in cultured fibroblasts from a patient with cholesteryl ester storage disease, *J. Biol. Chem.* 250, 8487-8495.

Guex, N., Peitsch, M.C., 1997. SWISS-MODEL and the Swiss-PdbViewer: An environment for comparative protein modelling. *Electrophoresis* 18, 2714-2723.

Guinea Pig Genome Sequencing Project, 2008. <http://www.broadinstitute.org/science/projects/mammals-models/guinea-pig/guinea-pig>.

Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT, *Nucleic Acids Symposium Series*, 41, 95-98.

Hirata, K., Dichek, H.L., Cioffi, J.A., Chooi, S.Y., Leeper, N.J., Quintana, L., Kronmal, G.S., Cooper, A.D., Quertermous, T., 1999. Cloning of a unique lipase from endothelial cells extends the lipase gene family, *J. Biol. Chem.* 274, 14170-14175.

Hoeg, J.M., Demosky, S.J., Pescovitz, O.H., Brewer, H.B., 1984. Cholesteryl ester storage disease and Wolman disease: phenotypic variants of lysosomal acid cholesteryl ester hydrolase deficiency, *Amer. J. Human Genet.*, 36, 1190-1203.

Horse Genome Project., 2008. <http://www.uky.edu/Ag/Horsemap/>

Horton, P., Nakai, K., 1997. Better prediction of cellular localization sites with the k nearest neighbors classifier. *Proc. Int. Conf. Intell. Syst. Mol. Biol.* 5, 147-152.

International Human Genome Sequencing Consortium., 2001. Initial sequencing and analysis of the human genome. *Nature* 409, 860-921.

- Jaye, M., Lynch, K.J., Krawiec, J., Marchadier, D., Maugeais, C., Doan, K., South, V., Amin, D., Perrone, M., & Daniel J. Rader, D.J., 1999. A novel endothelial-derived lipase that modulates HDL metabolism, *Nature Genetics* 21, 424-428.
- Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., Haussler, D., 2003. The human genome browser at UCSC. *Genome Res.* 12, 994-1006.
- Kimura, M., 1983. *The Neutral Theory of Molecular Evolution*, Cambridge University Press, Cambridge Mass.
- Kopp, J., Schwede, T., 2004. The SWISS-MODEL Repository of annotated three-dimensional protein structure homology models. *Nucleic Acids Res.* 32, D230-D234.
- Lohse, P., Chahrokh-Zadeh, S., Lohse, P., Seidel, D., 1997. Human lysosomal acid lipase/cholesteryl ester hydrolase and human gastric lipase: identification of the catalytically active serine, aspartic acid, and histidine residues. *J. Lipid Res.* 38, 892-903.
- Lowe M.E., Rosenblum J.L., Strauss A.W., 1989. Cloning and characterization of human pancreatic lipase cDNA. *J. Biol. Chem.* 264, 20042-20048.
- Martin, G.A., Busch, S.J., Meredith, G.D., Cardin, A.D., Blankenship, D.T., Mao, S.J.T., Reichtin, A.E., Woods, C.W., Racke, M.M., Schafer, M.P., Fitzgerald, M.C., Burke, D.M., Flanagan, M.A., Jackson, R.L., 1998. Isolation and cDNA sequence of human postheparin plasma hepatic triglyceride lipase, *J. Biol. Chem.* 263, 10907-10914.
- McGuffin, L.J., Bryson, K., Jones, D.T., 2000. The PSIPRED protein structure prediction server. *Bioinformatics* 16, 404-405.
- Mouse Genome Sequencing Consortium, 2002. Initial sequencing and comparative analysis of the mouse genome. *Nature* 420, 520-562.
- Orangutan Genome Project, 2007. http://genome.wustl.edu/genomes/view/pongo_abelii/
- Patrick, A.D., Lake, B.D., 1969. Deficiency of an acid lipase in Wolman's disease. *Nature* 222, 1067-1068.
- Rat Genome Sequencing Project Consortium., 2004. Genome sequence of the Brown Norway rat yields insights into mammalian evolution. *Nature* 428, 493-521.
- Roussel, A., Canaan, S., Egloff, M-P., Riviere, M., Dupuis, L., Verger, R., Cambillau, C., 1999. Crystal structure of human gastric lipase and model of lysosomal acid lipase, two lipolytic enzymes of medical interest, *J. Biol. Chem.* 274, 16995-17002.
- Saitou, N., Nei, N., 1987. The neighbour-joining method: a new method for reconstructing phylogenetic trees, *Mol. Biol. Evol.* 4, 406-426.
- Sleat, D.E., Zheng, H., Qian M, Lobel, P., 2006. Identification of sites of mannose 6-phosphorylation on lysosomal proteins, *Mol. Cell Proteomics* 5, 686-701.
- The MGC Project Team., 2004. The status, quality and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection. *Genome Res.* 14, 2121-2127.
- Thierry-Mieg, D., Thierry-Mieg, J., 2006. AceView: A comprehensive cDNA-supported gene and transcripts annotation. *Genome Biology* 7, S12 <http://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/index.html?human>
- Toulza, E., Mattiuzzo, N.R., Galliano, M.F., Jonca, N., Dossat, C., Jacob, D., de Daruvar, A., Wincker, P., Serre, G., Guerrin, M., 2007. Large-scale identification of human genes implicated in epidermal barrier function, *Genome Biol.* 8, R107.1-R107.23.

Van De Peer, Y., de Wachter, R., 1994. TreeCon for Windows: a software package for the construction and drawing of evolutionary trees for the Microsoft Windows environment, Comp. App. Science 10, 569-570.

Wang, F., Wang, W., Wahala, K., Adlercreutz, H., Ikonen, E., Tikkanen, M.J., 2008. Role of lysosomal acid lipase in the intracellular metabolism of LDL-transported dehydroepiandrosterone-fatty acyl esters, Amer. J. Physiol. Endocrinol. Metab., 295, E1455-61.

Wion, K.L., Kirchgessner, T.G., Lusi, A.G., Schotz, M.C., Lawn, R.M., 1987. Human lipoprotein lipase complementary DNA sequence, Science 235, 1638-1641.

Woodburne MO, Rich TH, Springer MS (2003) The evolution of tribospheny and the antiquity of mammalian clades. Mol Phylogenet Evol 28: 360-385.

Xenopus tropicalis Genome Project, 2005. <http://genome.jgi-psf.org/Xentr4/Xentr4.home.html>.

Figure 1: Amino acid sequence alignments for mouse (*Mus musculus*) acid lipase sequences.

See Table 1 for sources of acid lipase sequences; * identical residues; . 1 or 2 conservative substitutions; - 1 or 2 non-conservative substitutions; residues involved in processing at N-terminus (signal peptide); potential N-glycosylation sites including residues **NKT** (161-163) which serves as a lysosomal targeting sequence **^LTS**; active site residues **Ser174**; **Asp345**; and **His374**; disulfide bond C residues **Cys248** and **Cys257** for human LIPA; **helix** (human LIPA) or predicted helix; **Sheet** (human LIPA) or predicted sheet; numbered according to Roussel et al [37]; potential basic amino acid 'patch' for lysosomal targeting at LIPA C-terminus **■**; residues **Leu89** and **Gln175** contribute to the oxyanion 'hole' near active site [37]; and bold **underlined** font shows known or predicted exon junctions.

Figure 2: Predicted tertiary structures for mouse acid lipases.

The predicted structures for mouse LIPF, LIPK, LIPA, LIPM, LIPN and LIPO1 are based on the reported structure for human LIPF (Roussel et al., 1999). Predicted structures were obtained using the SWISS MODEL web site <http://swissmodel.expasy.org/workspace/index.php?>. See Table 1 for sources of these sequences. The rainbow color code describes the 3-D structures from the N- (blue) to C-termini (red color).

Figure 3: Gene structures and major splicing isoforms for mouse *Lipa*, *Lipf*, *Lipk*, *Lipm*, *Lipn* and *Lipol* genes.

From AceView website (Thierry-Mieg and Thierry-Mieg, 2006) <http://www.ncbi.nlm.nih.gov/IEB/Research/AceView/>
Mature isoform variants (designated as a or b isoform) are shown for each mouse gene transcript with capped 5'- and 3'-ends for the predicted mRNA sequences. Scales of base pairs of nucleotide sequences are shown. Flags identify validated endings: cap site on the 5' side, polyadenylation site on the 3' side. Filled flags correspond to frequent events while empty flags have lesser supporting cDNAs (all validated); at the 3' side, black flags are associated to the main AATAAA signal, blue flags to any single letter variant of the major sequence.

Figure 4: Mouse tissue gene expression ‘heat maps’ for acid lipase genes.

Taken from the mouse genome browser (<http://genome.ucsc.edu>) (NCBI37/mm9 2007 assembly) (Kent *et al.* 2003). GNF Expression Atlas 2 data using various expression chips for mouse acid lipase genes *Lipa*, *Lipf*, *Lipk*, *Lipm*, *Lipn* and *Lipo1* (using GenBank ID AI747699) (<http://biogps.gnf.org>). Comparative gene expression levels among mouse tissues: red (high); black, (intermediate); and green (low) expression levels.

Figure 5: Phylogenetic tree of mammalian acid lipase LIPF, LIPJ, LIPK, LIPM, LIPN and LIPO amino acid sequences with frog (*Xenopus tropicalis*) LIP A sequence.

The tree is labeled with the acid lipase gene name and the name of the mammal or frog. Note the major clusters for each of the 7 acid lipase gene families. The gene duplication events generating these distinct gene families (*LIPF*, *LIPJ*, *LIPK*, *LIPM*, *LIPN* and *LIPO*) is proposed to have occurred prior to the eutherian mammalian common ancestor estimated at ~ 100 million years ago (Woodburne *et al.*, 2003). A genetic distance scale is shown. The number of times a node (sequences common to a node or branch) occurred in the bootstrap replicates are shown. Only replicate values of 90 or more which are highly significant are shown. 100 bootstrap replicates were performed in each case. Of particular interest are the nodes (marked with an asterisk*) supporting the significant separation of each of the acid lipase gene families examined.

Acid Lipase Gene	Chromosome	Gene	Coding	Subunit	Amino	pI	GenBank	Vega Gene ID	Ensembl Protein	UNIPROT	Ref Seq ID	Other
(Expression) ¹	Coordinates	Size	Exons	MW	Acids		ID		Prediction	ID	*NCBI Reference	Name
Human	Chromosome 10	bps	Strand								*NCBI Predicted	Other ID
<i>Homo sapiens</i>	Gene order:J-F-K-N-M-A										*BLAT Predicted	
<i>LIPA</i> (4.9)	90,964,568-90,997,385	32,818	9 -ve	45,419	399	6.4	BC012287	OTTHUMG00000018716	ENST00000371829	P38571	NM_001127605	<i>LICH</i>
<i>LIPF</i> (2.5)	90,424,214-90,438,571	14,357	9 +ve	45,238	398	6.8	AK208334	OTTHUMG00000018695	ENST00000238983	P07098	NM_004190	
<i>LIPJ</i> (0.3)	90,346,519-90,366,732	20,213	9 +ve	42,388	366	6.1	BC031219	OTTHUMG00000018691	ENST00000371939	Q5W064	NM_001010939	
<i>LIPK</i> (0.05)	90,474,281-90,502,490	28,210	9 +ve	45,563	399	8.4	EF426482	OTTHUMG00000018693	ENST00000404190	Q5VXJ0	NM_001080518	
<i>LIPM</i> (0.04)	90,552,634-90,570,235	17,602	9 +ve	48,233	423	6.6	EF426484	OTTHUMG00000018698	ENST00000404743	Q5VYY2	NM_001128215	
<i>LIPN</i> (0.05)	90,511,143-90,527,979	16,834	9 +ve	45,534	399	6.4	EF426483	OTTHUMG00000018694	ENST00000404459	Q5VXI9	NM_001102469	
Mouse	Chromosome 19											MGI ID
<i>Mus musculus</i>	Gene order:o4-3-2-1-f-k-n-m-a											
<i>Lipa</i> (3.2)	34,568,473-34,599,332	30,860	9 -ve	45,435	397	8.2	BC058564		ENSMUST00000049572	Q9Z0M5	NM_021460	96789
<i>Lipf</i> (2.5)	34,035,738-34,051,303	15,566	9 +ve	44,637	395	6.4	BC061067	OTTMUSG00000028348	ENSMUST00000025680	Q9CPP7	NM_026334	1914967
<i>Lipk</i> (0.3)	34,082,780-34,122,287	39,508	9 +ve	45,243	398	8.7	BK055815		ENSMUST00000054260	Q8BM14	NM_172837	2679259
<i>Lipm</i> (0.8)	34,175,433-34,197,049	21,617	9 +ve	48,254	422	8.7	BC031933		ENSMUST00000025681	Q8K2A6	NM_023903	1926003
<i>Lipn</i> (0.13)	34,141,848-34,159,408	17,561	9 +ve	45,744	400	6.8	AK154333	OTTMUSG00000029762	ENSMUST00000025682	Q3U4B4	NM_027340	1917416
<i>Lipo1</i> (1.0)	33,851,027-33,861,942	10,916	9 (-ve)	44,668	399	6.1	AI747699	ENSMUSG00000079342	ENSMUST00000112506		*NP_001013792	2147592
<i>Lipo2</i> ³	33,795,278-33,825,842	30,565	9 -ve	44,856	399	6.5	AK139780	OTTMUSG00000033077	ENSMUST00000025694		*XM_994361	3644466
<i>Lipo3</i> ³	33,830,724-33,859,458	28,735	9 -ve	44,695	399	6.5	AK170332	ENSMUSG00000024766	ENSMUST00000112508			
<i>Lipo4</i> ³	33,573,637-33,592,260	18,624	9 -ve	44,771	398	6.3		OTTMUSG00000033071	ENSMUST00000112511		*XM_001477927	3779637
Rat	Chromosome 1											RGD ID
<i>Rattus norvegicus</i>	Gene order:o-f-k-n-m-a											
<i>Lipa</i> (0.5)	238,466,493-238,500,195	33,703	9 -ve	45,079	397	6.3	BC072532		ENSRNOT00000025845	Q64194	NM_012732	3008
<i>Lipf</i> (0.05)	237,841,539-237,856,579	15,041	9 +ve	44,588	395	6.1	X02309	ENSRNOG00000019448	ENSRNOT00000027965	P04634	NM_017341	708441
<i>Lipk</i> (0.01)	237,889,400-237,910,233	20,834	9 +ve	45,337	397	8.7		ENSRNOG00000019409	ENSRNOT00000026299		NM_001106374	1309724
<i>Lipm</i> ³	237,964,915-237,983,386	18,472	9 +ve	48,103	422	8.7		ENSRNOG00000019301	ENSRNOT00000026219		*XP_001079892	1304912
<i>Lipo</i> (0.03)	237,932,633-237,948,802	16,169	9 +ve	45,628	398	8.2		ENSRNOG00000019395	ENSRNOT00000026242		*XP_574655	1560354
<i>Lipo</i> ³	237,642,349-237,653,919	11,571	9 -ve	44,441	397	5.7		ENSRNOG00000025444	ENSRNOT00000035013		*XP_220070	1565682
Clawed Frog												
<i>Xenopus tropicalis</i>	Scaffold 150											Xenbase
<i>LIPA</i> ³	1,826,750-1,838,449	11,700	9 +ve	45,454	404	5.8	BC106353			Q3KQ76	NM_001015847	952976

Table 1: Mammalian acid lipase (*LIPA*, *LIPF*, *LIPJ*, *LIPK*, *LIPM*, *LIPN* and *LIPO*) and frog (*Xenopus tropicalis*) *LIPA* genes and proteins.

¹RefSeq: the reference amino acid sequence; ²predicted Ensembl amino acid sequence; and ³BLAT predicted amino acid sequences are shown (see <http://www.ncbi.nlm.nih.gov>); GenBank IDs are derived from NCBI sources

<http://www.ncbi.nlm.nih.gov/genbank/>; Vega gene ID was derived from the Vertebrate Genome Annotation (VEGA) database <http://vega.sanger.ac.uk>; Ensembl ID was derived from Ensembl genome database <http://www.ensembl.org>;

⁴relative level of gene expression was obtained from the AceView database for human, mouse and rat gene transcripts <http://www.ncbi.nlm.nih.gov/IEB/Research/AceView/index.html?mouse>; ⁵result not available; UNIPROT refers to

UniprotKB/Swiss-Prot IDs for individual acid lipases (see <http://kr.expasy.org>); bps refers to base pairs of nucleotide sequences; pI refers to theoretical isoelectric points; the number of coding exons are listed. Sources for acid lipase sequences were provided by the above sources (see Table 1).

Acid Lipase Gene	Human LIPA	Mouse LIPA	Human LIPF	Mouse LIPF	Human LIPJ	Human LIPK	Mouse LIPK	Human LIPM	Mouse LIPM	Human LIPN	Mouse LIPN	Mouse LIPD1	Rat LIPO
Human LIPA	100	74	57	55	53	55	52	59	59	53	55	47	47
Mouse LIPA	74	100	50	51	50	51	48	54	54	49	50	44	45
Human LIPF	57	50	100	76	57	62	60	52	53	50	51	50	51
Mouse LIPF	55	51	76	100	56	61	60	52	54	50	51	50	51
Human LIPJ	53	50	57	56	100	59	54	52	52	48	48	46	46
Human LIPK	55	51	62	61	59	100	81	52	52	50	52	48	50
Mouse LIPK	52	48	60	60	54	81	100	50	52	46	47	48	50
Human LIPM	59	54	52	52	52	52	50	100	87	52	52	44	46
Mouse LIPM	59	54	53	54	52	52	52	87	100	52	52	45	46
Human LIPN	53	49	50	50	48	50	46	52	52	100	81	41	46
Mouse LIPN	55	50	51	51	48	52	47	52	52	81	100	41	47
Mouse LIPD1	47	44	50	50	46	48	48	44	45	41	41	100	81
Rat LIPO	47	45	51	51	46	50	50	46	46	46	47	81	100
	Mouse LIPD1	Mouse LIPD2	Mouse LIPD3	Mouse LIPD4									
Mouse LIPD1	100	76	77	78									
Mouse LIPD2	76	100	96	91									
Mouse LIPD3	77	96	100	90									
Mouse LIPD4	78	91	90	100									

Table 2: Percentage identities for human, mouse and rat acid lipase amino acid sequences

Numbers show the percentage of amino acid sequence identities. Numbers in **bold** show higher sequence identities for mammalian acid lipase gene family members.

Mammal	Acid	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7	Site 8	Site 9	Site 10	Site 11	Potential	High	Lower
	Lipase												N-glycosylation	probability	Probability
													sites	sites (>0.75)	Sites (0.5-0.74)
														(confirmed)*	
Human	LIPA	36NYS	72NHS	101NSS	161NKT		273NMS		321NQS				6	4	2
Mouse	LIPA	34NYT		99NSS	159NKT		271NMS		319NQS				5	3	2
Rat	LIPA	34NYT		99NSS	159NKT		271NMS		319NQS				5	3	2
Human	LIPF	34NIS*		99NNS*		185NPS	271NTS*			327NVT*			5	4 confirmed*	0
Mouse	LIPF	33NYS		98NNS			270NYS						3	2	1
Rat	LIPF	33NIS	68NNS	98NNS		184NPT	270NYS						5	1	1
Human	LIPJ*	2NIS		68NNS			240NMS	268NST	288NQT	296NMT			6	2	2
Human	LIPK	34NIS		99NNS			271NMS			327NIT			4	2	1
Mouse	LIPK	33NIS		98NNS			270NMS			326NIS			4	1	2
Rat	LIPK	32NIS		97NNS			269NMS			325NIS			4	1	2
Human	LIPM	48NIS		113NNS			285NMS						3	0	2
Mouse	LIPM	48NYS		113NNS			285NMS						3	1	1
Rat	LIPM	48NYS		113NNS			285NMS						3	1	1
Human	LIPN	35NTS		100NGS	160NKT		272NQS		320NQS			413NLS	6	1	4
Mouse	LIPN	37NAS		102NGS	162NKT		274NMS		322NQS				5	0	4
Rat	LIPN	35NAS		100NGS	160NKT		272NMS		320NQS				5	0	4
Mouse	LIP01	32NYS	70NSS				268NTS		316NQT		357NLT		5	2	2
Mouse	LIP02	32NYS					268NKS		316NQT		357NLT		4	3	1
Mouse	LIP03	32NYS	70NSS			183NQT	268NKS		316NQT		357NLT		6	3	2
Mouse	LIP04	32NYS	70NSS				268NMS		316NQT		357NLT		5	2	3
Rat	LIP0	32NYS	70NSS		157NTT		268NTS		316NQT				5	1	3

Table 3: Predicted N-glycosylation sites for mammalian acid lipases.

Numbers refer to amino acids in the acid sequences, including N-asparagine; K-lysine; I-isoleucine; M-methionine; H-histidine; S-serine; R-arginine; T-threonine; Q-glutamine; and V-valine. Note that there are 11 potential sites identified, including 4 confirmed sites for human LIPF and a lysosomal targeting site (site 4) for human LIPA. High (in **bold**) and lower probability N-glycosylation sites were identified using the NetNGlyc 1.0 web server (<http://www.cbs.dtu.dk/services/NetNGlyc/>).

Acid Lipase Gene	Chromosome	Gene	Coding	Subunit	Amino	pI	Ensembl Protein	UNIPROT	Ref Seq ID
	Coordinates	Size	Exons	MW	Acids		Prediction	ID	¹ NCBI Reference
Chimpanzee <i>Pons troglodytes</i>	Chromosome 10								² NCBI Predicted
	Gene order: <i>J-F-K-N-M-A</i>								³ BLAT Predicted
<i>LIPA</i>	89,482,403-89,515,834	33,432	9 -ve	45,419	399	6.4	ENSPTRT00000005165		XP_521552 ¹
<i>LIPF</i>	88,917,921-88,929,291	11,371	9 +ve	45,300	398	6.6	ENSPTRT00000005140		XP_521545 ¹
<i>LIPJ</i>	88,837,829-88,854,222	16,394	9 +ve	42,383	366	6.1	ENSPTRT000000057646		XP_507899 ¹
<i>LIPK</i>	88,976,758-89,008,956	32,199	9 +ve	45,544	399	8.1	ENSPTRT000000057645		XP_001138639 ¹
<i>LIPM</i>	89,063,462-89,081,035	17,574	9 +ve	48,277	423	6.6	ENSPTRT00000005143		XP_001138977 ¹
<i>LIPN</i>	89,017,601-89,034,648	17,048	9 +ve	45,559	398	6.4	ENSPTRT000000057644		XP_521546 ¹
Orangutan <i>Pongo abelii</i>	Chromosome 10								
	Gene order: <i>N-K-F-J-M-A</i>								
<i>LIPA</i>	87,901,737-87,934,555	32,819	9 (-ve)	45,452	399	6.4	ENSPPYT00000002953		³ chr10.569.1
<i>LIPF</i>	85,710,697-85,722,068	11,372	9 (-ve)	45,240	398	7.2	ENSPPYT00000002935		³ chr10.550.1
<i>LIPJ</i>	85,796,879-85,816,972	20,094	9 (-ve)	42,239	366	6.1			⁵
<i>LIPK</i>	85,629,181-85,658,714	29,534	9 (-ve)	45,550	399	8.4	ENSPPYT00000002934		³ chr10.549.1
<i>LIPM</i>	87,474,905-87,491,040	16,136	9 (+ve)	48,275	423	6.6	⁵ ENSPPYT00000002946		³ chr10.562.1
<i>LIPN</i>	85,606,251-85,620,533	14,283	9 (-ve)	45,608	398	6.5	⁵ ENSPPYT00000002933		⁵
Rhesus Monkey <i>Macaca mulatta</i>	Chromosome 9								
	Gene order: <i>J-F-K-N-M-A</i>								
<i>LIPA</i>	88,805,255-88,839,453	34,199	9 (-ve)	45,480	399	6.4			² XP_001085160
<i>LIPF</i>	88,247,613-88,258,640	11,028	9 (+ve)	44,318	392	5.9			² XP_001082289
<i>LIPJ</i>	88,180,349-88,199,904	19,556	9 (+ve)	46,385	391	6.3			² XP_001082167
<i>LIPK</i>	88,298,630-88,329,151	30,522	9 (+ve)	46,506	399	8.9			² XP_001082335
<i>LIPM</i>	88,379,173-88,396,837	17,665	9 (+ve)	48,303	423	6.5			³
<i>LIPN</i>	88,337,780-88,353,937	16,158	9 (+ve)	45,651	398	6.4			³
Cow <i>Bos taurus</i>	Chromosome 26								
	Gene order: <i>J-F-K-M-A</i> ⁴ [N]								
<i>LIPA</i>	11,348,494-11,390,918	42,425	9 (-ve)	45,671	399	7.2	BC146075 (GenBank ID)		² NP_001096793
<i>LIPF</i>	10,779,450-10,793,506	14,047	9 (+ve)	45,194	397	6.5	ENSBTAT00000008453		NM_174103
<i>LIPJ</i>	10,632,628-10,651,794	19,167	9 (+ve)	45,574	398	8.3	ENSBTAT00000049487		² XP_593347
<i>LIPK</i>	10,849,060-10,866,363	17,304	9 (+ve)	45,453	396	8.8	ENSBTAT00000017773		² XP_584217
<i>LIPM</i>	10,937,349-10,957,185	19,837	9 (+ve)	48,505	423	8.3	ENSBTAT00000017778		² XP_595458
<i>LIPN</i>	⁴ 558-19,395 Unknown	34,223	⁴	⁴	⁴	⁴	⁴ ENSBTAT00000034704		² XP_602277
Horse <i>Equus caballus</i>	Chromosome 1								
	Gene order: <i>A-M-N-K-F-J</i>								
<i>LIPA</i>	39,069,159-39,103,381	34,223	9 (+ve)	45,327	409	7.3	ENSECAT00000011488		² XP_001503012
<i>LIPF</i>	39,769,658-39,780,294	10,637	9 (+ve)	45,214	398	7.7	ENSECAT00000010632		² XP_001503148
<i>LIPJ</i>	39,936,554-39,961,057	24,504	9 (-ve)	45,756	398	7.2	ENSECAT00000015158		² XP_001501589
<i>LIPK</i>	39,588,872-39,619,886	31,015	9 (-ve)	45,666	399	8.8	ENSECAT00000007415		² XP_001503125
<i>LIPM</i>	39,498,465-39,519,901	21,437	9 (-ve)	48,099	423	6.8	ENSECAT00000024102		² XP_001501533
<i>LIPN</i>	39,546,866-39,573,950	27,085	9 (-ve)	45,649	400	6.6	ENSECAT00000006452		² XP_001501541
Dog <i>Canis familiaris</i>	Chromosome 26								
	Gene order: <i>J-F-K-N-M-A</i>								
<i>LIPA</i>	41,958,963-41,981,595	22,633	9 (-ve)	45,063	398	6.7	ENSCAFT000000035174		² XP_853280
<i>LIPF</i>	41,541,401-41,555,137	17,737	9 (+ve)	45,103	398	6.3	ENSCAFT00000024834		² NP_001003209
<i>LIPJ</i> [<i>LIPL1</i>]	41,497,356-41,509,775	12,420	9 (+ve)	41,703	365	6.6	ENSCAFT00000036576		² XP_534779
<i>LIPK</i>	41,602,751-41,625,548	22,798	9 (+ve)	46,233	400	8.7	ENSCAFT00000036401		² XP_543591
<i>LIPM</i>	41,677,108-41,690,970	13,863	9 (+ve)	48,797	430	7.7	ENSCAFT00000036170		² XP_853107

LIPN	41,643,262-41,656,521	13,265	9 (+ve)	45,239	400	8.8	ENSCAFT00000036272	³
Guinea Pig	Scaffold 15							
Cavia porcellus	Gene order: F-K-N-M-A [J]							
LIPA	1,783,523-1,816,987	33,465	9 (-ve)	45,053	399	8.8	ENSCPOT00000000968	³ sc15.016.1
LIPF	1,254,927-1,269,927	15,001	9 (+ve)	45,087	398	6.1	ENSCPOT00000007411	³ sc15.009.1
LIPK	1,297,088-1,309,813	12,726	9 (+ve)	45,540	397	8.8	ENSCPOT00000007417	
LIPM	1,360,852-1,377,384	16,533	9 (+ve)	45,061	396	6.5	ENSCPOT00000001668	³ sc15.010.1
LIPN	1,327,922-1,340,779	12,858	9 (+ve)	45,765	398	6.4	ENSCPOT00000007424	

Supplementary Table 1: Other mammalian acid lipase (*LIPA*, *LIPF*, *LIPJ*, *LIPK*, *LIPM*, *LIPN* and *LIPO*) genes and proteins.

¹RefSeq: the reference amino acid sequence; ²predicted Ensembl amino acid sequence; and ³BLAT predicted amino acid sequences are shown (see <http://www.ncbi.nlm.nih.gov>); GenBank IDs are derived from NCBI sources <http://www.ncbi.nlm.nih.gov/genbank/>; Ensembl ID was derived from Ensembl genome database <http://www.ensembl.org>; ⁴unknown chromosomal location for cow LIPN gene; ⁵result not available; bps refers to base pairs of nucleotide sequences; pI refers to theoretical isoelectric points; the number of coding exons are listed. Sources for acid lipase sequences were provided by the above sources (see Table 1).