Title:

Impact of exercise selection on hamstring muscle activation

Authors:

Matthew N. Bourne^{1,2,3}, Morgan D. Williams⁴, David A. Opar⁵, Aiman Al Najjar⁶,
Graham K. Kerr^{1,2}, Anthony J. Shield^{1,2}.

¹School of Exercise and Nutrition Science, Faculty of Health, Queensland University of
Technology, Brisbane, Australia.

²Institute of Health and Biomedical Innovation, Queensland University of Technology,
Brisbane, Australia.

³Queensland Academy of Sport, Centre of Excellence for Applied Sport Science Research,
Brisbane, Australia.

⁴School of Health, Sport and Professional Practice, Faculty of Life Sciences and Education,
University of South Wales, Wales, United Kingdom

⁵School of Exercise Sciences, Australian Catholic University, Melbourne, Australia.

⁶Centre for Advanced Imaging, University of Queensland, Brisbane, Australia.

Corresponding Author:

Dr Anthony Shield

School of Exercise and Nutrition Sciences and the Institute of Health and Biomedical
Innovation,

Queensland University of Technology, Victoria Park Road, Kelvin Grove, 4059,
Brisbane, Queensland, Australia.

Email: aj.shield@qut.edu.au

Ph: +61 7 3138 5829

Fax: +61 7 3138 3980

Key Words

Injury prevention, rehabilitation, physical therapy

ABSTRACT

Objective: To determine the extent to which different strength training exercises selectively activate the commonly injured biceps femoris long head (BF_{LH}) muscle. **Methods:** This two-part observational study recruited 24 recreationally active males. **Part 1** explored the amplitudes and the ratios of lateral to medial hamstring (BF/MH) normalised electromyography (nEMG) during the concentric and eccentric phases of 10 common strength training exercises. **Part 2** used functional magnetic resonance imaging (fMRI) to determine the spatial patterns of hamstring activation during two exercises which i) most selectively, and ii) least selectively activated the BF in part 1. **Results:** Eccentrically, the largest BF/MH nEMG ratio was observed in the 45° hip extension exercise and the lowest was observed in the Nordic hamstring (NHE) and bent-knee bridge exercises. Concentrically, the highest BF/MH nEMG ratio was observed during the lunge and 45° hip extension and the lowest was observed for the leg curl and bent-knee bridge. fMRI revealed a greater BF_{LH} to semitendinosus activation ratio in the 45° hip extension than the NHE ($p < 0.001$). The T2 increase after hip extension for BF_{LH}, semitendinosus and semimembranosus muscles were greater than that for BF_{SH} ($p < 0.001$). During the NHE, the T2 increase was greater for the semitendinosus than for the other hamstrings ($p \leq 0.002$). **Conclusion:** This investigation highlights the non-uniformity of hamstring activation patterns in different tasks and suggests that hip extension exercise more selectively activates the BF_{LH} while the NHE preferentially recruits the semitendinosus. These findings have implications for strength training interventions aimed at preventing hamstring injury.

What are the new findings?

- The hamstrings are activated non-uniformly during hip- and knee-based exercises
- Hip extension exercise more evenly activates the three long heads of the hamstrings and the Nordic hamstring exercise preferentially recruits the semitendinosus (ST)

INTRODUCTION

Paragraph 1 Hamstring strain injuries (HSIs) are commonly experienced by athletes involved in running-based sports. They are the most prevalent injury in track and field,[1] Australian Rules football,[2] and soccer[3] and up to 30% recur within 12 months.[4] Upwards of 80% of HSIs involve the biceps femoris long head (BF_{LH}) muscle[5-7] and most injuries are thought to occur during the late swing phase of high-speed running.[8] During this phase of the gait cycle, the BF_{LH} reaches its peak length and develops maximal force while undergoing a forceful eccentric contraction to decelerate the shank for foot strike,[9] and it is thought that these conditions may at least partly explain its propensity for injury. It has also been reported that prior BF_{LH} injury is associated with a degree of neuromuscular inhibition[10 11] and prolonged atrophy[12], which suggests that current rehabilitation practices do not adequately restore function to this muscle.

Paragraph 2 It has been proposed that hamstring weakness is a risk factor for future strain injury[6 13 14] and interventions aimed at increasing strength, particularly eccentric knee flexor strength, have been effective in reducing HSI rates in several sports.[15-18] However, despite an increased focus on hamstring strength in prophylactic programs,[19] exercise selection is often implemented on the basis of clinical recommendations and assumptions rather than empirical evidence.[20 21] There is currently a very small body of work on the activation patterns of the hamstrings during commonly employed exercises. Studies using functional magnetic resonance imaging (fMRI) have shown that activation differs within and between hamstring muscles during different tasks.[11 22-24] For example, the semitendinosus (ST) appears to be selectively activated during the Nordic hamstring exercise (NHE)[11] and the eccentric prone leg curl,[24] while the semimembranosus (SM) is preferentially recruited during the stiff leg deadlift.[23] Surface electromyography (sEMG) has been used extensively in the analysis of hamstring exercises.[23-26] However, these

studies are sometimes contradictory and are often inconsistent with the results from fMRI.[11 22-25 27] The lack of complete agreement between fMRI and sEMG might reflect the different physiological basis of each technique.[28] Surface EMG amplitude is sensitive to the electrical activity generated by active motor units and is detected by electrodes overlying the skin.[29] This provides valuable information on the neural strategies involved during muscle activation with high temporal resolution, but is prone to cross talk[29] and cannot discriminate between closely approximated segments of muscles[30] such as the medial hamstrings (semimembranosus and semitendinosus). By contrast, fMRI is a relatively new[31] technique which reflects the metabolic activity associated with exercise.[28] Muscle activation is associated with a transient increase in the transverse (T2) relaxation time of tissue water, which can be detected from signal intensity changes in fMR images. These T2 shifts, which, like sEMG, increase in proportion to exercise intensity,[31 32] can be mapped in cross-sectional images of muscles and therefore provide significantly greater spatial clarity than sEMG.[28 30]

Paragraph 3 An improved understanding of the patterns of hamstring muscle activation during common strength training exercises may enable practitioners to make better informed decisions regarding exercise selection in injury prevention and rehabilitation programs. These data may also inform the design of training studies aimed at investigating the chronic adaptations induced by different exercises. The purpose of this two-part study was to determine the extent to which different exercises selectively activate the commonly injured BF_{LH}. Part 1 used sEMG to determine the amplitude and ratio of lateral to medial hamstring activation during 10 commonly employed exercises. Based on these findings, part 2 employed fMRI to map muscle activation during two exercises that appeared to a) most selectively; and b) least selectively activate the BF according to sEMG. We hypothesised that the patterns of hamstring muscle activation would be non-uniform between exercises and, on

the basis of previous work,[23] that more selective activation of the BF_{LH} would be observed during hip-extension exercise.

METHODS

Participants

Paragraph 4 Twenty-four recreationally active male athletes (age, 24.4 ± 3.3 years, height, 181.8 ± 6.1 cm, weight, 85.2 ± 13.4 kg) participated in this study. Eighteen athletes (age, 23.9 ± 3.1 , height, 180.6 ± 5.9 , weight, 86.0 ± 14.8) participated in part 1 and ten athletes (age, 24.6 ± 4.0 , height, 183.5 ± 7.0 , weight, 83.5 ± 8.7) participated in part 2. *A priori* sample size estimates were based on 1) the capacity to detect a 10% difference in the ratio of BF to MH (BF/MH) sEMG amplitude between exercises,[25] and 2) an effect size of 1.0 in the percentage change in T2 relaxation time between muscles,[11] at a power of 0.80 and with $p < 0.05$. Participants were free from soft tissue and orthopaedic injuries to the trunk, hips and lower limbs at the time of testing and had no known history of cardiovascular, metabolic or neurological disorders. Participants had no history of HSI in the previous 18 months and had never suffered an anterior cruciate ligament injury. Prior to testing, all participants completed a cardiovascular screening questionnaire to make sure it was safe for them to perform intense exercise and those who were involved in part 2 also completed a standard MRI screening questionnaire to ensure it was safe for them to enter the magnetic field. All participants provided written, informed consent for this study, which was approved by the Queensland University of Technology Human Research Ethics Committee and the University of Queensland Medical Research Ethics Committee.

Study Design

Paragraph 5 This cross-sectional study involved two parts. In the first we explored the sEMG amplitudes and ratios of BF to medial hamstring (MH) sEMG activity during ten

commonly employed strength training exercises. Based on these findings, part 2 involved an fMRI investigation of two exercises which appeared to a) most selectively, and b) least selectively activate the BF muscle during eccentric contractions.

PART 1

Paragraph 6 On a separate day prior to experimental testing, participants were familiarised with the exercises used in this investigation. All were shown a demonstration of each exercise (Figure 1) and performed several practice repetitions while receiving verbal feedback from the investigators. Once the participant could complete the exercise with appropriate technique, the loads were progressively increased until an approximate 12RM load was determined (unless the exercise was already supramaximal, ie. NHE and glute-ham-raise). On the day of testing, participants reported to the laboratory and were prepared for sEMG measurement. The testing session began with two maximal voluntary isometric contractions (MVICs) for the hamstrings. Subsequently, participants completed a single set of six repetitions of each exercise, each with the predetermined 12RM load, in randomised order. All data were sampled from a randomly selected limb (dominant or non-dominant), which was the exercised limb during all unilateral movements and all testing sessions were supervised by the same investigator (MNB) to ensure consistency of procedures.

Exercise Protocol

Paragraph 7 The 10 exercises were chosen based on a review of the scientific literature[23 25 27]. They included the bilateral and unilateral stiff-leg deadlift, hip hinge, lunge, unilateral bent and straight knee bridges, leg curl, 45° hip extension, glute-ham-raise and the NHE. Unless the exercise was explosive (hip hinge) or supramaximal and eccentric-only (NHE and

glute-ham raise) participants completed both the concentric and eccentric phases of each exercise using a 12-RM load at a constant pace (~2s up and ~2s down).

INSERT FIGURE 1

Electromyography

Paragraph 8 Bipolar pre-gelled Ag/AgCl sEMG electrodes (10mm diameter, 15mm interelectrode distance) (Ambu, BlueSensor N) were used to record electromyographical activity from the BF and MH. The skin of the participants was shaved, lightly abraded and cleaned with alcohol before electrodes were placed on the posterior thigh, midway between the ischial tuberosity and tibial epicondyles. Electrodes were oriented parallel to the line between these two landmarks, as per SENIAM guidelines,[33] and secured with tape to minimise motion artefact. The reference electrode was placed on the ipsilateral head of the fibula. Muscle bellies of the BF and MH were identified via palpation and correct placement was confirmed by observing active external and internal rotation of the knee in 90° of flexion.[10] During all exercise trials, hip and knee joint angles were measured simultaneously with sEMG data using two digital goniometers. The hip sensor's axis of rotation was aligned with the greater trochanter of the femur and the knee sensor was positioned superficial to the lateral femoral epicondyle.

Maximal voluntary contraction

Paragraph 9 Surface EMG activity was recorded during MVICs of the hamstrings using a custom-made device which was fitted with two uniaxial load cells.[34] Participants lay prone with their hips in 0° of flexion and knees fully extended (180°), with their ankles secured in

immoveable yokes and were asked to perform forceful knee flexion while investigators provided strong verbal encouragement. After 1-2 warm-up contractions, participants completed two 3-4s MVICs, with 30-sec of rest separating each attempt. The contraction that elicited the highest average amplitude for the BF and MH was used to represent the maximal EMG amplitude.

Data analysis

Paragraph 10 All sEMG and joint angle data were sampled at 1 kHz through a 16-bit PowerLab 26T AD unit (ADInstruments, New South Wales, Australia) (amplification = 1000; common mode rejection ratio = 110dB) and analysed using LabChart 8.0 (AD Instruments, New South Wales, Australia). Raw sEMG data were filtered using a Bessel filter (frequency bandwidth = 10-500Hz) and then full wave rectified. Joint angle data were used to determine the concentric and eccentric phases of each repetition for each exercise. For each phase, the filtered sEMG signal was normalised to values obtained during MVIC and these normalised sEMG (nEMG) values were averaged across the six repetitions.

Statistical analysis

Paragraph 11 Data were analysed using JMP version 10.02 (SAS Institute Inc, 2012). Descriptive statistics were calculated for mean nEMG amplitudes of BF and MH for the concentric and eccentric phases of each exercise and an activation ratio was determined by dividing the average BF nEMG amplitude by the average MH nEMG amplitude (BF/MH); ratios >1.0 indicated that the BF was more active than the MH muscles. For both the concentric and eccentric phases, repeated measures linear mixed models fitted with the restricted maximum likelihood method were used to determine differences between exercises.

For this analysis, *exercise* was the fixed factor and *participant identity* the random factor. When a significant main effect was observed for *exercise*, post hoc t-tests with Bonferroni corrections were used to identify the source and reported as mean differences with 95% CIs. For these analyses, the Bonferroni adjusted p value was set at <0.002.

PART 2

Paragraph 12 A cross-sectional design was used to map the spatial patterns of hamstring muscle activation during the 45° hip extension and NHE. These exercises were chosen because they a) most selectively (45° hip extension) and b) least selectively (NHE) activated the BF muscle during eccentric contractions according to sEMG. Participants completed two separate exercise sessions, separated by at least six days (14 ± 5 days), with each session involving one of the aforementioned exercises. Functional MRI scans of both thighs were acquired before and immediately after each exercise bout. All testing sessions were supervised by the same investigator (MNB).

Exercise Protocol

Paragraph 13 A depiction of the 45° hip extension and NHE can be found in Figure 1. All exercise was completed using the same equipment as that used in part 1. Participants completed five sets of 10 repetitions of each exercise with one-minute rest intervals between sets. The higher volume of exercise (compared to part 1) was necessary because transient T2 changes reflect fluid shifts associated with glycolysis and have a higher detection threshold than sEMG.[28] All subjects completed 50 repetitions successfully. During the rest periods, participants remained in a seated position (for the hip extension exercise) or lay prone (NHE) to minimise activation of the hamstrings. The 45° hip extension exercise was performed

unilaterally (with the limb chosen randomly) with a starting load corresponding to each participant's approximate 12-RM (median = 10kg; range = 10 to 20kg). However if the participant could no longer complete the exercise with the allocated load, the weight was gradually reduced by increments of 5kg until it could be completed at the desired speed (2sec up and 2 sec down), which was controlled by an electronic metronome. The NHE was performed bilaterally with body weight only. Participants received verbal support from the investigators throughout all exercise sessions to promote maximal effort. All participants were returned to the scanner immediately following the cessation of exercise and post-exercise scans began within $148.6 \pm 24\text{sec}$ (mean $\pm$ SD).

Functional muscle magnetic resonance imaging (fMRI)

Paragraph 14 All fMRI scans were performed using a 3-Tesla (Siemens TrioTim, Germany) imaging system with a spinal coil. The participant was positioned supine in the magnet bore with their knees fully extended and hips in neutral and straps were secured around both limbs to prevent any undesired movement. Consecutive T2-weighted axial images were acquired of both limbs beginning at the level of the iliac crest and finishing distal to the tibial plateau using a 180×256 image matrix. Images were acquired before and immediately after exercise using a Car-Purcel-Meiboom-Gill (CPMG) spin-echo pulse sequence and the following parameters: transverse relaxation time (TR) = 2540ms; echo time (TE) = 8, 16, 24, 32, 40, 48 and 56ms; number of excitations = 1; slice thickness = 10mm; interslice gap = 10mm; field of view = $400 \times 281.3\text{mm}$). The total acquisition time for each scan was 6min 24s. A localiser adjustment (20s) was applied prior to the first sequence of each scan to standardise the field of view and to align collected images between the pre- and post- exercise scans.[11] To minimise any inhomogeneity in MR images caused by dielectric resonances at 3T, a (B1) filter was applied to all scans; this is a post-processing image filter that improves the image signal intensity profile without affecting the image contrast. In addition, to ensure that the

signal intensity profile of T2-weighted images was not disrupted by anomalous fluid shifts, participants were instructed to avoid any exhaustive resistance training of the lower limbs in the week preceding testing, and were seated for a minimum of 15 minutes[23] before pre-exercise imaging.

Paragraph 15 For each exercise session, the T2 relaxation times of each hamstring muscle were measured in T2-weighted images acquired before and after exercise to evaluate the degree of muscle activation during exercise. All fMRI scans were transferred to a Windows computer in the digital imaging and communications in medicine (DICOM) file format. The T2 relaxation times of each hamstring muscle (BF_{LH}, BF_{SH}, ST and SM) were measured in five axial slices, corresponding to 30, 40, 50, 60 and 70% of thigh length; these values were determined relative to the distance between the inferior margin of the ischial tuberosity (0%) and the superior border of the tibial plateau (100%).[11 23] Image analysis software (Sante Dicom Viewer and Editor, Cornell University) was used to measure the signal intensity of each hamstring muscle in the exercised limb in both the pre- and post-exercise scans. The signal intensity was measured manually in each slice using a circular region of interest (ROI)[27] which was placed in a homogenous region of contractile tissue in each muscle belly (avoiding fat, aponeurosis, tendon, bone and blood vessels). The size of each ROI varied (0.2 to 5.6 cm²) based on the cross-sectional area and the amount of homogeneous tissue available in each slice. The signal intensity reflected the mean value of all pixels within the ROI and was measured across seven echo times (8, 16, 24, 32, 40, 48, 56ms). To calculate the T2 relaxation time for each ROI, the signal intensity value at each echo time was fitted to a mono-exponential decay model using a least squares algorithm:

$$[(SI= M \times \exp(\text{echo time} / T2))][23]$$

where SI is the signal intensity at a specific echo time, and M represents the pre-exercise fMRI signal intensity. To assess the extent to which each ROI was activated during exercise, the mean percentage change in T2 was calculated as:

$$[(\text{mean post-exercise T2} / \text{mean pre-exercise T2}) \times 100].$$

To provide a meaningful measure of whole-muscle activation, the percentage change in T2 relaxation time for each hamstring muscle was evaluated using ROIs from all five thigh levels. Previous studies have demonstrated excellent reliability of T2 relaxation time measures with intra-class correlation coefficients ranging from 0.87 to 0.94.[28 35]

Statistical analysis

Paragraph 16 Absolute T2 values before and after each exercise session were reported descriptively as mean $\pm$ SD. Repeated measures linear mixed models fitted with the restricted maximum likelihood (REML) method were used to determine the spatial activation patterns of the hamstring muscles during the 45° hip extension and NHE. The percentage change in T2 relaxation time was compared between each hamstring muscle (BF_{LH}, BF_{SH}, ST and SM) for both exercises. For this analysis, *muscle* was the fixed factor and both *participant identity* and *participant identity x muscle* the random factors. When a significant main effect was detected for *muscle*, post hoc t tests with Bonferroni corrections were used to determine the source; the adjusted alpha was set at $p < 0.008$. Given that the two examined exercises differed in intensity and contraction mode(s), it was not appropriate to directly compare the magnitude of the T2 shifts between exercises.[36] Instead, repeated measures linear mixed models fitted with the REML method were used to determine differences in the ratio of BF to ST (BF_{LH}/ST and BF_{SH}/ST) and SM to ST (SM/ST) percentage change in T2 relaxation time between exercises. For these analyses *exercise* was the fixed factor and *participant identity* the random factor. When a main effect was found for *exercise*, post hoc t tests were again used to

292 determine the source and reported as mean difference (and 95% CI). Alpha was set at $p < 0.05$
293 for these analyses.

294

295

296

297

298

299

300

RESULTS

Levels of hamstring muscle activation

Paragraph 17 Average BF muscle activity ranged from 21.4% (lunge) to 99.3% (unilateral straight knee bridge) MVIC during the concentric phase and 10.7% (hip hinge) to 71.9% (NHE) during the eccentric phase. Average MH muscle activity ranged from 18.1% (lunge) to 120.7% (leg curl) during the concentric phase and 11.6% (hip hinge) to 101.8% (NHE) during the eccentric phase.

Concentric biceps femoris to medial hamstring (BF:MH) activation ratio

Paragraph 18 The concentric BF/MH activation ratio for each exercise can be found in Figure 2a. A significant main effect was detected between exercises ($p < 0.001$) with post hoc t tests showing that the BF/MH ratio was greater during the lunge than the leg curl (mean difference = 0.8, 95% CI = 0.5 to 1.1, $p < 0.001$) and bent-knee bridge (mean difference = 0.7, 95% CI = 0.4 to 1.1, $p < 0.001$). Similarly, the BF/MH ratio was greater in the 45° hip extension exercise than the leg curl (mean difference = 0.6, 95% CI = 0.3 to 1.0, $p < 0.001$) and bent-knee bridge (mean difference = 0.6, 95% CI = 0.2 to 0.9, $p = 0.001$).

Eccentric biceps femoris to medial hamstring (BF:MH) activation ratio

Paragraph 19 The eccentric BF/MH activation ratio for each exercise can be found in Figure 2b. A significant main effect was observed for exercise ($p < 0.001$) with post hoc analyses revealing that the BF/MH ratio was significantly greater in the 45° hip extension than the NHE (mean difference = 0.7, 95% CI = 0.4 to 1.0, $p < 0.001$), bent-knee bridge (mean

difference = 0.7, 95% CI = 0.4 to 1.0, $p < 0.001$), leg curl (mean difference = 0.6, 95% CI = 0.3 to 0.9, $p < 0.001$) and the glute-ham raise (mean difference = 0.6, 95% CI = 0.3 to 0.9, $p < 0.001$). No other between-exercise differences were observed once adjusted for multiple comparisons ($p > 0.002$).

INSERT FIGURE 2

Percentage change in T2 relaxation time following the 45° hip extension exercise

Paragraph 20 A significant main effect was observed for muscle ($p < 0.001$) with post hoc t tests revealing that the exercise-induced T2 changes in the BF_{SH} were significantly lower than those observed for the BF_{LH} (mean difference = 60.7%, 95% CI = 41.3 to 80.1%, $p < 0.001$), ST (mean difference = 78.0%, 95% CI = 58.4 to 97.6%, $p < 0.001$) and SM muscles (mean difference = 49.8%, 95% CI = 30.1 to 69.5%, $p < 0.001$) (Figure 3). The T2 change for ST was significantly greater than SM (mean difference = 28.2%, 95% CI = 9.2 to 47.1%, $p = 0.005$) however, no difference was observed between the BF_{LH} and SM ($p = 0.245$) or between the BF_{LH} and ST muscles ($p = 0.067$). Absolute T2 values before and after the hip extension exercise are reported in Table 1.

INSERT FIGURE 3

Percentage change in T2 relaxation time following the Nordic hamstring exercise

Paragraph 21 A main effect was detected for muscle ($p < 0.001$). Post hoc analyses showed that the T2 changes induced by exercise within the ST were significantly larger than those

observed for the BF_{LH} (mean difference = 29.8%, 95% CI = 20.5 to 39.2%, p<0.001), BF_{SH} (mean difference = 16.2%, 95% CI = 6.4 to 26.0%, p=0.002) and SM (mean difference = 29.9%, 95% CI = 20.4 to 39.4%, p<0.001) muscles (Figure 4). In addition, the T2 increase observed for BF_{SH} was significantly greater than for the BF_{LH} (mean difference = 13.7%, 95% CI = 3.9 to 23.4%, p=0.008) and SM (mean difference = 13.8, 95% CI = 3.8 to 23.7, p=0.008) muscles. No difference was observed between the BF_{LH} and SM muscles (p=0.982). The absolute T2 values before and after the NHE are reported in Table 1.

INSERT FIGURE 4

Comparison of hamstring activation ratios between exercises

Paragraph 22 When comparing the BF_{LH}/ST ratio, a significant main effect was observed for exercise (p<0.001) with post hoc analyses revealing a significantly greater ratio during 45° hip extension exercise than during the NHE (mean difference = 0.7, 95% CI = 0.6 to 0.9, p<0.001) (Figure 5).

INSERT FIGURE 5

Paragraph 23 A significant main effect was also detected for exercise when comparing the BF_{SH}/ST ratio (p<0.001). Post hoc t tests demonstrated that this ratio was significantly greater during the NHE than during the 45° hip extension exercise (mean difference = 0.42, 95% CI = 0.24 to 0.62, p<0.001). When comparing the SM/ST ratio a significant main effect was

detected for exercise ($p<0.001$) with post hoc t tests showing relatively higher ratios during the 45° hip extension than during the NHE (mean difference = 0.51, 95% CI = 0.39 to 0.64, $p<0.001$).

Table 1. T2 relaxation time values measured before (T2 Pre) and immediately after (T2 Post) the 45° hip extension and Nordic hamstring exercise (NHE) sessions.

45° hip extension			NHE	
	T2 Pre (ms)	T2 Post (ms)	T2 Pre (ms)	T2 Post (ms)
BF_{LH}	35.61 (±8.13)	62.56 (±17.34)	35.84 (±8.83)	39.76 (±11.13)
BF_{SH}	31.03 (±3.85)	36.17 (±4.65)	32.36 (±6.05)	40.57 (±9.64)
ST	38.68 (±11.89)	74.18 (±20.06)	37.31 (±10.74)	53.04 (±19.97)
SM	44.26 (±13.23)	71.09 (17.48)	41.39 (±10.87)	45.4 (±11.67)

Data are presented as mean values (±SD). BF_{LH}, biceps femoris long head; BF_{SH}, biceps femoris short head; ST, semitendinosus; SM, semimembranosus.

DISCUSSION

Paragraph 24 The primary aim of this study was to determine movements that most selectively activate the commonly injured BF_{LH}. The results support the hypothesis that hamstring activation patterns differ markedly between exercises and provide evidence to suggest that hip extension exercise more selectively targets the BF_{LH} than the NHE.

Paragraph 25 The NHE has been shown, in a number of studies,[15 16 18] to be very effective at reducing HSIs in soccer players as long as compliance is adequate.[37] However, we[11] and others[22] have previously reported that the NHE preferentially activates the ST and this might be interpreted as evidence that the exercise is sub-optimal to protect against running-related strain injury. It is entirely possible that the NHE confers injury-preventive benefits via an improved load-bearing capacity of its agonist,[38] however, in this study, we have provided EMG evidence which shows, despite the relatively selective activation of the ST, that the lateral hamstrings were still strongly activated during the NHE. Indeed, BF nEMG was higher during the NHE than during the eccentric phase of any other exercise and the evidence for this exercise's protective effects[15 16 18] suggests that eccentric actions alone in a training program are sufficient to make the hamstrings more resistant to strain injury. High levels of BF nEMG during the NHE are consistent with previous investigations[25] and are the result of the supramaximal intensity of the exercise, which potentially explains why high levels of BF nEMG were also observed in the eccentric glute-ham-raise. High levels of BF nEMG in concentric actions were observed in several other exercises including the straight-knee bridge, leg curl and the 45° hip extension which corroborates previous observations.[25] However, the importance of hamstring activation patterns during concentric actions remains unclear from the perspective of injury prevention.

Paragraph 26 While high levels of nEMG are an important stimulus for improving strength and voluntary activation,[39] exercise selectivity may still have important implications for

404 rehabilitation. For example, inhibition of previously injured BF muscles during eccentric
405 actions has been reported many months after rehabilitation,[10 11 40] and it has been
406 proposed[41] that these deficits might partly explain observations of persistent eccentric knee
407 flexor weakness,[10] BF_{LH} atrophy[12] and a chronic shortening of BF_{LH} fascicles.[42] These
408 data[10-12 40 42] are consistent with the possibility that conventional rehabilitation strategies
409 may not adequately target the commonly injured BF_{LH}. Previous studies have shown that the
410 ratio of lateral to medial hamstring (BF/MH) sEMG varies with foot rotation[43] and differs
411 between exercises.[25] In the current study, the eccentric phase of the 45° hip extension
412 exercise exhibited the greatest BF/MH nEMG ratio (1.5 ± 0.1) while the NHE (0.8 ± 0.1) and
413 bent-knee bridge exercises (0.8 ± 0.1) displayed the lowest ratios. These observations were
414 confirmed in the subsequent fMRI analysis whereby the ratio of BF_{LH} to ST in the 45° hip
415 extension exercise (0.96 ± 0.09) was markedly higher than that observed for the NHE ($0.23 \pm$
416 0.08). It is also noteworthy that the eccentric phase of other hip-oriented exercises (straight-
417 knee bridge, unilateral and bilateral stiff-leg deadlift and hip hinge) displayed BF/MH nEMG
418 ratios >1.0 . In contrast, the eccentric phase of exercises that involved significant movement at
419 the knee (leg curl, glute-ham-raise, bent-knee bridge and NHE) had higher levels of medial
420 hamstring nEMG (BF/MH ratio <1.0). These data suggest that hamstring activation strategies
421 are partly dependent on the joints involved in each movement. During concentric
422 contractions, the most selective BF activation was observed in the lunge exercise which
423 corroborates a previous fMRI investigation.[27] However, it is important to consider that the
424 lunge also exhibited the lowest BF nEMG amplitude ($21.4 \pm 7.4\%$) of any exercise which
425 likely renders it an inadequate stimulus for improving strength or stimulating hypertrophy in
426 this muscle.[39] Interestingly, the exercise that least selectively activated the BF during
427 concentric contractions was the leg curl, which mimics the joint positions and hamstring
428 muscle-tendon lengths experienced in the NHE.

Paragraph 27 The mechanism for higher levels of BF_{LH} activity during hip extension-oriented movements remains unclear, however, it is possible that hamstring muscle moment arms play a role. For example, the BF_{LH} exhibits a larger moment arm at the hip than at the knee[44] and therefore possesses a greater mechanical advantage at this joint. As a result, the BF_{LH} undergoes significantly more shortening during hip extension than knee flexion.[44] By contrast, the ST displays a larger sagittal plane moment arm at the knee than both BF_{LH} and SM,[44] which may explain its preferential recruitment during movements at this joint, such as the NHE and leg curl exercises. It is also noteworthy that the ST is a fusiform muscle with long fibre lengths and many sarcomeres in series, which potentially makes it well-suited to forceful eccentric contractions[45] such as those experienced in the NHE. Further work is needed to clarify the mechanisms underpinning these unique strategies of hamstring activation during hip and knee movements.

Paragraph 28 The current findings are different to some others that have investigated hamstring activation patterns. For example, Zebis and colleagues[25] recently reported that both the NHE and the prone isokinetic leg curl were performed with very similar levels of ST and BF_{LH} nEMG. However, in the current investigation, the NHE and leg curl exercises resulted in more selective activation of the medial hamstrings and, in the case of the NHE, the fMRI results also suggest selective use of the ST muscle. Differences between these studies may conceivably be related to participant sex (females[25] versus males in the current study), electrode placement, and the fact that this earlier work did not differentiate between the concentric and eccentric phases of each exercise. However, it is also important to consider that sEMG does not have the spatial resolution of fMRI and cannot reliably distinguish between neighbouring muscles,[30] such as the long and short heads of BF or the ST and SM, which appear to display distinct activation magnitudes.[11 23 24] These data highlight the

limitations of relying exclusively on sEMG to infer strategies of hamstring muscle activation during exercise and suggest the need for more spatially robust methods in future work.

Paragraph 29 In interpreting the results of this study, it is important to consider that sEMG and fMRI techniques measure different aspects of muscle activity. The absence of T2 relaxation time changes in people with McCardle's disease[46] suggests that fMRI is sensitive to glycolysis and it is thought that the osmotic fluid shifts which persist after exercise and give rise to T2 changes are a consequence of the accumulation of glycolytic metabolites.[36] Fortunately, the proportion of Type II glycolytic fibres does not appear to vary across the hamstring muscles[47] so this is unlikely to be a confounding factor in this study. However, exercise induced changes in T2 will be influenced by contraction mode because concentric work is characterised by higher nEMG amplitudes[29] and is markedly less efficient than eccentric work against the same loads.[48] As a consequence, the differences in T2 relaxation time changes after the 45° hip extension exercise which involved concentric and eccentric actions and the almost entirely eccentric NHE do not reflect only the levels of voluntary muscle activation. Instead, fMRI can offer insights into the relative metabolic activity and reliance upon different hamstring muscles in each exercise. According to fMRI, the NHE involves preferential ST use with modest use of the other hamstrings, while the 45° hip extension exercise appears to heavily recruit both the BF_{LH} and ST muscles. These observations are largely consistent with the sEMG component of this study, which also suggested higher activation of the medial than lateral hamstrings in the NHE and more even activation of the medial and lateral hamstrings in the 45° hip extension.

Paragraph 30 Characterising the activation patterns of the hamstrings during different tasks is an important first step in identifying exercises worthy of further investigation, however, electrical or metabolic activity of muscles should not be the only factors considered in exercise selection. Indeed, despite the BF_{LH} being more active in hip extension, there is

currently no evidence to suggest that training with this exercise actually leads to a reduction in the risk of HSI. Further work is required to understand how the hamstrings adapt to this and other exercises and adaptation is influenced by a range of factors, such as contraction mode[49] and range of motion,[45] which were not a part of the current investigation. For example, there is little reason to believe that concentric or concentrically-biased exercise is effective in HSI prevention or rehabilitation programs.[17] Indeed, there is evidence that concentric training shortens BF_{LH} fascicles[49] and shifts knee flexor torque-joint angle relationships towards shorter muscle lengths[50] and neither of these adaptations are considered beneficial for HSI prevention.[7 51] Because eccentric and concentric training programs appear to have opposing effects on fascicle lengths,[49] it is possible that exercises combining contraction modes may have minimal or at least blunted effects on muscle architecture. Future studies are needed to assess the impact of certain exercises on known or proposed risk factors for HSI such as eccentric strength[6] and fascicle lengths,[7] and only then will there be sufficient evidence to justify use of those exercises in intervention studies aimed at reducing the risk of injury. Based on the current findings, for example, it seems logical to compare the effects of training programs including the NHE and the 45⁰ hip extension exercises on the abovementioned variables.

Limitations

Paragraph 31 Given the high cost of fMRI, it was not possible to include all participants in both parts of the experiment. Therefore, comparing the results of part 1 and 2 should be done with caution. Furthermore, all of our participants were recreationally active so it remains to be seen if these findings can be applied to more highly trained athletes. We have previously shown that recreationally active young men with a history of unilateral hamstring strain exhibited less T2 change in previously injured muscles than in their uninjured homologous muscles from the contralateral limbs after performing the NHE.[22] More work will be

needed to establish whether the patterns of selective muscle activation observed in the current study are also evident in athletes with a history of strain injury.[10 11 40] Lastly, it should be acknowledged that the T2 response to an exercise stimulus is highly dynamic and can be influenced by a range of factors such as the metabolic capacity and vascular dynamics of the active tissue.[28 36] We attempted to minimise this by recruiting only male participants with a similar age and training status.

Conclusion

Paragraph 32 The current study suggests that the patterns of hamstring muscle activation are heterogeneous across a range of different strength exercises. We have provided sEMG evidence to suggest that, during eccentric contractions, hip extension exercise more selectively activates the lateral hamstrings while knee flexion-oriented exercises preferentially recruit the medial hamstrings. However, despite being the least selective activator of the BF, the NHE still elicited higher levels of BF nEMG during eccentric actions than any other exercise which may help to explain how it confers HSI-preventive benefits.[15 16 18] The results of the fMRI investigation largely confirm our initial sEMG observations, showing that, relative to the ST, the BF_{LH} was ~4 times more active in hip extension than the NHE. However, they also show that the BF_{LH}, BF_{SH}, ST and SM display distinct patterns of muscle use during different tasks. Collectively, the results of this study highlight the limitations of relying on a single method to infer strategies of muscle activation and suggest that the hip extension exercise may be useful for improving strength and voluntary activation of the commonly injured BF_{LH}. Future work is needed to determine the effect of this and other exercises on hamstring architecture and morphology before we can justify their inclusion in interventions aimed at reducing the risk of HSI.

527
528
529
530
531
532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

How might it impact upon clinical practice in the future?

- Hamstring injury prevention and rehabilitation exercises can potentially be targeted to the site of injury.
- Hip extension exercise may be more useful than the NHE for selectively activating the commonly injured BF_{LH}.

ACKNOWLEDGEMENTS

We thank the Queensland Academy of Sport's Centre of Excellence for Applied Sport Science Research for funding this investigation. The authors also acknowledge the facilities, and the scientific and technical assistance of the National Imaging Facility at the Centre for Advanced Imaging, University of Queensland.

CONTRIBUTORS

MB was the principle investigator and was involved with study design, recruitment, analysis and manuscript write up. MW, DO, GK and TS were involved with the study design, analysis and manuscript preparation. AA was involved in fMRI data acquisition. All authors had full access to all of the data (including statistical reports and tables) in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis.

TRANSPARENCY DECLARATION

The lead author* (MB) affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained. * = The manuscript's guarantor.

COPYRIGHT DECLARATION

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, [a worldwide licence](#) to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third

576 *party material where-ever it may be located; and, vi) licence any third party to do any or all of the*
577 *above.*

578 **DATA SHARING**

579 Consent was not obtained for data sharing but the presented data are anonymised and risk of
580 identification is low.

581 **FUNDING**

582 This study was funded by the Queensland Academy of Sport's Centre of Excellence for Applied
583 Sports Science Research.

584 **COMPETING INTERESTS**

585 None declared. All authors have completed the Unified Competing Interest form
586 at www.icmje.org/doi_disclosure.pdf (available on request from the corresponding author) and declare
587 that (1) the Queensland Academy of Sport's Centre of Excellence for Applied Sports Science
588 Research funded this study; (2) MB, MW, DO, GK, AA and TS have no relationships with companies
589 that might have an interest in the submitted work in the previous 3 years; (3) their spouses, partners,
590 or children have no financial relationships that may be relevant to the submitted work; and (4) MB,
591 MW, DO, GK, AA and TS have no non-financial interests that may be relevant to the submitted work.

592

593 **ETHICAL CLEARANCE**

594 All participants provided written, informed consent for this study, which was approved by the
595 Queensland University of Technology Human Research Ethics Committee and the University of
596 Queensland Medical Research Ethics Committee.

597

598

599

600

601

602

603

REFERENCES

1. Opar DA, Drezner J, Shield A, et al. Acute hamstring strain injury in track-and-field athletes: A 3-year observational study at the Penn Relay Carnival. *Scand J Med Sci Sports* 2013; 24 (4): 254–259.
2. Orchard JW, Seward H, Orchard JJ. Results of 2 decades of injury surveillance and public release of data in the Australian Football League. *Am J Sports Med* 2013;41(4):734-41.
3. Ekstrand J, Hagglund M, Walden M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br J Sports Med* 2011;45(7):553-8.
4. Orchard J, Best TM. The management of muscle strain injuries: an early return versus the risk of recurrence. *Clin J Sport Med* 2002;12(1):3-5.
5. Koulouris G, Connell DA, Brukner P, et al. Magnetic Resonance Imaging Parameters for Assessing Risk of Recurrent Hamstring Injuries in Elite Athletes. *Am J Sports Med* 2007;35(9):1500-06.
6. Opar DA, Williams MD, Timmins RG, et al. Eccentric Hamstring Strength and Hamstring Injury Risk in Australian Footballers. *Med Sci Sports Exerc* 2014;47(4):857-65.
7. Timmins R, Bourne M, Shield A, et al. Biceps femoris long head architecture, eccentric knee flexor strength and hamstring injury risk in professional football (soccer): a prospective cohort study. *Medicine and science in sports and exercise. Br J Sports Med* In press.
8. Schache AG, Wrigley TV, Baker R, et al. Biomechanical response to hamstring muscle strain injury. *Gait Posture* 2009;29(2):332-8.
9. Chumanov ES, Heiderscheit BC, Thelen DG. Hamstring musculotendon dynamics during stance and swing phases of high-speed running. *Med Sci Sports Exerc* 2011;43(3):525-32.

10. Opar DA, Williams MD, Timmins RG, et al. Knee flexor strength and bicep femoris electromyographical activity is lower in previously strained hamstrings. *J Electromyogr Kinesiol* 2013; 23(3):696-703.
11. Bourne M, Opar,DA, Williams,MD, Al Najjar, A, Shield, AJ Muscle activation patterns in the Nordic hamstring exercise: Impact of prior strain injury. *Scand J Med Sci Sports* 2015 doi: 10.1111/sms.12494.
12. Silder A, Heiderscheit B, Thelen D, et al. MR observations of long-term musculotendon remodeling following a hamstring strain injury. *Skeletal Radiol* 2008;37:1101-09.
13. Croisier JL, Ganteaume S, Binet J, et al. Strength imbalances and prevention of hamstring injury in professional soccer players. *Am J Sports Med* 2008;36(8):1469.
14. Thorborg K. What are the most important risk factors for hamstring muscle injury? *Clin J Sport Med* 2014;24(2):160-1.
15. Arnason A, Andersen TE, Holme I, et al. Prevention of hamstring strains in elite soccer: an intervention study. *Scand J Med Sci Sports* 2008;18(1):40-8.
16. Petersen J, Thorborg K, Nielsen MB, et al. Preventive Effect of Eccentric Training on Acute Hamstring Injuries in Men's Soccer: A Cluster-Randomized Controlled Trial. *Am J Sports Med* 2011;39(11):2296-303.
17. Askling CM, Tengvar M, Tarassova O, et al. Acute hamstring injuries in Swedish elite sprinters and jumpers: a prospective randomised controlled clinical trial comparing two rehabilitation protocols. *Br J Sports Med* 2014;48(7):532-9.
18. van der Horst N, Smits DW, Petersen J, et al. The preventive effect of the nordic hamstring exercise on hamstring injuries in amateur soccer players: a randomized controlled trial. *Am J Sports Med* 2015;43(6):1316-23.

19. Heiderscheit BC, Sherry MA, Silder A, et al. Hamstring strain injuries: Recommendations for diagnosis, rehabilitation and injury prevention. *J Orthop Sports Phys Ther* 2010;40(2):67.
20. Malliaropoulos N, Mendiguchia J, Pehlivanidis H, et al. Hamstring exercises for track and field athletes: injury and exercise biomechanics, and possible implications for exercise selection and primary prevention. *Br J Sports Med* 2012;46(12):846-51.
21. Guex K, Millet GP. Conceptual framework for strengthening exercises to prevent hamstring strains. *Sports Med* 2013;43(12):1207-15.
22. Mendiguchia J, Arcos AL, Garrues MA, et al. The use of MRI to evaluate posterior thigh muscle activity and damage during Nordic hamstring exercise. *J Strength Cond Res* 2013;27 (12):3426–3435.
23. Ono T, Higashihara A, Fukubayashi T. Hamstring functions during hip-extension exercise assessed with electromyography and magnetic resonance imaging. *Res Sports Med* 2011;19(1):42-52.
24. Ono T, Okuwaki T, Fukubayashi T. Differences in activation patterns of knee flexor muscles during concentric and eccentric exercises. *Res Sports Med* 2010;18(3):188-98.
25. Zebis MK, Skotte J, Andersen CH, et al. Kettlebell swing targets semitendinosus and supine leg curl targets biceps femoris: an EMG study with rehabilitation implications. *Br J Sports Med* 2013;47(18):1192-8.
26. Ditroilo M, De Vito G, Delahunt E. Kinematic and electromyographic analysis of the Nordic Hamstring Exercise. *J Electromyogr Kinesiol* 2013;23(5):1111-8.
27. Mendiguchia J, Garrues MA, Cronin JB, et al. Non-uniform changes in MRI measurements of the thigh muscles following two hamstring strengthening exercises. *J Strength Cond Res* 2013; 27(3):574-81

28. Cagnie B, Elliott JM, O'Leary S, et al. Muscle functional MRI as an imaging tool to evaluate muscle activity. *J Orthop Sports Phys Ther* 2011;41(11):896-903.
29. Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG. *J Appl Physiol (1985)* 2004;96(4):1486-95.
30. Adams GR, Duvoisin MR, Dudley GA. Magnetic resonance imaging and electromyography as indexes of muscle function. *J Appl Physiol (1985)* 1992;73(4):1578-83.
31. Fleckenstein JL, Canby RC, Parkey RW, et al. Acute effects of exercise on MR imaging of skeletal muscle in normal volunteers. *AJR Am J Roentgeno* 1988;151(2):231-7.
32. Fisher MJ, Meyer RA, Adams GR, et al. Direct relationship between proton T2 and exercise intensity in skeletal muscle MR images. *Invest Radiol* 1990;25(5):480-5.
33. Hermens HJ, Freriks B, Disselhorst-Klug C, et al. Development of recommendations for SEMG sensors and sensor placement procedures. *J Electromyogr Kinesiol* 2000;10:361-74.
34. Opar DA, Piatkowski T, Williams MD, et al. A novel device using the Nordic hamstring exercise to assess eccentric knee flexor strength: a reliability and retrospective injury study. *J Orthop Sports Phys Ther* 2013;43(9):636-40.
35. Cagnie B, Dickx N, Peeters I, et al. The use of functional MRI to evaluate cervical flexor activity during different cervical flexion exercises. *J Appl Physiol (1985)* 2008;104(1):230-5.
36. Patten C, Meyer RA, Fleckenstein JL. T2 mapping of muscle. *Semin Musculoskelet Radiol* 2003;7(4):297-305.
37. Goode AP, Reiman MP, Harris L, et al. Eccentric training for prevention of hamstring injuries may depend on intervention compliance: a systematic review and meta-analysis. *J Orthop Sports Phys Ther* 2015;49(6):349-56.

38. Schuermans J, Van Tiggelen D, Danneels L, et al. Biceps femoris and semitendinosus-teammates or competitors? New insights into hamstring injury mechanisms in male football players: a muscle functional MRI study. *Br J Sports Med* 2014;48(22):1599-606.
39. Kraemer WJ, Adams K, Cafarelli E, et al. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Ex* 2002;34(2):364-80.
40. Opar DA, Williams MD, Timmins RG, et al. Rate of torque and electromyographic development during anticipated eccentric contraction is lower in previously strained hamstrings. *Am J Sports Med* 2013;41(1):116-25.
41. Fyfe JJ, Opar DA, Williams MD, et al. The role of neuromuscular inhibition in hamstring strain injury recurrence. *J Electromyogr Kinesiol.* 2013;23(3):523-30.
42. Timmins RG, Shield AJ, Williams MD, et al. Biceps femoris long-head architecture: a reliability and retrospective injury study. *Med Sci Sports Ex* 2015;47(5):905-13.
43. Lynn SK, Costigan PA. Changes in the medial-lateral hamstring activation ratio with foot rotation during lower limb exercise. *J Electromyogr Kinesiol* 2009;19(3):e197-205.
44. Thelen DG, Chumanov ES, Hoerth DM, et al. Hamstring muscle kinematics during treadmill sprinting. *Med Sci Sports Ex* 2005;37(1):108-14.
45. Lieber RL, Friden J. Functional and clinical significance of skeletal muscle architecture. *Muscle Nerve* 2000;23(11):1647-66.
46. Fleckenstein JL, Haller RG, Lewis SF, et al. Absence of exercise-induced MRI enhancement of skeletal muscle in McArdle's disease. *J Appl Physiol* (1985) 1991;71(3):961-9.
47. Garrett WE, Jr., Califf JC, Bassett FH. Histochemical correlates of hamstring injuries. *Am J Sports Med* 1984;12(2):98-103.

48. Shellock FG, Fukunaga T, Mink JH, et al. Acute effects of exercise on MR imaging of skeletal muscle: concentric vs eccentric actions. *AJR Am J Roentgenol* 1991;156(4):765-8.
49. Timmins RG, Ruddy JD, Presland J, et al. Architectural changes of the biceps femoris after concentric or eccentric training. *Med Sci Sports Ex* 2015 [Epub ahead of print].
50. Kilgallon M, Donnelly AE, Shafat A. Progressive resistance training temporarily alters hamstring torque-angle relationship. *Scand J Med Sci Sports* 2007;17(1):18-24.
51. Brockett CL, Morgan DL, Proske U. Predicting hamstring strain injury in elite athletes. *Med Sci Sports Ex* 2004;36(3):379-87.

Figure 1. The 10 examined exercises. (a) bilateral stiff-leg deadlift, (b) hip hinge, (c) unilateral stiff-leg deadlift, (d) lunge, (e) unilateral bent knee bridge, (f) unilateral straight knee bridge, (g) leg curl, (h) 45° hip extension, (i) glute-ham-raise, (j) Nordic hamstring exercise (NHE).

Figure 2. Biceps femoris (BF) to medial hamstring (MH) normalised EMG (nEMG) relationship for the (a) concentric and (b) eccentric phases of each exercise. (SDL) Bilateral stiff-leg deadlift, (HH) hip hinge, (USDL) unilateral stiff-leg deadlift, (L) lunge, (bKb) unilateral bent knee bridge, (SKB) unilateral straight knee bridge, (LC) leg curl, (HE) 45° hip extension, (GHR) glute-ham-raise, (NHE) Nordic hamstring exercise. Exercises to the left of and above the 45° line exhibited higher levels of BF than MH nEMG and exercises to the right and below the line displayed higher levels of MH than BF nEMG.

Figure 3. Percentage change in fMRI T2 relaxation times of each hamstring muscle following the 45° hip extension exercise. Values are expressed as mean percentage change compared to values at rest. ** indicates significantly different from ST, BF_{LH} and SM (p<0.001). * indicates significantly different from ST (p=0.005). Error bars depict standard error. BF_{LH}, biceps femoris long head; BF_{SH}, biceps femoris short head; ST, semitendinosus; SM, semimembranosus.

Figure 4. Percentage change in fMRI T2 relaxation times of each hamstring muscle following the Nordic hamstring exercise. Values are expressed as mean percentage change compared to values at rest. ** indicates significantly different from BF_{LH}, BF_{SH} and SM (p≤0.002). * indicates significantly different from BF_{LH} and SM (p=0.008) Error bars depict standard error. BF_{LH}, biceps femoris long head; BF_{SH}, biceps femoris short head; ST, semitendinosus; SM, semimembranosus.

768 **Figure 5.** Ratio of biceps femoris long head (BF_{LH}) to semitendinosus (ST) (BF_{LH}/ST)
769 percentage change in fMRI T2 relaxation times following the 45° hip extension and the
770 Nordic hamstring exercise (NHE). * indicates a significant difference between exercises
771 ($p<0.001$). Error bars depict standard error.

772

773