Note. This article will be published in a forthcoming issue of the *International Journal of Sports Physiology and Performance*. The article appears here in its accepted, peer-reviewed form, as it was provided by the submitting author. It has not been copyedited, proofread, or formatted by the publisher.

Section: Original Investigation

Article Title: Optimising Pre-Season Training Loads in Australian Football

Authors: David L. Carey¹, Justin Crow^{1,2}, Kok-Leong Ong³, Peter Blanch⁴, Meg E. Morris¹, Ben J. Dascombe¹, and Kay M. Crossley¹

Affiliations: ¹La Trobe Sport and Exercise Medicine Research Centre, College of Science, Health and Engineering, La Trobe University, Melbourne, Victoria, Australia. ²Essendon Football Club, Melbourne, Victoria, Australia. ³SAS Analytics Innovation Lab, La Trobe Business School, La Trobe University, Melbourne, Victoria, Australia. ⁴School of Allied Health Sciences, Griffith University, Gold Coast, Queensland, Australia.

Journal: *International Journal of Sports Physiology and Performance*

Acceptance Date: May 10, 2017

©2017 Human Kinetics, Inc.

DOI: <https://doi.org/10.1123/ijsp.2016-0695>

Title: Optimising pre-season training loads in Australian football

Submission type: Original investigation

Authors:

David L. Carey¹, Justin Crow^{1,2}, Kok-Leong Ong³, Peter Blanch⁴, Meg E. Morris¹,
Ben J. Dascombe¹, Kay M. Crossley¹

Affiliations:

¹La Trobe Sport and Exercise Medicine Research Centre, College of Science, Health and Engineering, La Trobe University, Melbourne, Victoria, Australia.

²Essendon Football Club, Melbourne, Victoria, Australia.

³SAS Analytics Innovation Lab, La Trobe Business School, La Trobe University, Melbourne, Victoria, Australia.

⁴School of Allied Health Sciences, Griffith University, Gold Coast, Queensland, Australia.

Corresponding author:

David L. Carey

d.carey@latrobe.edu.au

La Trobe Sport and Exercise Medicine Research Centre,
La Trobe University,
Plenty Road and Kingsbury Drive,
Melbourne, Australia 3083.

Running head: Optimising training loads in AFL

Abstract word count: 247

Text-only word count: 2806

References: 15

Number of figures: 3

Number of tables: 0

Abstract

Purpose: To investigate whether pre-season training plans for Australian football can be computer generated using current training load guidelines to optimise injury risk reduction and performance improvement. **Methods:** A constrained optimisation problem was defined for daily total and sprint distance, using the pre-season schedule of an elite Australian football team as a template. Maximising total training volume and maximising Banister model projected performance were both considered as optimisation objectives. Cumulative workload and acute:chronic workload ratio constraints were placed on training programs to reflect current guidelines on relative and absolute training loads for injury risk reduction. Optimisation software was then used to generate pre-season training plans. **Results:** The optimisation framework was able to generate training plans that satisfied relative and absolute workload constraints. Increasing the off-season chronic training loads enabled the optimisation algorithm to prescribe higher amounts of ‘safe’ training and attain higher projected performance levels. Simulations showed that using a Banister model objective led to plans that included a taper in training load prior to competition in order to minimise fatigue and maximise projected performance. In contrast, when the objective was to maximise total training volume, more frequent training were prescribed in order to accumulate as much load as possible. **Conclusions:** Feasible training plans that maximise projected performance and satisfy injury risk constraints can be automatically generated by an optimisation problem for Australian football. The optimisation methods allow for individualised training plan design and the ability to adapt to changing training objectives and different training load metrics.

Keywords: AFL; injury; performance; workload ratio

INTRODUCTION

Training load prescription in team sport athletes is a balance between performance improvement^{1,2} and injury risk reduction³⁻⁶. The manipulation of training intensity, duration and frequency in order to induce improvements in athletic performance is a fundamental objective of training plan prescription⁷. To inform this process, mathematical models of the relationship between training loads and performance have been proposed for multiple athletic populations^{1,7,8}. Banister et al.¹ modelled the response to a training dose using two time-decaying functions representing fitness and fatigue. This allowed performance to be projected at a later time by taking the difference between the modelled fitness and fatigue functions. While the accuracy of these models in predicting performance has been limited^{7,9}, they provide a generalised basis for training prescription. Studies modelling team sport performance are fewer, possibly due to difficulties in quantifying individual performance within a team environment and mixed training methods.

Training load has been identified as a risk factor for injury in recent reviews, with both absolute and relative loads needing to be considered when assessing injury risk^{4,6}. The acute:chronic workload ratio quantifies an athletes' relative amount of short term (acute) to long term (chronic) training, and is an injury risk factor in a number of team sports^{4-6,10}. Additionally, there is evidence that cumulative absolute workloads can influence injury risk in Australian football³.

Currently, physical preparation staff are tasked with balancing the training guidelines associated with injury risk reduction and performance improvement when prescribing training loads. Mathematical optimisation is a method which may assist this process, particularly as more data on training load monitoring becomes available^{5,6,10-12}. Optimisation is the task of finding a set of values (decision variables) that maximise an objective function (goal) and satisfy a set of constraints. Optimising training loads has been explored in studies of tapering⁸

and to generate training plans for performance improvement¹³. No study has explored optimisation models that have incorporated training guidelines for injury prevention based on cumulative loads and workload ratios in team sport athletes.

This study aimed to: (i) determine the extent to which current training load guidelines (for relative and absolute training loads) can be used to generate optimised pre-season training plans in Australian football and (ii) investigate the effects of varying optimisation targets and load constraints on the computer generated plans.

METHODS

The task of planning pre-season training loads was posed as an optimisation problem. The decision variables were the amount of training prescribed each day and constraints were defined based on recommended acute:chronic workload ratio and cumulative load limits for injury risk reduction.

The fixture for an elite Australian football club was used as a template. Players were scheduled to play three practice matches; 98, 104 and 112 days from the start of pre-season, and their first competitive match on day 125. Outside of these matches, training loads were able to be freely prescribed on each day.

Decision variables

The goal of the optimisation procedure was to generate a full pre-season training plan by specifying the workload value (w) each day.

$$w_i = \text{training load on day } i; i \in \{1, 2, \dots, 125\}$$

In this study, total distance and sprint distance (SD) were explored since most of the research to date has examined the relationship between these variables and injury risk in team sport^{3,5,10}. However, in general, the workload variable could be substituted for any method of training load quantification.

For clarity, the following sections describe the complete optimisation formulation for total session distance and expands to then present the simple modifications needed to adapt the method to SD distance. This illustrates how the methodology can be generalised to different training load representations.

Constraints

The optimisation model required training plans to satisfy five constraints:

(i) Daily training loads were constrained to be between 0-50,000m (a generous upper bound based on unpublished data from previous seasons).

$$0 \leq w_i \leq 50,000$$

(ii) The acute:chronic workload ratio (r) was calculated on a daily basis, using a 6-day acute and 24-day chronic time window, found to be appropriate for this cohort¹⁰.

$$r_i = \sum_{j=i-6}^{i-1} \frac{w_j}{6} \bigg/ \sum_{j=i-24}^{i-1} \frac{w_j}{24}$$

Training plans were constrained to keep daily workload ratios in the previously described ‘safe zone’ for injury risk reduction ($0.6 < r_i < 1.3$)^{5,10}.

(iii) 3-weekly cumulative distance loads of between 73,721-86,662m were reported to increase pre-season injury risk in Australian footballers (OR: 5.49)³. To account for this, a rolling 21-day cumulative load (C_i) was calculated and constrained to not exceed 73,721m.

$$C_i = \sum_{j=i-21}^{i-1} w_j$$

$$C_i < 73,721$$

(iv) Rest days ($w_i = 0$) need to be considered when planning training in professional team sport due to contractual entitlements. Rest days were included by replicating the proposed rest schedule at the football club. Whilst it can be argued that the timing of rest may be a

component of designing an optimal plan, considerations around public holidays, weekends and player requests were considered beyond the scope of mathematical constraints.

(v) Match demands were taken from the 2015 AFL GPS Report¹⁴ and were incorporated by constraining the total match distance to be 13,200m ($w_{125} = 13,200$). This is an average value and may not reflect the largest loads seen in matches or the differences between playing positions. By increasing the match demand constraint the method can be adapted to prepare for higher match loads.

Pre-season matches in the AFL are subject to altered rules which generally involve more interchange players and shorter match durations. Unpublished data collected within the participating club suggested that total distances covered were approximately 15% lower in pre-season matches when compared to in-season matches ($w_{98,104,112} = 11,220$).

Calculating cumulative workloads and acute:chronic workload ratios at the beginning of pre-season requires knowledge of off-season chronic loads. At the participating club, players are generally given, and expected to follow, an off-season training program, but are not monitored with GPS devices due to league restrictions. As such there is an inherent assumption that players are completing their off-season training. Two levels of off-season chronic load were considered (representing typically prescribed loads); 14km/week and 21km/week.

Objective

Two objective functions (f_A, f_B) were considered. Objective A was to maximise the total amount of training distance in the pre-season. Representing the simple goal of allowing players to complete as much training as possible without violating injury risk constraints (assumed desirable for team sports).

$$f_A(\mathbf{w}) = \sum_{i=1}^{125} w_i$$

Objective B was to maximise the Banister model¹ projected performance on match-day. This objective was chosen as it included a consideration of the fatiguing effects of training as well as a realistic goal of trying to maximise player preparation before their first match. Banister model parameters were adapted from a study of middle distance runners¹⁵, as no such research has been undertaken in Australian football.

$$p_i = p_0 + k_1 \sum_{j=1}^{i-1} w_j e^{\frac{-(i-j)}{t_1}} - k_2 \sum_{j=1}^{i-1} w_j e^{\frac{-(i-j)}{t_2}}$$

$$k_1 = 1, k_2 = 2, t_1 = 45, t_2 = 11$$

$$f_B(\mathbf{w}) = p_{125}$$

Modifications for sprint distance

Adapting the optimisation problem outlined above to SD (defined as the distance covered above 75% of an athletes recorded top speed³) requires only a few parameter changes.

- The 3-weekly cumulative load constraint was changed to ($C_i < 1,453$) to reflect findings on increased injury risk by Colby et al³.
- Regular and pre-season match demands were changed to 268m and 200m of SD respectively³.
- Off-season chronic loads were considered at 150m/week and 225m/week of SD to reflect levels typically prescribed by the participating club.

These example values are taken from a study of one team, and may not represent the demands of other athletes. It is recommended that constraints be tailored to suit team specific demands when adapting this methodology.

Simulations

Training plans were initialised by random sampling from a normal distribution [mean (standard deviation) = 3km (1km) and 30m (10m) for distance and SD respectively]. Starting

from the random training plan the optimisation software sought to find a solution that satisfied all of the constraints and maximised the objective function (Supplementary movie 1).

Optimisation was performed using the MATLAB software package (MATLAB 8.1, The MathWorks Inc., 2013), the nonlinear programming solver and the sequential quadratic programming algorithm. Default step and function convergence tolerances were used (10^{-6}). Twenty simulations were run for distance and SD under each combination of objective and off-season chronic load.

RESULTS

The optimisation approach produced solutions that were able to satisfy both the acute:chronic workload ratio and cumulative workload constraints. Each simulated pre-season training program converged to an optimal objective value within a similar number of iterations (Figure 1), suggesting that the optimisation problem formulation, and the algorithms used were appropriate. Increasing the off-season chronic load parameter resulted in higher total pre-season training loads under optimisation objective A (Figure 1[a & c]) and higher projected performance under objective B (Figure 1[b & d]).

The distribution of prescribed session distances and session SD loads for each objective is shown in Figure 2. Each simulation had slightly different distributions (i.e. the lines in Figure 2 do not perfectly overlap), meaning it was possible for the optimisation software to achieve the same objective value using slightly different training plans. Changing the objective of the optimisation problem resulted in different load distributions in the generated training plans. In general, plans constructed with the objective of maximising Banister projected performance (objective B) prescribed more short sessions (distance < 5,000m and SD < 100m) as well as more long sessions (distance ~ 7,000-11,000m and SD > 100m).

Sample pre-season training plans (for distance and SD) generated by the optimisation procedure are shown in Figure 3. Plans for total session distance (Figure 3[a & b]) were

generally similar under the different objective functions. The longest prescribed sessions were around 15km and scheduled approximately 30 days into pre-season. A notable difference was observed near the end of the pre-season before the first regular season match. Plans aiming to maximise total distance (Figure 3[a]) prescribed frequent training up to and around matches whereas those generated using the Banister model refrained from prescribing any training in the 2-3 days preceding the first competitive match (Figure 3[b]). This difference can likely be attributed to the fatigue component of the Banister objective function used. Refraining from training before the first regular season match allows for the fatigue component to decay towards zero, and the projected performance level to be maximised¹.

SD plans (Figure 3[c & d]) showed a different load progression than distance plans. A gradual increase in SD load was prescribed by the optimisation software, with maximal loads not reached until around 90 days into pre-season. Similar to distance plans, using the Banister model to guide training led to a reduction in SD load leading into the first match (Figure 3[d]).

DISCUSSION

This study aimed to determine whether training load guidelines could be used to generate optimised pre-season training plans in Australian football, and how varying optimisation targets and load constraints influenced the computer generated plans. The results demonstrated that the total and SD distances across pre-season training programs could be generated using an optimisation approach. Cumulative load and acute:chronic workload ratio guidelines for injury risk reduction^{3-5,10} were able to be satisfied on each day of the pre-season plan when they were defined as mathematical constraints. The theoretical approach taken, based on match demands and workload constraints, generated pre-season plans (Figure 1 & 3) comparable to those previously reported in professional Australian football teams (mean total distance = 314-411km, mean total SD = 2.7-8.9km)³. This research extends the Schaefer et al.¹³

approach by including constraints based on injury risk factors and putting the approach in a sport specific context.

Choice of optimisation objective

The choice of optimisation objective influenced the distribution of training sessions prescribed by the model (Figure 2). The generated plans were generally similar in how they progressed loads until the latter stages of the pre-season, prior to the first competitive match (Figure 3; days 112-125). Maximising projected performance with a Banister model (objective B) was achieved by reducing training frequency leading into competitive matches. This reduction allows for the fatigue component of the Banister model to decay towards zero¹, maximising the projected fitness level. This aligns with previous theoretical results from Fitz-Clarke et al.⁸, where a taper before competition maximised projected performance. Maximising total pre-season volume (objective A) was accomplished by prescribing more frequent moderately sized sessions. More frequent training is a way to accumulate more load (the goal of objective A) without breaking acute:chronic workload constraints. No taper is included since objective A does not include consideration for fatigue accumulation, suggesting a Banister model objective may be more appropriate.

Effect of off-season chronic load

Modifying the off-season chronic load parameter changed the amounts of prescribed training and projected performance in the generated plans (Figure 1). A higher off-season chronic load (21 km/week for distance or 225 m/week for SD) enabled the optimiser to prescribe larger ‘safe’ training volumes (Figure 1[a & c]) and achieve higher projected performance (Figure 1[b & d]). These findings highlight the potential benefit of prescribing and adhering to training plans during off-season periods to promote follow on positive effects. The importance of high chronic workloads aligns with the findings of Malone et al.¹¹ that they may be protective against injury in Gaelic football.

Ability to customise plans

The methodological framework outlined in this study allows for customisation and individualisation of training plans depending on the preferences of the practitioner and needs of the athlete and team. For example, more aggressive or conservative training plans could be generated by simply modifying the workload ratio constraint; lowering it to 1.1 for a safer plan or raising it to 1.5 if the user was willing to accept higher injury risk⁵. Individualised planning could be incorporated by changing the off-season chronic load parameter or cumulative load constraint. For example, first year players, or others returning from injury, could be assigned a lower off-season chronic load or a lower acceptable cumulative load – and have their plans represent this lower load capacity.

The framework also allows for customisation to suit different team objectives. For example a team may want to employ a game style that requires a higher amount of SD. This could be incorporated into plans by increasing the match demand constraint for SD from 268m to 350m (or whatever the desired level may be). The timeframes used in this study (125 day pre-season with practise matches on days 98, 104 and 112) can be modified for teams or sports with different schedules. This could be accomplished by moving the timing of the match demand constraints (e.g. for a 30 day shorter pre-season: $w_{95} = 13,200$ and $w_{68,74,82} = 11,220$).

The optimisation objective can also be adapted to suit different goals. Instead of maximising projected performance (peaking) at round 1, the user may prefer to have athletes peaking for each pre-season practise match as well, or a match later in the season. A plan with this goal could be generated by changing the objective function to be the sum of Banister projected performance on each practise match day, or on a day later in the season.

In general, the optimisation approach described can be used to generate training plans that consider a number of combinations of the above modifications, allowing for rapid troubleshooting of different training strategies.

Limitations

This study presented a method for optimised training plan generation in the context of a standard Australian football pre-season. As such, the parameters considered were specific to the cohort of interest. A full evaluation of the effects of varying model parameters to reflect different possible training philosophies and timelines was considered beyond the scope of this study. The intent of the study was to determine if training plans could be generated using an optimisation approach. As such, there was no data available on the implementation of a generated plan to allow for comparison between planned and actual loads. Future studies are needed to evaluate the effects of an optimisation approach on injury occurrence and performance.

A Banister impulse-response model was used to model athlete responses to training loads¹. A discussion of the merits of using a Banister model for team sport athletes was considered beyond the scope of this study and it is possible that other models may be more appropriate.

PRACTICAL APPLICATIONS

The methodology outlined in this report provides an adaptable framework for physical preparation staff to quickly create athletic training plans that objectively optimise training goals whilst satisfying injury risk and life-balance constraints (i.e. days off) without exposing their plans to subjective bias. Practical applications include individualised training plan design and adaptability to changing training objectives. The framework described also provides theoretical scope for testing different training strategies and assumptions (e.g. how much more total training volume could athletes attain if acute:chronic workload ratio limits are increased to 1.5?

Or, are athletes able to reach match fitness levels if their off-season chronic loads are reduced by 50%?).

CONCLUSION

Feasible pre-season training plans for Australian football can be automatically generated using an optimisation approach that maximises performance whilst being constrained by injury risk guidelines. Training plans generated for athletes that enter pre-season with higher off-season chronic loads prescribed larger total training volumes. This allowed larger projected performance improvements, whilst theoretically avoiding exposure to high risk training patterns. The methodology described allows for individualised training plan design and the ability to adapt to changing training objectives.

REFERENCES

1. Banister E, Calvert T, Savage M, Bach T. A systems model of training for athletic performance. *Aust J Sports Med.* 1975;7(3):57-61.
2. Mujika I, Chatard J-C, Busso T, Geyssant A, Barale F, Lacoste L. Effects of training on performance in competitive swimming. *Canadian Journal of Applied Physiology.* 1995;20(4):395-406.
3. Colby MJ, Dawson B, Heasman J, Rogalski B, Gabbett TJ. Accelerometer and GPS-derived running loads and injury risk in elite Australian footballers. *J Strength Cond Res.* 2014;28(8):2244-2252.
4. Soligard T, Swellnus M, Alonso J-M, et al. How much is too much?(Part 1) International Olympic Committee consensus statement on load in sport and risk of injury. *British Journal of Sports Medicine.* 2016;50(17):1030-1041.
5. Gabbett TJ. The training—injury prevention paradox: should athletes be training smarter and harder? *British journal of sports medicine.* 2016;50(5):273-280.
6. Drew MK, Finch CF. The Relationship Between Training Load and Injury, Illness and Soreness: A Systematic and Literature Review. *Sports Medicine.* 2016;46(6):861-883.
7. Borresen J, Lambert MI. The quantification of training load, the training response and the effect on performance. *Sports Med.* 2009;39(9):779-795.
8. Fitz-Clarke JR, Morton RH, Banister EW. Optimizing athletic performance by influence curves. *J Appl Physiol (1985).* 1991;71(3):1151-1158.
9. Hellard P, Avalos M, Lacoste L, Barale F, Chatard J-C, Millet GP. Assessing the limitations of the Banister model in monitoring training. *Journal of sports sciences.* 2006;24(05):509-520.
10. Carey DL, Blanch P, Ong KL, Crossley KM, Crow J, Morris ME. Training loads and injury risk in Australian football-differing acute: chronic workload ratios influence match injury risk. *Br J Sports Med.* 2016.
11. Malone S, Roe M, Doran DA, Gabbett TJ, Collins K. High chronic training loads and exposure to bouts of maximal velocity running reduce injury risk in elite Gaelic football. *Journal of Science and Medicine in Sport.* 2016.
12. Akenhead R, Nassis GP. Training load and player monitoring in high-level football: current practice and perceptions. *International journal of sports physiology and performance.* 2016;11(5):587-593.
13. Schaefer D, Asteroth A, Ludwig M. Training plan evolution based on training models. Paper presented at: Innovations in Intelligent SysTems and Applications (INISTA), 2015 International Symposium on 2015.
14. Wisbey B, Montgomery PG. *Quantifying AFL Player Demands Using GPS Tracking – 2015 Season.* AFL Research Board Report 2015.

15. Morton RH, Fitz-Clarke JR, Banister EW. Modeling human performance in running.
Journal of applied physiology. 1990;69(3):1171-1177.

Figure 1: Convergence of 20 simulated pre-season training plans for: (a) distance under objective A, (b) distance under objective B, (c) SD under objective A and (d) SD under objective B. Objective A is to maximise total pre-season load and Objective B is to maximise Banister projected performance at round 1.

Figure 2: Distribution of session loads in optimal pre-season plans for: (a) distance and (b) SD. Objective A is to maximise total pre-season load and Objective B is to maximise Banister projected performance at round 1 (off-season chronic loads: 14km/week distance and 150m/week SD).

Figure 3: Computer generated optimal pre-season training plans for: (a) distance under objective A, (b) distance under objective B, (c) SD under objective A and (d) SD under objective B. Objective A is to maximise total pre-season load and Objective B is to maximise Banister projected performance at round 1 (off-season chronic loads: 14km/week distance and 150m/week SD).