Balancing biodiversity outcomes and pollution management in urban stormwater treatment wetlands

Robin Hale^{1*}, Stephen E Swearer¹, Michael Sievers^{1,+}, Rhys Coleman^{2,3}

¹School of BioSciences, University of Melbourne, Parkville, Victoria, Australia 3010

²Melbourne Water Corporation, Docklands, Victoria, Australia 3008

³School of Ecosystem and Forest Sciences, University of Melbourne, Parkville, Victoria, Australia 3010

* Email: robin.hale@unimelb.edu.au

⁺ Current address: Australian Rivers Institute, Griffith University

Role of the funding source

We acknowledge funding from the Australian Research Council (LP140100343) and Melbourne Water. R. Coleman is an employee of Melbourne Water and contributed to the development and preparation of the article, and the decision to submit for publication.

Competing interests

We have no conflict of interest to declare.

Abstract

Wetlands are increasingly being constructed to mitigate the effects of urban stormwater, such as altered hydrological regimes and reduced water quality, on downstream aquatic ecosystems. While the primary purpose of these wetlands is to manage stormwater, they also attract animals whose growth, survival and breeding (i.e. 'fitness') may be compromised. Such deleterious effects will be exacerbated if animals are caught in 'ecological traps', mistakenly preferring wetlands with unsuitable environmental conditions. Alternatively, wetlands that offer suitable habitat conditions for animals could be beneficial, especially in fragmented urban landscapes. Consequently, a thorough understanding of the potential ecological impacts of stormwater treatment wetlands is critical for managing unintended consequences to urban biodiversity.

To help facilitate this understanding, we draw upon findings from a four-year research program conducted in the city of Melbourne in south-eastern Australia as a case study. First, we summarise our research demonstrating that some stormwater wetlands can be ecological traps for native frogs and fish in the study region, whilst others likely provide important habitat in areas where few natural waterbodies remain. We use our work to highlight that while stormwater wetlands can be ecological traps, their effects can be properly managed. We propose the need for a better understanding of the ecological consequences of changes to wetland quality and their population-level impacts across the landscape. We hope that this study will generate discussions about how to most effectively manage constructed wetlands in urban landscapes and more research for a better understanding of the issues and opportunities regarding potential ecological traps.

Keywords: animal behaviour, ecological trap, urban planning, water sensitive urban design

1. Introduction

In 2015, 54% of the world's population lived in urban areas, and this is predicted to be closer to 70% by 2050 (WHO, 2017). The development of infrastructure to support this growth places considerable demand on aquatic ecosystems (Chin, 2006), and changes the nature and distribution of wetlands around cities (Kentula et al., 2004). Urbanisation causes a raft of effects on the biological, chemical and physical characteristics of aquatic environments (Walsh et al., 2005). One of the major changes is an increase in stormwater runoff from impervious surfaces (e.g. roads, buildings) connected to waterways. Stormwater runoff can alter the hydrological regime in receiving waters with subsequent impacts to stream geomorphology (Vietz et al., 2016b), and contain a diverse range of pollutants (e.g. nutrients, metals, herbicides and hydrocarbons: Malaviya and Singh, 2012).

One of the most common methods to manage the impact of urban run-off on receiving environments is to construct stormwater treatment wetlands (Malaviya and Singh, 2012), with tens of thousands of these built in residential, commercial and industrial urban areas throughout the world (Tixier et al., 2011). These wetlands have been shown to effectively treat pollutants in stormwater both in specific case studies (e.g. Al-Rubaei et al., 2017; Schulz et al., 2003) and more general assessments (e.g. for the UK; Lucas et al., 2015). Although not the primary function of these treatment systems, it is common for animals to colonise stormwater wetlands (Hassall and Anderson, 2015; Tixier et al., 2011). In some circumstances, stormwater treatment wetlands can be important habitats (e.g. Holtmann et al., 2018; Holtmann et al., 2017), especially since one of the major challenges facing animals in urban landscapes is habitat loss and fragmentation. Recognising the importance of unconventional habitats like these wetlands may be an important component of urban biodiversity conservation (Soanes et al., In press).

Urban stormwater wetlands can suitable habitats for some animals, with comparable biodiversity and community structure to natural lakes (Stephansen et al., 2016). While the pollutants that accumulate as part of the stormwater treatment function may not have deleterious effects on some species (Søberg et al., 2016), they have the potential to cause a range of lethal and sub-lethal effects for others. For instance, experiments in the US have shown that exposure to pond sediments with elevated heavy metal and salt concentrations can kill frog embryos, and cause sub-lethal effects such as reduced size at metamorphosis in other species (Gallagher et al., 2014; e.g. Snodgrass et al., 2008). A recent global assessment indicated that while measures of abundance and species richness in stormwater wetlands can be comparable to natural wetlands, the fitness (i.e. growth, survival, reproduction) of animals is often decreased (Sievers et al., 2018a). There appears to be a conflict between the engineered purpose of these wetlands as stormwater treatment systems, and their incidental use as habitats by wildlife.

Understanding the potential ecological costs and benefits of urban stormwater wetlands depends on knowledge about how the presence and fitness of animals that inhabit them varies. If poor fitness outcomes are associated with occupying certain wetlands, the consequences will be exacerbated if animals do not avoid these habitats. From an evolutionary perspective, animals should choose habitats where their fitness is high, but some mistakenly choose habitats where their fitness is reduced – a situation known as an ecological trap (Robertson and Hutto, 2006). Many animals use indirect cues associated with likely future habitat quality to select habitats (e.g. birds use vegetation to predict future food resources or predator densities; Cody, 1985). Ecological traps arise when these cues are poor predictors of habitat quality (Robertson and Chalfoun, 2016; Robertson and Hutto, 2006; Robertson et al., 2013). Perhaps the clearest example of an ecological trap is when insects are

attracted to polarised light reflected from artificial surfaces (e.g. street lamps, sides of buildings, roads) (Horvath, 1995; Horvath et al., 1998).

Ecological traps can also arise as unintended consequences of management activities and pose a serious but largely unexplored conservation risk (Battin, 2004; Hale et al., 2015a; Hale and Swearer, 2017). Traps could cause local extinctions if animals that colonise them are unable to breed or survive. Traps may also increase the risk of regional extinction by attracting animals away from high quality sites and into those where their fitness is reduced (Hale et al., 2015b).

There is a strong conceptual basis for predicting that some stormwater wetlands are ecological traps given the potential conflict between their intended stormwater treatment purpose and their use by animals (Hale et al., 2015a; Tilton, 1995). There has, however, been limited work examining how urban stormwater treatment wetlands function as habitats for animals, and less on whether they are ecological traps. Being able to identify wetlands where animals have low fitness and especially those that are traps is critical if impacts to susceptible animals are to be managed. It is equally important to know where high quality wetlands occur to help target conservation and management efforts, since these wetlands may enhance connectivity and increase biodiversity in urban areas where few natural waterbodies remain.

It may be possible to plan management activities to prevent ecological traps arising in the first place, but in instances when ecological traps have arisen as unintended consequences of management actions there are several steps to identify and mitigate their impact (Hale et al., 2015a). The first step is to assess the risks that management activities pose to animals (e.g. how will environmental conditions change and where will changes take place relative to the distribution of animals) and implementing preventative measures such as alternative treatment systems that are less likely to attract animals (e.g. biofilters, raingardens, rainwater

tanks) or providing off-line habitat more suitable for animals. The second step involves testing if an ecological trap has formed, requiring information about how the fitness of animals is affected by habitat changes, and the habitat preferences of animals (Robertson and Hutto, 2006). If an ecological trap has formed, the third step is to mitigate its effects. This could involve improving habitat quality (e.g. remediating sediments to reduce pollutant loads) or reducing the attractiveness of habitats (e.g. removing cues, such as particular types of vegetation).

For the past four years, we have been conducting research to test how stormwater treatment wetlands perform as habitats for native aquatic fauna around the city of Melbourne, in south-eastern Australia, and to examine if some might be functioning as ecological traps (step 2, above). Our aim here is to use our empirical findings to highlight key considerations for agencies involved in stormwater management in terms of managing these wetlands for biodiversity outcomes in relation to steps 1 and 3, above. Given that stormwater wetlands are being constructed in most cities globally, we hope this case study and the considerations that have arisen subsequently will fuel discussions and research on the critical knowledge gaps surrounding the ecological impacts and opportunities of constructed wetlands in urban landscapes.

2. Urban stormwater wetlands around Melbourne, south-eastern Australia

2.1. Study system

Melbourne is the capital city of the state of Victoria in south-eastern Australia, with a population of 4.85 million people (ABS, 2018), and is an ideal case study system to examine how stormwater wetlands might perform as habitats for animals for several reasons. First, as Melbourne has expanded over the past 30 years, the increase in stormwater wetlands has been

substantial, from fewer than 50 in the mid-1980s to over 700 currently (Hale et al., 2015a; Sharley et al., 2017). Therefore, stormwater wetlands are a major feature of the landscape, and one of the most common habitats available for animals. Second, animals are found in these wetlands (e.g. Murray et al., 2013). Third, wetlands range considerably in terms of their likely habitat quality for animals, and many have sediment pollutant concentrations that are above published sediment quality guidelines e.g. heavy metals (Sharley et al., 2017) and pesticides (Allinson et al., 2015). Stormwater wetlands in Melbourne have also been built in areas with a diverse range of land uses including rural, urban growth, residential, industrial and commercial. We therefore predicted that animals could inhabit some locations that are potentially suitable habitats but also others (e.g. those that are highly polluted) where deleterious impacts on fitness are likely.

2.2. Are stormwater wetlands ecological traps for native frogs?

Stormwater pollutants can impact amphibians by reducing survival and impairing growth (Gallagher et al., 2014; Snodgrass et al., 2008), causing physical abnormalities (Reeves et al., 2010) and changing behaviour (Moore et al., 2015). We surveyed amphibian assemblages at 67 urban wetlands spanning gradients in pollutant concentrations, to determine if frogs were found in highly polluted wetlands where they likely suffer fitness costs (Sievers et al., In press). The presence of most frog species was related to emergent and fringing vegetation, which is consistent with surveys from other urban ponds in the region (Hamer and Parris, 2011), but contaminant levels were largely unimportant. Frogs readily inhabited wetlands with concentrations of heavy metals and pesticides capable of causing lethal- and sub-lethal effects (Sievers et al., 2018b; Sievers et al., In press; Sievers et al., 2018c).

Our surveys highlighted that some stormwater wetlands may be ecological traps for frogs, so we experimentally tested if this was the case (Sievers et al., 2018c), focussing on a species that commonly breeds in urban wetlands, the spotted marsh frog (*Limnodynastes tasmaniensis*) (Ficken and Byrne, 2013; Hamer and Parris, 2011). We tested if survival and metamorphosis-related measures of frogs differed within mesocosms that simulated six wetlands with different contaminant levels, and if tadpoles raised in more contaminated conditions were less able to avoid cues from predators in laboratory trials. We paired this work with a field-based oviposition (egg laying) experiment to assess breeding-site preferences. Our results showed that stormwater wetlands can be ecological traps for frogs: survival was lower and tadpoles were less responsive to predator cues when raised in polluted conditions, but adults laid equivalent numbers of eggs across wetlands (Sievers et al., 2018c).

Animals often respond to environmental changes by altering their behaviour, which can determine how they cope in changing environments (Tuomainen and Candolin, 2011; Wong and Candolin, 2015). Behaviour can be directly linked to fitness, especially in terms of behavioural responses to contaminants (Zala and Penn, 2004) and might serve as a sensitive ‘early warning’ indicators of deleterious effects (Zala and Penn, 2004). In addition, animals are likely to experience multiple contaminants simultaneously, and determining how these potential stressors interact to impact behaviour is critical to understanding responses under natural conditions (Hale et al., 2017; Halfwerk and Slabbekoorn, 2015). We exposed spotted marsh frog tadpoles to two pollutants, copper and imidacloprid, at concentrations observed in wetlands around Melbourne (Allinson et al., 2015), and conducted laboratory choice experiments to document responses to predator olfactory (sense of smell) cues and tail prodding (simulated ‘attack’). Imidacloprid is a neonicotinoid insecticide and whereas exposure to other neurotoxic insecticides can affect amphibian behaviour (Brunelli et al., 2009), its effects on frogs have not been investigated. We found (Sievers et al., 2018b) that

tadpoles suffered high mortality rates when exposed to imidacloprid, and that only tadpoles that were not exposed to either contaminant avoided predator cues. In addition, imidacloprid and copper interacted to increase rates of erratic swimming, and exposed tadpoles were less able to escape simulated attacks. Collectively, we have shown that frogs are found in many stormwater treatment wetlands across Melbourne that vary in terms of the level of pollutants that are present, and that some wetlands are likely ecological traps for spotted marsh frogs.

2.3. Are stormwater wetlands ecological traps for native fish?

We tested if stormwater wetlands can be ecological traps for a native fish, the dwarf galaxias (*Galaxiella pusilla*), which is listed as vulnerable on both the IUCN Red List (IUCN, 2015) and the Australian *Environment Protection and Biodiversity Conservation Act 1999*. *G. pusilla* is a small (generally < 40 mm) fish native to south-eastern Australia that occurs in slow-flowing or still, shallow habitats (e.g. swamps, drains, backwaters, wetlands) often containing dense macrophytes and emergent plants (Coleman, 2014; Coleman et al., 2015). Habitat loss and invasive species are identified as major threats (Saddler et al., 2010). We examined habitat selection behaviour of *G. pusilla*, using laboratory choice experiments to show that fish were not responsive to a range of cues (e.g. from conspecifics or vegetation), and did not avoid water from stormwater wetlands (Hale et al., 2018). However, in a field experiment, survival was lower at stormwater wetlands than paired, nearby non-stormwater sites, and fish suffered delayed ovarian maturation, potentially due to a lack of food (zooplankton). We also conducted a series of additional laboratory and field experiments to examine the effects of the invasive mosquitofish (*Gambusia holbrooki*), which commonly occurs with *G. pusilla* (Coleman et al., 2017) and has been linked to the decline of numerous native fish and amphibians (Pyke, 2008). We found that *G. holbrooki* heavily predated *G. pusilla* larvae, and disrupted adult spawning (Brown et al., 2018). Despite these fitness costs,

G. pusilla did not avoid chemical and visual cues from the invasive *G. holbrooki* (Brown et al., 2018).

Taken together, these stormwater wetlands (separate to the ones studied for the frog experiments) may be ‘equal-preference’ ecological traps for *G. pusilla*, with fish equally preferring poorer quality options (Robertson and Hutto, 2006). Our results also suggest that invasive species in stormwater wetlands can cause ecological traps. *G. holbrooki* is frequently observed in high densities within stormwater wetlands across Melbourne (R. Coleman personal observations). Much of the work on managing ecological traps has focussed at the site scale such as improving habitat quality or reducing the attractiveness of poor quality habitats. Our results illustrate that for species like *G. pusilla* that are highly dispersive but not selective about habitats, a whole-of-landscape approach is necessary. This requires that we consider where wetlands are constructed relative to existing populations, which may be important for frogs also.

3. Key considerations for managing stormwater wetlands for biodiversity

3.1. Stormwater wetlands can be ecological traps

The potential for stormwater wetlands to be ecological traps was first raised in the mid-1990s (Tilton, 1995), but our research provides the first empirical evidence that this is the case for both fish and frogs. Furthermore, the mechanisms causing these ecological traps were different for the two taxonomic groups. Measures of frog fitness (e.g. tadpole survival) was lower at more polluted sites (Sievers et al., 2018c), but other environmental characteristics, in particular vegetation and macrophyte cover, zooplankton (food) densities, and the invasive eastern gambusia were more important determinants of the fitness of *G. pusilla* (Brown et al., 2018; Hale et al., 2018). This highlights that stormwater wetland

management needs to reduce the unintended consequences to native animals but also that these requirements will likely vary between species and life history stages. This is a well-known part of wetland management (e.g. Hamer et al., 2012) but is not routine as part of stormwater wetland planning and design.

To gain a better understanding of whether the effects we observed are widespread, similar assessments are needed at a greater number of sites. Ideally, these sites would span a range of potentially influential environmental contexts, such as pollutant levels which vary according to catchment land-use (Sharley et al., 2017). It is also possible that what is an ecological trap for one species may be suitable habitat for others. For instance, there might be variability in how different species are affected by heavy metals, with some stream invertebrates (Hale et al., 2014) being more tolerant of heavy metals than frogs (Sievers et al., 2018c). Clearly there will be pollution levels that adversely affect all species but impacts on fitness at other levels is likely to be species-specific.

Only recently have studies attempted to examine inter-specific variability in the effects of ecological traps (Robertson et al., 2018), and future work exploring how similar conditions affect the fitness of different species would provide guidance to wetland managers about the need for targeted interventions for the most susceptible animals (e.g. those that are rare or threatened, that have very specific habitat requirements, or those of strategic importance such as conservation-listed species).

3.2. Managing the effects of ecological traps in the context of urban stormwater wetlands

Management is often based on human perceptions of good habitats for animals, typically in terms of structural elements such as land cover or vegetation type (Van Dyck, 2012).

Managing ecological traps depends on thinking more about (1) what represents good habitat from the perspective of animals, as well as (2) if, and how, do animals recognise and prefer good habitats. Therefore, ecological traps can be managed by using this information to either improve habitat quality or reduce the attractiveness of habitats.

One of the main aspects of habitat quality in relation to stormwater wetlands is the accumulation of pollutants, which can have clear deleterious effects on frogs (Sievers et al., 2018b; Sievers et al., 2018c) and other animals. To ensure the functionality of stormwater wetlands for stormwater treatment, a key aspect is to develop schedules for inspection and maintenance (EPA, 2009b). The aim of maintenance is to preserve a desired level of performance and efficiency, and to extend the life-span of the wetland (Erickson et al., 2013). Determining the frequency that accumulated polluted sediment must be removed is important, and this can be based on sedimentation rates and the remaining capacity of treatment cells such as inlet sediment pond, or published guidelines such as the EPA Victoria Industrial waste resource guidelines for soils and sediments (EPA, 2009a). The frequency of maintenance is an important decision, given the typically high cost of sediment disposal (Sharley et al., 2017). Managing wetlands to decrease the likelihood of unintended consequences to biodiversity may require increased maintenance regimes, and so a key avenue for future work is to develop optimal stormwater wetland maintenance regimes that reduce the risk of pollutant accumulation to biodiversity i.e. what is the ideal frequency of maintenance to maintain pollutant levels below critical thresholds for certain animals? If more frequent maintenance is required, how do we assess and balance the ecological risks with the cost of sediment disposal?

A mechanistic view of animal behaviour can help identify management levers that can be used to modify behaviour and ultimately population-level processes (Blumstein and Berger-Tal, 2015). Understanding the habitats animals prefer and why, is an important knowledge

gap to prevent ecological traps, but habitat selection behaviour of animals is not routinely considered in stormwater wetland management (Sievers et al., 2018a). Principles from cognition have been proposed as a useful tool to manipulate problematic behaviours by animals (Greggor et al., 2014) based on identifying the cue that animals use, and the cognitive processes underpinning this behaviour, before targeting those processes, within the constraints of the animal and the cue, and this approach may be fruitful for managing stormwater wetlands that are ecological traps.

If we can identify the specific elements of habitats that animals respond to, it may be possible to decrease the attractiveness of habitats to ameliorate ecological traps. For example, Robertson (2012) developed a set of management prescriptions to disarm an ecological trap for the Olive-sided fly catcher (*Contopus cooperi*) based on understanding which habitat features this species responds most strongly to. Alternatively, we may be able to attract animals into suitable habitats by changing characteristics of the environment to alter habitat selection behaviour. For example, playbacks from calling conspecifics can be used to attract birds and amphibians to breeding sites (Ward and Schlossberg, 2004). This may be difficult to implement on a broader scale but could be useful at particular wetlands. At broader scales, other measures such as planting or removing certain types of vegetation in and around the wetland may be more feasible.

Stormwater wetlands are often constructed as a series of ponds (CSIRO, 2005), with inlets initially leading to deeper sediment ponds where many pollutants settle before water moves through a series of other pools containing vegetation. Pollutant levels are likely to vary spatially throughout this series of ponds, so one possibility is to attract animals into areas where exposure to pollutants is reduced. This has been proposed as an option at landscape scales to reduce animals using particular wetlands (Sharley et al., 2017) but may also be possible within the wetland treatment train. For example, by minimising emergent or

fringing vegetation or substituting it with alternative vegetation (e.g. deep submerged) it may discourage wildlife occupation (Sievers et al., In press) without compromising the performance of the wetland. This increased flexibility in wetland design is worth further investigation.

3.3. Better understanding the ecological consequences of changes to wetlands

While our work shows that some stormwater wetlands can be ecological traps, most studies examining how animals respond to changes in wetland quality rarely collect information about fitness and behaviour (Sievers et al., 2018a). Community and population-level metrics that are commonly used to assess wetlands provide information about whether animals are present but not whether they are able to survive, grow and reproduce. Community and population metrics are also likely to be less sensitive than measures of individual fitness (Sievers et al., 2018a). While previous work has shown that species richness and the abundance of animals can be high in stormwater wetlands (e.g. Holtmann et al., 2018), there is a need to also assess the fitness of these animals i.e. to understand the potential short-term versus long-term benefits or impacts.

Stormwater wetlands can contain a suite of different pollutants, as two recent studies from Melbourne demonstrate. In the first, Allinson et al. (2015) collected sediment samples from 24 wetlands and screened for 90 different pesticides, with 23 of these detected at least once. In the second, Sharley et al. (2017) collected sediment samples from 98 wetlands, with 13 metals recorded at all sites, and petrochemical hydrocarbons recorded at 94% of sites. Almost a third of sites had zinc values above published sediment quality thresholds. There has been extensive research on the ecotoxicology of stormwater pollutants and more work in this area is needed, especially in terms of potential interactions between different stressors, and also

less studied chemicals. We observed lethal effects of imidacloprid at concentrations well below published LC_{50} values, and showed that exposed tadpoles did not avoid cues from predators (Sievers et al., 2018b). More research is needed to understand the effects of pollutants on a wide range of species, life history stages (we might predict early stages like eggs and larvae to be more susceptible for example) and potential endpoints (e.g. mortality vs. sub-lethal effects like changes in behaviour). Given that pollutants co-occur at wetlands and could interact in complex ways (e.g. Piggott et al., 2015), looking at potential interactive effects between stressors is also critical.

3.4. Considering landscape context in wetland management

Although it is important to demonstrate where ecological traps are present, evaluating their population level consequences is pertinent for the sustained management of urban stormwater wetlands for wildlife. For example, do ecological traps mean populations of native animals are less likely to persist? Presently, this question has been addressed mainly using models (e.g. Delibes et al., 2001; Hale et al., 2015b) rather than empirical data collection. To evaluate the ultimate ecological costs and benefits of stormwater wetlands, it is necessary to explore whether observed fitness costs affect the likelihood of species persistence.

From a regional biodiversity perspective, it is critical to understand whether the presence of stormwater wetlands within urban landscapes (that may be polluted, poor quality habitats) provides greater benefits than no or few aquatic habitats. This is an important question across cities where habitat loss and fragmentation commonly occur. Poor quality habitats may still play an important role in terms of providing connectivity to better quality habitats throughout the landscape or maintaining a persistent but sub-optimal metapopulation. It is important to

understand the local-scale effects of stormwater wetlands (i.e. if animals colonise them, can they survive and breed?) but also how individual wetlands contribute to landscape connectivity (i.e. do they provide dispersers that can move elsewhere?). Conservation priorities are often based on the contribution of individual patches to landscape connectivity (Rubio et al., 2015), and it is important to consider stormwater wetlands in this regard. Simulation modelling has demonstrated that if habitats vary temporally in terms of quality, it is possible that regional populations can persist even if all patches are of low quality overall (population growth rate < 1) i.e. all patches are low quality when considered over longer time scales but at any one time point there is sufficient variability so that some are suitable (Schreiber, 2010). This could be important for stormwater wetlands, where pollutant loads may be higher after rainfall events (Walsh et al., 2016).

For species that are highly dispersive, it will also be important to consider where stormwater wetlands are located relative to other habitats while considering the movement capacity and behaviour of different species. For instance, *G. pusilla* are highly dispersive and appear to have low habitat selectivity (Hale et al., 2018). Should we therefore consider building alternative high-quality habitats nearby to stormwater wetlands for the persistence of this nationally significant species? Or avoid building stormwater wetlands when possible in areas where this species occurs?

Landscape context is also an important management consideration. For example, catchment land use is likely to be a key determinant of the pollution levels within wetlands (Sharley et al., 2017), so actions undertaken at the catchment-scale to address the underlying causes of stormwater run-off might be more effective than actions at the site level (Vietz et al., 2016a). While such catchment-scale management attempts are rare and require interdisciplinary collaborations that consider the complex social-ecological aspects of urban catchments, some examples do exist (e.g. Walsh et al., 2015). For example, retaining more

water in the upstream catchment (e.g. via water sensitive urban design methods such as biofilters, raingardens and rainwater tanks to reduce and treat run-off) reduces stormwater pollutant loads into wetlands. Alternatively, diversifying the types of wetlands that are built to include adjacent offline habitats for animals that are less likely to accumulate pollutants for animals when designing stormwater treatment wetlands.

Conclusions

Stormwater run-off is a major environmental issue for aquatic ecosystems in urban areas, and stormwater treatment wetlands can be an effective pollution management tool. However, stormwater wetlands are often used incidentally as habitats for animals. This can be beneficial for the persistence of some species in fragmented urban environments. However, we have demonstrated that there is the potential that animals that colonise these wetlands suffer deleterious impacts such as from accumulated pollutants or the presence of invasive species. To guide planning and maintenance decision-making, we have highlighted some of the key issues and management options for these wetlands so that impacts to biodiversity may be reduced. We acknowledge the preliminary nature of these recommendations and strongly advocate for a deeper understanding of the benefits and impacts of urban stormwater treatment wetlands across a broader suite of animals across both site- and regional-scales.

4. Acknowledgements

This work was supported by Melbourne Water and the Australian Research Council under project LP140100343. We thank the following people for attending a workshop to discuss this project, and for providing feedback on an earlier draft of this manuscript: Alison Rickard, David Reginato, Bronwen Hutchison, William Steele, Trish Grant, Birgit Jordan, Rachael

407 Bathgate, Daniella Gerente, Arezou Houshmand, Leon Metzeling, Chris Walsh, Dennis
408 Corbett and Michael Scroggie. Thanks to Rachael Bathgate for organising and coordinating
409 the workshop.

410

411 **References**

412 ABS, 2018. <http://www.abs.gov.au/ausstats/abs@.nsf/mf/3218.0>.

413 Al-Rubaei, A.M., Engstrom, M., Viklander, M., Blecken, G.T., 2017. Effectiveness of a 19-
414 Year Old Combined Pond-Wetland System in Removing Particulate and Dissolved
415 Pollutants. *Wetlands* 37, 485-496.

416 Allinson, G., Zhang, P., Bui, A., Allinson, M., Rose, G., Marshall, S., Pettigrove, V., 2015.
417 Pesticide and trace metal occurrence and aquatic benchmark exceedances in surface waters
418 and sediments of urban wetlands and retention ponds in Melbourne, Australia. *Environmental*
419 *Science and Pollution Research* 22, 10214-10226.

420 Battin, J., 2004. When good animals love bad habitats: Ecological traps and the conservation
421 of animal populations. *Conserv Biol* 18, 1482-1491.

422 Blumstein, D.T., Berger-Tal, O., 2015. Understanding sensory mechanisms to develop
423 effective conservation and management tools. *Curr. Opin. Behav. Sci.* 6, 13-18.

424 Brown, T.R., Coleman, R., Swearer, S.E., Hale, R., 2018. Behavioural responses to, and
425 fitness consequences from, an invasive species are life-stage dependent in a threatened native
426 fish. *Biological Conservation* 228, 10-16.

427 Brunelli, E., Bernabò, I., Berg, C., Lundstedt-Enkel, K., Bonacci, A., Tripepi, S., 2009.
428 Environmentally relevant concentrations of endosulfan impair development, metamorphosis
429 and behaviour in *Bufo bufo* tadpoles. *Aquatic Toxicology* 91, 135-142.

430 Chin, A., 2006. Urban transformation of river landscapes in a global context. *Geomorphology*
 431 79, 460-487.

432 Cody, M.L., 1985. *Habitat selection in birds*. Academic Press, Orlando, FL, USA.

433 Coleman, R.A., 2014. Conservation of the dwarf galaxias, *Galaxiella pusilla* (Mack 1936)
 434 (Teleostei: Galaxiidae), a threatened freshwater fish from south-eastern Australia, School of
 435 BioSciences. PhD Dissertation, University of Melbourne, Australia.

436 Coleman, R.A., Hoffmann, A.A., Raadik, T.A., 2015. A review of *Galaxiella pusilla* (Mack)
 437 (Teleostei: Galaxiidae) in south-eastern Australia with a description of a new species.
 438 *Zootaxa* 4021, 243-281.

439 Coleman, R.A., Raadik, T.A., Pettigrove, V., Hoffmann, A.A., 2017. Taking advantage of
 440 adaptations when managing threatened species within variable environments: the case of the
 441 dwarf galaxias, *Galaxiella pusilla* (Teleostei, Galaxiidae). *Marine and Freshwater Research*
 442 68, 175-186.

443 CSIRO, 2005. *WSUD Engineering Procedures : Stormwater*. CSIRO Publishing, Victoria,
 444 Australia.

445 Delibes, M., Gaona, P., Ferreras, P., 2001. Effects of an attractive sink leading into
 446 maladaptive habitat selection. *The American Naturalist* 158, 277-285.

447 EPA, 2009a. *Soil hazard categorisation and management, industrial waste resource*
 448 *guidelines*. Environmental Protection Authority, Melbourne.

449 EPA, 2009b. *Stormwater Wet Pond and Wetland Management Guidebook*.

450 Erickson, A.J., Weiss, P.T., Gulliver, J.S., 2013. Maintenance of stormwater treatment
451 practices, Optimizing Stormwater Treatment Practices. Springer, pp. 265-283.

452 Ficken, K., Byrne, P.G., 2013. Heavy metal pollution negatively correlates with anuran
453 species richness and distribution in south-eastern Australia. Austral Ecology 38, 523-533.

454 Gallagher, M.T., Snodgrass, J.W., Brand, A.B., Casey, R.E., Lev, S.M., Van Meter, R.J.,
455 2014. The role of pollutant accumulation in determining the use of stormwater ponds by
456 amphibians. Wetlands Ecology and Management 22, 551-564.

457 Greggor, A.L., Clayton, N.S., Phalan, B., Thornton, A., 2014. Comparative cognition for
458 conservationists. Trends Ecol. Evol. 29, 489-495.

459 Hale, R., Coleman, R., Pettigrove, V., Swearer, S.E., 2015a. Identifying, preventing and
460 mitigating ecological traps to improve the management of urban aquatic ecosystems. Journal
461 of Applied Ecology 52, 928-939.

462 Hale, R., Coleman, R., Sievers, M., Brown T, Swearer, S., 2018. Using conservation
463 behaviour to manage ecological traps for a threatened freshwater fish. Ecosphere 9, e02381.

464 Hale, R., Marshall, S., Jeppe, K., Pettigrove, V., 2014. Separating the effects of water
465 physicochemistry and sediment contamination on *Chironomus tepperi* (Skuse) survival,
466 growth and development: A boosted regression tree approach. Aquatic Toxicology 152, 66-
467 73.

468 Hale, R., Piggott, J.J., Swearer, S.E., 2017. Describing and understanding behavioral
469 responses to multiple stressors and multiple stimuli. Ecol Evol 7, 38-47.

470 Hale, R., Swearer, S.E., 2017. When good animals love bad restored habitats: how
 471 maladaptive habitat selection can constrain restoration. *Journal of Applied Ecology* 54, 1478-
 472 1486.

473 Hale, R., Treml, E.A., Swearer, S.E., 2015b. Evaluating the metapopulation consequences of
 474 ecological traps. *Proc. R. Soc. Lond. B Biol. Sci.* 282, 20142930.

475 Halfwerk, W., Slabbekoorn, H., 2015. Pollution going multimodal: the complex impact of the
 476 human-altered sensory environment on animal perception and performance. *Biology Letters*
 477 11, 20141051.

478 Hamer, A.J., Parris, K.M., 2011. Local and landscape determinants of amphibian
 479 communities in urban ponds. *Ecol. Appl.* 21, 378-390.

480 Hamer, A.J., Smith, P.J., McDonnell, M.J., 2012. The importance of habitat design and
 481 aquatic connectivity in amphibian use of urban stormwater retention ponds. *Urban*
 482 *Ecosystems* 15, 451-471.

483 Hassall, C., Anderson, S., 2015. Stormwater ponds can contain comparable biodiversity to
 484 unmanaged wetlands in urban areas. *Hydrobiologia* 745, 137-149.

485 Holtmann, L., Juchem, M., Bruggeshemke, J., Mohlmeyer, A., Fartmann, T., 2018.
 486 Stormwater ponds promote dragonfly (Odonata) species richness and density in urban areas.
 487 *Ecol. Eng.* 118, 1-11.

488 Holtmann, L., Philipp, K., Becke, C., Fartmann, T., 2017. Effects of habitat and landscape
 489 quality on amphibian assemblages of urban stormwater ponds. *Urban Ecosystems* 20, 1249-
 490 1259.

491 Horvath, G., 1995. Reflection polarization patterns at flat water surfaces and their relevance
 492 for insect polarization vision. *Journal of Theoretical Biology* 175, 27-37.

493 Horvath, G., Bernath, B., Molnar, G., 1998. Dragonflies find crude oil visually more
 494 attractive than water: Multiple-choice experiments on dragonfly polarotaxis.
 495 *Naturwissenschaften* 85, 292-297.

496 IUCN, 2015. The IUCN Redlist for Threatened Species. International Union of Conservation
 497 of Nature, www.iucnredlist.org.

498 Kentula, M.E., Gwin, S.E., Pierson, S.M., 2004. Tracking changes in wetlands with
 499 urbanization: Sixteen years of experience in Portland, Oregon, USA. *Wetlands* 24, 734-743.

500 Lucas, R., Earl, E.R., Babatunde, A.O., Bockelmann-Evans, B.N., 2015. Constructed
 501 wetlands for stormwater management in the UK: a concise review. *Civil Engineering and*
 502 *Environmental Systems* 32, 251-268.

503 Malaviya, P., Singh, A., 2012. Constructed wetlands for management of urban stormwater
 504 runoff. *Critical Reviews in Environmental Science and Technology* 42, 2153-2214.

505 Moore, H., Chivers, D.P., Ferrari, M.C., 2015. Sub-lethal effects of Roundup™ on tadpole
 506 anti-predator responses. *Ecotoxicology and environmental safety* 111, 281-285.

507 Murray, C.G., Kasel, S., Loyn, R.H., Hepworth, G., Hamilton, A.J., 2013. Waterbird use of
 508 artificial wetlands in an Australian urban landscape. *Hydrobiologia* 716, 131-146.

509 Piggott, J.J., Townsend, C.R., Matthaei, C.D., 2015. Reconceptualizing synergism and
 510 antagonism among multiple stressors. *Ecology and Evolution* 5, 1538-1547.

511 Pyke, G.H., 2008. Plague minnow or mosquito fish? A review of the biology and impacts of
512 introduced *Gambusia* species. Annual Review of Ecology Evolution and Systematics 39,
513 171-191.

514 Reeves, M.K., Jensen, P., Dolph, C.L., Holyoak, M., Trust, K.A., 2010. Multiple stressors
515 and the cause of amphibian abnormalities. Ecological Monographs 80, 423-440.

516 Robertson, B.A., 2012. Investigating targets of avian habitat management to eliminate an
517 ecological trap. Avian Conservation and Ecology 7.

518 Robertson, B.A., Chalfoun, A.D., 2016. Evolutionary traps as keys to understanding
519 behavioral maladaptation. Curr. Opin. Behav. Sci. 12, 12-17.

520 Robertson, B.A., Hutto, R.L., 2006. A framework for understanding ecological traps and an
521 evaluation of existing evidence. Ecology 87, 1075-1085.

522 Robertson, B.A., Keddy-Hector, I.A., Shrestha, S.D., Silverberg, L.Y., Woolner, C.E.,
523 Hetterich, I., Horváth, G., 2018. Susceptibility to ecological traps is similar among closely
524 related taxa but sensitive to spatial isolation. Animal Behaviour 135, 77-84.

525 Robertson, B.A., Rehage, J.S., Sih, A., 2013. Ecological novelty and the emergence of
526 evolutionary traps. Trends in Ecology and Evolution 28, 552-560.

527 Rubio, L., Bodin, Ö., Brotons, L., Saura, S., 2015. Connectivity conservation priorities for
528 individual patches evaluated in the present landscape: how durable and effective are they in
529 the long term? Ecography 38, 782-791.

530 Saddler, S., Jackson, J., Hammer, M., 2010. National Recovery Plan for the Dwarf Galaxias
531 *Galaxiella pusilla*. Department of Sustainability and Environment, Melbourne, Australia.

532 Schreiber, S.J., 2010. Interactive effects of temporal correlations, spatial heterogeneity and
533 dispersal on population persistence. *Proc. R. Soc. Lond. B Biol. Sci.*, rspb20092006.

534 Schulz, C., Gelbrecht, J., Rennert, B., 2003. Treatment of rainbow trout farm effluents in
535 constructed wetland with emergent plants and subsurface horizontal water flow. *Aquaculture*
536 217, 207-221.

537 Sharley, D.J., Sharp, S.M., Marshall, S., Jeppe, K., Pettigrove, V.J., 2017. Linking urban land
538 use to pollutants in constructed wetlands: Implications for stormwater and urban planning.
539 *Landscape and Urban Planning* 162, 80-91.

540 Sievers, M., Hale, R., Parris, K.M., Swearer, S.E., 2018a. Impacts of human-induced
541 environmental change in wetlands on aquatic animals. *Biological Reviews* 93, 529-554.

542 Sievers, M., Hale, R., Swearer, S.E., Parris, K.M., 2018b. Contaminant mixtures interact to
543 impair predator-avoidance behaviours and survival in a larval amphibian. *Ecotoxicology and*
544 *Environmental Safety* 161, 482-488.

545 Sievers, M., Hale, R., Swearer, S.E., Parris, K.M., In press. Occupancy of polluted wetlands
546 is a risk to conserving frogs in urban landscapes. *Conserv Biol.*

547 Sievers, M., Parris, K.M., Swearer, S.E., Hale, R., 2018c. Stormwater wetlands can function
548 as ecological traps for urban frogs. *Ecol. Appl.* 28, 1106-1115.

549 Snodgrass, J.W., Casey, R.E., Joseph, D., Simon, J.A., 2008. Microcosm investigations of
550 stormwater pond sediment toxicity to embryonic and larval amphibians: variation in
551 sensitivity among species. *Environmental Pollution* 154, 291-297.

552 Soanes, K., Sievers, M., Chee, Y.E., Williams, N.S.G., Bhardwaj, M., Marshall, A.J., Parris,
553 K.M., In press. Correcting common misconceptions to inspire conservation action in urban
554 environments. *Conserv Biol.*

555 Søbørg, L.C., Vollertsen, J., Blecken, G.-T., Nielsen, A.H., Viklander, M., 2016.
556 Bioaccumulation of heavy metals in two wet retention ponds. *Urban Water Journal* 13, 697-
557 709.

558 Stephansen, D.A., Nielsen, A.H., Hvitved-Jacobsen, T., Pedersen, M.L., Vollertsen, J., 2016.
559 Invertebrates in stormwater wet detention ponds—Sediment accumulation and
560 bioaccumulation of heavy metals have no effect on biodiversity and community structure.
561 *Science of the Total Environment* 566, 1579-1587.

562 Tilton, D.L., 1995. Integrating wetlands into planned landscapes. *Landscape and Urban*
563 *Planning* 32, 205-209.

564 Tixier, G.T., Lafont, M., Grapentine, L., Rochfort, Q., Marsalek, J., 2011. Ecological risk
565 assessment of urban stormwater ponds: literature review and proposal of a new conceptual
566 approach providing ecological quality goals and the associated bioassessment tools.
567 *Ecological Indicators* 11, 1497-1506.

568 Tuomainen, U., Candolin, U., 2011. Behavioural responses to human-induced environmental
569 change. *Biological Reviews* 86, 640-657.

570 Van Dyck, H., 2012. Changing organisms in rapidly changing anthropogenic landscapes: the
571 significance of the Umwelt'-concept and functional habitat for animal conservation. *Evol.*
572 *Appl.* 5, 144-153.

573 Vietz, G.J., Rutherford, I.D., Fletcher, T.D., Walsh, C.J., 2016a. Thinking outside the
 574 channel: Challenges and opportunities for protection and restoration of stream morphology in
 575 urbanizing catchments. *Landscape and Urban Planning* 145, 34-44.

576 Vietz, G.J., Walsh, C.J., Fletcher, T.D., 2016b. Urban hydrogeomorphology and the urban
 577 stream syndrome: Treating the symptoms and causes of geomorphic change. *Progress in*
 578 *Physical Geography* 40, 480-492.

579 Walsh, C.J., Booth, D.B., Burns, M.J., Fletcher, T.D., Hale, R.L., Hoang, L.N., Livingston,
 580 G., Rippey, M.A., Roy, A.H., Scoggins, M., Wallace, A., 2016. Principles for urban
 581 stormwater management to protect stream ecosystems. *Freshwater Science* 35, 398-411.

582 Walsh, C.J., Fletcher, T.D., Bos, D.G., Imberger, S.J., 2015. Restoring a stream through
 583 retention of urban stormwater runoff: a catchment-scale experiment in a social–ecological
 584 system. *Freshwater Science* 34, 1161-1168.

585 Walsh, C.J., Roy, A.H., Feminella, J.W., Cottingham, P.D., Groffman, P.M., Morgan, R.P.,
 586 2005. The urban stream syndrome: current knowledge and the search for a cure. *Journal of*
 587 *the North American Benthological Society* 24, 706-723.

588 Ward, M.P., Schlossberg, S., 2004. Conspecific attraction and the conservation of territorial
 589 songbirds. *Conserv Biol* 18, 519-525.

590 WHO, 2017.
 591 http://www.who.int/gho/urban_health/situation_trends/urban_population_growth/en/.

592 Wong, B.B.M., Candolin, U., 2015. Behavioral responses to changing environments. *Behav.*
 593 *Ecol.* 26, 665-673.

594 Zala, S.M., Penn, D.J., 2004. Abnormal behaviours induced by chemical pollution: a review
595 of the evidence and new challenges. *Animal Behaviour* 68, 649-664.

596