1 Multimodal nature of GTPS

2
3 1 **Individuals with persistent greater trochanteric pain syndrome exhibit impaired pain**
4
5 2 **modulation, as well as poorer physical and psychological health than pain free**
6
7 3 **individuals: a cross sectional study**
8
9

10
11 4 Melanie Louise Plinsinga¹, PhD, Brooke Kaye Coombes², PhD, Rebecca Mellor¹, PhD, Bill
12
13 5 Vicenzino¹, PhD
14
15

16 6
17
18
19 7 **Affiliations**
20

21
22 8 ¹ School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane,
23
24 9 QLD 4072, Australia
25

26
27 10 ² School of Allied Health Sciences, Griffith University, QLD 4222, Australia
28
29

30 11
31
32
33 12 **Corresponding author**
34

35
36 13 Bill Vicenzino
37
38
39 14 School of Health and Rehabilitation Sciences Building 84A, The University of Queensland,
40
41 15 Brisbane, QLD 4072, Australia; +61 7 336 52781; b.vicenzino@uq.edu.au
42
43

44 16
45
46
47 17 **Conflicts of Interest and Source of Funding:** This study was supported by the National
48
49 18 Health and Medical Research Council (NHMRC) Program Grant (#631717). MLP is
50
51 19 supported by the International Postgraduate Research Scholarship (IPRS)/University of
52
53 20 Queensland Centennial Scholarship (UQcent). Authors have no conflicts of interest to
54
55 21 declare.
56
57
58
59 22

Multimodal nature of GTPS

24 **Abstract**

25 **Objectives:** To compare physical, sensory and psychosocial factors between individuals with
26 greater trochanteric pain syndrome and controls, and to explore factors associated with pain
27 and disability.

28 **Design:** Cross-sectional study.

29 **Setting:** General community.

30 **Subjects:** Patients with persistent, clinically diagnosed greater trochanteric pain syndrome
31 and healthy controls.

32 **Methods:** Participants completed tests of thermal and pressure pain threshold; conditioned
33 pain modulation; temporal summation; muscle strength, physical function, physical activity,
34 psychological factors and health related quality of. Standardised mean differences between
35 groups were calculated, and multiple linear regression identified factors associated with pain
36 and disability.

37 **Results:** Forty patients (95% female, average (standard deviation) age 51 (9) years), and 58
38 controls (95% female, age 53 (11) years) were included. Heat pain threshold, temporal
39 summation and pain catastrophizing were not different between groups. Compared to
40 controls, patients displayed significantly poorer quality of life (standardized mean difference
41 -2.66), lower pressure pain threshold locally (-1.47, remotely (-0.57)), poorer health status (-
42 1.22), impaired physical function (range 0.64 to 1.20), less conditioned pain modulation (-
43 1.01), weaker hip abductor/extensor strength (-1.01 and -0.59), higher depression (0.72) and
44 anxiety (0.61) levels, lower cold pain threshold locally (-0.47, (remotely (-0.39)) and less
45 time spent in (vigorous) physical activity (-0.43 to -0.39). 26% of pain and disability was
46 explained by depression, hip abductor strength and time to complete stairs.

1 Multimodal nature of GTPS

2

3 47 **Conclusions:** Patients with greater trochanteric pain syndrome exhibited poorer health

4

5 48 related quality of life, physical impairments, widespread hyperalgesia, and greater

6

7 49 psychological distress than healthy controls. Physical and psychological factors were

8

9 50 associated with pain and disability.

10

11

12

13 51 **Key words:** Tendinopathy, musculoskeletal pain, chronic pain.

14

15

16 52

17

18

19 53

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Multimodal nature of GTPS

54 **Introduction**

55 Greater trochanteric pain syndrome (GTPS) presents as lateral hip pain on activities that load
56 the structures at the greater trochanter. Most common in middle-aged females, it is the most
57 prevalent lower limb tendinopathy to present to primary care (4.22 per 1000 per year) (1).
58 GTPS often persists, and high levels of pain and disability and poor quality of life are
59 reported (2). Pain self-efficacy, anxiety and depression are associated with other persistent
60 musculoskeletal conditions (3, 4) and tendinopathies (5), however, the presence of
61 psychological factors in persistent GTPS when compared to controls has not been studied.

62

63 There is evidence of local and widespread mechanical and thermal hyperalgesia in upper limb
64 tendon studies, with discrepancies seen in lower limb studies (6-9). These lower limb studies
65 are limited by low sample size and high methodological heterogeneity including testing
66 methods, sites of assessment and equipment (7). Currently, there is one cross-sectional study
67 available in GTPS that reports local and widespread mechanical hyperalgesia in GTPS (n =
68 18) compared to controls (n = 18) (8). Altered endogenous pain inhibitory mechanisms have
69 been reported in several chronic musculoskeletal conditions like osteoarthritis (10), low back
70 pain (11, 12), Achilles tendinopathy (9), and recently in GTPS (13). The latter study reported
71 a negative condition pain modulation in 34.7% of GTPS participants, although there was no
72 comparator group included. Investigation of a more comprehensive battery of sensory tests in
73 people with persistent greater trochanteric pain is warranted.

74 The gluteal muscles are crucial for functions like walking and stair climbing (14). Hip
75 abductor weakness is observed unilaterally (15-17) and bilaterally (15, 16) in GTPS, but hip
76 extensor muscle strength has received limited attention.

77

1 Multimodal nature of GTPS

2
3 78 Our primary aim was to quantify differences in physical, sensory, and psychosocial outcome
4
5 79 measures in participants with GTPS compared to pain-free controls. Based on the current
6
7
8 80 literature, we hypothesized that impaired physical strength and function, mechanical
9
10 81 hyperalgesia and psychological distress would be found in persistent GTPS compared to
11
12 82 pain-free controls. Our secondary aim was to identify factors associated with pain and
13
14 83 disability in participants with GTPS. We hypothesized that pain and disability would be
15
16
17 84 explained by multidimensional patient factors.

20 85

22 86

24 87 **Methods**

26 88 Study design

28
29 89 This cross-sectional study compared individuals with GTPS and age and sex matched
30
31 90 controls. Reporting follows the Strengthening the Reporting of Observational studies in
32
33 91 Epidemiology (STROBE) guidelines for cross-sectional studies (18). Ethical approval was
34
35 92 granted by The University of Queensland Human Research Ethics Committee (approval
36
37
38 93 #2015000219). All participants provided verbal and written consent.

41 94

44 95

47 96 Setting

49
50 97 Between 16 May 2017 and 11 September 2018, participants were recruited from the Brisbane
51
52 98 metropolitan area via newsletters and social media. A three-stage eligibility screening process
53
54 99 was adopted: online screening, telephone interview and physical screening. Physical
55
56
57 100 screening was performed by a registered post-graduate qualified musculoskeletal
58
59 101 physiotherapist with >20 years clinical experience and baseline measures were conducted
60

Multimodal nature of GTPS

during one session of approximately 2.5 hours in a sound and temperature controlled room at

The University of Queensland.

Participants

Volunteers were aged 18-70 years. Individuals with GTPS were eligible if they experienced lateral hip pain rated $\geq 2/10$ on an 11-point Numeric Rating Scale (NRS) (0 = no pain; 10 = worst pain imaginable) at its worst in the past week; pain duration ≥ 3 months; pain on palpation of the tendon insertion on the greater trochanter (19), plus reproduction of trochanteric pain on at least one of the following clinical tests: (1) 30s single leg stance, (2) the Hip FADER test, (3) static muscle test in the FADER position, (4) the FABER test, (5) Ober's test, (6) a static muscle contraction in the Ober's test position (19). GTPS participants were excluded if they experienced groin pain on quadrant testing $> 3/10$ PNRS, received a corticosteroid injection in the previous six months, experienced major trauma in the past 12 months, or had lower limb or back pain that was worse than their hip pain, required treatment or prevented usual work or leisure activities in the previous six months. Controls were excluded if they had lateral hip pain or other areas of pain requiring treatment or preventing work or leisure activities in the preceding six months. Pregnancy, systemic inflammatory or neurological disorders, uncontrolled diabetes, and fibromyalgia were general exclusion criteria.

Outcome measures

1 Multimodal nature of GTPS

2
3 125 Data collection was performed by one researcher (MP) not blinded to the condition of the
4
5 126 participants.

6
7
8 127

9
10
11 128 *Pain and disability*

12
13
14 129 Average pain, worst pain and pain during activity in the past week were rated using a PNRs.

15
16 130 Pain and disability was measured using the Victorian Institute of Sports Assessment – GTPS
17
18 (VISA-G) questionnaire (20), consisting of eight items. Scores range from 0 to 100, with
19 131
20
21 132 higher scores indicating less pain and better function. The VISA-G demonstrates good test-retest
22
23 133 reliability (ICC 0.827; 95% CI 0.638-0.923), internal consistency (Cronbach’s alpha 0.809; 95% CI
24
25 134 0.709-0.934) and construct validity, compared to the Harris Hip Score and the Oswestry Disability
26
27 135 Index (20).

28
29
30 136

31
32
33 137 *Quantitative sensory testing*

34
35
36 138 Somatosensory profiles were assessed unilaterally at a local hip site (the most painful spot in
37
38 139 GTPS; 1-2cm posterior to the greater trochanter in controls) and a remote site (the lateral
39
40 140 elbow, side randomly allocated by a coin toss). In participants with bilateral symptoms, the
41
42 141 most affected side was measured. The order of quantitative sensory tests (QSTs) was
43
44 142 standardized as follows.

45
46
47 143

48
49
50
51 144 Pain thresholds were determined by asking participants to ‘press a button at the first onset of
52
53 145 pain’. Cold pain threshold (CPT) and heat pain threshold (HPT) were assessed using a
54
55 146 Thermotest Analyser (Somedic AB, Farsta, Sweden). From a baseline temperature of 32°C,
56
57 147 incremental increases/decreases in temperature (1°C /s) were applied and tests were
58
59
60

Multimodal nature of GTPS

terminated when the participant pressed the button or if upper (50°C) or lower (5°C) cut-off temperatures were reached. Pressure pain threshold (PPT) was assessed by placing a 1cm² probe of a handheld pressure algometer (Somedic, Sweden) on the skin and gradually increasing the pressure at a rate of 30kPa/s, until the participant pressed the button. A maximum pressure of 900 kPa could be applied. Average pain thresholds were calculated from three consecutive measurements with 30-second rest intervals.

Temporal summation was measured as the difference in perceived intensity (PNRS) between a single pinprick stimuli (256 mN pinprick) and the average of 10 repetitive pinprick stimuli applied within an area of 1cm² at 1/s. Five series were applied with 30-second rest intervals.

Conditioned pain modulation (CPM) was assessed by immersing the contralateral hand in a cold-water bucket (conditioning stimulus) and measurement of PPT over the lateral hip (test stimulus) before, during and immediately after immersion of the hand (21). The starting water temperature was 10°C (range 9.5°C – 10.5°C). Water temperature was adjusted by adding warm or cold water until a pain intensity of 4-6/10 on PNRS was reported. The highest PPT score recorded during or immediately post immersion of the hand was subtracted from the PPT score pre-immersion to calculate the CPM response. A positive CPM response (increase) reflects a hypo-algesic response, whereas a negative response (decrease) reflects a hyper-algesic response.

Strength and Physical measures

1 Multimodal nature of GTPS

2

3 170 Hip abductor and extensor muscle strength were measured using a handheld dynamometer

4

5 171 (Nicholas, Lafayette, IN47903 USA) (Figure 1). The center of the dynamometer was placed 5

6

7

8 172 cm proximal to the lateral malleolus for testing hip abductor strength, and in line with the

9

10 173 posterior aspect of the thigh, 5 cm proximal from the kneecap for testing hip extensor

11

12 174 strength (22). One practice and three experimental trials of maximal voluntary effort were

13

14

15 175 conducted with 30-second rest intervals. Strength was reported as torque standardized to

16

17 176 body weight (Nm/kg).

18

19

20 177

21

22

23 178 >>Insert Figure 1<<

24

25

26 179

27

28

29 180 Physical function was measured through balance (Star Excursion Balance Test (SEBT)) (23)

30

31 181 and time to complete a 20-meter walk test (20MWT), five times sit to stand test (STS), and a

32

33 182 flight of stairs. For the SEBT, participants were asked to maintain single leg stance while

34

35

36 183 maximally reaching with the contralateral leg in anterior, posterolateral, and posteromedial

37

38 184 directions (24). The 20MWT and flight of stairs were performed on a 10-meter course and a

39

40 185 set of 11 stairs that participants were asked to respectively complete/ ascend and descend as

41

42 186 fast as they were able to, without running (25). STS was performed from a seated position,

43

44

45 187 with arms across the chest, and participants stood up and down five times as quickly as

46

47 188 possible (26). One practice trial and two testing trials with 3-minute rests were performed,

48

49 189 and the fastest trial was used for analysis.

50

51

52 190

53

54

55 191 Physical activity time was determined with the Active Australia Survey (27). Participants

56

57

58 192 reported minutes spent in walking, and performing moderate, and vigorous activity during the

59

60

Multimodal nature of GTPS

previous week. Total physical activity was calculated as the sum of minutes spent in these activities.

Self-reported psychosocial outcomes and health related quality of life

Quality of life was measured with the EuroQoL questionnaire (28). Participants rated their current level of disability (severe, moderate or none) in five dimensions (mobility, self-care, usual activities, pain/discomfort, anxiety/depression). Responses were transformed into a single index score ranging from 0 to 1, with higher scores reflecting greater quality of life (28). Health status was measured on a visual analogue scale of the EuroQol, ranging from 0 (worst health you can imagine), to 100 (best health you can imagine). The EuroQol has demonstrated moderate validity and reliability (29, 30).

Pain catastrophizing was quantified with the self-reported Pain Catastrophizing Scale (PCS), that has high reliability (Cronbach's alpha = 0.91) and moderate construct validity compared to the Inventory of Negative Thoughts in Response to Pain (31). Participants indicated the degree to which they experience each of 13 thoughts or feelings when in pain on a 5-point scale, ranging from 0 (not at all), to 4 (all the time) (32). Total scores range from 0-52, higher scores indicating higher levels of pain catastrophizing (32). A total score ≥ 30 was defined as clinically relevant (32).

Anxiety and depression were measured with the Hospital Anxiety and Depression Scale (HADS) (33). Total anxiety and depression scores ranged from 0-21, higher scores indicating higher levels of anxiety and depression. Cases of anxiety and depression were defined by a

1 Multimodal nature of GTPS
2
3 216 total score ≥ 8 (33). The HADS has good reliability (Cronbach’s alpha 0.82 and 0.83) and
4
5 217 validity (33).
6
7
8 218
9
10
11 219 Fear of movement (kinesiophobia) was measured with the Tampa Scale of Kinesiophobia
12
13 220 (TSK) (34), asking participants to rate their agreement on a 4-point Likert scale, ranging from
14
15 221 1 (strongly disagree), to 4 (strongly agree). Total scores range from 17 to 68, with higher
16
17 222 scores indicating greater kinesiophobia. High fear was defined as a total score >37 , low fear
18
19 223 as a total score ≤ 37 (34). The TSK has reported good reliability (Cronbach’s alpha 0.74 -
20
21 224 0.87) and validity (35).
22
23
24
25
26 225
27
28
29 226 Pain self-efficacy was measured with the Pain Self-Efficacy Questionnaire (PSEQ), a
30
31 227 questionnaire used to assess confidence in performing activities while in pain (36).
32
33 228 Participants rate how confidently they can perform activities described on a 7-point Likert
34
35 229 scale, ranging from 0 (not at all confident), to 6 (completely confident). Total scores range
36
37 230 from 0 to 60, where higher scores reflect stronger self-efficacy beliefs (36). Scores $<40/60$
38
39 231 were identified as participants having “less sustainable gains,” as opposed to scores ≥ 40
40
41 232 identifying participants that “could maintain functional gains” (36). The PSEQ is shown to be
42
43 233 a valid and reliable measure (37).
44
45
46
47 234
48
49
50
51 235
52
53
54 236 Study Size
55
56 237 Sample size calculations were done a-priori with G*Power (Universitat Dusseldorf) v3.1(38),
57
58 238 based on values of the remote PPT measure in lateral elbow tendinopathy (39). Based on a
59
60

Multimodal nature of GTPS

moderate effect size of 0.67 (39), power of 0.8, alpha of 0.05, and a 2:1 allocation ratio, we estimated a total sample size of 36 GTPS and 72 control participants.

Statistical Method

Statistical analyses were performed in SPSS version 25.0 (IBM, New York, NY, USA).

Normality was assessed visually (histograms, quantile-quantile plots).

Differences between GTPS and controls were assessed using parametric tests for continuous outcome measures (independent t-tests) and non-parametric tests for ordinal and nominal data (Mann-Whitney U test, Chi-square test). CPM efficacy was analysed using ANCOVA, including baseline hip PPT, water temperature and pain ratings of the conditioning stimulus prior to PPT testing as covariates. Significance was determined with the Benjamini-Hochberg procedure to control for possible false positives caused by multiple comparisons (40). The false discovery rate was set at 0.10 (10%).

Standardized mean differences (SMDs) and 95% confidence intervals (CI) were calculated to facilitate comparison for outcomes with different measurement units (RevMan V5.0; random effect model). Effect sizes were considered as: 0.2 small, 0.6 moderate, 1.2 large and ≥ 2.0 very large (41).

1 Multimodal nature of GTPS

2
3 260 Correlations between pain intensity, number of pain regions and all outcome measures were
4
5 261 calculated to evaluate the degree of association of these commonly regarded co-factors in
6
7 262 presentation of musculoskeletal pain. Pearson correlation coefficients of (negative) 0.0 to 0.3
8
9
10 263 were regarded as negligible, 0.30 to 0.50 low, 0.5 to 0.7 moderate, 0.7 – 0.9 high, and 0.9 to
11
12 264 1.0 very high.

13
14
15 265
16
17
18 266 Multiple linear regression analysis (‘Enter’ method) was performed with variables that were
19
20 267 (a) continuous, (b) significantly impaired compared to controls (variables with highest effect
21
22 268 sizes were chosen if they measured similar constructs), (c) significantly correlated with
23
24 269 VISA-G. Symptom duration and pain intensity during activity were considered possible
25
26 270 confounders. Independent variables were sequentially excluded from the model based on
27
28 271 magnitude of significance. Multicollinearity reflected by a VIF of <2 was considered
29
30 272 respectable.

31
32
33
34
35 273
36
37 274

38
39 275 **Results**

40 276 Forty participants with GTPS and 58 controls were recruited (Figure 2). GTPS and controls
41
42 277 were comparable in age and sex, the majority being female (Table 1). GTPS participants had
43
44 278 a significantly greater BMI, waist girth and hip circumference compared to controls (Table
45
46 279 1). Within the GTPS group, 35% had a normal BMI, 32.5% were classified overweight,
47
48 280 17.5% class I obesity, 7.5% class II obesity, and 7.5% class III obesity compared to 3.4%
49
50 281 underweight, 63.8% normal, 22.4% overweight and 10.3% class I obesity in controls (p<0.01,
51
52 282 Table 1)(42).

53
54
55
56
57 283
58
59 284 >>Insert Figure 2<<
60

Multimodal nature of GTPS

>>Insert Table 1<<

Outcome measures

Sensory, physical and psychological outcomes, and mean differences (95% CI) are presented in Table 2 and Supplementary Material 1 outlines significant p-values.

>>Insert Table 2<<

Quantitative Sensory Testing

Compared to controls, GTPS displayed large differences in local PPT and small differences in remote PPT. Small differences in local and remote CPT (GTPS displaying cold hyperalgesia) were noted. HPT and temporal summation did not differ between groups.

CPM was measured for 39 GTPS and 41 controls (Figure 3). One GTPS and three control participants did not consent to CPM testing, three controls could not be measured due to time constraints or skin condition, and nine controls reached >900kPa during PPT or CPM testing. Moderate differences were found between groups, with GTPS displaying smaller changes in PPT (Table 2, Figure 4).

Low associations were found for worst pain and local HPT ($r = -0.37$, $p = 0.02$; as was average pain), local temporal summation ($r = 0.48$, $p < 0.01$) and local PPT ($r = -0.37$, $p = 0.02$). No significant associations were found between the number of pain regions and any of the QST measures including temporal summation and CPM.

Multimodal nature of GTPS

>>Insert Figure 3 and Figure 4<<

Strength and Physical measures

GTPS showed large differences in STS performance and moderate differences for unilateral hip abductor strength, 20MWT, flight of stairs, and SEBT performance (all directions) compared to controls. A small reduction in hip extensor strength was observed (Table 2, Figure 4).

Significant but small group differences were found for total physical activity time and vigorous physical activity in the previous week (Table 2, Figure 4).

Low associations were found for worst pain and SEBT posteromedial ($r = -0.35$, $p = 0.04$; as was average pain), flight of stairs ($r = 0.47$, $p < 0.01$; as was average pain $r = 0.41$, $p = 0.01$), STS ($r = 0.41$, $p = < 0.01$), and extensor strength ($r = -0.36$, $p = 0.03$). No significant associations were found between the number of pain regions and any strength or physical measures.

Self-reported psychological distress and quality of life

Compared to controls, GTPS displayed large differences in quality of life, health status and moderate differences in anxiety and depression scores. No significant group differences were found for pain catastrophisation (Table 2, Figure 4). Within the GTPS group, 25% were classified with anxiety, 5% with depression, compared to 5.2% and 3.4% of controls respectively. (Table 2). Within the GTPS group, 87.5% were able to ‘maintain functional gains’ (PSEQ scores), whereas 57.5% had a high degree of kinesiophobia (Table 2). These measures were not collected in the control group.

Multimodal nature of GTPS

Low associations were found for worst pain and anxiety ($r = -0.33$, $p = 0.04$), health status ($r = -0.48$, $p < 0.001$; as was average pain $r = -0.49$, $p < 0.001$), and for the number of pain sites and pain catastrophising ($r = 0.43$, $p < 0.01$). High associations were found for worst pain and quality of life ($r = -0.75$, $p < 0.001$; as was average pain $r = -0.73$, $p < 0.001$).

Factors associated with pain and disability (VISA-G)

The final multivariate regression model explained 26% of the variance of VISA-G. Cases with evidence of marginal significance were retained in the final model. Based on the non-standardised coefficients, one unit increase in depression on the HADS is likely to decrease VISA-G by 1.67 units, one N.m.kg⁻¹ increase in abductor strength is likely to increase VISA-G by 12.64 units, and one second increase in time to complete stairs is likely to decrease VISA-G by 0.53 units (Supplementary Material 2).

Discussion

This study aimed to quantify differences in physical, sensory, and psychosocial outcome measures in participants with GTPS compared to pain-free controls, and to explore the factors associated with pain and disability. Large effects were found for health-related quality of life, hip PPT, and STS; moderate effects for CPM, hip abductor strength, all other functional measures, anxiety and depression; small effects for elbow PPT, hip and elbow CPT, hip extensor strength and physical activity time. Notably we did not find any differences in temporal summation of pain between groups. VISA-G scores were explained by a combination of depression, hip abductor strength and time to complete stairs.

1 Multimodal nature of GTPS

2
3 356 This study found evidence of widespread mechanical and cold hyperalgesia. The degree of
4
5 357 impairment in PPT was greater at the local tendon (SMD -1.49) than the elbow (SMD -0.58),
6
7 358 and small for local and remote CPT (SMD -0.47 and -0.39). Previous literature reported
8
9 359 evidence of *local* mechanical and thermal hyperalgesia in some, but not all lower
10
11 360 tendinopathy studies (7-9, 43). *Widespread* mechanical and cold hyperalgesia have been
12
13 361 reported in Achilles and lateral elbow tendinopathy (9, 39), but not in other studies (43). A
14
15 362 recent cross-sectional study in persistent GTPS reported significant group differences in PPT
16
17 363 locally, bilaterally and remotely, inferring primary and secondary mechanical hyperalgesia in
18
19 364 GTPS (8). Comparison of our study results with previous literature shows inconsistencies in
20
21 365 findings. High quality methodological studies with a larger sample size that explore the
22
23 366 involvement of pain sensitivity in persistent tendinopathies like GTPS are warranted.
24
25
26
27
28

29 367
30
31
32 368 CPM is an established paradigm used to study endogenous pain inhibitory mechanisms. It is
33
34 369 thought to reflect the efficacy of the diffuse noxious inhibitory control system (DNIC) (44), a
35
36 370 spinal-medullary-spinal pathway that triggers pain inhibition when two painful stimuli are
37
38 371 applied simultaneously (45). In humans, CPM may reflect peripheral neural mechanisms,
39
40 372 such as chronic inflammation and hyper-excitability (46) as well as higher cortical centres,
41
42 373 which are also affected by age, sex, menstruation cycle, and psychological features (45). We
43
44 374 found moderate differences in CPM efficacy, possibly reflecting alterations in DNIC,
45
46 375 however we did not find evidence of altered facilitatory ascending mechanisms in GTPS (i.e.,
47
48 376 temporal summation was not different between groups). We could speculate that persistent
49
50 377 pain causes ongoing activation of pain inhibitory mechanisms, leaving patients unable to
51
52 378 further activate these systems, resulting in a ceiling effect. This has been previously proposed
53
54 379 in other persistent pain conditions (47).
55
56
57
58
59
60

Multimodal nature of GTPS

380

381 The presence of widespread pain sites is often considered linked with central nervous system
382 sensitization. We did not find this in our cohort, which is consistent with a study of knee
383 osteoarthritis (48), but not others. A recent cross-sectional study including 49 participants
384 with GTPS reported significant associations of widespread pain distribution with less CPM
385 and increased temporal summation (13). The association between widespread pain and less
386 CPM has also been reported in low back pain (11, 49). It might be that a subgroup of
387 individuals who have GTPS and widespread pain may be more likely to experience central
388 nervous system sensitization. Future research should further assess if subgroups of
389 individuals with different severity of GTPS vary according to their pain distribution. This
390 may lead to improved patient specific pain management.

391

392 The VISA-G is a self-reported questionnaire concerning pain and functional disability
393 specifically designed for GTPS (20). Linear regression revealed a strong association between
394 VISA-G and depression, implying depression levels to be an indicator of severity of self-
395 reported pain and disability. High quality evidence shows depression levels to be associated
396 with many other chronic musculoskeletal pain conditions (50, 51), and reported depression to
397 be a possible moderator in the relationship with chronic pain (52). Further, positive
398 behaviours like resilience, optimism and pain acceptance are thought to positively affect
399 chronic pain (53, 54). Thus, the interpretation of self-reported pain and disability
400 questionnaires, such as VISA-G, should take this association with levels of depression into
401 account.

402

Multimodal nature of GTPS

The limitations of this study include the unblinded assessor, although a standardized protocol was rigorously followed. Only 80% of the calculated sample size for controls was recruited due to time constraints. Further, our population was highly educated (>65% of both groups had at least graduated from University or polytechnic) and predominantly female, which should be taken into account when generalizing findings. QST requires operator skill, however, members of this research lab have considerable experience in QST studies. Future research should aim to identify the prognostic significance of physical, sensory and psychological characteristics on pain and disability in persistent GTPS before targeted treatments can be developed. In addition, the impact of social factors and environment should be further investigated to understand the patient characteristics and prognosis of individuals with GTPS.

Conclusions

This study shows that compared to controls, people with persistent GTPS exhibit impaired health related quality of life, physical strength and functioning, widespread hyperalgesia, and altered psychological factors. No differences were seen in heat hyperalgesia, temporal summation or pain catastrophizing. Self-reported pain and disability was associated with depression, hip abductor strength and time to complete stairs. The connections between biopsychological features and persistent pain in GTPS should be considered in order to and inform management approaches.

Acknowledgments and Conflict of Interest

This study was supported by the National Health and Medical Research Council (NHMRC) Program Grant (#631717). MLP is supported by the International Postgraduate Research

Multimodal nature of GTPS

427 Scholarship (IPRS)/University of Queensland Centennial Scholarship (UQcent). Authors

428 have no conflicts of interest to declare.

429

For Review Only

Multimodal nature of GTPS

References

1. Albers IS, Zwerver J, Diercks RL, Dekker JH, Van den Akker-Scheek I. Incidence and prevalence of lower extremity tendinopathy in a Dutch general practice population: a cross sectional study. *BMC Musculoskelet Disord.* 2016;17:16.

2. Fearon AM, Cook JL, Scarvell JM, Neeman T, Cormick W, Smith PN. Greater trochanteric pain syndrome negatively affects work, physical activity and quality of life: a case control study. *J Arthroplasty.* 2014;29(2):383-6.

3. Brosschot JF. Cognitive-emotional sensitization and somatic health complaints. *Scandinavian Journal of Psychology.* 2002;43:113-21.

4. Das De S, Vranceanu AM, Ring DC. Contribution of kinesophobia and catastrophic thinking to upper-extremity-specific disability. *J Bone Joint Surg Am.* 2013;95(1):76-81.

5. Mallows A, Debenham J, Walker T, Littlewood C. Association of psychological variables and outcome in tendinopathy: a systematic review. *Br J Sports Med.* 2016;51(9):743-8.

6. Heales LJ, Lim EC, Hodges PW, Vicenzino B. Sensory and motor deficits exist on the non-injured side of patients with unilateral tendon pain and disability--implications for central nervous system involvement: a systematic review with meta-analysis. *Br J Sports Med.* 2014;48(19):1400-6.

7. Plinsinga ML, Brink MS, Vicenzino B, van Wilgen P. Evidence of Nervous System Sensitization in Commonly Presenting and Persistent Painful Tendinopathies: A Systematic Review. *J Orthop Sports Phys Ther.* 2015;45(11):1-34.

8. French HP, Jong CC, McCallan M. Do features of central sensitisation exist in Greater Trochanteric Pain Syndrome (GTPS)? A case control study. *Musculoskelet Sci Pract.* 2019;43:6-11.

9. Tompra N, van Dieen JH, Coppieters MW. Central pain processing is altered in people with Achilles tendinopathy. *Br J Sports Med.* 2015;0:1-6.

10. Arendt-Nielsen L, Nie H, Laursen MB, Laursen BS, Madeleine P, Simonsen OH, et al. Sensitization in patients with painful knee osteoarthritis. *Pain.* 2010;149(3):573-81.

11. Gerhardt A, Eich W, Treede RD, Tesarz J. Conditioned pain modulation in patients with nonspecific chronic back pain with chronic local pain, chronic widespread pain, and fibromyalgia. *Pain.* 2017;158(3):430-9.

12. Mlekusch S, Neziri AY, Limacher A, Juni P, Arendt-Nielsen L, Curatolo M. Conditioned Pain Modulation in Patients With Acute and Chronic Low Back Pain. *Clin J Pain.* 2016;32(2):116-21.

13. Ferrer-Pena R, Munoz-Garcia D, Calvo-Lobo C, Fernandez-Carnero J. Pain Expansion and Severity Reflect Central Sensitization in Primary Care Patients with Greater Trochanteric Pain Syndrome. *Pain Med.* 2019;20(5):961-70.

14. Grimaldi A, Mellor R, Hodges P, Bennell K, Wajswelner H, Vicenzino B. Gluteal Tendinopathy: A Review of Mechanisms, Assessment and Management. *Sports Med.* 2015;45(8):1107-19.

15. Allison K, Vicenzino B, Wrigley TV, Grimaldi A, Hodges PW, Bennell KL. Hip Abductor Muscle Weakness in Individuals with Gluteal Tendinopathy. *Med Sci Sports Exerc.* 2015;48(3):346-52.

16. Fearon A, Neeman T, Smith P, Scarvell J, Cook J. Pain, not structural impairments may explain activity limitations in people with gluteal tendinopathy or hip osteoarthritis: A cross sectional study. *Gait Posture.* 2017;52:237-43.

17. Ganderton C, Pizzari T, Harle T, Cook J, Semciw A. A comparison of gluteus medius, gluteus minimus and tensor fascia latae muscle activation during gait in post-menopausal women with and without greater trochanteric pain syndrome. *J Electromyogr Kinesiol.* 2017;33:39-47.

18. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *The Lancet.* 2007;370:1453-7.

19. Grimaldi A, Mellor R, Nicolson P, Hodges P, Bennell K, Vicenzino B. Utility of clinical tests to diagnose MRI-confirmed gluteal tendinopathy in patients presenting with lateral hip pain. *Br J Sports Med.* 2016;0:1-7.

Multimodal nature of GTPS

20. Fearon AM, Ganderton C, Scarvell JM, Smith PN, Neeman T, Nash C, et al. Development and validation of a VISA tendinopathy questionnaire for greater trochanteric pain syndrome, the VISA-G. *Man Ther.* 2015;20(6):805-13.
21. Arendt-Nielsen L, Andresen T, Malver LP, Oksche A, Mansikka H, Drewes AM. A Double-blind, Placebo-controlled Study on the Effect of Buprenorphine and Fentanyl on Descending Pain Modulation. A Human Experimental Study. 2012.
22. Ireland ML, Wilson JD, Ballantyne BT, Davies IM. Hip Strength in Females With and Without Patellofemoral Pain. *Journal of Orthopaedic & Sports Physical Therapy.* 2003;33(11):671-6.
23. Gribble PA, Hertel J, Plisky P. Using the Star Excursion Balance Test to assess dynamic postural-control deficits and outcomes in lower extremity injury: a literature and systematic review. *J Athl Train.* 2012;47(3):339-57.
24. Robinson R, Gribble P. Kinematic Predictors of Performance on the Star Excursion Balance Test. *Journal of Sport Rehabilitation.* 2008.
25. Simonsick EM, Gardner AW, Poehlman ET. Assessment of physical function and exercise tolerance in older adults: reproducibility and comparability of five measures. *Aging Clin Exp Res.* 2000;12:274-80.
26. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *Journal of Gerontology.* 1994;49(2):M85-M94.
27. Australian Institute of Health and Welfare. The Active Australia Survey: A guide and manual for implementation, analysis and reporting; 2004.
28. Reenen van M, Oppe M. EQ-5D-3L UserGuide. Rotterdam: EuroQol Group. 2015.
29. Fransen M, Edmonds J. Reliability and validity of the EuroQol in patients with osteoarthritis of the knee. *Rheumatology.* 1999;38:807-13.
30. Janssen MF, Pickard AS, Golicki D, Gudex C, Niewada M, Scalone L, et al. Measurement properties of the EQ-5D-5L compared to the EQ-5D-3L across eight patient groups: a multi-country study. *Qual Life Res.* 2013;22(7):1717-27.
31. Osman A, Barrios FX, Kopper BA, Hauptmann W, Jones J, O'Neill E. Factor structure, reliability, and validity of the pain catastrophizing scale. *Journal of Behavioral Medicine.* 1997;20(6):589-605.
32. Sullivan MJL, Bishop J. The Pain Catastrophizing Scale: Development and Validation. *American Psychological Association.* 1995;7(4):524-32.
33. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *Journal of Psychosomatic Research.* 2002;52:69-77.
34. Vlaeyen JWS, Kole-Snijders AMJ, Boeren RGB, van Eek H. Fear of movement/(re)injury in chronic low back pain and its relation to behavioral performance. *Pain.* 1995;62:363-72.
35. Larsson C, Hansson EE, Sundquist K, Jakobsson U. Psychometric properties of the Tampa Scale of Kinesiophobia (TSK-11) among older people with chronic pain. *Physiother Theory Pract.* 2014;30(6):421-8.
36. Nicholas MK. The pain self-efficacy questionnaire: Taking pain into account. *Eur J Pain.* 2007;11(2):153-63.
37. Tonkin L. The pain self-efficacy questionnaire. *Aust J Physiother.* 2008;54(1):77.
38. Faul F, Erdfelder E, Lang A, Buchner A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods.* 2007;39(2):175-91.
39. Coombes BK. Thermal Hyperalgesia Distinguishes Those With Severe Pain and Disability in Unilateral Lateral Epicondylalgia. *Clin J Pain.* 2012;28:595-601.
40. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of Royal Statistical Society* 1995;57(1):289-300.

Multimodal nature of GTPS

41. Hopkins WG. A new view of statistics. 2013 [cited 11/08/2018]; Available from: <http://www.sportsci.org/resource/stats/index.html>

42. WHO. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee.; 1995.

43. Plinsinga ML. Patellar and Achilles tendinopathies are predominantly peripheral pain states: a blinded case control study of somatosensory and psychological profile. *Br J Sports Med*. 2018;52(5):284-91.

44. Lewis GN, Rice DA, McNair PJ. Conditioned pain modulation in populations with chronic pain: a systematic review and meta-analysis. *J Pain*. 2012;13(10):936-44.

45. van Wijk G, Veldhuijzen DS. Perspective on diffuse noxious inhibitory controls as a model of endogenous pain modulation in clinical pain syndromes. *J Pain*. 2010;11(5):408-19.

46. Nir RR, Yarnitsky D. Conditioned pain modulation. *Curr Opin Support Palliat Care*. 2015;9(2):131-7.

47. Lannersten L, Kosek E. Dysfunction of endogenous pain inhibition during exercise with painful muscles in patients with shoulder myalgia and fibromyalgia. *Pain*. 2010;151(1):77-86.

48. Lluch Girbes E, Duenas L, Barbero M, Falla D, Baert IAC, Meeus M, et al. Expanded distribution of pain as a sign of central sensitization in individuals with symptomatic knee osteoarthritis. *Physical Therapy*. 2016;96(8):1196-207.

49. Gerhardt A, Eich W, Janke S, Leisner S, Treede RD, Tesarz J. Chronic Widespread Back Pain is Distinct From Chronic Local Back Pain: Evidence From Quantitative Sensory Testing, Pain Drawings, and Psychometrics. *Clin J Pain*. 2016;32(7):568-79.

50. MacLachlan LR, Collins NJ, Matthews MLG, Hodges PW, Vicenzino B. The psychological features of patellofemoral pain: a systematic review. *Br J Sports Med*. 2017;0:1-11.

51. Stubbs B, Aluko Y, Myint PK, Smith TO. Prevalence of depressive symptoms and anxiety in osteoarthritis: a systematic review and meta-analysis. *Age Ageing*. 2016;45(2):228-35.

52. Vadivelu N, Kai AM, Kodumudi G, Babayan K, Fones M, Burg MM. Pain and psychology - A reciprocal relationship. *Ochsner Journal*. 2017;17:173-80.

53. Ahmed SA, Shantharam G, Eltorai AEM, Hartnett DA, Goodman A, Daniels AH. The effect of psychosocial measures of resilience and self-efficacy in patients with neck and lower back pain. *Spine J*. 2019;19(2):232-7.

54. Ferreira VM, Sherman AM. The relationship of optimism, pain and social support to well-being in older adults with osteoarthritis. *Aging Ment Health*. 2007;11(1):89-98.

55. Crawford JO. The Nordic Musculoskeletal Questionnaire. *Occupational Medicine*. 2007;57(4):300-1.

Multimodal nature of GTPS

Tables

Table 1. Participant characteristics of the GTPS and the control group. Data represent mean (SD) for each group unless otherwise indicated, with group mean differences (95% CI; p-value). Significant group differences are indicated by *.

	GTPS group n = 40	Control group n = 58	Group difference
Age years	51 (9)	53 (11)	-1.2 (-5.5, 3.1; p=0.59)
Sex female n, (%) [*]	38 (95%)	55 (95%)	p = 0.97
Height cm	165.7 (6.8)	165.7 (5.7)	0 (-0.0, 0.0; 1.0)
Body mass kg	78.3 (18.5)	66.2 (12.8)	12 (5.7, 18.3; p<0.001*)
Body Mass Index kg/m ²	28.5 (6.2)	24.0 (4.1)	4.4 (2.3, 6.5; p<0.001*)
Underweight (BMI <18.5)	0%	3.4%	p <0.01*†
Normal (18.5–24.9)	35%	63.8%	
Overweight (25.0–29.9)	32.5%	22.4%	
Obese class I (30.0–34.9)	17.5%	10.3%	
Obese class II (35.0–39.9)	7.5%	-	
Obese class III (≥40)	7.5%	-	
Waist girth cm	94.7 (14.0)	84.12 (13.3)	10.6 (5.1, 16.2; p<0.001*)
Hip circumference cm	110.5 (13.9)	101.4 (10.2)	9.1 (4.3, 14.0; p<0.001*)
Highest education†			p = 0.28
Less than 3y high school n, (%)	1 (2.5)	1 (1.7%)	
3y or more high school n, (%)	5 (12.5%)	2 (8.6%)	
Some tertiary training n, (%)	8 (20%)	9 (15.5%)	
Graduated from University or polytechnic n, (%)	11 (27.5%)	19 (32.8%)	
Any post-graduate study n, (%)	15 (37.5%)	24 (41.4%)	
Work status†			p = 0.91
Employed for wages n, (%)	26 (65%)	35 (60.3%)	
Self-employed n, (%)	4 (10%)	3 (5.2%)	
Out of work n, (%)	1 (2.5%)	2 (3.4%)	
A homemaker n, (%)	3 (7.5%)	1 (1.7%)	
Student n, (%)	6 (15%)	4 (6.9%)	

Multimodal nature of GTPS

Retired n, (%)	0	13 (22.4%)	
Hip pain interference with sleep quality			-
Not at all n, (%)	4 (10%)	-	
Occasionally n, (%)	9 (22.5%)	-	
Some of the time n, (%)	16 (40%)	-	
Most of the time n, (%)	7 (17.5%)	-	
All of the time n, (%)	4 (10%)	-	
Duration of symptoms months, median (IQR)	18 (15)	-	-
Unilateral n, (%)	28 (70%)	-	-
VISA-G (0-100)	63 (11)	-	-
Pain severity in past week (PNRS 0-10)			
Average, median (IQR)	4 (3)	-	-
Worst, median (IQR)	7 (3)	-	-
During activity, median (IQR)	5 (4)	-	-
Number of pain regions, median (IQR)‡	3 (2)		

Abbreviations: BMI, body mass index; GT, gluteal tendinopathy; IQR, interquartile range; PNRS, pain numeric rating scale; SD, standard deviation; VISA-G, Victorian Institute of Sports Assessment – Gluteal tendinopathy; 3y, 3 years.

*significant p level set on p = 0.05

†Group differences calculated with the χ^2 test.

‡Number of pain regions recorded with The Nordic Musculoskeletal Questionnaire (55).

Multimodal nature of GTPS

Table 2. Mean (SD) for quantitative sensory testing, strength and physical tests, psychological self-reported outcome measures, and quality of life in the GTPS and control groups, with group mean differences (95% CI; p-value).

	GTPS group n = 40	Control group n = 58	Group differences
QST measures			
Cold pain threshold hip °C	10 (7.52)	7.02 (5.24)	2.97 (0.41, 5.53; 0.03*)
Cold pain threshold elbow °C	10.40 (7.77)	7.79 (5.85)	2.60 (0.13, 5.53; 0.08*)
Heat pain threshold hip °C	45.09 (2.84)	46.09 (2.76)	-1.00 (-2.14, 0.14; 0.09)
Heat pain threshold elbow °C	46.05 (3.59)	46.82 (2.80)	-0.78 (-2.06, 0.50; 0.23)
Pressure pain threshold hip kPa	203.25 (71.99)	395.88 (155.59)	-192.63 (-245, -140.26; <0.001*)
Pressure pain threshold elbow kPa	317.83 (133.81)	396.48 (134.96)	-78.66 (-133.53, -23.79; <0.01*)
Temporal summation hip /10 PNRS	1.33 (1.01)	1.16 (1.33)	-0.17 (-0.67 to 0.34; 0.51)
Temporal summation elbow /10 PNRS	0.58 (0.69)	0.73 (1.08)	0.15 (-0.21 to 0.50; 0.42)
Conditioned pain modulation hip kPa†	48.96 (51.48)	117.98 (79.31)	-59.79 (-98.22, -21.36; <0.01*)
Strength and physical measures			
Hip abductor strength N.m.kg ⁻¹	0.70 (0.25)	1.00 (0.31)	-0.30 (-0.42, -0.18; <0.001*)
Hip extensor strength N.m.kg ⁻¹	0.42 (0.20)	0.58 (0.30)	-0.16 (-0.27, -0.05; <0.01*)
Star Excursion Balance Test			
Anterior	0.52 (0.07)	0.58 (0.07)	-0.05 (-0.08, -0.03; <0.001*)
Posterolateral	0.78 (0.12)	0.86 (0.13)	-0.08 (-0.13, -0.03; <0.01*)
Posteromedial	0.66 (0.13)	0.77 (0.14)	-0.11 (-0.16, -0.05; <0.001*)
Sit to stand test s	11.98 (3.15)	8.76 (2.23)	3.22 (2.15, 4.30; <0.001*)
20 meter walk test s	9.84 (2.87)	7.42 (1.26)	2.42 (1.58, 3.27; <0.001*)
Flight of stairs test s	20.94 (5.47)	16.93 (3.64)	4.01 (2.18, 5.84; <0.001*)
Total physical activity min/week	465.38 (472.25)	677.16 (502.27)	211.78 (11.76, 411.80; 0.04*)
Vigorous physical activity min/week	98.25 (163.13)	166.98 (182.09)	68.73 (-2.54, 140.01; 0.06*)
Psychosocial self-reported outcome measures			
Pain catastrophizing scale			

Multimodal nature of GTPS

Total score /52	7.23 (6.56)	5.55 (7.0)	1.67 (-1.10, 4.45; 0.23)
Clinical level n, (%)‡	1 (2.5%)	1 (1.7%)	p = 0.79
Non clinical level n, (%)‡	39 (97.5%)	57 (98.3%)	

Hospital Anxiety and Depression

Scale: Anxiety	5.38 (3.43)	3.43 (2.96)	1.94 (0.66, 3.23; <0.01*)
Total score /21	10 (25%)	3 (5.2%)	p<0.01
Cases n, (%)‡	30 (75%)	55 (94.8%)	
Non-cases n, (%)‡			

Hospital Anxiety and Depression

Scale: Depression	3.15 (2.476)	1.52 (2.096)	1.63 (0.71, 2.55; 0.001*)
Total score /21	2 (5%)	2 (3.4%)	p = 0.70
Cases n, (%)‡	38 (95%)	56 (96.6%)	
Non-cases n (%)‡			

Pain self-efficacy

Total score /60	49.58 (8.258)	-	-
Less sustainable gains n, (%)	5 (12.5%)	-	-
Maintenance of function n, (%)	35 (87.5%)	-	-

Tampa Scale of Kinesiophobia

Total score /68	35.93 (4.599)	-	-
High degree of kinesiophobia n, (%)	23 (57.5%)	-	-
Low degree of kinesiophobia n, (%)	17 (42.5%)		

EQ-5D-3L

Index score /1	0.72 (0.14)	0.98 (0.06)	-0.26 (-0.30, -0.22; <0.001*)
Self-reported health /100 VAS	68.68 (15.50)	84.21 (10.16)	-15.53 (-20.67, -10.39; <0.001*)

Abbreviations: GT, gluteal tendinopathy; PNRs, pain numeric rating scale SD, standard deviation; VAS, visual analogue scale.

*Significance calculated with the Benjamini-Hochberg procedure (Supplementary Material 1).

‡Group differences calculated with an ANCOVA with GT n = 39 and HC n = 41.

‡Group differences calculated with the χ^2 test.

Multimodal nature of GTPS

Figure legends

Figure 1. Positioning of the participant during measurements of hip abductor muscle strength (upper image) and hip extensor muscle strength (lower image). Hip abductor muscle strength was measured with the test knee extended and the hip at 10° abduction and 0° flexion, and the opposite leg flexed at the hip (45°) and knee (90°). Hip extensor strength was measured with both hips in 30° flexion and both knees in 45° flexion. In both tests, the pelvis was stabilized with a seatbelt and towel, strapped to the plinth, and the participant held the side(s) of the plinth with both hands.

Figure 2. Flowchart of recruitment.

Figure 3. Individual data points of the greater trochanteric pain syndrome (GTPS) group (blue colour scheme; n = 39) and control group (grey colour scheme; n = 41) and the group average (black line) for the pressure pain threshold (PPT) measure pre conditioning stimulus and the PPT max (maximum of PPT during/after).

Figure 4. Forest plot including standardized mean differences (SMD) and 95% confidence intervals (CI) for all outcome measures. Quantitative sensory testing on the hip and elbow (cold pain threshold (CPT), heat pain threshold (HPT), pressure pain threshold (PPT) and conditioned pain modulation (CPM)); strength (abductor and extensor), physical function (star excursion balance test (SEBT), time to complete 20 meter walk test, sit to stand, flight of stairs) and physical activity (Active Australia Survey (AAS)); psychological outcome measures (the Hospital Anxiety and Depression Scale (HADS)), and the Pain Catastrophizing Scale (PCS)); health related quality of life.

Multimodal nature of GTPS

Figures

Figure 1.

Multimodal nature of GTPS

Figure 2.

Multimodal nature of GTPS

Figure 3.

Figure 4.