Title

Acknowledging cancer as a ‘family disease’: A systematic review of family research during cancer.

Abstract

Purpose

Family members are a part of the team to improve the outcomes of the person with cancer.

Families require support and information to optimise their care, however, their needs are often unacknowledged and within clinical areas there is a lack of family focused interventions. Studies highlight families’ needs but lack a family representation. The aim was to explore research with family as the unit-of-care during cancer treatment.

Method

The Pickering systematic quantitative literature review method; a 15-step process from searching, database development and analysis was followed. Research published 2008-2019 within databases: MEDLINE, SCOPUS, PsycINFO, Cochrane, CINAHL; key words, ‘family* or caregiver*, and cancer*, neoplasm* and coping*, distress* in November 2019. Quality assessment completed using Mixed Methods Appraisal Tool, descriptive quantitative analysis and thematic analysis.

Results

Studies involving patients and family members were included in the review (N=73). The analysis identified participants had a mean age of 58 years and primarily lung, breast or prostate cancer. Over 80% were from America and European countries; 93% had a dyad sample, quantitative

studies (76%). There was eight intervention studies between four to sixteen weeks long, focused on family wellbeing. Themes described the impact of cancer on the whole family, the importance of communication between family members, and resources for family members.

Conclusion

The review identified four main scales and optimum intervention styles. Family research in the adult cancer needs to focus on intervention studies, increase international focus and inclusion of other family members such as children, friends and older adults.

Tweetable abstract: Acknowledging cancer as a family disease, analysing the current research, highlighting gaps and best practice

Key words: Cancer, Family, Family Nursing, Interventions, Literature review, Nursing

Introduction

A diagnosis of cancer will influence the person with cancer and their family over a long period of time (Finley, 2018). The supportive role of family in the care of person with cancer is well recognized, including both physical and psychosocial support (Gilliss et al., 2019), and may begin even before the cancer diagnosis is made, and continue through different trajectories of the cancer journey (Coyne et al., 2017; Rha et al., 2015). Family caregivers include partners, close or extended family members and friends (Romito et al., 2013). Family caregiving varies according to severity of symptoms, treatments required, stage of disease, or closeness to death; taking up to 14-24 months during the active treatment phase; averaging 8.8 hours/day, and the family caregiving process may last up to 4 years (Given et al., 2012). The increase in survival of oncology patients can result in longer term of caregiving for family members.

As cancer care has transitioned to have more of a focus on outpatient and home-based care, the family caregiver has become the main provider of support to the person with cancer (Ferrell & Wittenberg, 2017). Research has highlighted, that cancer will influence both the person with cancer and their family equally with significant burden from depression, physical health changes, declines in social functioning and reduction in quality of life (QoL) (Dieperink et al., 2018; Ferrell & Wittenberg, 2017). An individual caregiver's perception of caregiving demand, and use of coping mechanisms are influenced by their own needs and also the person with cancer's symptoms and dependence on them as family caregiver (Stamataki et al., 2014). The needs of family caregivers vary according to the age, sex, ethnicity, education and the person's disease progression and are connected to the person they are caring for (Andersen et al., 2019; Johansen et al., 2018). The consequences of caregiving may include physical problems such as fatigue, sleep disturbances, loss of weight and appetite, headaches, and psychological

distress including anxiety and depression, but caregiving can be both a positive and negative experience with the family having changes to social activities and relationships (Rha et al., 2014, 2015; Romito et al., 2013).

The definition of family caregiver, often referred as informal caregiver, varies across studies (Bell & Wright, 2015). A universal definition of a family caregiver or informal caregiver is the person or persons who are not paid for the care they provide (Australian Commission on Safety and Quality in Health Care, 2018; Shajani & Snell, 2019). The type of support provided by caregivers includes emotional, medical, tangible and practical support (Romito et al., 2013). The support provided involves different aspects of person with cancer's care such as daily life assistance, symptom management, monitoring treatment complications, providing accommodation, scheduling visits, housekeeping, meal preparation and financial support (Rha et al., 2015). The family caregiving role has received increased attention in the last decade related not only to psychosocial aspects of care but also economic influences (Girgis & Lambert, 2017; Zaleta et al., 2018).

Recent research suggests caregivers and patients should be conceptualized as a family unit rather than caregivers as individuals separate from patients (Bell & Wright, 2015). Caregiver-patient congruence and reciprocal influence highlight the effects of patients and family on each other (Fletcher et al., 2011; K. S. Lyons et al., 2016). To understand the complexity and implications of patient and caregivers' appraisal of cancer management, a family as the unit care is the best approach (Coyne et al., 2017). Some researchers suggest that consideration of interpersonal relationships and complexity of family networks should be taken into account when conducting research, developing interventions and supporting persons with cancer (Illingworth et al., 2009).

Systematic reviews have explored aspects of family research in chronic disease including healthcare interventions and caregiver-mediated interventions (Deek et al., 2016; Fiest et al., 2018), and a recent scoping review on family research for older patients with cancer is published (Konradsen et al., 2020). The role of family in chronic disease management is established (Gilliss et al., 2019; Whitehead et al., 2018), however, there has not been a mapping of research specific to adults with cancer including their family members. The aim of this review is to explore studies with a focus of family as the unit of care during cancer treatment for all adult age groups including, prevention, treatment, supportive care, rehabilitation and early palliative care and to identify research gaps.

The systematic quantitative literature review approach

To fully explore and understand the scope of research available a systematic quantitative literature review was conducted. This type of review differs from a traditional systematic review as it allows for the inclusion and analysis of both quantitative and qualitative studies in the review (Pickering & Byrne, 2014). It also differs from an integrative review by mapping the data from the studies within a database. To guide this process a systematic quantitative literature review framework was used (Pickering & Byrne, 2014). The Pickering method guides the development of a database of variables from the included articles allows for comparison of variables within the database and trends in data (Pickering et al., 2015). The steps in a systematic quantitative review are to define the topic, inclusion and exclusion criteria, identify databases, complete literature searches, download into endnote, complete study assessment, finalise articles for inclusion, prepare database, enter first ten percent of papers and refine database, headings, complete binary data entry of all articles, complete quantitative analysis and qualitative analysis (Pickering & Byrne, 2014).

Method

The following method for the systematic quantitative literature review is described according to the Pickering framework (Pickering & Byrne, 2014):

Pickering framework Step 1 – identify the topic, Step 2 Formulate research question,

A modified PICO table [population, intervention, comparator, outcome] was developed to capture the range of terms used for family and disease states (Carman et al., 2013). Research question: What are the types and common themes of research with adults with cancer and their family members as participants?

Population: Research that included data from both adult patients with cancer and their family caregivers were considered for the study. A definition of family research was developed to capture the range of research available. Family research was defined as data collected from both patients and family members (Ferrell & Wittenberg, 2017; Shajani & Snell, 2019).

Intervention: All interventions used in previous studies that focused on the family unit were included. *Comparator:* All comparator types identified in original studies were included. Descriptive studies without comparators were also included. *Outcomes:* The outcomes of this review included the family research methods, sample, research instruments used and common themes across these studies. The search focussed on research during cancer treatment [within first year] but excluded end of life as much research is on managing supportive care for the person with cancer rather than the family as unit of care (Oakley et al., 2019).

Pickering framework Step 3 – identify key words, Step 4 – identify database and Step 5 – complete searches

Searches were conducted by two researchers (JJ, EC) across MEDLINE, SCOPUS (Web of Science), PsycINFO, Cochrane, CINAHL databases using the key words, ‘family* or caregiver*, and cancer*, neoplasm* and coping*, distress* in November 2019. Searches were limited to English language and published between 2008 - 2019. Ganong (2011) conducted a review of family nursing research and highlighted the development of theory and interventions, the current time span enables a view of current trends in family research in cancer building from previous literature reviews (Bell, 2014). Key words were developed in consultation with a health librarian to ensure relevant articles were included. The reference lists of included articles and literature reviews were searched (Moher et al., 2009). Inclusion criteria were primary research, peer reviewed, full text, written in English and had a sample of adult patients with cancer and their family. Exclusion was paediatric patients, or end of life focus.

The initial search was downloaded in EndNote X9 into database groups, duplicates were removed. A stepped process of assessment within EndNote using key word search ‘family / caregiver and cancer’ within title and abstract. A stepped approach of review was taken, first the abstract and then full article was read to ensure participants were both patient and family / caregiver (Hopia et al., 2016). Full texts were screened by two authors (EC, NH) and disputes were resolved through consultation with all authors. When several articles across the same group of participants were found, if the articles presented different aspects of the study they were included (Pickering & Byrne, 2014). We followed the systematic review guidelines outlined by the Preferred Reporting Items for Systematic Reviews (PRISMA) (Moher, Liberati, Tetzlaff, & Altman, 2009). Figure 1 depicts the PRISMA flowchart for the search.

Data extraction

Pickering framework Step 6- prepare database, Step 7 – enter first ten papers, Step 8 – refine database, Step 9- enter all papers as binary data

An excel spreadsheet was used to collect the binary data from included articles to allow quantitative data analysis. Variables such as journal, country, methodology, method, participants was entered as binary data [0 or 1] to enable a spreadsheet of data which can be used to generate tables and figures. Content of the database was developed and refined as data was extracted (Pickering & Byrne, 2014). For example, columns for different scale measurements were added as they were identified providing a clear picture of the range of tools used within family research.

Pickering framework Step 12 - complete quantitative analysis and qualitative analysis

The ‘Mixed Methods Appraisal Tool (MMAT) – Version 2018’ framework for appraisal of systematic mixed study literature reviews was used (Hong et al., 2018; Pace et al., 2012). Two researchers [EC, JJ] independently completed MMAT appraisals and compared to identify differences in scores. Differences were resolved through discussion with all authors. All articles were included irrespective of their quality appraisal, as the only aspect of missing quality was response rate below 50 which highlighted the question of was the sample representative of the target population (Hong et al., 2018).

A descriptive analysis of the quantitative data set was completed and graphs developed to present the data for easy review (Pickering & Byrne, 2014). Means of ratio data were completed for age and sample size, frequency of other variables was completed. The analysis is presented as graphs and tables to supplement the results (Pickering & Byrne, 2014).

Using a thematic qualitative analysis approach an analysis across all the articles was completed to identify common themes within the included articles. Each member of the research team was allocated a group of the included articles to review and identify themes within the results and discussion section of the articles (Braun & Clarke, 2014). These themes were collated within the database.

Results

Pickering framework Step 13-15 prepare the results and manuscript

A total of 814 articles were identified through the databases with 41 additional articles from reference lists. After removing duplicated 730 titles were screened within Endnote against the inclusion criteria and 132 full text articles screened for eligibility with 73 articles included in the final synthesis. Insert Table 1 Summary of articles and themes

Descriptive characteristics

Included studies were conducted across six countries/geographical areas with the majority from the United States of America (n = 45, 62%) or Europe (n = 20, 27%). Predominantly articles were published in ten major oncology or nursing journals with most published in Supportive Care in Cancer (n = 18, 25%). Most studies used a dyad approach, including one caregiver, mainly the spouse (n = 68, 93%). There was no statistical difference between the genders of the patients and however there were 15% lower male caregivers within the samples. Three studies identified patient and spouse dyads with gender diversity (K. Lyons et al., 2016; Manne et al., 2012; Robbins et al., 2014), and three studies excluded gender diverse couples (Morgan et al., 2011; Rottmann et al., 2015; Zwahlen et al., 2011). See Figure 2: The frequency of articles published grouped by country/geographical area.

Five studies included a range of family members which was referred to in their results (Ashouri et al., 2018; Coyne et al., 2017; Dieperink et al., 2018; Kissil et al., 2013; Robinson et al., 2010). The mean age of participants across all studies was 58 (range 25-88, SD = 9). Twenty-two articles (29%) studied cancer in general without targeting individuals with a specific cancer type. Lung cancer was the most frequent cancer studied (n = 13, 18%). The mean sample size in quantitative studies was 327 participants (range 48 - 2007, SD = 329) and 28 participants (range 12 - 60, SD = 14) in qualitative studies. The majority of studies (n = 45, 62%) had a response rate above 50% and 22 (30%) had a response rate above 75%. See Figure 3: Frequency of the types of cancer focused on in each study.

Study designs

Over three quarters (n = 56, 76%) of studies used quantitative research methods. Intervention designs including random control trial were used for eight studies. Interviews was the most common method of qualitative study designs (n = 13, 18%), focus groups and observation studies were also qualitative methods used to explore family response to adult cancer. Four studies were mixed method using survey and interview or focus group to collect family data. Figure 4 presents the frequency of quantitative and qualitative study design types used in articles included in this review.

Study interventions and duration

Eight studies including six RCTs tested interventions, which were developed for the person with cancer and family members and were delivered by a variety of modalities. Modalities included telephone or videophone (n = 3) (Badger et al., 2013; Mosher et al., 2018; Porter et al., 2011), a combination of telephone and face-to-face methods (n = 2) (Harden et al.,

2009; Northouse et al., 2013), face-to-face only sessions (n = 1) (Dockham et al., 2016), DVD (n = 1) (Collinge et al., 2013), mobile phone applications or computer platforms (n = 1) (Kubo et al., 2018). Figure 5 presents the intervention types used in each experimental study. The interventions mainly focused on improving dyad or family communication. Two interventions provided an educational approach (Dockham et al., 2016; Porter et al., 2011), one focused on coping (Mosher et al., 2018), and two combined education and coping (Harden et al., 2009; Northouse et al., 2013). Another two studies provided psychological support (Badger et al., 2013; Kubo et al., 2018) and one used massage as a relaxation technique (Collinge et al., 2013).

The interventions delivered in acute diagnostic phase were between four weeks and three months. Twenty-seven quantitative studies (37%) collected data prospectively over several time points. Research follow up occurred at 4 or 6 months (n = 12) (Badger et al., 2013; Badr et al., 2008; Badr & Carmack Taylor, 2008; Harden et al., 2009; Manne et al., 2012; Milbury et al., 2012; Newberry et al., 2012; Northouse et al., 2013; Porter et al., 2011; Richardson et al., 2015; Segrin & Badger, 2013, 2014), with study length ranging from one month (Collinge et al., 2013; Hendrix et al., 2016; Mosher et al., 2018) to 24 months (Eisemann et al., 2014). Figure 6 presents the duration of studies collecting prospective data.

Outcomes measured quantitatively

In the included studies, three outcomes were commonly measured for both person with cancer and family members including: functional assessment, health and wellbeing and relationship assessment, with most studies (n=53, 73%) measuring health and wellbeing. Within functional assessment the most commonly used scale was the Brief Symptom Inventory (n= 15, 20%) which measured symptom management of person with cancer and Hospital Anxiety and Depression Scale (HADS) (n =11 15%). Physical health was measured using the Medical

Outcomes Study 36-Item Short-Form Health Survey to assess health and wellbeing of both persons with cancer and family members (n = 8, 11%). Within relationship assessment marital relationships were most commonly measured using the Dyadic Adjustment Scale for both person with cancer and family members (n = 11, 15%). See Table 2: Presentation of scales used in the quantitative research.

Themes

The thematic qualitative analysis revealed three main themes in the included research. These related to ‘Who cancer affects’, ‘Communication within the family’ and ‘Family resources used’.

Who cancer affects. Current research explored how the patient and family experienced cancer. Assessment of functional changes such as fatigue, depression, distress and symptom burden were measured for both patient and family highlighting that functional changes were not isolated to the patient and that family members experienced changes which were often similar or worse than those of the patient. Health and wellbeing such as QoL, physical and psychological health were measured for both patient and family member with a collective finding that cancer influenced both. Caregiver distress was often higher than patient distress. There was a positive relationship between patient and family burden, meaning the lower the patient burden, the lower the family burden. Longitudinal studies explored distress and burden over time finding that distress was still present years after diagnosis. Several studies found that the level of distress at diagnosis and functioning of the family influenced distress over time. Another finding was that distress and burden influenced QoL. The analysis identified that QoL was linked to social determinants of health such as education and access to health services.

Communication within the family. Communication within the family was key to better patient and family outcomes. Communication processes such as being open and sharing concerns, talking about intimacy and the burden of cancer empowered the family to access resources and use positive adaptation strategies. Research interventions were designed to improve communication strategies and appraisal of the cancer within the family, such as health professional facilitated discussion, therapeutic communication and intimacy strategies. Our analysis identified that families who could share concerns and normalise the cancer experience were more likely to obtain tailored support and achieve better outcomes. The patients' stage of illness influenced how they communicated within the family and was related to their sense of burden and distress.

Family resources used. The theme of family resources related to tailoring resources for the patient and family and how this changed over time as their needs changed. Assessment of distress, needs and priorities was important to ensure resources provided met the family's needs. A range of interventions provided information resources in printed booklets, pamphlets, telephone and DVD. Across the studies, recommendations suggested the need to tailor the resource to the individuals' learning preferences. The research interventions varied in length however short family focused interventions, tailored to the family stage of life and cancer stage, were identified as the best way to achieve better patient and family outcomes. Patients and family identified that clinical interventions that facilitated coping with cancer and how to support each other were helpful.

Discussion

This style of quantitative systematic review is the first to explore current family research where participants are persons with cancer and their family. This style of review enables

information from included studies to be synthesised using a quantitative approach. The analysis of binary data provides descriptive statistics with the development of graphs and tables to present data. This review was able to collect data from 73 studies across eleven years providing a view of family research and highlighting commonly used interventions, scales, sample sizes and outcomes.

Although over 50% of the studies were conducted in the United States there was a range of studies across all continents highlighting the international extent of research exploring the family as the unit of care. The need for more family research across diverse areas such as Africa and Asia is needed however, to illustrate how caregivers from different cultures and potentially underdeveloped regions can be better supported (Gabriel et al., 2020). As global migration continues to change how families provide care (Whitehead et al., 2018), the ability to support caregivers of diverse backgrounds also needs to be informed by evidence. The literature review found that there was several research groups who had conducted a range of studies often longitudinal in design or across cancer groups (Badger et al., 2013; Badr, 2016; Mosher et al., 2018; Northouse et al., 2012). These studies enable an understanding of family over time and how different interventions work to improve the quality of life and coping strategies for families.

Initially this review was structured to include research focusing on family as a unit of care, however there was only a few studies involving a range of family members such as adult children, close friends or other relatives (Ashouri et al., 2018; Coyne et al., 2017; Dieperink et al., 2018; Kissil et al., 2013; Robinson et al., 2010). Knowing the complexity of family and the influence of relationships on personal well-being, conducting research with multiple family members should be a consideration as we move to support the family as a unit of care to improve the outcomes for the person with cancer (Illingworth et al., 2009; Van Parys et al., 2017). There

is a need to understand the connectedness and roles within the family to enable interventions and translation of research to practice to be fully family-centred incorporating cancer as a family problem.

A key gap in the literature that was identified in this review was the mean age of the participants; 58 years old. The lack of research for the families with older cancer patients is also reported by (Konradsen et al., 2020). With the increasing age of the general population and higher risk of cancer for adults over the age of sixty, more research with a focus on the older adult may be warranted to develop family focused interventions which can support the younger family carers who may be caring for older parents as well as their own children (Magnuson et al., 2018). The response rate and sample size differed across studies highlighting the difficulty of recruiting family members (Coyne, 2013; Coyne et al., 2019), however the results reflected quality research outcomes and transferability of ideas across studies (Kitson & Harvey, 2016).

The studies were conducted across most cancer groups with the expected focus on lung, breast and prostate cancer as these are the most prevalent cancer types worldwide. These studies provided information on the experience, interventions and outcomes which may be useful across other cancer groups. However there were some missing cancer groups such as cervical cancer (Caster et al., 2017). Cancer type can identify the gender of the person with cancer, and as demonstrated by the large number of partner/spouse dyads in this review, cancer type may therefore influence the family caregiver gender. The analysis of the included articles identified little significant difference in the caregiving relationship between genders. Previous research exploring gender and age differences in caregiving for persons with dementia identified that more than half of caregivers were female, who also had other caregiving burdens related to children (Xiong et al., 2018). This review identified the scarcity of research involving patients

and spouses of gender diversity. There are currently few studies available that include gender diverse couples and little is known about whether the unmet needs and support required by gender diverse couples is comparable to heterosexual couples (Gibson et al., 2017; Harding et al., 2012).

This review explored differences in interventions identifying that most were delivered via telephone with a face to face aspect as well. This method of delivery was common for intervention studies aimed at improving communication and increasing coping capacity within the family and specifically the carer dyad. A Cochrane Systematic Review exploring psychosocial interventions for caregivers after cancer diagnosis highlighted a lack of high level evidence for the interventions (Treanor et al., 2019). There was evidence to support that an intervention with a family focus was likely to improve quality of life outcomes for both the client and the family (Treanor et al., 2019). This is a similar finding to our review which found that interventions with a focus on communication showed improved family quality of life and ability to cope. Improved communication provided avenues for supporting family members but also seeking help from health professionals (Dieperink et al., 2018). One of the concerns identified was the lack of research to explore longitudinal outcomes. Although interventions contributed to improved quality of life and coping these outcomes were not sustained over time and maintained within the survival period (Northouse et al., 2013).

The review identified outcome measures related to functional assessment, health and wellbeing and relationship assessment. These measures provided a view of how the person with cancer and family members were coping with the cancer diagnosis and treatment and how their relationship as a couple or family was influenced by cancer. The relationship changes related to open sharing of the burden of cancer, meeting each other's needs and learning new ways to cope

with the diagnosis and lifestyle changes. Previous reviews have highlighted the suitability of online resources in reducing unmet needs of family members when there is limited health professionals to address the unmet needs (Monton et al., 2019). Online health resources are being developed which enable family members to access information and support via Smartphone apps, however, resources need to be at the appropriate health literacy level as they are used without health professional support (Wittenberg et al., 2019). There is still a need for health professionals to support the family as they work through changes as a family unit, focusing on improving family communication and access to tailored support (Goldsmith et al., 2019).

There was scant research on socio-economic impact of cancer on the patient and family. It is well known that disparities of access to treatment and ability to maintain family functioning are influenced by socio-economic status (Girgis & Lambert, 2017; Salehi et al., 2019). The cost of caregiving warrants more research to really understand the family contribution to support and how best to ensure health professionals enable the family to sustain support to the person with cancer.

Many of the studies explored who was most affected by the cancer finding that it was often the family members with a higher level of distress and unmet needs. The complexity of family interaction and provision of information at the appropriate health literacy level was highlighted. Hence the need to understand the family and who they are and tailor information and interventions to meet their specific needs is warranted (Shajani & Snell, 2019). Intimacy and relationships was a focus within the dyad research exploring the effects of cancer and the couples' intimacy and how this influenced their coping with the cancer. Interventions aimed at improving family communication either by teaching supportive behaviours such as massage, spirituality, or general coping skills identified positive outcomes for the family and person with

cancer. Teaching couples and family communication skills to encourage open the discussion about how the cancer influenced them was found to improve overall quality of life outcomes (Wittenberg et al., 2019).

This literature review revealed the need for resources to be accessible, culturally correct, support the family self-efficacy and provide a connection to health profession or community support. The level of distress at diagnosis influenced the person with cancer and the family ability to maintain family functioning suggesting that family assessment and tailoring supportive care for the family would improve family outcomes (Fiest et al., 2018). The delivery of information and resources was preferred as telephone or technology based, this reduced the travel and costs to attend information and support sessions. Across chronic disease there has been a move to deliver information and tailor support using technology enhanced devices enabling the accessibility to be enhanced (Heynsbergh et al., 2018).

Strength & Limitations

While all efforts were made to identify relevant studies, it is possible that some were excluded due lack of clear identification of patient and family participants in the title or abstract. However, a systematic quantitative literature review includes a diverse range of study designs and has the advantage to draw a quantified map of the research within a specific time frame, and point out gaps for further investigation. Only studies registered in English databases were accessed and this may have narrowed the number of studies included, especially those studies from middle and low-income countries and Asian countries. The included studies used various methodologies, outcome measures and sample sizes, and it was challenging to conduct the analysis but the review method enables the analysis of research.

Recommendations

Future research should aim to include a range of family participants and higher response rates to understand the depth of family response and how to improve outcomes. The development of family interventions need to be tailored to the family as a unit of care. Incorporating a family assessment to truly tailor information for family members and meet their needs is required. As society becomes increasingly connected with technology, health care must move to incorporate this but not lose the personal touch and value of listening to the family. Using video telephone support and other technology will enhance face-to-face health professional support (Petersson et al., 2020; Vestergaard et al., 2019). Additionally, there is a need for research among older family members, this review identified that, family research has focused on caregivers with a mean age of 58. As our populations continue to age there is an undisputed need for evidence on how to better support family in the outpatient setting to ensure resources are suitable and acceptable to caregivers.

In current practice, nurses are at the forefront in having the opportunity to assess, respond to and evaluate caregivers' needs. Improved communication between nurses and caregivers can be driven by nurses and is a vital step in ensuring that caregivers' needs are acknowledged and resources provided.

Conclusion

This review has explored research which had patients with cancer and family members as a sample. Research gaps include support for older family caregivers and providing individualised support to caregivers and the need for research involving non spouse family caregivers. There is a research gap in interventions studies and in studies from countries outside US and Europe. This

review proposes that family focused interventions delivered during the acute treatment phase using technology, over a six month period have the highest possibility of sustained influence for the family functioning.

Reference List

- Akin, S., & Durna, Z. (2013). A comparative descriptive study examining the perceptions of cancer patients, family caregivers, and nurses on patient symptom severity in Turkey. *European Journal of Oncology Nursing*, 17(1), 30-37. <https://doi.org/10.1016/j.ejon.2012.02.002>
- Andersen, N., Nielsen, C., Danbjørg, D., Møller, P., & Dieperink, K. (2019). Cancer & Caregivers. Caregivers' needs for support during outpatient cancer settings. *Oncology Nursing Forum*, 46(6), 757-767. <https://doi.org/doi: 10.1188/19.ONF.757-767>
- Ashouri, E., Taleghani, F., Memarzadeh, M., Saburi, M., & Babashahi, F. (2018). The perceptions of nurses, patients and family members regarding nurses' empathetic behaviours towards patients suffering from cancer: a descriptive qualitative study. *Journal of Research in Nursing*, 23(5), 428-443. <https://doi.org/10.1177/1744987118756945>
- Australian Commission on Safety and Quality in Health Care. (2018). *National safety and quality health service standards*. ACSQHC Retrieved 30th Oct from <https://www.safetyandquality.gov.au/our-work/clinical-communications/>
- Badger, T., Segrin, C., Pasvogel, A., & Lopez, A. M. (2013). The effect of psychosocial interventions delivered by telephone and videophone on quality of life in early-stage breast cancer survivors and their supportive partners. *J Telemed Telecare*, 19(5), 260-265. <https://doi.org/10.1177/1357633x13492289>
- Badr, H. (2016). Psychosocial Interventions for Patients With Advanced Cancer and Their Families. *American Journal of Lifestyle Medicine*, 10(1), 53-63. <https://doi.org/10.1177/1559827614530966>
- Badr, H., Acitelli, L., & Carmack Taylor, C. (2008). Does talking about their relationship affect couples' marital and psychological adjustment to lung cancer? *J Cancer Surviv*, 2(1), 53-64. <https://doi.org/10.1007/s11764-008-0044-3>
- Badr, H., & Carmack Taylor, C. (2008). Effects of relationship maintenance on psychological distress and dyadic adjustment among couples coping with lung cancer. *Health Psychol*, 27(5), 616-627. <https://doi.org/10.1037/0278-6133.27.5.616>
- Badr, H., Gupta, V., Sikora, A., & Posner, M. (2014). Psychological distress in patients and caregivers over the course of radiotherapy for head and neck cancer. *Oral Oncol*, 50(10), 1005-1011. <https://doi.org/10.1016/j.oraloncology.2014.07.003>

- Bell, J. M. (2014, February 1, 2014). Knowledge translation in family nursing: Gazing into the promised land. *Journal of Family Nursing*, 20(1), 3-12. <https://doi.org/10.1177/1074840714521731>
- Bell, J. M., & Wright, L. M. (2015). The Illness Beliefs Model: advancing practice knowledge about illness beliefs, family healing, and family interventions. *Journal of Family Nursing*, 21(2), 179. <http://griffith.summon.serialssolutions.com>
- Braun, V., & Clarke, V. (2014). What can "thematic analysis" offer health and wellbeing researchers? *International Journal of Qualitative Studies on Health and Well-being*, 9(1), 26152-26152. <https://doi.org/10.3402/qhw.v9.26152>
- Carmack Taylor, C., Badr, H., Lee, J., Fossella, F., Pisters, K., Gritz, E., & Schover, L. (2008). Lung cancer patients and their spouses: psychological and relationship functioning within 1 month of treatment initiation. *Ann Behav Med*, 36(2), 129-140. <https://doi.org/10.1007/s12160-008-9062-7>
- Carman, M. J., Wolf, L. A., Henderson, D., Kamienski, M., Koziol-McLain, J., Manton, A., & Moon, M. D. (2013). Developing Your Clinical Question: The Key to Successful Research. *Journal of Emergency Nursing*, 39(3), 299-301. <https://doi.org/org/10.1016/j.jen.2013.01.011>
- Castellanos, E. H., Dietrich, M. S., Bond, S. M., Wells, N., Schumacher, K., Ganti, A. K., & Murphy, B. A. (2019). Impact of patient symptoms and caregiving tasks on psychological distress in caregivers for head and neck cancer (HNC). *Psychooncology*, 28(3), 511-517. <https://doi.org/10.1002/pon.4968>
- Caster, M. M., Norris, A. H., Butao, C., Carr Reese, P., Chemey, E., Phuka, J., & Turner, A. N. (2017). Assessing the Acceptability, Feasibility, and Effectiveness of a Tablet-Based Cervical Cancer Educational Intervention. *Journal of Cancer Education*, 32(1), 35-42. <https://doi.org/10.1007/s13187-015-0953-6>
- Collinge, W., Kahn, J., Walton, T., Kozak, L., Bauer-Wu, S., Fletcher, K., Yarnold, P., & Soltysik, R. (2013). Touch, Caring, and Cancer: Randomized controlled trial of a multimedia caregiver education program. *Supportive Care in Cancer*, 21(5), 1405-1414. <https://doi.org/10.1007/s00520-012-1682-6>
- Coyne, E. (2013). The strengths and resources used by families of young women with breast cancer. *Australian Journal of Cancer Nursing*, 14(2), 10-16. <http://trove.nla.gov.au/version/192031675>
- Coyne, E., Dieperink, K., Østergaard, B., & Creedy, D. (2017). Strengths and resources used by Australian and Danish adult patients and their family caregivers during treatment for cancer. *European Journal of Oncology Nursing*, 29, 53-59. <https://doi.org/10.1016/j.ejon.2017.05.005>

- Coyne, E., Frommolt, V., & Salehi, A. (2019). The experience and challenges of rural persons with cancer and their families. *Collegian*, 26(6), 609-614. <https://doi.org/10.1016/j.colegn.2019.10.002>
- Deek, H., Hamilton, S., Brown, N., Inglis, S. C., Digiacomio, M., Newton, P. J., Nouredine, S., MacDonald, P. S., Davidson, P. M., Investigators, F. P., & the, F. P. I. (2016). Family-centred approaches to healthcare interventions in chronic diseases in adults: a quantitative systematic review. *Journal of Advanced Nursing*, 72(5), 968-979. <https://doi.org/10.1111/jan.12885>
- Dieperink, K., Coyne, E., Creedy, D., & Ostergaard, B. (2018). Family functioning and perceived support from nurses during cancer treatment among Danish and Australian patients and their families *Journal of Clinical Nursing*, 27(1-2), e154-e161. <https://doi.org/10.1111/jocn.13894>
- Dieperink, K. B., Mark, K., & Mikkelsen, T. B. (2016). Marital rehabilitation after prostate cancer – a matter of intimacy. *International Journal of Urological Nursing*, 10(1), 21-29. <https://doi.org/10.1111/ijun.12091>
- Dockham, B., Schafenacker, A., Yoon, H., Ronis, D., Kershaw, T., Titler, M., & Northouse, L. (2016). Implementation of a psychoeducational program for cancer survivors and family caregivers at a cancer support community affiliate: A pilot effectiveness study. *Cancer Nursing*, 39(3), 169-180. <https://doi.org/10.1097/NCC.0000000000000311>
- Eisemann, N., Waldmann, A., Rohde, V., & Katalinic, A. (2014). Quality of life in partners of patients with localised prostate cancer. *Qual Life Res*, 23(5), 1557-1568. <https://doi.org/10.1007/s11136-013-0588-1>
- Elliott, K. E., Scott, J. L., Monsour, M., & Nuwayhid, F. (2015). Profiles of dyadic adjustment for advanced prostate cancer to inform couple-based intervention. *Psychol Health*, 30(11), 1259-1273. <https://doi.org/10.1080/08870446.2015.1043301>
- Ellis, J., Lloyd Williams, M., Wagland, R., Bailey, C., & Molassiotis, A. (2013, Jan). Coping with and factors impacting upon the experience of lung cancer in patients and primary carers. *Eur J Cancer Care (Engl)*, 22(1), 97-106. <https://doi.org/10.1111/ecc.12003>
- Ellis, K., Janevic, M., Kershaw, T., Caldwell, C., Janz, N., & Northouse, L. (2017). The influence of dyadic symptom distress on threat appraisals and self-efficacy in advanced cancer and caregiving. *Supportive Care in Cancer*, 25(1), 185-194.
- Ferrell, B., & Wittenberg, E. (2017). A review of family caregiving intervention trials in oncology. *CA: A Cancer Journal for Clinicians*, 67(4), 318-325. <https://doi.org/10.3322/caac.21396>

- Fiest, K. M., McIntosh, C. J., Demiantschuk, D., Leigh, J. P., & Stelfox, H. T. (2018). Translating evidence to patient care through caregivers: a systematic review of caregiver-mediated interventions. *BMC MEDICINE*, 16(1), 105. <https://doi.org/10.1186/s12916-018-1097-4>
- Finley, J. P. (2018). Caregiver Café: Providing education and support to family caregivers of patients with cancer. *Clinical Journal of Oncology Nursing*, 22(1), 91-96. <https://doi.org/10.1188/18.CJON.91-96>
- Fletcher, B. S., Miaskowski, C., Given, B., & Schumacher, K. (2011). The cancer family caregiving experience: An updated and expanded conceptual model. *European Journal of Oncology Nursing*, 16(4), 387-398. <https://doi.org/10.1016/j.ejon.2011.09.001>
- Gabriel, I., Creedy, D., & Coyne, E. (2020). A systematic review of psychosocial interventions to improve quality of life of patients with cancer and their family caregivers. *Nursing Open*, 7, 1299-1312. <https://doi.org/10.1002/nop2.543>
- Ganong, L. (2011). Return of the "Intimate Outsider": Current trends and issues in family nursing research revisited. *Journal of Family Nursing*, 17(4), 416-440. <https://doi.org/10.1177/1074840711425029>
- Gansler, T., Kepner, J., Willacy, E., Soloe, C., Rupert, D., Jarblum, M., Driscoll, D., Orr, A., Fitzgerald, T., & Esparza, A. (2010). Evolving Information Priorities of Hematologic Cancer Survivors, Caregivers, and Other Relatives. *Journal of Cancer Education*, 1-10. <https://doi.org/10.1007/s13187-009-0034-9>
- Gibson, A. W., Radix, A. E., Maingi, S., & Patel, S. (2017). Cancer care in lesbian, gay, bisexual, transgender and queer populations. *Future Oncology*, 13(15), 1333-1344. <https://doi.org/10.2217/fon-2017-0482>
- Gilliss, C. L., Pan, W., & Davis, L. L. (2019). Family Involvement in Adult Chronic Disease Care: Reviewing the Systematic Reviews. *Journal of Family Nursing*, 25(1), 3-27. <https://doi.org/10.1177/1074840718822365>
- Girgis, A., & Lambert, S. (2017). Cost of informal caregiving in cancer care. *Cancer Forum*, 41(2), 16-22.
- Given, B. A., Given, C. W., & Sherwood, P. (2012). The Challenge of Quality Cancer Care for Family Caregivers. *Seminars in Oncology Nursing*, 28(4), 205-212. <https://doi.org/10.1016/j.soncn.2012.09.002>
- Goldsmith, J. V., Wittenberg, E., Terui, S., Kim, H., & Umi, S. (2019). Providing Support for Caregiver Communication Burden: Assessing the Plain Language Planner Resource As a Nursing

- Intervention. *Seminars in Oncology Nursing*, 35(4), 354-358.
<https://doi.org/https://doi.org/10.1016/j.soncn.2019.06.007>
- Goldzweig, G., Baider, L., Andritsch, E., Pfeffer, R., & Rottenberg, Y. (2017). A Dialogue of Depression and Hope: Elderly Patients Diagnosed with Cancer and Their Spousal Caregivers. *J Cancer Educ*, 32(3), 549-555. <https://doi.org/10.1007/s13187-015-0975-0>
- Götze, H., Friedrich, M., Brähler, E., Romer, G., Mehnert, A., & Ernst, J. (2017). Psychological distress of cancer patients with children under 18 years and their partners-a longitudinal study of family relationships using dyadic data analysis. *Supportive Care in Cancer*, 25(1), 255-264.
<https://link.springer.com/content/pdf/10.1007%2Fs00520-016-3411-z.pdf>
- Gröpper, S., van der Meer, E., Landes, T., Bucher, H., Stickel, A., & Goerling, U. (2016). Assessing cancer-related distress in cancer patients and caregivers receiving outpatient psycho-oncological counseling. *Supportive Care in Cancer*, 24(5), 2351-2357. <https://doi.org/10.1007/s00520-015-3042-9>
- Haj Mohammad, N., Walter, A., van Oijen, M., Hulshof, M., Bergman, J., Anderegg, M., van Berge Henegouwen, M., Henselmans, I., Sprangers, M., & van Laarhoven, H. (2015). Burden of spousal caregivers of stage II and III esophageal cancer survivors 3 years after treatment with curative intent. *Supportive Care in Cancer*, 23(12), 3589-3598. <https://doi.org/10.1007/s00520-015-2727-4>
- Harden, J., Falahee, M., Bickes, J., Schafenacker, A., Walker, J., Mood, D., & Northouse, L. (2009). Factors associated with prostate cancer patients' and their spouses' satisfaction with a family-based intervention. *Cancer Nurs*, 32(6), 482-492. <https://doi.org/10.1097/NCC.0b013e3181b311e9>
- Harden, J., Northouse, L., Cimprich, B., Pohl, J., Liang, J., & Kershaw, T. (2008). The influence of developmental life stage on quality of life in survivors of prostate cancer and their partners. *Journal of Cancer Survivorship*, 2(2), 84-94. <https://doi.org/10.1007/s11764-008-0048-z>
- Harding, R., Epiphaniou, E., & Chidgey-Clark, J. (2012). Needs, Experiences, and Preferences of Sexual Minorities for End-of-Life Care and Palliative Care: A Systematic Review. *Journal of Palliative Medicine*, 15(5), 602-611. <https://doi.org/10.1089/jpm.2011.0279>
- Hasson-Ohayon, I., Goldzweig, G., Braun, M., & Galinsky, D. (2010). Women with advanced breast cancer and their spouses: diversity of support and psychological distress. *Psycho-Oncology*, 19(11), 1195-1204. <https://doi.org/10.1002/pon.1678>
- Hendriksen, E., Williams, E., Sporn, N., Greer, J., DeGrange, A., & Koopman, C. (2015). Worried together: a qualitative study of shared anxiety in patients with metastatic non-small cell lung cancer and

- their family caregivers. *Support Care Cancer*, 23(4), 1035-1041. <https://doi.org/10.1007/s00520-014-2431-9>
- Hendrix, C., Bailey, D., Steinhäuser, K., Olsen, M., Stechuchak, K., Lowman, S., Schwartz, A., Riedel, R., Keefe, F., Porter, L., & Tulsky, J. (2016). Effects of enhanced caregiver training program on cancer caregiver's self-efficacy, preparedness, and psychological well-being. *Supportive Care in Cancer*, 24(1), 327-336. <https://doi.org/10.1007/s00520-015-2797-3>
- Heynsbergh, N., Heckel, L., Botti, M., & Livingston, P. M. (2018). Feasibility, useability and acceptability of technology-based interventions for informal cancer carers: a systematic review [journal article]. *BMC Cancer*, 18(1), 244. <https://doi.org/10.1186/s12885-018-4160-9>
- Hong, Q. N., Gonzalez-Reyes, A., & Pluye, P. (2018). Improving the usefulness of a tool for appraising the quality of qualitative, quantitative and mixed methods studies, the Mixed Methods Appraisal Tool (MMAT). *Journal of Evaluation in Clinical Practice*, 24(3), 459-467. <https://doi.org/10.1111/jep.12884>
- Hopia, H., Latvala, E., & Liimatainen, L. (2016). Reviewing the methodology of an integrative review. *Scandinavian Journal of Caring Sciences*, 30(4), 662-669. <https://doi.org/10.1111/scs.12327>
- Hsieh, C. C., Yu, C. J., Chen, H. J., Chen, Y. W., Chang, N. T., & Hsiao, F. H. (2019). Dispositional mindfulness, self-compassion, and compassion from others as moderators between stress and depression in caregivers of patients with lung cancer. *Psycho-Oncology*, 28(7), 1498-1505. <https://doi.org/10.1002/pon.5106>
- Illingworth, N., Forbat, L., Hubbard, G., & Kearney, N. (2009). The importance of relationships in the experience of cancer: A re-working of the policy ideal of the whole-systems approach. *European Journal of Oncology Nursing*, 14(1), 23-28. <https://doi.org/10.1016/j.ejon.2009.06.006>
- Johansen, S., Cvancarova, M., & Ruland, C. (2018). The Effect of Cancer Patients' and Their Family Caregivers' Physical and Emotional Symptoms on Caregiver Burden. *Cancer Nurs*, 41(2), 91-99. <https://doi.org/10.1097/NCC.0000000000000493>
- Kauffman, R., Bitz, C., Clark, K., Loscalzo, M., Kruper, L., & Vito, C. (2016). Addressing psychosocial needs of partners of breast cancer patients: a pilot program using social workers to improve communication and psychosocial support. *Supportive Care In Cancer: Official Journal Of The Multinational Association Of Supportive Care In Cancer*, 24(1), 61-65. <https://doi.org/10.1007/s00520-015-2721-x>
- Kershaw, T., Mood, D., Newth, G., Ronis, D., Sanda, M., Vaishampayan, U., & Northouse, L. (2008). Longitudinal analysis of a model to predict quality of life in prostate cancer patients and their spouses. *Ann Behav Med*, 36(2), 117-128. <https://doi.org/10.1007/s12160-008-9058-3>

- Kim, Y., Carver, C. S., Spillers, R. L., Crammer, C., & Zhou, E. S. (2011). Individual and dyadic relations between spiritual well-being and quality of life among cancer survivors and their spousal caregivers. *Psychooncology*, 20(7), 762-770. <https://doi.org/10.1002/pon.1778>
- Kim, Y., Kashy, D. A., Wellisch, D. K., Spillers, R. L., Kaw, C. K., & Smith, T. G. (2008). Quality of life of couples dealing with cancer: dyadic and individual adjustment among breast and prostate cancer survivors and their spousal caregivers. *Ann Behav Med*, 35(2), 230-238. <https://doi.org/10.1007/s12160-008-9026-y>
- Kim, Y., van Ryn, M., Jensen, R., Griffin, J., Potosky, A., & Rowland, J. (2015). Effects of gender and depressive symptoms on quality of life among colorectal and lung cancer patients and their family caregivers. *Psycho-Oncology*, 24(1), 95-105. <https://doi.org/10.1002/pon.3580>
- Kissil, K., Niño, A., Ingram, M., & Davey, M. (2013). "I Knew From Day One That I'm Either Gonna Fight This Thing or Be Defeated": African American Parents' Experiences of Coping With Breast Cancer. *Journal of Family Nursing*, 20(1), 98-119. <https://doi.org/10.1177/1074840713504035>
- Kitson, A. L., & Harvey, G. (2016). Methods to Succeed in Effective Knowledge Translation in Clinical Practice: Effective Knowledge Translation. *Journal of Nursing Scholarship*, 48(3), 294-302. <https://doi.org/10.1111/jnu.12206>
- Konradsen, H., Brødsgaard, A., Østergaard, B., Svavarsdóttir, E., Dieperink, K. B., Imhof, L., Luttik, M. L., Mahrer-Imhof, R., & García-Vivar, C. (2020). Health practices in Europe towards families of older patients with cancer: a scoping review. *Scandinavian Journal of Caring Sciences*. <https://doi.org/10.1111/scs.12855>
- Kubo, A., Altschuler, A., Kurtovich, E., Hendlish, S., Laurent, C., Kolevska, T., Li, Y., & Avins, A. (2018). A Pilot Mobile-based Mindfulness Intervention for Cancer Patients and their Informal Caregivers. *Mindfulness*, 9(6), 1885-1894. <https://doi.org/10.1007/s12671-018-0931-2>
- La, I. S., & Yun, E. K. (2017). Effects of stress appraisal on the quality of life of adult patients with multiple myeloma and their primary family caregivers in Korea. *Psycho-Oncology*, 26(10), 1640-1646. <https://doi.org/10.1002/pon.4348>
- Lageman, S. K., Brown, P. D., Anderson, S. K., Lachance, D. H., Yan, E., Laack, N. N. I., & Cerhan, J. H. (2015). Exploring primary brain tumor patient and caregiver needs and preferences in brief educational and support opportunities. *Supportive Care in Cancer*, 23(3), 851-859. <https://doi.org/10.1007/s00520-014-2413-y>
- Lambert, S. D., Girgis, A., Turner, J., Regan, T., Candler, H., Britton, B., Chambers, S., Lawsins, C., & Kayser, K. (2013). "You need something like this to give you guidelines on what to do": patients' and

- partners' use and perceptions of a self-directed coping skills training resource. *Supportive Care in Cancer*, 21(12), 3451-3460. <https://doi.org/10.1007/s00520-013-1914-4>
- Li, Q., Lin, Y., Xu, Y., & Zhou, H. (2018). The impact of depression and anxiety on quality of life in Chinese cancer patient-family caregiver dyads, a cross-sectional study. *Health Qual Life Outcomes*, 16(1), 230. <https://doi.org/10.1186/s12955-018-1051-3>
- Liang, J., Lee, S. J., Storer, B. E., Shaw, B. E., Chow, E. J., Flowers, M. E., Krakow, E. F., Bar, M., Syrjala, K. L., Salit, R. B., Kurukulasuriya, C. E., & Jim, H. S. L. (2019). Rates and Risk Factors for Post-Traumatic Stress Disorder Symptomatology among Adult Hematopoietic Cell Transplant Recipients and Their Informal Caregivers. *Biol Blood Marrow Transplant*, 25(1), 145-150. <https://doi.org/10.1016/j.bbmt.2018.08.002>
- Libert, Y., Merckaert, I., Slachmuylder, J. L., & Razavi, D. (2013). The ability of informal primary caregivers to accurately report cancer patients' difficulties. *Psychooncology*, 22(12), 2840-2847. <https://doi.org/10.1002/pon.3362>
- Locher, J. L., Robinson, C. O., Bailey, F. A., Carroll, W. R., Heimbürger, D. C., Saif, M. W., Tajeu, G., & Ritchie, C. S. (2010). Disruptions in the organization of meal preparation and consumption among older cancer patients and their family caregivers. *Psycho-Oncology*, 19(9), 967-974. <https://doi.org/10.1002/pon.1656>
- Lyons, K., Lee, C., Bennett, J., Nail, L., Fromme, E., Hiatt, S., & Sayer, A. (2014). Symptom incongruence trajectories in lung cancer dyads. *J Pain Symptom Manage*, 48(6), 1031-1040. <https://doi.org/10.1016/j.jpainsymman.2014.02.004>
- Lyons, K., Miller, L., & McCarthy, M. (2016). The Roles of Dyadic Appraisal and Coping in Couples With Lung Cancer. *Journal of Family Nursing*, 22(4), 493-514. <https://doi.org/10.1177/1074840716675976>
- Lyons, K. S., Miller, L. M., & McCarthy, M. J. (2016). The Roles of Dyadic Appraisal and Coping in Couples With Lung Cancer. *Journal of Family Nursing*, 22(4), 493-514. <https://doi.org/10.1177/1074840716675976>
- Magnuson, A., Lemelman, T., Pandya, C., Goodman, M., Noel, M., Tejani, M., Dougherty, D., Dale, W., Hurria, A., Janelins, M., Lin, F. V., Heckler, C., & Mohile, S. (2018). Geriatric assessment with management intervention in older adults with cancer: a randomized pilot study. *Supportive Care in Cancer*, 26(2), 605-613. <https://doi.org/10.1007/s00520-017-3874-6>
- Manne, S., Badr, H., & Kashy, D. (2012). A longitudinal analysis of intimacy processes and psychological distress among couples coping with head and neck or lung cancers. *J Behav Med*, 35(3), 334-346. <https://doi.org/10.1007/s10865-011-9349-1>

- Manne, S., Kissane, D., Zaider, T., Kashy, D., Lee, D., Heckman, C., & Virtue, S. (2015). Holding back, intimacy, and psychological and relationship outcomes among couples coping with prostate cancer. *J Fam Psychol*, 29(5), 708-719. <https://doi.org/10.1037/fam0000096>
- Milbury, K., Badr, H., & Carmack Taylor, C. (2012). The role of blame in the psychosocial adjustment of couples coping with lung cancer. *Ann Behav Med*, 44(3), 331-340. <https://doi.org/10.1007/s12160-012-9402-5>
- Mishra, S. I., Rishel Brakey, H., Kano, M., Nedjat-Haiem, F. R., & Sussman, A. L. (2018). Health related quality of life during cancer treatment: Perspectives of young adult (23-39 years) cancer survivors and primary informal caregivers [Article]. *European journal of oncology nursing : the official journal of European Oncology Nursing Society*, 32, 48-54. <https://doi.org/10.1016/j.ejon.2017.11.007>
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*, 339(7716), 332-336. <https://doi.org/10.1136/bmj.b2535>
- Molassiotis, A., Wilson, B., Blair, S., Howe, T., & Cavet, J. (2011). Unmet supportive care needs, psychological well-being and quality of life in patients living with multiple myeloma and their partners. *Psychooncology*, 20(1), 88-97. <https://doi.org/10.1002/pon.1710>
- Monton, O., Lambert, S., Belzile, E., & Mohr-Elzeki, D. (2019). An evaluation of the suitability, readability, quality, and usefulness of online resources for family caregivers of patients with cancer. *Patient Education and Counseling*, 102(10), 1892-1897. <https://doi.org/https://doi.org/10.1016/j.pec.2019.05.010>
- Morgan, M. A., Small, B. J., Donovan, K. A., Overcash, J., & McMillan, S. (2011). Cancer patients with pain: the spouse/partner relationship and quality of life. *Cancer Nurs*, 34(1), 13-23. <https://doi.org/10.1097/NCC.0b013e3181efed43>
- Moschopoulou, E., Hutchison, I., Bhui, K., & Korszun, A. (2018). Post-traumatic stress in head and neck cancer survivors and their partners [Article in Press]. *Supportive Care in Cancer*, 1-9. <https://doi.org/10.1007/s00520-018-4146-9>
- Mosher, C., Secinti, E., Johns, S., O'Neil, B., Helft, P., Shahda, S., Jalal, S., & Champion, V. (2018). Examining the effect of peer helping in a coping skills intervention: a randomized controlled trial for advanced gastrointestinal cancer patients and their family caregivers. *Quality of life research*, 27(2), 515-528. <https://doi.org/10.1007/s11136-017-1620-7>

- Mosher, C. E., Adams, R. N., Helft, P. R., O'Neil, B. H., Shahda, S., Rattray, N. A., & Champion, V. L. (2016). Family caregiving challenges in advanced colorectal cancer: patient and caregiver perspectives. *Supportive Care In Cancer: Official Journal Of The Multinational Association Of Supportive Care In Cancer*, 24(5), 2017-2024. <https://doi.org/10.1007/s00520-015-2995-z>
- Mosher, C. E., Ott, M. A., Hanna, N., Jalal, S. I., & Champion, V. L. (2015). Coping with physical and psychological symptoms: a qualitative study of advanced lung cancer patients and their family caregivers. *Supportive Care In Cancer: Official Journal Of The Multinational Association Of Supportive Care In Cancer*, 23(7), 2053-2060. <https://doi.org/10.1007/s00520-014-2566-8>
- Newberry, A., Kuo, J., Donovan, H., Given, B., Given, C. W., Schulz, R., & Sherwood, P. (2012). Identifying family members who are likely to perceive benefits from providing care to a person with a primary malignant brain tumor. *Oncology Nursing Forum*, 39(3), E226-E232. <https://doi.org/10.1188/12.ONF.E226-E232>
- Northouse, L., Mood, D. W., Schafenacker, A., Kalemkerian, G., Zalupski, M., LoRusso, P., Hayes, D., Hussain, M., Ruckdeschel, J., Fendrick, A., Trask, P., Ronis, D., & Kershaw, T. (2013). Randomized clinical trial of a brief and extensive dyadic intervention for advanced cancer patients and their family caregivers. *Psycho-Oncology*, 22(3), 555-563. <https://doi.org/10.1002/pon.3036>
- Northouse, L., Williams, A., M, Given, B., & McCorkle, R. (2012). Psychosocial care for family caregivers of patients with cancer. *Journal of clinical Oncology*, 30(11), 1227-1234. <https://doi.org/10.1200/JCO.2011.39.5798>
- Nund, R. L., Rumbach, A. F., Debattista, B. C., Goodrow, M. N. T., Johnson, K. A., Tupling, L. N., Scarinci, N. A., Cartmill, B., Ward, E. C., & Porceddu, S. V. (2015). Communication changes following non-glottic head and neck cancer management: The perspectives of survivors and carers. *International Journal of Speech-Language Pathology*, 17(3), 263-272. <https://doi.org/10.3109/17549507.2015.1010581>
- Oakley, S., Grealish, L., El Amouri, S., & Coyne, E. (2019). The lived experience of expatriate nurses providing end of life care to Muslim patients in a Muslim country: An integrated review of the literature. *International Journal of Nursing Studies*, 94, 51-59. <https://doi.org/https://doi.org/10.1016/j.ijnurstu.2019.03.002>
- Ovayolu, Ö., Ovayolu, N., Aytaç, S., Serçe, S., & Sevinc, A. (2015). Pain in cancer patients: pain assessment by patients and family caregivers and problems experienced by caregivers [Article]. *Supportive Care in Cancer*, 23(7), 1857-1864. <https://doi.org/10.1007/s00520-014-2540-5>
- Pasipanodya, E. C., Parrish, B. P., Laurenceau, J. P., Cohen, L. H., Siegel, S. D., Graber, E. C., & Belcher, A. J. (2012). Social constraints on disclosure predict daily well-being in couples coping with early-stage breast cancer. *J Fam Psychol*, 26(4), 661-667. <https://doi.org/10.1037/a0028655>

- Petersson, N. B., Løvendahl, A. J., Danbjørg, D. B., & Dieperink, K. B. (2020). Video-consulted rounds with caregivers: The experience of patients with cancer. *European Journal of Oncology Nursing*, 101763. <https://doi.org/10.1016/j.ejon.2020.101763>
- Petruzzi, A., Finocchiaro, C. Y., Lamperti, E., & Salmaggi, A. (2013). Living with a brain tumor : reaction profiles in patients and their caregivers. *Supportive Care in Cancer*, 21(4), 1105-1111. <https://doi.org/10.1007/s00520-012-1632-3>
- Pickering, C., & Byrne, J. (2014). *The benefits of publishing systematic quantitative literature reviews for PhD candidates and other early-career researchers* (Vol. 33). <https://doi.org/10.1080/07294360.2013.841651>
- Pickering, C., Grignon, J., Steven, R., Guitart, D., & Byrne, J. (2015). Publishing not perishing: how research students transition from novice to knowledgeable using systematic quantitative literature reviews. *Studies in Higher Education*, 40(10), 1756-1769. <https://doi.org/10.1080/03075079.2014.914907>
- Porter, L. S., Keefe, F. J., Garst, J., Baucom, D. H., McBride, C. M., McKee, D. C., Sutton, L., Carson, K., Knowles, V., Rumble, M., & Scipio, C. (2011). Caregiver-assisted coping skills training for lung cancer: results of a randomized clinical trial. *J Pain Symptom Manage*, 41(1), 1-13. <https://doi.org/10.1016/j.jpainsymman.2010.04.014>
- Porter, L. S., Keefe, F. J., Garst, J., McBride, C. M., & Baucom, D. (2008). Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: associations with symptoms and distress. *Pain*, 137(2), 306-315. <https://doi.org/10.1016/j.pain.2007.09.010>
- Posluszny, D. M., Dougall, A. L., Johnson, J. T., Argiris, A., Ferris, R. L., Baum, A., Bovbjerg, D. H., & Dew, M. A. (2015). Posttraumatic stress disorder symptoms in newly diagnosed patients with head and neck cancer and their partners. *Head Neck*, 37(9), 1282-1289. <https://doi.org/10.1002/hed.23760>
- Rha, S. Y., Park, Y., Song, S. K., Lee, C. E., & Lee, J. (2014). Caregiving burden and health-promoting behaviors among the family caregivers of cancer patients. *European Journal of Oncology Nursing*, 19(2), 174-181. <https://doi.org/10.1016/j.ejon.2014.09.003>
- Rha, S. Y., Park, Y., Song, S. K., Lee, C. E., & Lee, J. (2015). Caregiving burden and the quality of life of family caregivers of cancer patients: the relationship and correlates. *European Journal of Oncology Nursing*, 19(4), 376-382. <https://doi.org/10.1016/j.ejon.2015.01.004>

- Richardson, A. E., Morton, R., & Broadbent, E. (2015). Psychological support needs of patients with head and neck cancer and their caregivers: A qualitative study. *Psychology & Health, 30*(11), 1288-1305. <https://doi.org/10.1080/08870446.2015.1045512>
- Robbins, M. L., Lopez, A. M., Weihs, K. L., & Mehl, M. R. (2014). Cancer conversations in context: naturalistic observation of couples coping with breast cancer. *J Fam Psychol, 28*(3), 380-390. <https://doi.org/10.1037/a0036458>
- Robinson, C. A., Bottorff, J. L., Smith, M. L., & Sullivan, K. M. (2010). "Just Because You've Got Lung Cancer Doesn't Mean I Will": Lung Cancer, Smoking, and Family Dynamics. *Journal of Family Nursing, 16*(3), 282-301. <https://doi.org/10.1177/1074840710370747>
- Romito, F., Goldzweig, G., Cormio, C., Hagedoorn, M., & Andersen, B. L. (2013). Informal caregiving for cancer patients: Informal Caregiving. *Cancer, 119*, 2160-2169. <https://doi.org/10.1002/cncr.28057>
- Rottmann, N., Hansen, D. G., Larsen, P. V., Nicolaisen, A., Flyger, H., Johansen, C., & Hagedoorn, M. (2015). Dyadic coping within couples dealing with breast cancer: A longitudinal, population-based study. *Health Psychol, 34*(5), 486-495. <https://doi.org/10.1037/hea0000218>
- Salehi, A., Frommolt, V., & Coyne, E. (2019). Factors affecting provision of care services for patients with cancer living in the rural area: An integrative review. *Australian Journal of Cancer Nursing, 20*(1), 3-13. <https://doi.org/10.33235/ajcn.20.1.3-13>
- Segrin, C., & Badger, T. A. (2013). Interdependent psychological distress between Latinas with breast cancer and their supportive partners. *Journal of Latina/o Psychology, 1*(1), 21-34. <https://doi.org/10.1037/a0030345>
- Segrin, C., & Badger, T. A. (2014). Psychological and physical distress are interdependent in breast cancer survivors and their partners. *Psychology, Health & Medicine, 19*(6), 716-723. <https://doi.org/10.1080/13548506.2013.871304>
- Shajani, Z., & Snell, D. (2019). *Wright & Leahey's Nurses and Families: A Guide to Family Assessment and Intervention* (7 ed.). F A Davies.
- Song, L., Northouse, L. L., Zhang, L., Braun, T. M., Cimprich, B., Ronis, D. L., & Mood, D. W. (2012). Study of dyadic communication in couples managing prostate cancer: a longitudinal perspective. *Psychooncology, 21*(1), 72-81. <https://doi.org/10.1002/pon.1861>
- Stamatakis, Z., Ellis, J. E., Costello, J., Fielding, J., Burns, M., & Molassiotis, A. (2014). Chronicles of informal caregiving in cancer: using 'The Cancer Family Caregiving Experience' model as an

- explanatory framework. *Supportive Care in Cancer*, 22(2), 435-444.
<https://doi.org/10.1007/s00520-013-1994-1>
- Treanor, C. J., Santin, O., Prue, G., Coleman, H., Cardwell, C. R., O'Halloran, P., & Donnelly, M. (2019). Psychosocial interventions for informal caregivers of people living with cancer. *Cochrane Database of Systematic Reviews*(6). <https://doi.org/10.1002/14651858.CD009912.pub2>
- Van Parys, H., Provoost, V., De Sutter, P., Pennings, G., & Buysse, A. (2017). Multi family member interview studies: a focus on data analysis. *Journal of Family Therapy*, 39(3), 386-401.
<https://doi.org/10.1111/1467-6427.12169>
- Vestergaard, L. V., Ostervang, C., Danbjorg, D. B., & Dieperink, K. B. (2019). Video-Based Patient Rounds for Caregivers of Patients With Cancer. *Oncology Nursing Forum*, 46(4), 485-492.
<https://doi.org/10.1188/19.ONF.485-492>
- Whitehead, L., Jacob, E., Towell, A., Abu-qamar, M. e., & Cole-Heath, A. (2018). The role of the family in supporting the self-management of chronic conditions: A qualitative systematic review. *Journal of Clinical Nursing*, 27(1-2), 22-30. <https://doi.org/10.1111/jocn.13775>
- Wittenberg, E., Xu, J., Goldsmith, J., & Mendoza, Y. (2019). Caregiver Communication About Cancer: Development of a mhealth resource to support family caregiver communication burden. *Psycho-Oncology*, 28(2), 365-371. <https://doi.org/10.1002/pon.4950>
- Xiong, C., Biscardi, M., Nalder, E., & Colantonio, A. (2018). Sex and gender differences in caregiving burden experienced by family caregivers of persons with dementia: a systematic review protocol. *BMJ Open*, 8(8), e022779. <https://doi.org/10.1136/bmjopen-2018-022779>
- Yoo, S. H., Kim, M., Yun, Y. H., Keam, B., Kim, Y. A., Kim, Y. J., Shim, H. J., Song, E. K., Kang, J. H., Kwon, J. H., Lee, J. L., Lee, S. N., Kim, S. Y., Kang, E. J., Do, Y. R., Choi, Y. S., & Jung, K. H. (2018). Attitudes toward early palliative care in cancer patients and caregivers: a Korean nationwide survey [Article]. *Cancer Medicine*, 7(5), 1784-1793. <https://doi.org/10.1002/cam4.1441>
- Zaleta, A., McManus, S., Miller, M., Johnson, J., & Buzaglo, J. (2018). Preparedness for care and formal training of family cancer caregivers. *Psycho-Oncology*, 27, 77. <https://doi.org/10.1002/pon.4623>
- Zwahlen, D., Hagenbuch, N., Jenewein, J., Carley, M. I., & Buchi, S. (2011). Adopting a family approach to theory and practice: measuring distress in cancer patient-partner dyads with the distress thermometer. *Psychooncology*, 20(4), 394-403. <https://doi.org/10.1002/pon.1744>

Highlights

- Family caregivers such as children or friends are under-represented in the literature
- Support for older adult family caregivers has not been satisfactory established
- Optimum length for intervention was identified as ten weeks
- Interventions which are multimodal improved family outcomes

Table 1 Summary of studies and themes

Author year Country	Design	Title	Themes	Theme Category
(Akin & Durna, 2013) Turkey	Survey	A comparative descriptive study examining the perceptions of cancer patients, family caregivers, and nurses on patient symptom severity in Turkey	Carers report patients' symptoms as more severe than what patients report.	Who is affected
(Ashouri et al., 2018) Iran	Interview	The perceptions of nurses, patients and family members regarding nurses' empathetic behaviours towards patients suffering from cancer: a descriptive qualitative study.	Need for patient nurse communication	Communication
(Badger et al., 2013) USA	RCT	The effect of psychosocial interventions delivered by telephone and videophone on quality of life in early-stage breast cancer survivors and their supportive partners	Technology is a preferential way for carers and patients to seek support however the preferred mode of technology differs. Counselling can have positive improvements in social wellbeing of patients and carers	Resources
(Badr et al., 2014) USA	Survey	Psychological distress in patients and caregivers over the course of radiotherapy for head and neck cancer.	Patient physical distress influences caregiver distress	Who is affected

(Badr et al., 2008) USA	Survey	Does talking about their relationship affect couples' marital and psychological adjustment to lung cancer?	Communication can impact on marital satisfaction and distress	Communication
(Badr & Carmack Taylor, 2008) USA	Survey	Effects of relationship maintenance on psychological distress and dyadic adjustment among couples coping with lung cancer	Relationship management improves overall outcomes	Communication
(Carmack Taylor et al., 2008) USA	Survey	Lung Cancer Patients and Their Spouses: Psychological and Relationship Functioning Within 1 Month of Treatment Initiation	Counselling improves communication	Communication
(Castellanos et al., 2019) USA	Survey	Impact of patient symptoms and caregiving tasks on psychological distress in caregivers for head and neck cancer (HNC)	Caregiver distress associated with burden	Who is affected
(Collinge et al., 2013) USA	RCT	Touch, Caring, and Cancer: Randomized controlled trial of a multimedia caregiver education program.	Building quality time- specially with physical touch in a relationship improves relationship and reduces patient distress and increases the caregivers self-efficacy in caregiving	Resources communication
(Coyne et al., 2017) Aus	Survey	Strengths and resources used by Australian and Danish adult patients and their family caregivers during treatment for cancer	Work with the family to promote connection to right external resources	Resources
(Dieperink et al., 2018) Denmark	Survey	Family functioning and perceived support from nurses during cancer treatment among	Lack of communication between carer and patient. Lack of acknowledgement of the role of the carer	Communication

		Danish and Australian patients and their families	Social support can help to normalise cancer experience.	
(Dieperink et al., 2016) Denmark	Focus group	Marital rehabilitation after prostate cancer - a matter of intimacy	Program improved communication and intimacy	Communication
(Dockham et al., 2016) USA	Survey	Implementation of a Psychoeducational Program for Cancer Survivors and Family Caregivers at a Cancer Support Community Affiliate: A Pilot Effectiveness Study	Facilitating improved communication between family members improved overall self-efficacy and empowered family so they had higher quality of life	Communication
(Eisemann et al., 2014) USA	Survey	Quality of life in partners of patients with localised prostate cancer	Cancer diagnosis affects both patient and family over longer term	Who is affected
(Elliott et al., 2015) Aus	Survey Interview	Profiles of dyadic adjustment for advanced prostate cancer to inform couple-based interventions	Shared burden of cancer promotion of communication improves outcomes	Communication
(Ellis et al., 2013) USA	Interview	Coping with and factors impacting upon the experience of lung cancer in patients and primary carers	Caring is a complex task which constantly changes. Carers adopt coping strategies to manage the illness period.	Who is affected
(Ellis et al., 2017) USA	Survey	The influence of dyadic symptom distress on threat appraisals and self-efficacy in advanced cancer and caregiving	Improving communication and support improved overall dyad outcomes	Communication

(Gansler et al., 2010) USA	Interview	Evolving information priorities of hematologic cancer survivors, caregivers and other relatives	Changes in information needs and priorities over time	Who is affected
(Goldzweig et al., 2017) Israel	Survey	A Dialogue of Depression and Hope: Elderly Patients Diagnosed with Cancer and Their Spousal Caregivers	Improving communication and support improved overall outcomes	Communication
(Götze et al., 2017) Germany	Survey	Psychological distress of cancer patients with children under 18 years and their partners—a longitudinal study of family relationships using dyadic data analysis	Dyads react as an emotional system rather than individual. Patients and partners similar levels of distress. Unemployment linked to lower social wellbeing. High social support leads to decreased distress for patient and partner	Who is affected
(Gröpper et al., 2016) Germany	Survey	Assessing cancer-related distress in cancer patients and caregivers receiving outpatient psycho-oncological counselling	Caregiver stress equal or higher than patients	Who is affected
(Haj Mohammad et al., 2015) Netherlands	Survey	Burden of spousal caregivers of stage II and III oesophageal cancer survivors 3 years after treatment with curative intent	Caregiver burden related to patients symptoms. Important to assess caregivers in risk	Who is affected
(Harden et al., 2009) USA	RCT	Factors Associated with Prostate Cancer Patients' and their Spouses' Satisfaction with a Family-based Intervention	Carers and patients require support to learn how to cope with cancer. Family based face-to-face support programs are acceptable to carers and patients.	Resources

(Harden et al., 2008) USA	Survey	The influence of developmental life stage on quality of life in survivors of prostate cancer and their partners	Quality of life influenced by life stage	Who is affected
(Hasson-Ohayon et al., 2010) Israel	Survey	Women with advanced breast cancer and their spouses: diversity of support and psychological distress	Patients and carer experience distress during cancer diagnosis. Carers perceive fewer social supports than patients. Social support can have a positive impact on distress.	Who is affected
(Hendriksen et al., 2015) USA	Interview	Worried together: a qualitative study of shared anxiety in patients with metastatic non-small cell lung cancer and their family caregivers	Patients and carers experience anxiety and fear related to uncertainty and death and after death. Change in roles within the partnership and change in what "normal" life is.	Who is affected
(Hendrix et al., 2016) Germany	Survey	Effects of enhanced caregiver training program on cancer caregiver's self-efficacy, preparedness, and psychological well-being	Short term improvement in self-efficacy and stress management from support program	Resources
(Hsieh et al., 2019) Taiwan	Survey	Dispositional mindfulness, self-compassion, and compassion from others as moderators between stress and depression in caregivers of patients with lung cancer	Caregiver mindfulness protective of distress	Who is affected
(Johansen et al., 2018) Norway	Survey	The Effect of Cancer Patients' and Their Family Caregivers' Physical and Emotional Symptoms on Caregiver Burden.	Caregiver burden related to patients symptoms. Important to assess caregivers in risk	Who is affected

(Kauffmann et al., 2016) USA	Survey	Addressing psychosocial needs of partners of breast cancer patients: a pilot program using social workers to improve communication and psychosocial support	Partners are equally likely to feel unsupported by their partner (patient) and feel down or depressed.	Who is affected
(Kershaw et al., 2008) USA	Survey	Longitudinal Analysis of a Model to Predict Quality of Life in Prostate Cancer Patients and their Spouses	Coping mechanisms have an impact on quality of life in carers and patients. Personal characteristics and illness related factors can impact on quality of life.	Resources
(Kim et al., 2008) USA	Survey	Quality of Life of Couples Dealing with Cancer: Dyadic and Individual Adjustment among Breast and Prostate Cancer Survivors and Their Spousal Caregivers	Carers and patients respond to each other's levels of distress. Distress can predict quality of life.	Who is affected
(Kim et al., 2011) USA	Survey	Individual and dyadic relations between spiritual well-being and quality of life among cancer survivors and their spousal caregivers	Spiritual wellbeing impacts on carers and patients quality of life. Wellbeing of one person in the carer/patient dyad impacts on the wellbeing of the other person in the relationship.	Who is affected
(Kim et al., 2015) USA	Survey	Effects of gender and depressive symptoms on quality of life among colorectal and lung cancer patients and their family caregivers.	Couple communication improves outcomes	Communication
(Kissil et al., 2013) USA	Focus group	I knew from day one that Im either gonna fight this thing or be defeated': African American parents experience of coping with breast cancer	Culturally aware support groups are beneficial	Resources

(Kubo et al., 2018) USA	Survey Interview	A Pilot Mobile-based Mindfulness Intervention for Cancer Patients and their Informal Caregivers.	Mindfulness app feasible to reduce distress	Resources
(La & Yun, 2017) Korea	Survey	Effects of stress appraisal on the quality of life of adult patients with multiple myeloma and their primary family caregivers in Korea	Interventions designed to reconstruct negative perspectives and improve self-efficacy may help both patients and caregivers to improve their quality of life	Resources
(Lageman et al., 2015) USA	Survey	Exploring primary brain tumor patient and caregiver needs and preferences in brief educational and support opportunities	Targeted interventions that are short improve outcomes	Resources
(Lambert et al., 2013) Aus	Interview	"You need something like this to give you guidelines on what to do": patients' and partners' use and perceptions of a self-directed coping skills training resource	Books brochures to information sources preferred booklets rather than DVD	Resources
(Li et al., 2018) China	Survey	The impact of depression and anxiety on quality of life in Chinese cancer patient-family caregiver dyads, a cross-sectional study.	Caregiver dyad distress influenced overall distress	Who is affected
(Liang et al., 2019) USA	Survey	Rates and Risk Factors for Post-Traumatic Stress Disorder Symptomatology among Adult Hematopoietic Cell Transplant Recipients and Their Informal Caregivers.	Caregiver experienced greater distress	Who is affected

(Libert et al., 2013) Belgium	Survey	The ability of informal primary caregivers to accurately report cancer patients' difficulties	Caregiver reported patient difficulties may not always be accurate	Who is affected
(Locher et al., 2010) USA	Interview	Disruptions in the organization of meal preparation and consumption among older cancer patients and their family caregivers.	Spouse experience mental health worse than patients View of unit of care, family cohesion influence mental health	Who is affected
(Lyons et al., 2014) USA	Survey	Symptom Incongruence Trajectories in Lung Cancer Dyads	Patients' symptoms are reported as more severe by carers than patients self-reports.	Who is affected
(Lyons et al., 2016) USA	Survey	The Roles of Dyadic Appraisal and Coping in Couples With Lung Cancer	Importance of food preparation and diet in reducing distress in daily living	Resources
(Manne et al., 2012) USA	Survey	A longitudinal analysis of intimacy processes and psychological distress among couples coping with head and neck or lung cancers	Communication can impact on distress and intimacy in carers and patients	Communication
(Manne et al., 2015) USA	Survey	Holding Back, Intimacy, and Psychological and Relationship Outcomes Among Couples Coping With Prostate Cancer	Carer and patients have reluctance to communicate sexual concerns during cancer diagnosis. Hesitancy to communicate has negative impacts on intimacy.	Communication
(Milbury et al., 2012)	Survey	The Role of Blame in the Psychosocial Adjustment of Couples Coping with Lung Cancer	Smoking status in patients and carers correlates with blame associated with lung cancer diagnosis.	Who is affected

(Mishra et al., 2018) USA	Interview	Health related quality of life during cancer treatment: Perspectives of young adult (23-39 years) cancer survivors and primary informal caregivers.	Support specific to life phase	Who is affected
(Molassiotis et al., 2011) UK	Survey	Unmet supportive care needs, psychological well-being and quality of life in patients living with multiple myeloma and their partners	Carer experience additional unmet needs to patients which relate to practical tasks of caring. Carer and patients experience negative mental wellbeing up to 5 years after diagnosis	Who is affected
(Morgan et al., 2011) USA	Survey	Cancer Patients with Pain: The Spouse/Partner Relationship and Quality of Life	The quality of the relationship between carer and patient can impact on quality of life	Who is affected
(Moschopoulou et al., 2018) UK	Survey	Post-traumatic stress in head and neck cancer survivors and their partners	Distress years after diagnosis	Who is affected
(Mosher et al., 2018) USA	RCT	Examining the effect of peer helping in a coping skills intervention: a randomized controlled trial for advanced gastrointestinal cancer patients and their family caregivers.	Patient and caregiver outcomes did not change overtime	Who is affected
(Mosher et al., 2016) USA	Interview	Family caregiving challenges in advanced colorectal cancer: patient and caregiver perspectives	Assessment of concerns Challenges related to physical aspects of caregiving	Who is affected

(Mosher et al., 2015) USA	Interview	Coping with physical and psychological symptoms: a qualitative study of advanced lung cancer patients and their family caregivers.	Social network often used for emotional and practical assistance	Resources
(Newberry et al., 2012) USA	Survey	Identifying family members who are likely to perceive benefits from providing care to a person with a primary malignant brain tumor	Level of distress at diagnosis influenced changes overtime focus on assessment to tailor care	Who is affected
(Northouse et al., 2013) USA	RCT	Randomized clinical trial of a brief and extensive dyadic intervention for advanced cancer patients and their family caregivers	Improving communication and support improved overall dyad outcomes	Communication
(Nund et al., 2015) Aus	Interview	Communication changes following non-glottic head and neck cancer management: The perspectives of survivors and carers.	Communication key influencing factor for adjustment	Communication
(Ovayolu et al., 2015) Turkey	Survey	Pain in cancer patients: pain assessment by patients and family caregivers and problems experienced by caregivers.	Inclusion of family in teaching pain management improves patient outcomes	Resources
(Pasipanodya et al., 2012) USA	Survey	Social Constraints on Disclosure Predict Daily Well-Being in Couples Coping With Early-Stage Breast Cancer	Poor communication about cancer related topics impacts on self-esteem, willingness to share positive and negative events, intimacy and happiness	Communication
(Petruzzi et al., 2013) Italy	Survey	Cross sectional analysis of quality of life of dyads living with brain tumour.	Caregivers need ongoing support to adapt to the patient's illness	Who is affected

(Porter et al., 2008) USA	Survey	Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: Associations with symptoms & distress	<p>Patients feel better about to managing their disease then carers</p> <p>Carers' self-efficacy is influenced by patient's self-management of symptoms.</p>	Who is affected
(Porter et al., 2011) USA	RCT	Caregiver-assisted coping skills training for lung cancer: results of a randomized clinical trial.	Influence of patient caregiver adjustment and self-efficacy	Communication
(Posluszny et al., 2015) USA	Survey	Posttraumatic Stress Disorder (PTSD) Symptoms in Newly Diagnosed Head and Neck Cancer Patients and their Partners	Carers are more likely to experience PTSD related to a cancer diagnosis than patients	Who is affected
(Richardson et al., 2015) NZ	Survey	Psychological support needs of patients with head and neck cancer and their caregivers: A qualitative study.	More psychological support is needed for patients and caregivers	Who is affected
(Robbins et al., 2014) USA	Survey Focus group	Cancer Conversations in Context: Naturalistic Observation of Couples Coping with Breast Cancer	Carers and patients are unlikely to talk about cancer. Carer and patients communicate about different aspects of cancer diagnosis and to different people. Communication type	Communication
(Robinson et al., 2010) Canada	Interview	Just Because You've Got Lung Cancer Doesn't Mean I Will": Lung Cancer, Smoking, and Family Dynamics.	Family influence not always present in cancer diagnosis	Who is affected

(Rottmann et al., 2015) Denmark	Survey	Dyadic Coping Within Couples Dealing With Breast Cancer: A Longitudinal, Population-Based Study	Depression and relationship quality can be impacted on by couples coping skills	Communication
(Segrin & Badger, 2013) USA	Survey	Interdependent Psychological Distress Between Latinas With Breast Cancer and Their Supportive Partners	Carers and patients respond to each other's psychological wellbeing in cultural sub-groups	Communication
(Segrin & Badger, 2014) USA	Survey	Psychological and physical distress are interdependent in breast cancer survivors and their partners	Patients and caregivers psychological wellbeing can impact on each other	Who is affected
(Song et al., 2012) USA	Survey	Study of dyadic communication in couples managing prostate cancer: a longitudinal perspective	Dyadic communication about cancer decreases over the illness trajectory. The stage of illness and role in the person in the dyad impact on communication patterns over time.	Communication
(Yoo et al., 2018) Korea	Survey	Attitudes toward early palliative care in cancer patients and caregivers: a Korean nationwide survey	Perception of palliative care can impact on uptake.	Communication
(Zwahlen et al., 2011) Switzerland	Survey	Adopting a family approach to theory and practice: measuring distress in cancer patient-partner dyads with the distress thermometer	Female carers experience more distress than male carers and patients of both gender.	Who is affected

- Akin, S., & Durna, Z. (2013). A comparative descriptive study examining the perceptions of cancer patients, family caregivers, and nurses on patient symptom severity in Turkey. *European Journal of Oncology Nursing*, 17(1), 30-37. <https://doi.org/10.1016/j.ejon.2012.02.002>
- Ashouri, E., Taleghani, F., Memarzadeh, M., Saburi, M., & Babashahi, F. (2018). The perceptions of nurses, patients and family members regarding nurses' empathetic behaviours towards patients suffering from cancer: a descriptive qualitative study. *Journal of Research in Nursing*, 23(5), 428-443. <https://doi.org/10.1177/1744987118756945>
- Badger, T., Segrin, C., Pasvogel, A., & Lopez, A. M. (2013). The effect of psychosocial interventions delivered by telephone and videophone on quality of life in early-stage breast cancer survivors and their supportive partners. *J Telemed Telecare*, 19(5), 260-265. <https://doi.org/10.1177/1357633x13492289>
- Badr, H., Acitelli, L., & Carmack Taylor, C. (2008). Does talking about their relationship affect couples' marital and psychological adjustment to lung cancer? *J Cancer Surviv*, 2(1), 53-64. <https://doi.org/10.1007/s11764-008-0044-3>
- Badr, H., & Carmack Taylor, C. (2008). Effects of relationship maintenance on psychological distress and dyadic adjustment among couples coping with lung cancer. *Health Psychol*, 27(5), 616-627. <https://doi.org/10.1037/0278-6133.27.5.616>
- Badr, H., Gupta, V., Sikora, A., & Posner, M. (2014). Psychological distress in patients and caregivers over the course of radiotherapy for head and neck cancer. *Oral Oncol*, 50(10), 1005-1011. <https://doi.org/10.1016/j.oraloncology.2014.07.003>
- Carmack Taylor, C., Badr, H., Lee, J., Fossella, F., Pisters, K., Gritz, E., & Schover, L. (2008). Lung cancer patients and their spouses: psychological and relationship functioning within 1 month of treatment initiation. *Ann Behav Med*, 36(2), 129-140. <https://doi.org/10.1007/s12160-008-9062-7>
- Castellanos, E. H., Dietrich, M. S., Bond, S. M., Wells, N., Schumacher, K., Ganti, A. K., & Murphy, B. A. (2019). Impact of patient symptoms and caregiving tasks on psychological distress in caregivers for head and neck cancer (HNC). *Psychooncology*, 28(3), 511-517. <https://doi.org/10.1002/pon.4968>

- Collinge, W., Kahn, J., Walton, T., Kozak, L., Bauer-Wu, S., Fletcher, K., Yarnold, P., & Soltysik, R. (2013). Touch, Caring, and Cancer: Randomized controlled trial of a multimedia caregiver education program. *Supportive Care in Cancer*, 21(5), 1405-1414. <https://doi.org/10.1007/s00520-012-1682-6>
- Coyne, E., Dieperink, K., Østergaard, B., & Creedy, D. (2017). Strengths and resources used by Australian and Danish adult patients and their family caregivers during treatment for cancer. *European Journal of Oncology Nursing*, 29, 53-59. <https://doi.org/10.1016/j.ejon.2017.05.005>
- Dieperink, K., Coyne, E., Creedy, D., & Ostergaard, B. (2018). Family functioning and perceived support from nurses during cancer treatment among Danish and Australian patients and their families *Journal of Clinical Nursing*, 27(1-2), e154-e161. <https://doi.org/10.1111/jocn.13894>
- Dieperink, K. B., Mark, K., & Mikkelsen, T. B. (2016). Marital rehabilitation after prostate cancer – a matter of intimacy. *International Journal of Urological Nursing*, 10(1), 21-29. <https://doi.org/10.1111/ijun.12091>
- Dockham, B., Schafenacker, A., Yoon, H., Ronis, D., Kershaw, T., Titler, M., & Northouse, L. (2016). Implementation of a psychoeducational program for cancer survivors and family caregivers at a cancer support community affiliate: A pilot effectiveness study. *Cancer Nursing*, 39(3), 169-180. <https://doi.org/10.1097/NCC.0000000000000311>
- Eisemann, N., Waldmann, A., Rohde, V., & Katalinic, A. (2014). Quality of life in partners of patients with localised prostate cancer. *Qual Life Res*, 23(5), 1557-1568. <https://doi.org/10.1007/s11136-013-0588-1>
- Elliott, K. E., Scott, J. L., Monsour, M., & Nuwayhid, F. (2015). Profiles of dyadic adjustment for advanced prostate cancer to inform couple-based intervention. *Psychol Health*, 30(11), 1259-1273. <https://doi.org/10.1080/08870446.2015.1043301>
- Ellis, J., Lloyd Williams, M., Wagland, R., Bailey, C., & Molassiotis, A. (2013, Jan). Coping with and factors impacting upon the experience of lung cancer in patients and primary carers. *Eur J Cancer Care (Engl)*, 22(1), 97-106. <https://doi.org/10.1111/ecc.12003>

- Ellis, K., Janevic, M., Kershaw, T., Caldwell, C., Janz, N., & Northouse, L. (2017). The influence of dyadic symptom distress on threat appraisals and self-efficacy in advanced cancer and caregiving. *Supportive Care in Cancer*, 25(1), 185-194.
- Gansler, T., Kepner, J., Willacy, E., Soloe, C., Rupert, D., Jarblum, M., Driscoll, D., Orr, A., Fitzgerald, T., & Esparza, A. (2010). Evolving Information Priorities of Hematologic Cancer Survivors, Caregivers, and Other Relatives. *Journal of Cancer Education*, 1-10. <https://doi.org/10.1007/s13187-009-0034-9>
- Goldzweig, G., Baider, L., Andritsch, E., Pfeffer, R., & Rottenberg, Y. (2017). A Dialogue of Depression and Hope: Elderly Patients Diagnosed with Cancer and Their Spousal Caregivers. *J Cancer Educ*, 32(3), 549-555. <https://doi.org/10.1007/s13187-015-0975-0>
- Götze, H., Friedrich, M., Brähler, E., Romer, G., Mehnert, A., & Ernst, J. (2017). Psychological distress of cancer patients with children under 18 years and their partners-a longitudinal study of family relationships using dyadic data analysis. *Supportive Care in Cancer*, 25(1), 255-264. <https://link.springer.com/content/pdf/10.1007%2Fs00520-016-3411-z.pdf>
- Gröpper, S., van der Meer, E., Landes, T., Bucher, H., Stickel, A., & Goerling, U. (2016). Assessing cancer-related distress in cancer patients and caregivers receiving outpatient psycho-oncological counseling. *Supportive Care in Cancer*, 24(5), 2351-2357. <https://doi.org/10.1007/s00520-015-3042-9>
- Haj Mohammad, N., Walter, A., van Oijen, M., Hulshof, M., Bergman, J., Anderegg, M., van Berge Henegouwen, M., Henselmans, I., Sprangers, M., & van Laarhoven, H. (2015). Burden of spousal caregivers of stage II and III esophageal cancer survivors 3 years after treatment with curative intent. *Supportive Care in Cancer*, 23(12), 3589-3598. <https://doi.org/10.1007/s00520-015-2727-4>
- Harden, J., Falahee, M., Bickes, J., Schafenacker, A., Walker, J., Mood, D., & Northouse, L. (2009). Factors associated with prostate cancer patients' and their spouses' satisfaction with a family-based intervention. *Cancer Nurs*, 32(6), 482-492. <https://doi.org/10.1097/NCC.0b013e3181b311e9>
- Harden, J., Northouse, L., Cimprich, B., Pohl, J., Liang, J., & Kershaw, T. (2008). The influence of developmental life stage on quality of life in survivors of prostate cancer and their partners. *Journal of Cancer Survivorship*, 2(2), 84-94. <https://doi.org/10.1007/s11764-008-0048-z>

- Hasson-Ohayon, I., Goldzweig, G., Braun, M., & Galinsky, D. (2010). Women with advanced breast cancer and their spouses: diversity of support and psychological distress. *Psycho-Oncology*, 19(11), 1195-1204. <https://doi.org/10.1002/pon.1678>
- Hendriksen, E., Williams, E., Sporn, N., Greer, J., DeGrange, A., & Koopman, C. (2015). Worried together: a qualitative study of shared anxiety in patients with metastatic non-small cell lung cancer and their family caregivers. *Support Care Cancer*, 23(4), 1035-1041. <https://doi.org/10.1007/s00520-014-2431-9>
- Hendrix, C., Bailey, D., Steinhauer, K., Olsen, M., Stechuchak, K., Lowman, S., Schwartz, A., Riedel, R., Keefe, F., Porter, L., & Tulsky, J. (2016). Effects of enhanced caregiver training program on cancer caregiver's self-efficacy, preparedness, and psychological well-being. *Supportive Care in Cancer*, 24(1), 327-336. <https://doi.org/10.1007/s00520-015-2797-3>
- Hsieh, C. C., Yu, C. J., Chen, H. J., Chen, Y. W., Chang, N. T., & Hsiao, F. H. (2019). Dispositional mindfulness, self-compassion, and compassion from others as moderators between stress and depression in caregivers of patients with lung cancer. *Psycho-Oncology*, 28(7), 1498-1505. <https://doi.org/10.1002/pon.5106>
- Johansen, S., Cvancarova, M., & Ruland, C. (2018). The Effect of Cancer Patients' and Their Family Caregivers' Physical and Emotional Symptoms on Caregiver Burden. *Cancer Nurs*, 41(2), 91-99. <https://doi.org/10.1097/NCC.0000000000000493>
- Kauffmann, R., Bitz, C., Clark, K., Loscalzo, M., Kruper, L., & Vito, C. (2016). Addressing psychosocial needs of partners of breast cancer patients: a pilot program using social workers to improve communication and psychosocial support. *Supportive Care In Cancer: Official Journal Of The Multinational Association Of Supportive Care In Cancer*, 24(1), 61-65. <https://doi.org/10.1007/s00520-015-2721-x>
- Kershaw, T., Mood, D., Newth, G., Ronis, D., Sanda, M., Vaishampayan, U., & Northouse, L. (2008). Longitudinal analysis of a model to predict quality of life in prostate cancer patients and their spouses. *Ann Behav Med*, 36(2), 117-128. <https://doi.org/10.1007/s12160-008-9058-3>
- Kim, Y., Carver, C. S., Spillers, R. L., Crammer, C., & Zhou, E. S. (2011). Individual and dyadic relations between spiritual well-being and quality of life among cancer survivors and their spousal caregivers. *Psychooncology*, 20(7), 762-770. <https://doi.org/10.1002/pon.1778>

- Kim, Y., Kashy, D. A., Wellisch, D. K., Spillers, R. L., Kaw, C. K., & Smith, T. G. (2008). Quality of life of couples dealing with cancer: dyadic and individual adjustment among breast and prostate cancer survivors and their spousal caregivers. *Ann Behav Med*, 35(2), 230-238. <https://doi.org/10.1007/s12160-008-9026-y>
- Kim, Y., van Ryn, M., Jensen, R., Griffin, J., Potosky, A., & Rowland, J. (2015). Effects of gender and depressive symptoms on quality of life among colorectal and lung cancer patients and their family caregivers. *Psycho-Oncology*, 24(1), 95-105. <https://doi.org/10.1002/pon.3580>
- Kissil, K., Niño, A., Ingram, M., & Davey, M. (2013). "I Knew From Day One That I'm Either Gonna Fight This Thing or Be Defeated": African American Parents' Experiences of Coping With Breast Cancer. *Journal of Family Nursing*, 20(1), 98-119. <https://doi.org/10.1177/1074840713504035>
- Kubo, A., Altschuler, A., Kurtovich, E., Hendlish, S., Laurent, C., Kolevska, T., Li, Y., & Avins, A. (2018). A Pilot Mobile-based Mindfulness Intervention for Cancer Patients and their Informal Caregivers. *Mindfulness*, 9(6), 1885-1894. <https://doi.org/10.1007/s12671-018-0931-2>
- La, I. S., & Yun, E. K. (2017). Effects of stress appraisal on the quality of life of adult patients with multiple myeloma and their primary family caregivers in Korea. *Psycho-Oncology*, 26(10), 1640-1646. <https://doi.org/10.1002/pon.4348>
- Lageman, S. K., Brown, P. D., Anderson, S. K., Lachance, D. H., Yan, E., Laack, N. N. I., & Cerhan, J. H. (2015). Exploring primary brain tumor patient and caregiver needs and preferences in brief educational and support opportunities. *Supportive Care in Cancer*, 23(3), 851-859. <https://doi.org/10.1007/s00520-014-2413-y>
- Lambert, S. D., Girgis, A., Turner, J., Regan, T., Candler, H., Britton, B., Chambers, S., Lawsin, C., & Kayser, K. (2013). "You need something like this to give you guidelines on what to do": patients' and partners' use and perceptions of a self-directed coping skills training resource. *Supportive Care in Cancer*, 21(12), 3451-3460. <https://doi.org/10.1007/s00520-013-1914-4>
- Li, Q., Lin, Y., Xu, Y., & Zhou, H. (2018). The impact of depression and anxiety on quality of life in Chinese cancer patient-family caregiver dyads, a cross-sectional study. *Health Qual Life Outcomes*, 16(1), 230. <https://doi.org/10.1186/s12955-018-1051-3>

- Liang, J., Lee, S. J., Storer, B. E., Shaw, B. E., Chow, E. J., Flowers, M. E., Krakow, E. F., Bar, M., Syrjala, K. L., Salit, R. B., Kurukulasuriya, C. E., & Jim, H. S. L. (2019). Rates and Risk Factors for Post-Traumatic Stress Disorder Symptomatology among Adult Hematopoietic Cell Transplant Recipients and Their Informal Caregivers. *Biol Blood Marrow Transplant*, 25(1), 145-150. <https://doi.org/10.1016/j.bbmt.2018.08.002>
- Libert, Y., Merckaert, I., Slachmuylder, J. L., & Razavi, D. (2013). The ability of informal primary caregivers to accurately report cancer patients' difficulties. *Psychooncology*, 22(12), 2840-2847. <https://doi.org/10.1002/pon.3362>
- Locher, J. L., Robinson, C. O., Bailey, F. A., Carroll, W. R., Heimbarger, D. C., Saif, M. W., Tajeu, G., & Ritchie, C. S. (2010). Disruptions in the organization of meal preparation and consumption among older cancer patients and their family caregivers. *Psycho-Oncology*, 19(9), 967-974. <https://doi.org/10.1002/pon.1656>
- Lyons, K., Lee, C., Bennett, J., Nail, L., Fromme, E., Hiatt, S., & Sayer, A. (2014). Symptom incongruence trajectories in lung cancer dyads. *J Pain Symptom Manage*, 48(6), 1031-1040. <https://doi.org/10.1016/j.jpainsymman.2014.02.004>
- Lyons, K., Miller, L., & McCarthy, M. (2016). The Roles of Dyadic Appraisal and Coping in Couples With Lung Cancer. *Journal of Family Nursing*, 22(4), 493-514. <https://doi.org/10.1177/1074840716675976>
- Manne, S., Badr, H., & Kashy, D. (2012). A longitudinal analysis of intimacy processes and psychological distress among couples coping with head and neck or lung cancers. *J Behav Med*, 35(3), 334-346. <https://doi.org/10.1007/s10865-011-9349-1>
- Manne, S., Kissane, D., Zaider, T., Kashy, D., Lee, D., Heckman, C., & Virtue, S. (2015). Holding back, intimacy, and psychological and relationship outcomes among couples coping with prostate cancer. *J Fam Psychol*, 29(5), 708-719. <https://doi.org/10.1037/fam0000096>
- Milbury, K., Badr, H., & Carmack Taylor, C. (2012). The role of blame in the psychosocial adjustment of couples coping with lung cancer. *Ann Behav Med*, 44(3), 331-340. <https://doi.org/10.1007/s12160-012-9402-5>

- Mishra, S. I., Rishel Brakey, H., Kano, M., Nedjat-Haiem, F. R., & Sussman, A. L. (2018). Health related quality of life during cancer treatment: Perspectives of young adult (23-39 years) cancer survivors and primary informal caregivers [Article]. *European journal of oncology nursing : the official journal of European Oncology Nursing Society*, 32, 48-54. <https://doi.org/10.1016/j.ejon.2017.11.007>
- Molassiotis, A., Wilson, B., Blair, S., Howe, T., & Cavet, J. (2011). Unmet supportive care needs, psychological well-being and quality of life in patients living with multiple myeloma and their partners. *Psychooncology*, 20(1), 88-97. <https://doi.org/10.1002/pon.1710>
- Morgan, M. A., Small, B. J., Donovan, K. A., Overcash, J., & McMillan, S. (2011). Cancer patients with pain: the spouse/partner relationship and quality of life. *Cancer Nurs*, 34(1), 13-23. <https://doi.org/10.1097/NCC.0b013e3181efed43>
- Moschopoulou, E., Hutchison, I., Bhui, K., & Korszun, A. (2018). Post-traumatic stress in head and neck cancer survivors and their partners [Article in Press]. *Supportive Care in Cancer*, 1-9. <https://doi.org/10.1007/s00520-018-4146-9>
- Mosher, C., Secinti, E., Johns, S., O'Neil, B., Helft, P., Shahda, S., Jalal, S., & Champion, V. (2018). Examining the effect of peer helping in a coping skills intervention: a randomized controlled trial for advanced gastrointestinal cancer patients and their family caregivers. *Quality of life research*, 27(2), 515-528. <https://doi.org/10.1007/s11136-017-1620-7>
- Mosher, C. E., Adams, R. N., Helft, P. R., O'Neil, B. H., Shahda, S., Rattray, N. A., & Champion, V. L. (2016). Family caregiving challenges in advanced colorectal cancer: patient and caregiver perspectives. *Supportive Care In Cancer: Official Journal Of The Multinational Association Of Supportive Care In Cancer*, 24(5), 2017-2024. <https://doi.org/10.1007/s00520-015-2995-z>
- Mosher, C. E., Ott, M. A., Hanna, N., Jalal, S. I., & Champion, V. L. (2015). Coping with physical and psychological symptoms: a qualitative study of advanced lung cancer patients and their family caregivers. *Supportive Care In Cancer: Official Journal Of The Multinational Association Of Supportive Care In Cancer*, 23(7), 2053-2060. <https://doi.org/10.1007/s00520-014-2566-8>
- Newberry, A., Kuo, J., Donovan, H., Given, B., Given, C. W., Schulz, R., & Sherwood, P. (2012). Identifying family members who are likely to perceive benefits from providing care to a person with a primary malignant brain tumor. *Oncology Nursing Forum*, 39(3), E226-E232. <https://doi.org/10.1188/12.ONF.E226-E232>

- Northouse, L., Mood, D. W., Schafenacker, A., Kalemkerian, G., Zalupski, M., LoRusso, P., Hayes, D., Hussain, M., Ruckdeschel, J., Fendrick, A., Trask, P., Ronis, D., & Kershaw, T. (2013). Randomized clinical trial of a brief and extensive dyadic intervention for advanced cancer patients and their family caregivers. *Psycho-Oncology*, 22(3), 555-563. <https://doi.org/10.1002/pon.3036>
- Nund, R. L., Rumbach, A. F., Debattista, B. C., Goodrow, M. N. T., Johnson, K. A., Tupling, L. N., Scarinci, N. A., Cartmill, B., Ward, E. C., & Porceddu, S. V. (2015). Communication changes following non-glottic head and neck cancer management: The perspectives of survivors and carers. *International Journal of Speech-Language Pathology*, 17(3), 263-272. <https://doi.org/10.3109/17549507.2015.1010581>
- Ovayolu, Ö., Ovayolu, N., Aytaç, S., Serçe, S., & Sevinc, A. (2015). Pain in cancer patients: pain assessment by patients and family caregivers and problems experienced by caregivers [Article]. *Supportive Care in Cancer*, 23(7), 1857-1864. <https://doi.org/10.1007/s00520-014-2540-5>
- Pasipanodya, E. C., Parrish, B. P., Laurenceau, J. P., Cohen, L. H., Siegel, S. D., Graber, E. C., & Belcher, A. J. (2012). Social constraints on disclosure predict daily well-being in couples coping with early-stage breast cancer. *J Fam Psychol*, 26(4), 661-667. <https://doi.org/10.1037/a0028655>
- Petruzzi, A., Finocchiaro, C. Y., Lamperti, E., & Salmaggi, A. (2013). Living with a brain tumor : reaction profiles in patients and their caregivers. *Supportive Care in Cancer*, 21(4), 1105-1111. <https://doi.org/10.1007/s00520-012-1632-3>
- Porter, L. S., Keefe, F. J., Garst, J., Baucom, D. H., McBride, C. M., McKee, D. C., Sutton, L., Carson, K., Knowles, V., Rumble, M., & Scipio, C. (2011). Caregiver-assisted coping skills training for lung cancer: results of a randomized clinical trial. *J Pain Symptom Manage*, 41(1), 1-13. <https://doi.org/10.1016/j.jpainsymman.2010.04.014>
- Porter, L. S., Keefe, F. J., Garst, J., McBride, C. M., & Baucom, D. (2008). Self-efficacy for managing pain, symptoms, and function in patients with lung cancer and their informal caregivers: associations with symptoms and distress. *Pain*, 137(2), 306-315. <https://doi.org/10.1016/j.pain.2007.09.010>
- Posluszny, D. M., Dougall, A. L., Johnson, J. T., Argiris, A., Ferris, R. L., Baum, A., Bovbjerg, D. H., & Dew, M. A. (2015). Posttraumatic stress disorder symptoms in newly diagnosed patients with head and neck cancer and their partners. *Head Neck*, 37(9), 1282-1289. <https://doi.org/10.1002/hed.23760>

- Richardson, A. E., Morton, R., & Broadbent, E. (2015). Psychological support needs of patients with head and neck cancer and their caregivers: A qualitative study. *Psychology & Health, 30*(11), 1288-1305. <https://doi.org/10.1080/08870446.2015.1045512>
- Robbins, M. L., Lopez, A. M., Weihs, K. L., & Mehl, M. R. (2014). Cancer conversations in context: naturalistic observation of couples coping with breast cancer. *J Fam Psychol, 28*(3), 380-390. <https://doi.org/10.1037/a0036458>
- Robinson, C. A., Bottorff, J. L., Smith, M. L., & Sullivan, K. M. (2010). "Just Because You've Got Lung Cancer Doesn't Mean I Will": Lung Cancer, Smoking, and Family Dynamics. *Journal of Family Nursing, 16*(3), 282-301. <https://doi.org/10.1177/1074840710370747>
- Rottmann, N., Hansen, D. G., Larsen, P. V., Nicolaisen, A., Flyger, H., Johansen, C., & Hagedoorn, M. (2015). Dyadic coping within couples dealing with breast cancer: A longitudinal, population-based study. *Health Psychol, 34*(5), 486-495. <https://doi.org/10.1037/hea0000218>
- Segrin, C., & Badger, T. A. (2013). Interdependent psychological distress between Latinas with breast cancer and their supportive partners. *Journal of Latina/o Psychology, 1*(1), 21-34. <https://doi.org/10.1037/a0030345>
- Segrin, C., & Badger, T. A. (2014). Psychological and physical distress are interdependent in breast cancer survivors and their partners. *Psychology, Health & Medicine, 19*(6), 716-723. <https://doi.org/10.1080/13548506.2013.871304>
- Song, L., Northouse, L. L., Zhang, L., Braun, T. M., Cimprich, B., Ronis, D. L., & Mood, D. W. (2012). Study of dyadic communication in couples managing prostate cancer: a longitudinal perspective. *Psychooncology, 21*(1), 72-81. <https://doi.org/10.1002/pon.1861>
- Yoo, S. H., Kim, M., Yun, Y. H., Keam, B., Kim, Y. A., Kim, Y. J., Shim, H. J., Song, E. K., Kang, J. H., Kwon, J. H., Lee, J. L., Lee, S. N., Kim, S. Y., Kang, E. J., Do, Y. R., Choi, Y. S., & Jung, K. H. (2018). Attitudes toward early palliative care in cancer patients and caregivers: a Korean nationwide survey [Article]. *Cancer Medicine, 7*(5), 1784-1793. <https://doi.org/10.1002/cam4.1441>

Zwahlen, D., Hagenbuch, N., Jenewein, J., Carley, M. I., & Buchi, S. (2011). Adopting a family approach to theory and practice: measuring distress in cancer patient-partner dyads with the distress thermometer. *Psychooncology*, 20(4), 394-403. <https://doi.org/10.1002/pon.1744>

Table 2 *Presentation of scales used in the quantitative research*

Variables		Measurements	
Functional assessment	Fatigue Pain Sleep	Lee Fatigue Scale	(Johansen et al., 2018)
		Brief Fatigue Inventory	(Kubo et al., 2018; Porter et al., 2008)
		Multidimensional Fatigue Inventory (MFI)	(Segrin & Badger, 2013)
		Fatigue scale of the Functional Assessment in Chronic Illness Therapy (FACIT) measures	(Lyons et al., 2016)
		Functional Assessment of Chronic Illness Therapy measure.	(Lyons et al., 2014)
		Brief Pain Inventory	(Lyons et al., 2016; Morgan et al., 2011; Porter et al., 2008)
		Visual Analogue Scale	(Ovayolu et al., 2015)
		General Sleep Disturbance Scale	(Johansen et al., 2018)
		Pittsburg Sleep Quality Index (PSQI)	(Kubo et al., 2018)
	Depression	Centre for Epidemiological Studies-Depression Scale	(Badger et al., 2013; Hendrix et al., 2016; Johansen et al., 2018; Kim et al., 2015; Robbins et al., 2014; Rottmann et al., 2015; Segrin & Badger, 2014)
		Beck Depression Inventory (BDI)	(Porter et al., 2008)
		Hospital Anxiety and Depression Scale (HADS)	(Ashouri et al., 2018; Götze, Friedrich, Brähler, Romer, Mehnert, & Ernst, 2017; Haj Mohammad et al., 2015; Kubo et al., 2018; Li et al., 2018; Libert et al., 2013; Molassiotis et al., 2011; Moschopoulou et al., 2018; Petruzzi et al., 2013; Posluszny et al., 2015; Segrin & Badger, 2013)
		State Trait Anxiety inventory (STAI)	(Porter et al., 2008; Segrin & Badger, 2013, 2014)
		PTSD Checklist-Civilian Version (PCL-C)	(Liang et al., 2019; Moschopoulou et al., 2018; Posluszny et al., 2015)
	Distress	Distress Thermometer	(Badr et al., 2014; Goldzweig et al., 2017; Zwahlen et al., 2011)
		Risk for Distress Scale (RFD)	(Ellis et al., 2017; Northouse et al., 2013)

		Profile of Mood States (POMS)	(Kim et al., 2008; Porter et al., 2008)
		Impact of Events Scale (IES)	(Robbins et al., 2014)
		Perceived Stress Scale (PSS)	(Collinge et al., 2013; Dockham et al., 2016; Segrin & Badger, 2013)
		Cancer and Treatment Distress Scale (CTXDS)	(Liang et al., 2019)
		Inventory of Current Concerns (ICC) Scale	(Harden et al., 2008)
Symptom management		Edmonton Symptoms Assessment Scale	(Akin & Durna, 2013)
		Memorial Symptom Assessment Scale (MSAS)	(Johansen et al., 2018; Morgan et al., 2011)
		General Symptom Distress Scale	(Badger et al., 2013; Gröpper et al., 2016; Segrin & Badger, 2013, 2014)
		Shortness of Breath Questionnaire (SOBQ)	(Lyons et al., 2014)
		Functional Assessment of Cancer Therapy-General	(Collinge et al., 2013; Dockham et al., 2016; Harden et al., 2009; Lyons et al., 2014; Lyons et al., 2016; Mosher et al., 2018; Northouse et al., 2013; Petruzzi et al., 2013; Porter et al., 2011; Porter et al., 2008)
		Cancer Self Efficacy Scale	(Dockham et al., 2016; Harden et al., 2009; Harden et al., 2008; Johansen et al., 2018; Kershaw et al., 2008; La & Yun, 2017; Li et al., 2018)
		Expanded Prostate Cancer Index Composite (EPIC) scale	(Harden et al., 2008; Song et al., 2012)
		Brief Symptom Inventory BSI	(Badr et al., 2008; Badr & Carmack Taylor, 2008; Carmack Taylor et al., 2008; Hasson-Ohayon et al., 2010; Kubo et al., 2018; La & Yun, 2017; Lyons et al., 2016; Manne et al., 2012; Milbury et al., 2012; Morgan et al., 2011; Northouse et al., 2013; Ovayolu et al., 2015; Porter et al., 2008; Rottmann et al., 2015; Segrin & Badger, 2013)
Health and wellbeing	QoL	Patient Reported Outcomes Measurement System (PROMS)	(Castellanos et al., 2019; Hendrix et al., 2016; Kubo et al., 2018; Morgan et al., 2011; Mosher et al., 2018; Porter et al., 2011; Porter et al., 2008)
		Cancer Rehabilitation Evaluation System (CARES)	(Libert et al., 2013; Manne et al., 2012)
		Functional Assessment of Cancer Therapy – Lung Cancer (FACT-L)	(Porter et al., 2008)
		Cancer Survivors' Unmet Needs measure (CaSUN)	(Molassiotis et al., 2011)
		Karnofsky Performance Status scale (KPS)	(Petruzzi et al., 2013)
		Zarit Burden Interview	(Elliott et al., 2015; La & Yun, 2017; Mosher et al., 2018)

	Linear Analog Self-Assessment (LASA)	(Lageman et al., 2015)
Physical health	Medical Outcomes study 12-item short form health survey	(Harden et al., 2009; Harden et al., 2008; Kershaw et al., 2008; Kim et al., 2015; Li et al., 2018)
	Medical Outcomes Study 36-Item Short-Form Health Survey	(Eisemann et al., 2014; Kim et al., 2011; Kim et al., 2008; La & Yun, 2017; Lyons et al., 2014; Lyons et al., 2016; Morgan et al., 2011; Newberry et al., 2012; Petruzzi et al., 2013)
	Omega Screening Questionnaire (OSQ) Symptom Scale	(Kershaw et al., 2008; Song et al., 2012)
	Quality of Life in Chronic Illness: FACT-G	(Hendrix et al., 2016)
	OARS Multidimensional Functional Assessment Questionnaire (OMFAQ)	(Hendrix et al., 2016)
	European Organization for Research and Treatment of Cancer Core Quality of Life Questionnaire EORTIC QOL	(Eisemann et al., 2014; Haj Mohammad et al., 2015; Hsieh et al., 2019; Molassiotis et al., 2011; Moschopoulou et al., 2018)
	Benefits of Illness Scale	(Dockham et al., 2016)
	Self-Efficacy Scale	(La & Yun, 2017; Porter et al., 2008)
	Adult Comorbidity Evaluation-27	(Kim et al., 2015)
Psychological health	Mastery Scale	(Newberry et al., 2012)
	Spiritual wellbeing	(Badger et al., 2013)
	Functional Assessment of Chronic Illness Therapy-Spiritual Well-Being Scale (FACIT-Sp)	(Mosher et al., 2018)
	Mental Health Inventory	(Manne et al., 2012; Manne et al., 2015)
	Positive and Negative Affect Schedule (PANAS)	(Pasipanodya et al., 2012; Segrin & Badger, 2013, 2014)
	Satisfaction Questionnaires	(Harden et al., 2009)
	Goldberg Adjective Scale	(Newberry et al., 2012)
	Measure of Current Status (MOCS)	(Mosher et al., 2018)
	Cancer Behaviour Inventory (CBI)	(Johansen et al., 2018)
	Inventory of Complicated Grief	(Lyons et al., 2014)

	COPE	(Carmack Taylor et al., 2008)
	Brief Cope	(Harden et al., 2009; Kershaw et al., 2008; Northouse et al., 2013)
Appraisal of illness	Beck Hopelessness Scale	(Eisemann et al., 2014; Harden et al., 2009; Kershaw et al., 2008; Northouse et al., 2013; Porter et al., 2011; Porter et al., 2008; Song et al., 2012)
	Appraisal of Illness Scale	(Ellis et al., 2017; Harden et al., 2009; Harden et al., 2008; Kershaw et al., 2008; Northouse et al., 2013)
	Appraisal of Caregiving Scale (ACS)	(Harden et al., 2009; Harden et al., 2008; Kershaw et al., 2008; Northouse et al., 2013)
	Mishel Uncertainty in Illness Scale	(Harden et al., 2009; Hsieh et al., 2019; Song et al., 2012)
	Cancer Self-Efficacy Scale (CASE scale)	(Dockham et al., 2016; Ellis et al., 2017; Harden et al., 2009; Harden et al., 2008; Johansen et al., 2018; La & Yun, 2017; Northouse et al., 2013; Porter et al., 2011)
Marital	Lewis Mutuality and Interpersonal Sensitivity Scale	(Dockham et al., 2016; Harden et al., 2009; Lyons et al., 2016; Northouse et al., 2013; Song et al., 2012)
	Dyadic adjustment	(Badr et al., 2008; Badr & Carmack Taylor, 2008; Carmack Taylor et al., 2008; Götze, Friedrich, Brähler, Romer, Mehnert, Ernst, et al., 2017; Manne et al., 2012; Manne et al., 2015; Milbury et al., 2012; Morgan et al., 2011)
	Dyadic Coping Inventory (DCI)	(Rottmann et al., 2015)
	Quality of Marriage Index	(Pasipanodya et al., 2012)
	Marital Adjustment Scale	(Newberry et al., 2012)
	Communication Patterns Questionnaire (CPQ)	(Manne et al., 2012)
	Marital assessment	(Haj Mohammad et al., 2015)
	Personal Assessment of Intimacy in Relationships (PAIR)	(Manne et al., 2012; Manne et al., 2015)
	Emotional-Intimacy Disruptive Behavior (EIDB)	(Lyons et al., 2016)
	Relationship Maintenance Strategies Measure	(Badr & Carmack Taylor, 2008)
General support	Supportive Care Needs Survey (SCNS)	(Haj Mohammad et al., 2015)
	Medical Outcomes Study-Social Support Scale	(Carmack Taylor et al., 2008; Harden et al., 2009; Johansen et al., 2018; Milbury et al., 2012)

	Interpersonal Support Evaluation List (ISEL)	(Newberry et al., 2012)
	Iceland-Expressive Family Functioning Questionnaire (ICE-FFQ)	(Dieperink et al., 2017)
	Iceland Family Perceived Support Questionnaire (ICE-FPSQ)	
	Cancer Support Inventory (CSI)	(Manne et al., 2012)
	Personal Resource Questionnaire	(Song et al., 2012)
	Cancer Perceived Agents of Social Support (CPASS)	(Goldzweig et al., 2017; Hasson-Ohayon et al., 2010; Johansen et al., 2018)
	Social Constraints Scale (SCS)	(Pasipanodya et al., 2012)
	Social Support Scale	(Dockham et al., 2016; Götze, Friedrich, Brähler, Romer, Mehnert, & Ernst, 2017; Kershaw et al., 2008; Moschopoulou et al., 2018; Northouse et al., 2013)
Caregiver	Caregiver reaction assessment Scale	(Collinge et al., 2013; Dockham et al., 2016; Hendrix et al., 2016; Johansen et al., 2018; Petruzzi et al., 2013) Preparedness for Caregiving scale (Hendrix et al., 2016)
	Caregiver Demands Scale (CDS)	(Morgan et al., 2011)
	Caregiver Strain Index	(Porter et al., 2008)
	Social Problem Solving Inventory-Revised (SPSI-R)	(Morgan et al., 2011)
	Self-efficacy Scale for Cancer Caregivers	(Hendrix et al., 2016)
	Positive Aspects of Caregiving scale	(Newberry et al., 2012)
	General Self-Efficacy Scale	(Johansen et al., 2018)
	Family Hardiness Index	(Coyne et al., 2017)
	Family Crisis Orientated Personal Evaluation Scales	

Figure 1

Literature searches, screening and selection of studies

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit www.prisma-statement.org.

Figures

Figure 2

The frequency of articles published grouped by country/geographical area.

Figure 3

Frequency of the types of cancer focused on in each study

Figure 4

Frequency of quantitative and qualitative study design types used in articles included in this review

Figure 5

The intervention types used in each experimental study

Figure 6

The duration of studies collecting prospective data (n = 27)

Search terms within Title and abstract

(family* or caregiver*, and cancer*, neoplasm* and coping*, distress*)

((family and coping and cancer) not paed* Ped* child* not review).ab.

(Family* caregiver* coping; distress; cancer) not paed not review)

(TS=(family) AND TS=(cancer) AND TS=(coping) NOT

TS=(ped* paed* child*) AND

TS=(caregiver)) AND LANGUAGE: (English) AND DOCUMENT

TYPES: (Article)