Non-Invasive Approaches to Functional Recovery after Spinal Cord Injury: Therapeutic Targets and Multimodal Device Interventions

Claudio Pizzolato^{1,2}, Mehmet A. Gunduz⁷, Dinesh Palipana^{2,3,4,5}, Jingnan Wu⁷, Gary Grant⁶, Susan Hall⁶, Rachel Dennison⁷, Ross D. Zafonte⁷, David G. Lloyd^{1,2}, Yang D. Teng^{7*}

¹ School of Allied Health Sciences, Griffith University, Gold Coast, QLD, Australia,

² Griffith Centre of Biomedical and Rehabilitation Engineering, Menzies Health Institute Queensland, and Advanced Design and Prototyping Technologies Institute (ADAPT), Griffith University, Gold Coast, QLD, Australia

³ The Hopkins Centre, Menzies Health Institute Queensland, Griffith University, Gold Coast, QLD, Australia

⁴ Gold Coast Hospital and Health Service, Gold Coast, QLD, Australia

⁵ School of Medicine, Griffith University, Gold Coast, QLD, Australia

⁶ School of Pharmacy and Pharmacology, Griffith University, Gold Coast, Australia

⁷ Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Mass General Brigham, and Harvard Medical School, Boston, MA, United States

*Corresponding author: Yang D. Teng (yang_teng@hms.harvard.edu)

Journal: Experimental Neurology

Keywords: brain-computer interface; multimodal device; neurobiology; neuromodulation; non-invasive therapy; pharmacology; prosthesis; rehabilitation; serotonin; spinal cord injury

Abstract

This paper is an interdisciplinary narrative review of efficacious non-invasive therapies that are becoming increasingly used to restore function in people with chronic spinal cord injuries (SCI). First presented are the secondary injury cascade set in motion by the primary lesion and highlights in therapeutic developments for mitigating the acute pathophysiologic process. Then summarized are the pharmacological strategies for modulation of the noradrenergic, serotonergic, and dopaminergic neurotransmissions to enhance recovery in bench and clinical studies of subacute and chronic SCI. Last examined are how neuromechanical devices (i.e., electrical stimulation, robotic assistance, brain-computer interface, and augmented sensory feedback) could be comprehensively engineered to engage efferent and afferent motosensory pathways to induce neuroplasticity-based neural pattern generation. Emerging evidence shows that computational models of the human neuromusculoskeletal system (i.e., human digital twins) could serve as functionalized anchors to integrate different neuromechanical and pharmacological interventions into a single multimodal prosthesis. The system, if appropriately built, may cybernetically optimize therapies via coordination of heterogeneous biosensory, system output, and control signals. Overall, these rehabilitation protocols involved neuromodulation to evoke beneficial adaptive changes within spared supraspinal, intracord, and peripheral/neuromuscular circuits to elicit neurological improvement. Therefore, qualitatively advancing the theoretical understanding of spinal cord neurobiology and neuromechanics is pivotal to designing new ways to reinstate locomotion after SCI. Future research efforts should concentrate on personalizing combined strategies consisting of pharmacological adjuncts, targeted neurobiological and neuromuscular repairs, and brain-computer interfaces, which follow multimodal neuromechanical strategies.

Table of Contents

Background	4
Contemporary managements for acute SCI	7
Developing a standardized drug treatment for acute SCI.....	7
Other drugs investigated for secondary injury amelioration	8
Adrenergic compounds.....	8
Therapeutic agents to diminish gliosis	9
Axon repair and regeneration agents	10
Stem cell-based approaches.....	11
Efficacious treatments for chronic SCI.....	12
Pharmacological Agents of Neuromodulation for chronic SCI	13
Neuromechanical prostheses to restore function for chronic SCI.....	18
Conclusions.....	23

Background

The debilitating sequelae of spinal cord injuries (SCI) have, until very recently, been considered irreversible. Life expectancies of those with SCI were limited before improvements in acute clinical management emerged in the middle of the twentieth century (Middleton et al., 2012; Savic et al., 2017). Today, people with SCI are successfully reintegrating into society and, to some extent, individuals with incomplete injuries can live independently with nearly normal lifespans (Pershouse et al., 2012). Present reintegration, however, largely relies on compensatory tools and assistive devices rather than sustainable functional recovery (Côté et al., 2017). Therefore, restoration of volitional arm-hand function and/or locomotion remains a top priority (Holanda et al., 2017; Maciejasz et al., 2014). Lack of functional improvement can substantially increase rates of severe complications (e.g., respiratory failure, lower urinary tract disorder, and musculoskeletal defect; Hall et al., 2019).

The initial insult to a spinal cord, whether a contusion, compression, laceration, or penetration, instantly produces physical damage (i.e., the primary injury). Its pathological features depend mainly on the lesion's scale and anatomical location (Ahuja et al., 2017; Freund et al., 2019). Immediately afterwards, cascades of biochemical and pathophysiological events set secondary injury processes in motion, which last for minutes, days, weeks, months, or even years, depending on the specific mechanisms involved, to exacerbate structural and functional losses (Ahuja et al., 2017). Typical secondary biochemical events include excessive release of norepinephrine (Vise et al., 1974) and glutamate or aspartate (i.e., excitotoxicity; Bell et al., 2017; Liu et al., 1991), reduced intraparenchymal oxygen (O_2), raised lactate level and lactate/pyruvate ratios (Bell et al., 2017; Farooque et al., 1996), ionic imbalance (e.g., extra- or/and intracellular Na^+ , K^+ , H^+ , Ca^{2+} , Cl^- , OH^- disturbances; Agrawal and Fehlings, 1996; Teng and Wrathall, 1997), surges of reactive oxygen (ROS) and nitrogen species (RNS; Visavadiya et al., 2016; Yu et al., 2009), mitochondrial dysfunctions (e.g., respiration reduction, $\uparrow$ ROS, and cytochrome *c* release), and caspase activation (Rabchevsky et al., 2020; Teng et al., 2004).

These perturbations produce acute (minutes to hours: e.g., < 48 hours post injury [p.i.] in humans) and subacute (days to weeks: clinically, 48 hours – 14 days p.i.) excitotoxicity (Liu et al., 1999; Wrathall et al., 1992), hemorrhagic necrosis and vascular/blood circulation disruptions (Matsushita et al., 2015; Mautes et al., 2000), ischemia (Brown et al., 2014), acidosis (Li et al., 2021; Teng and Wrathall, 1997), action potential conduction failure (James

et al., 2011), and oxidative lesions (Christie et al., 2008; Liu et al., 2020; Visavadiya et al., 2016). Also occurring in these stages are neural cell necrosis, apoptosis, necroptosis, pyroptosis and parthanatos (Liu et al., 2020; Kuzhandaivel et al., 2010; Springer et al., 1999), axon and myelin destructions (James et al., 2011; Rosenberg et al., 1999a, b), leukocyte infiltration (Fleming et al., 2006), neural and systemic inflammation (Bloom et al., 2020; de Rivero Vaccari et al., 2016), reactive astrogliosis (Anderson et al., 2016), and cystic microcavity formation (Losey et al., 2014; Miyao et al., 2020). Other pathophysiological events are endoplasmic reticulum stress (Saraswat Ohri et al., 2018b) and abnormal autophagy (Saraswat Ohri et al., 2018a).

The events jointly produce neural and vascular cell loss and disrupt ascending and descending neuropathways, neuromuscular communication, and the central pattern generation (CPG) network. These damages result in sensorimotor deficits below the level of injury and acute systemic disorders including respiratory dysfunctions, cardiovascular disorders, gastrointestinal abnormality and gut dysbiosis (Aarabi et al., 2012; Furlan and Fehlings, 2008; Kabatas et al., 2008; O'Connor et al., 2018) (**Figure 1**). The body, challenged by the primary and secondary assaults, simultaneously activates its repair responses represented by transient neural stem cell (NSC) proliferation, beneficial inflammation, endocannabinoid production, and tissue healing. But the spontaneous responses are mostly futile attempts to restore neural function (**Figure 1**); however, they continually take place during all post-SCI stages, offering potential therapeutic targets (Arevalo-Martin et al., 2012; Dias et al., 2018; Teng, 2019a).

In the intermediate (weeks to months: 14 days – 6 months p.i. in humans) and chronic phases of SCI (months to years: clinically, > 6 months p.i.), accumulating secondary outcomes, from a particular lesion level and severity combination, gradually reach an equilibrium state. This is built via the interaction between the pathophysiological events and the body's endogenous healing mechanisms (see details above). Such an equilibrium defines a chronic clinical profile unique to each individual SCI case (Krishna et al., 2014). Without effective therapies, any early chronic stage “equilibrium” will inevitably deteriorate. The downfall is caused by a slowly worsened interstitial environment, characterized by buildup of trace amines and their metabolites (Gozal et al., 2014; Li et al., 2017), pro-neuroinflammatory milieu, and a degenerating neuromusculoskeletal network (**Figure 1**) (Fleming et al., 2006; Clark and Findlay, 2017). For example, trace amines produced by pericytes were shown to activate receptors in the absence of monoamine neurotransmitters in

the lesioned cord, which, depending on dose levels, could hinder motosensory function by inducing hypoxia and vasoconstriction (Gozal et al., 2014; Li et al., 2017).

The chronic pathology is characterized by axon demyelination (Guest et al., 2005), syrinx formation and enlargement (Krebs et al., 2016), astroglial scar aging (Yoon et al., 2018), and serotonin (5-hydroxytryptophan: 5HT) deprivation in the below-lesion cord (Ganzer et al., 2018). These are coupled with the maladaptive plasticity of the brain (Freund et al., 2011), spinal cord (Cadotte et al., 2012; Strain et al., 2019), and peripheral nerve and organs (Chariker et al., 2019; Kern et al., 2017; van De Meent et al., 2010). The impairments of muscles and bones jeopardize rehabilitation opportunities (Frontera et al., 2006; Clark and Findlay, 2017; Morse et al., 2008). Serious complications commonly arise in the respiratory (van Silfhout et al., 2016), lower urinary tract (Chan et al., 2018; Kanai et al., 2011), gastrointestinal (Holmes and Blanke, 2019; Jogia and Ruitenberg, 2020), cardiovascular (e.g., autonomic dysreflexia. (Dorton et al., 2020; Eldahan et al., 2020), and reproductive systems (Stoffel et al., 2018) (**Figure 1**).

In the USA, ~45.4% of patients were discharged with cervical SCI; the next most frequent injury levels were T12 (~6.1%) and L1 (~4.7%) (i.e., thoracolumbar SCI; 2019 data of The National Spinal Cord Injury Statistical Center: www.nscisc.uab.edu). Global data had similar distribution features (Lee et al., 2014). To consistently describe SCI, the International Standards for Neurological Classification of Spinal Cord Injury (ISNCSCI) was first published in 1982 by the American Spinal Injury Association (ASIA; Chicago, IL; 1982 [Google Scholar]). With periodic revisions, ISNCSCI (i.e., the ASIA Impairment Scale: AIS-A to E) has since become the most widely used SCI classification system (Kirshblum et al., 2020; Maynard et al., 1997). Based on the 2019 NSCISC data, at acute admission, 44.3% of individuals were diagnosed as AIS-A (complete), 12% as AIS-B (sensory only), 13.9% as AIS-C (non-functional motor), 19% as AIS-D (functional motor), and 10.9% as AIS-E (recovered).

Clinically, trauma to the spinal cord usually does not completely sever it. People with acute SCI who exhibit residual neurological signs below the lesion level were likely to have better prognoses for some natural or rehabilitation-induced recovery (Scivoletto et al., 2014). For example, there were 19.0% and 11.2% AIS-D participants at acute and rehabilitation admissions, respectively, which improved to 32.2% at system discharge (NSCISC: 2019

data). These facts have laid down the foundation for developing pharmacological, cell, bioengineering, gene, rehabilitation, and multimodal therapies to preserve, protect, regenerate, and reactivate the available neural matrix in the injured spinal cord to recover function.

Contemporary management of acute SCI

The management of acute SCI has successfully evolved from basic life support and wound care to comprehensive approaches. Prior to the end of the second world war, SCI was the most lethal encounter of all traumas and invariably resulted in a dismal prognosis. Marked progress has since been made to qualitatively improve outcomes (Shank et al., 2019). Present common protocols involve surgical decompression and stabilization, with or without neuroprotective or anticoagulative pharmacological interventions, treatment or prevention of infection and complication, and optimization of patients' options for rehabilitation (Côté et al., 2017; Furlan et al., 2011).

Decompression and stabilization, albeit with lack of consensus about timing of intervention, can minimize acute impairments to the cord that would otherwise aggravate pre-existing conditions and jeopardize clinical prognosis (Fehlings et al., 2012; Kim et al., 2018). Importantly, decompression via laminectomy and myelotomy was first investigated by Dr. Alfred Reginald Allen in 1911, in which the intervention improved neural outcomes in an experimental model of spinal cord contusion (Allen, 1911). Whether myelotomy has tangible clinical efficacy for acute SCI remains to be further verified (Koyanagi et al., 1989; Telemacque et al., 2020). Thus, for standardizing surgical treatment protocols, using an evidence-based decision-making approach is crucial. To achieve better clinical outcomes, combinational use of pharmacological treatment is imperative (Fehlings et al., 2017).

Developing a standardized drug treatment for acute SCI:

Bench-side pharmacological studies in the early 1980s uncovered the effects of methylprednisolone (MP; a glucocorticoid) on inhibition of lipid peroxidation in cell membranes, reduction of intraparenchymal ischemia, and enhancement of aerobic energy metabolism in animal models (Hall and Braughler, 1981; Hall et al., 1984; Young and Flamm, 1982). The findings, together with the clinical anti-inflammatory and antiedemic effects of MP, facilitated the launch of the Second National Acute Spinal Cord Injury Study (NASCIS II; Bracken et al., 1990). In an earlier trial, which was termed NASCIS I *post hoc*, a smaller dose regimen of MP given over 10 days post SCI produced no significant therapeutic

results (Bracken et al., 1984; Bracken et al., 1985). In distinction, evaluating MP, naloxone, and placebo, the NASCIS II trial revealed significant benefits of high-dose MP (30 mg/kg, i.v. bolus, plus a 23-hour infusion at 5 mg/kg/hour) when given within 8 hours p.i. (Bracken et al., 1990). Sustained effects were observed in neurologically complete or incomplete recipients. However, adverse effects typically associated with steroid therapy were recorded, including gastrointestinal bleeding, wound infections, and delayed wound healing, although the reported incident frequency was not statistically higher than that of placebo-treated patients (Bracken et al., 1990).

In the early 1990s, the aforementioned results enabled the Upjohn Company, the MP supplier, to register the drug for treating acute SCI in Canada, several Western European countries, and most Far Eastern countries. In contrast, the Food and Drug Administration (FDA) of the USA, in approving any application of a drug for the intended population, required tangible evidence of efficacy from two adequately controlled clinical trials (Darrow et al., 2020). This criterion made filing a new drug application for the MP treatment of acute SCI infeasible based solely on data from NASCIS II. NASCIS III, although validating MP's benefits, had no placebo control group due to ethical concerns about the exclusion of treating SCI patients with an MP formula holding reported advantages. Also, the evidence for claimed replications of NASCIS II's outcomes from other countries was deemed insufficient (Otani et al., 1994; Petitjean et al., 1998).

The regimen, nevertheless, was subsequently adopted as the standard of care for acute SCI, because, worldwide, MP had been in clinical use for many other indications (Bracken et al., 1997). Notably, in more recent clinical studies, the efficacy of high-dose MP appeared questionable (Suberviola et al., 2008). The treatment was also linked with significantly higher rates of severe complications (e.g., pneumonia, decubitus ulcer, and thrombosis; Chikuda et al., 2014). As such, many institutions decided to stop treating SCI with systemic high-dose MP (Chikuda et al., 2014; Felleiter et al., 2012). Since then, the field has not come up with another internationally accepted drug treatment for acute SCI.

Other drugs investigated for secondary injury amelioration:

Adrenergic compounds: Intraparenchymal hemorrhagic necrosis has been a classic drug target due to its aggravating effect on other secondary events (e.g., sizes of lesion volume and gliosis) and therapeutic opportunities (e.g., cell or biomaterial transplantation; Hong et al., 2017). Early studies aimed at impeding the formation of hypoxic hemorrhagic necrosis and

lesion size tested norepinephrine synthesis blocker alpha methyl tyrosine (or alpha receptor antagonists phenoxybenzamine and mellaril; Osterholm, 1974; Osterholm and Mathews, 1972). The treatment yielded some positive pilot results that were, however, not replicable (Gurden and Feringa, 1974). Conversely, administration of clenbuterol, a β_2 -adrenoceptor agonist, significantly ameliorated ischemia/reperfusion injury-induced spinal motor neuron loss and locomotion deficit in rabbits (Chen et al., 2013). Epinephrine (agonist of both alpha and beta adrenoceptors) and isoprenaline (nonselective agonist of beta adrenoceptors) treatment significantly suppressed cytotoxic edema of astrocytes *in vitro* and in acutely injured rat spinal cords by activating beta adrenoceptors *in vivo* (Vardjan et al., 2016). Taken together, future investigations to validate and translate these findings should determine the role of each adrenoceptor subtype in triggering specific pathophysiological events (e.g., vasogenic edema after ischemia/reperfusion versus cytotoxic edema post contusion) in a time-dependent manner.

Therapeutic agents to diminish gliosis: The proliferation of reactive astrocytes and microglia generates a glial scar that has long been postulated to be a physical barrier to axonal regeneration. Aiming to overcome this obstacle, trypsin, hyaluronidase, and elastase were systematically investigated for the purpose of dissolving the glial scar to promote axon regeneration. It was eventually concluded that enzymatic digestion of the glial scar tissue cannot induce axon regrowth and locomotion recovery (Guth et al., 1980).

The post-SCI glial scar (i.e., the glial cell border surrounding the lesion core) is formed by multiple types of cells (e.g., astrocyte, microglia, and NG2 glia). The extracellular matrix (ECM) in the core region contains a variety of proteoglycans or proteoglycan-like molecules such as chondroitin sulphate proteoglycans (CSPGs) and cytotactin/tenascin that were found to be inhibitors of axonal growth (Carlstedt et al., 1989; McKeon et al., 1991). The focus was thereby shifted towards developing chondroitinase ABC (ChABC), which removes glycosaminoglycan chains on the CSPGs, into a therapeutic agent (Bradbury et al., 2002). Numerous studies showed the effect of ChABC on shrinking the scale of glial scars or axon dieback and promoting corticospinal tract regeneration in different SCI models (Busch et al., 2009; Houle et al., 2006).

Importantly, the regrowth of serotonergic axons may not be affected by the presence of the astroglial scar (Hawthorne et al., 2011). Recent papers also described that SCI animals receiving genetic ablation of astrocytic scar had higher degrees of axon dieback and no

regrowth of corticospinal and sensory tracts across the lesion site when compared to control treatment (Anderson et al., 2016). In post-T9-10 transection rats that recovered locomotion under epidural electrical stimulation (EES), quipazine (a 5HT agonist) treatment, and training, the total amount of CSPG expression significantly increased with time after injury (Al'joboori et al., 2020). Hence, deeper understanding of the divergent impacts produced by different cell types and ECM components in the glial scar will facilitate identification of effective therapeutic targets for spinal cord repair (Fawcett, 2020).

Axon repair and regeneration agents: Promoting axonal function and regeneration has been another primary focus in therapeutic developments for SCI. Reviewed here were three different approaches. First, a voltage dependent potassium channel antagonist, 4-aminopyridine (4-AP), was tested for enhancing action potential propagation in demyelinated axons of injured spinal cords to improve locomotion (Gruner and Yee, 1999; Segal and Brunnemann, 1998). Although gait and locomotor function were improved in many studies, benefits were variable and certain side effects (e.g., vasospasm) occurred, preventing it from being formally licensed to treat clinical SCI (DeForge et al., 2004; Grijalva et al., 2003; Segal et al., 1999). Encouragingly, 4-AP (Dalfampridine/Ampyra[®]) received FDA approval in 2010 to strengthen ambulation in patients with multiple sclerosis (MS) after multiple phase II and III trials had demonstrated its enhancement of both walking speed and function in about one third of MS patients (Dunn and Blight, 2011).

Second, blocking Nogo, an inhibitor of axonal outgrowth, is another strategy to treat SCI. For this purpose, both Nogo Receptor-66 antagonists or antibodies against Nogo-A have been intensively investigated. *In vivo* animal studies showed that these agents stimulated axonal sprouting of the corticospinal, rubrospinal, and serotonergic axons into and across the lesion area, and promoted gait and motor recovery (Freund et al., 2007; GrandPré et al., 2002; Merkler et al., 2001). Despite some replicative studies failing to find significant benefit of a Nogo-66 antagonist, peptide NEP1-40 (Steward et al., 2008), the anti-Nogo-A antibody AT1355 showed tolerability by participants when evaluated in the first phase I study, permitting the trial to advance into phase II (Kucher et al., 2018).

Third, developing agents to disrupt the RhoA/Rho kinase pathway, which reportedly had many negative effects after SCI, including neurite growth inhibition, cell death, demyelination, inflammation, and neuropathic pain (Monnier et al., 2003). Two types of drugs to block this pathway have been extensively studied: RhoA inactivators and Nogo

inhibitors (Monnier et al., 2003). BA-210 (Cethrin[®]), a RhoA inhibitor, was evaluated in a small number of participants and found to have variable outcomes. In a cohort of 48 patients with complete SCI (cervical: n=16, thoracic: n=32), BA-210 treatment was found to be more beneficial for individuals with cervical injuries than thoracic injuries, but there was high inter-patient variability (Fehlings et al., 2011).

In general, drug development is a complex, lengthy, and costly process. In the case of BA-210/Cethrin[®], the drug took a circuitous route to enter a phase IIb/III trial under the name of VX-210 (Fehlings et al., 2018), following even earlier studies in 2000 and the clinical evaluation run by Alseres Pharmaceuticals in 2007 (Baptiste et al., 2009). Because interim analyses failed to show efficacy, Vertex Pharmaceuticals Inc., the trial sponsor, decided to halt development of VX-210 in 2019. Clinical trials like these underscored some of the major difficulties in developing therapies for SCI, regarding how to select therapeutic candidates, interpret results, determine appropriate control data, use adequate statistical analyses, and assess proper cause and effect relationships (Friede et al., 2018).

Stem cell-based approaches: Propelled by advances in stem cell biology and material science research, new opportunities to invent neural repair strategies comprising stem cells, with or without biomaterial scaffolding or encapsulation, have been actively pursued (Teng et al., 2002; Vismara et al., 2017). Nationally, the Japanese government in 2019 approved a therapy of systemically administering human mesenchymal stromal stem cells (hMSCs) for patients with acute SCI. This instantly triggered concerns from the international community, mainly about the clinical trial's design that was the basis for the approval (e.g., small sample size, lack of a double-blinded approach, no inclusion of a sham operation group, etc.; Cyranoski, 2019a, b). Japan's ministry of health formally responded to the criticisms, reiterating impracticality to run a double-blinded study on an autologous cell therapy and the ethical dilemma of sham surgeries for neurotrauma patients (Miyamoto, 2019). Clearly, more laboratory and clinical investigations are needed before drawing conclusions according to internationally agreeable criteria.

Nevertheless, in the Japanese protocol, the impact of hMSCs (i.v.) was largely derived from their ability to secrete trophic factors and other cytokines to suppress inflammation (and other secondary injury events) and stimulate beneficial plasticity. These mechanisms are different from conventional neural repair roles of stem cells (e.g., neural cell replacement and promotion of long-distance axon regeneration). Characterized more recently was the innate

“functional multipotency” of stem cells (e.g., MSCs, NSCs, etc.), enabling them to exert neurotrophic, anti-inflammatory, immunoregulatory, and homeostatic influence (Osaka et al., 2010; Teng, 2019a). The classic definition of totipotency, pluripotency and multipotency of stem cells principally touches on their capacity to differentiate into different phenotypes of lineage-designated cells. In distinction, functional multipotency reveals a more enriched repertoire of signaling, epigenetic, genetic, and cellular events that allow stem cells to build homeostasis during development and adulthood to maintain function of an organ system (Teng, 2019b).

Implementing this stem cell theory has played an important role in correctly investigating stem cells to advance basic biology and therapeutic translation. Our studies helped to define *recovery neurobiology* (Ropper et al., 2017; Teng, 2019a), which describes the ability of the injured adult spinal cord, under proper treatment, to deploy polysynaptic neural circuits different from normal neurophysiological pathways to enable postinjury restoration of function. Activation of the essential components of recovery neurobiology (i.e., propriospinal projection network, serotonergic neuromodulation, neuromuscular junction, and CPG) has effectively reinstated function for both experimental and clinical SCI (Gill et al., 2018; Ropper et al., 2017).

Efficacious treatments for chronic SCI

Primary challenges of chronic SCI are worsening disability, lack of spontaneous improvement below injury level, complications, and neuromuscular/musculoskeletal deterioration (**Figure 1**) (Adams and Hicks, 2005; Huie et al., 2017). Yet, compared to limited progress made in therapeutic management of acute SCI, there have been remarkable achievements in developing clinical rehabilitation therapies for chronic SCI (Gagnon et al., 2018; Rupp et al., 2015). This is attributable to newly gained understanding of neuroplastic mechanisms and their potentiators in both experimental and clinical settings, which led to breakthroughs in designing functional recovery strategies targeting the reactivation of the CPG circuitry (Behrman et al., 2006; Donati et al., 2016; Gill et al., 2018; Harkema et al., 2011; Hubscher et al., 2018).

Appropriate electrical stimuli that trigger CPG activation have been shown to induce functional recovery in animal models and individuals with even severe SCI (Gerasimenko et al., 2008; Harkema et al., 2011; Megía García et al., 2020;). As described above, clinically classified complete injuries rarely destroy the entire CPG network that is mainly in the

lumbar cord (Harnie et al., 2019; O'Shea et al., 2017). Unlike the corticospinal tract, the segmental spinal neural circuits do not mediate volitional movement. They, however, can elicit spinal locomotion pattern generation, serving as a target for developing pharmacological neuromodulation, computer interfacing, functional stimulation, and comprehensively integrated rehabilitation systems (e.g., the human digital twin; see below; Krucoff et al., 2016; McPherson et al., 2015; Pizzolato et al., 2019; Sasada et al., 2014).

Pharmacological Neuromodulation for Chronic SCI:

Based on the pathophysiological characteristics of chronic SCI (**Figure 1**), the late phase functional deficit severity of each patient is determined by “the equilibrium” interactively reached between the primary lesion plus secondary damages and the endogenous repair process. To approach a higher functional state in the dynamic system over time p.i., it is preferable to therapeutically optimize the recovery activities (Teng, 2019a). It has been demonstrated that pharmacological neuromodulations, when used as part of a multifaceted strategy, are efficacious in refining neuroplasticity, pain management, neurologic function, and complication reduction. These treatments are believed to work mainly by enhancing communications of the CPG network, spinal cord learning circuitry, and sensorimotor, neuromuscular-musculoskeletal, and inter-neurotransmission systems (Grau et al., 2020; Knikou et al., 2017; Rodgers et al., 2019).

Modulation of monoamine neurotransmitters in the descending noradrenergic, serotonergic, and dopaminergic pathways have surfaced as the leading pharmacological foci. For instance, based on the distinct spectrum of kinematic, kinetic, and clonus electromyogram (EMG) characteristics that each monoaminergic receptor modulated, optimal combinations of pharmacological treatments and electrical spinal cord stimulation were designed. The multimodal monoaminergic therapy restored coordinated hindlimb locomotion with nearly normal weight bearing without supraspinal innervation in paralyzed rats after spinal cord transection (Musienko et al., 2011). In general, studies have examined the mechanisms of drug action on direct activation of the (1) noradrenergic/adrenergic, (2) serotonergic, and (3) dopaminergic neurotransmissions in the spinal cord regarding plastic changes in the CNS, enhanced proprioceptive feedback, and improvements in muscular and/or cardiovascular fitness (see details in **Table 1**).

<Insert Table 1 about here>

(1) Adrenoceptor (α_2 and β_2) agonists: Clonidine, acting as an α_2 -adrenoceptor agonist, potentiated locomotor recovery in chronic spinalized models by augmenting crossed extensor responses via stimulating CPG neurons (Barbeau and Rossignol, 1991; Frigon et al., 2012; Naftchi, 1982). Clinically, with incorporation of progressive weight support and treadmill training, two participants with stabilized spastic paraplegia of SCI after receiving a coadministration of clonidine and cyproheptadine exhibited less clonus and more phasic EMG activity in their lower limbs during overground locomotion with the aid of Canadian crutches. Cyproheptadine is a first-generation antihistamine with anticholinergic, antiserotonergic, and local anesthetic properties (Fung et al., 1990). However, when used alone, clonidine improved locomotion in spastic paretic condition but not in paraplegic SCI (Stewart et al., 1991). The data suggested that neuromodulation for sensorimotor improvement after SCI involved multiple neurotransmitter and neuromuscular systems. Noticeably, all published human studies were in case report format and mostly performed in the 1990s. More investigations on this topic are needed, especially those designed to determine the mechanism of clonidine's impact on spasm and locomotion, to prevent serious side effects (e.g., bradycardia; Rosenblum 1993), and to achieve additive or synergistic outcomes when applied with other classes of drugs and gait training (Domingo et al., 2012; Frigon et al., 2012).

Clenbuterol, at low dose, is a long acting β_2 -adrenoceptor agonist ($T_{1/2}$: 25-39 h). In earlier experimental reports, clenbuterol markedly reduced muscle atrophy and neuromuscular scoliosis by retarding denervation-induced loss of muscle contractility and bone mineralization in models of peripheral nerve lesion and SCI, respectively (Zeman et al., 1987; Zeman et al., 1997). It also spared white matter and enhanced hindlimb function in SCI rats, likely by increasing the expression of nerve growth factor (NGF) and basic fibroblast growth factor (bFGF/FGF2; Hayes et al., 1995; Zeman et al., 1999). However, since 2006, clenbuterol has not been included as an ingredient in any therapeutic drug approved by the FDA because of its adverse effects and abuse potential (www.drugbank.ca/drugs/DB01407).

In contrast, salbutamol, an FDA-approved β_2 -adrenoceptor agonist for bronchodilation, used in short term (i.e., 14 or 21 days) administration increased voluntary muscle strength, including the maximum static inspiratory mouth pressure in man. However, there were variations in the magnitude and duration of this effect between muscle groups (Martineau et al., 1992). Prolonged use of metaproterenol, another β_2 -agonist, strengthened skeletal

muscles in participants with chronic SCI (Signorile et al., 1995). In a recent randomized, placebo-controlled, cross-over study (n = 12 trained men), salbutamol (6 × 4 mg, p.o.) augmented protein turnover rates in skeletal muscle 0.5-5 h after resistance exercise, through concomitant cAMP/PKA and Akt2 signaling (Hostrup et al., 2018). It is worth noting that although newer generation β_2 -agonists (e.g., formoterol) experimentally induced anabolic responses in skeletal muscle following denervation at doses linked with fewer side-effects on the heart, compared to clenbuterol and fenoterol (Mayne et al., 2015), their potential to harm the cardiovascular system has not been completely abolished (Hostrup et al., 2020).

(2) Serotonergic (5HT: 5-hydroxytryptamine) agonists: The potency of 5HT_{1A} activation for functional restoration after SCI was first demonstrated by the reversal of respiratory abnormalities resulting from either T8 contusion or C5 hemicontusion in rats following treatment with buspirone and 8-OH-DPAT (Choi et al., 2005; Teng et al., 2003). The effect was mediated through 5-HT_{1A} receptors because both buspirone (BuSpar[®]: a 5HT_{1A} partial agonist) and 8-OH-DPAT (a 5HT_{1A/7} full agonist, depending on species) improved post-SCI respiration. Furthermore, pretreatment of animals with p-MPPI, a 5-HT_{1A} selective antagonist, abolished the beneficial impact of either drug (Choi et al., 2005; Teng et al., 2003). The data suggested that activation of 5HT_{1A} receptors could be an effective strategy for recovering respiratory function and other types of patterned motor activities p.i. (Teng et al., 2003).

In vitro, N-methyl-D-aspartate (NMDA) induced fixative locomotion in the neonatal rat spinal cord. Adding 5-HT to the preparation improved left/right and flexor/extensor alterations that were recorded from the 3rd and 5th lumbar ventral roots, respectively. Although the effect could only be partly reproduced by stimulating 5-HT_{2A/2c} receptors, daily administration of p-chloro-phenylalanine (PCPA) to suppress 5-HT synthesis (starting on the day of birth) or application of either a 5-HT₂ or 5-HT₇ receptor antagonist disorganized the NMDA-triggered locomotion in the preparation. After adding 5-HT to the bath, locomotion-like activity was restored in spinal cords collected from animals pretreated with PCPA. Thus, 5-HT appeared to be pivotal in enhancing the locomotor-related alternations (i.e., CPG activity) in neonatal rat spinal cords *in vitro* (Pearlstein et al., 2005). Contrarily, stimulation of 5-HT_{1A} and 5-HT₇ receptors by 8OH-DPAT was able to produce CPG activity in mice with T9-10 spinal cord transection *in vivo* (Landry et al., 2006). When combined with EES and locomotor training, buspirone treatment activated and recruited the lumbar CPG circuitry

via affecting sensory input and functionally remodeling the pace-making pathways in rats with T7 spinal cord transection (Courtine et al., 2009). Administration of buspirone strengthened locomotion (e.g., improving the number of steps taken, between hindlimbs coupling strength, angular excursion of the hip joint, and paw positioning) in mice with either T8 transection or a dual lesion paradigm of T7 hemisection and T8 transection that was done at the end of a 3-week training period after T7 hemisection. Also, buspirone treatment achieved long-term improvement of symmetry, paw positioning at contact, and paw drag when combined with locomotor training in mice with dual T7 and T8 lesions (Jeffrey-Gauthier et al., 2018).

In clinical rehabilitation, buspirone treatment augmented muscular coordination and the effort level that a participant with AIS-B generated in the exoskeleton while stepping, producing a refined continuous, smooth stepping motion in the exoskeleton in addition to autonomic improvements (cardiovascular function and thermoregulation; Gad et al., 2017). Buspirone co-administered with transcutaneous electrical stimulation (TES) in chronic AIS-B participants (C5 or above; motor complete) significantly increased mean hand strength (Freyvert et al., 2018). Newly published results demonstrated that buspirone (7.5-15 mg, b.i.d., p.o.), not trazodone (100 mg/day; a 5HT uptake inhibitor) or placebo, significantly reduced chronic SCI-caused susceptibility to induced hypocapnic central sleep apnea by reducing chemosensitivity and increasing CO₂ reserve (Maresh et al., 2020).

The effects of use and non-use of buspirone were examined in participants ($n = 21$; < 2 years of SCI from C4 to T3) who underwent 6 months of equal exercise training facilitated by functional electrical stimulation (FES), i.e., direct skeletal muscle stimulation to generate muscle contraction and movement. The participants on buspirone ($n = 10$; 29 ± 17 mg/day), compared to well-matched individuals without buspirone ($n = 11$), had significantly greater increases in peak oxygen consumption (VO_{2peak}) and peak ventilation (VE_{peak}). While changes in VO_{2peak} and VE_{peak} were correlated across all patients ($r = 0.63$, $p < 0.01$), participants on buspirone showed the strongest correlation ($r = 0.85$, $p < 0.01$). Only in the buspirone group were changes in respiratory function significantly correlated with increased peak tidal volume ($r > 0.66$, $p < 0.05$). The data demonstrated that buspirone improved cardiorespiratory adaptations to FES-exercise training in individuals with chronic high-level SCI. These close associations between augmented ventilatory and aerobic capacities

indicated that improved respiratory function was the primary mechanism (Vivodtzev et al., 2020).

The majority of serotonin in the spinal cord originates from the raphe nuclei, which makes this neuromodulation innervation (i.e., there is no intrinsic 5HT synthesis in the adult mammalian spinal cord) vulnerable to neurotrauma (Perrier and Cotel, 2015). In mammalian spinal cords, 5-HT_{1A} receptors, which are coupled to different messenger pathways via association with G α isoforms or $\beta\gamma$ signaling in different cell types (Rojas and Fiedler, 2016), are primarily located on the soma and proximal dendrites of serotonergic-innervated locomotor-activated neurons in Rexed laminae I-IV, VII, VIII and X of thoraco-lumbar segments of cats (Giroux et al., 1999; Noga et al., 2009). In human lumbar spinal cords, 5-HT_{1A} receptors were mostly distributed in Rexed lamina II of the dorsal horn and in the ventral horn, participating in afferent, sensorimotor, and CPG regulation (Perrin et al., 2011). Presently, buspirone is one of the most widely investigated and clinically used neuromodulation drugs. It agonizes inhibitory presynaptic and postsynaptic 5HT_{1A} receptors (e.g., somatodendritic autoreceptors, postsynaptic heteroreceptors, etc.), while also having possible other weaker interactions (Loane and Politis, 2012). While discussion of other receptor subtypes of 5HT is beyond the scope of this review, it should be mentioned that intrathecal antagonism of spinal 5-HT₇ receptors disrupted spontaneous locomotion in intact animals, but not fictive locomotion induced by stimulation of the mesencephalic locomotor region. Since locomotor coordination in adult mammals requires the sensory input, the results suggested a 5-HT₇ receptor-mediated role in modulating afferent pathways to operate CPG (Cabaj et al., 2017). Thus, targeting 5HT_{1A}, 5-HT_{2A/2C}, and 5HT₇ receptors for neuromodulation may constitute a fruitful strategy to restoring locomotion after SCI.

(3) Dopaminergic modulation: Descending dopaminergic pathways from the brain, specifically that originating from the A11 area of the diencephalon, may be the major source of dopamine within the mammalian spinal cord. These tracts play crucial roles in sensorimotor integration, locomotion regulation, and pain modification (Barraud et al., 2010; Sharples et al., 2014). In early laboratory studies, adding DOPA to chronic spinal preparations liberated catecholamines (i.e., dopamine and noradrenaline). The data suggested that DOPA augmented the synthesis and terminal release of neurotransmitters from the descending noradrenergic pathway (Andén et al., 1966). Administration of DOPA together with the monoamine oxidase inhibitor nialamide (i.v.) generated rhythmic motor output in the

absence of phasic peripheral feedback in models of spinalization at different levels (i.e., T12, and L5-S1; Grillner and Zangger, 1979).

To modulate multiple neurotransmissions governing the CPG, a tri-therapy of buspirone, carbidopa, and levodopa was administered to mice with T9-10 spinal cord transection. The treatment enabled the mice to manifest episodes of steady weight-bearing stepping (Guertin et al., 2010). In a subsequent study, spinalized mice given the tri-therapy with or without clenbuterol, plus regular treadmill training for 8 weeks, exhibited locomotor movement and gradually increased muscle mass (Ung et al., 2012). Furthermore, delivery of a combined recipe of NMDA, 5-HT, and dopamine produced fictive locomotion in rodent spinal cords. This was disrupted via activation of the bradykinin receptor B2 by the inflammatory nonapeptide bradykinin, a secondary injury molecule of SCI (Mandadi et al., 2016).

The laboratory data suggested that dopaminergic stimulation co-enhanced by other neurotransmissions could be an effective formula for locomotor recovery. Supporting this notion, a double-blind, placebo-controlled, randomized phase I/IIa study on safety and efficacy of treatment with buspirone/levodopa/carbidopa (SpinalonTM) revealed that locomotor-like EMG patterns were displayed only in participants (AIS-A/B) who received SpinalonTM treatment (Radhakrishna et al., 2017). Notably, buspirone, clenbuterol, and levodopa can stimulate production of neurotrophic factors and antioxidants by neural cells (Hayes et al., 1995; Zhang et al., 2006), and this type of neurotrophic support is missing in lesioned adult spinal cords and essential for neural repair and plasticity-based functional recovery (Teng, 2019a). Therefore, if multiple neurotransmissions are pharmacologically modulated, locomotion is likely inducible in clinical chronic SCI. The efficacy can be further increased if the drug treatment is combined with rehabilitation training, especially those integrated by multimodal neuromechanical prostheses (see details below).

Neuromechanical prostheses to restore function for chronic SCI:

Neuromechanical prostheses encompass assistive and rehabilitative devices that interface with the individual's sensorimotor system (Pizzolato et al., 2019). Biological signals acquired via neuromechanical prosthesis include a range of electrical biopotentials (e.g., electroencephalograms [EEG], EMG, electrocardiogram [ECG]), biomechanical variables (e.g., limb movement and forces, hand gestures), and respiratory parameters (e.g., tidal volume, rate, and minute ventilation). These signals, if appropriately interpreted, can be used

to understand a person's motor intention and, accordingly, control electrical (e.g., electrical stimulation) and mechanical (e.g., exoskeletons) actuators to generate movement.

Neuromechanical prostheses can compensate for the SCI-caused loss of afferent feedback by redirecting and augmenting sensory input from disrupted pathways to residual ascending tracts representing the affected peripheral target, in order to reengage the somatotopic area remapped by SCI (Leemhuis et al., 2019) (**Figure 2**). Successful neuromechanical prostheses should provide electrical and mechanical support that is commensurate with the person's own neuromuscular capabilities, and consistent with the excitations of still intact neural pathways.

(1) Electrical stimulation: Generation of movement via FES and/or mechanical means, which also appropriately triggers peripheral mechanoreceptors, may reinstate lower limb motor synergies. FES gained heightened attention after a seminal study by McDonald et al. (2002), showing that long-term training using FES for cycling in one individual with cervical SCI improved the scale from AIS-A to AIS-C within 3 years. FES cycling also improved motor function in people with complete or incomplete SCI after a 10-week intervention. However, in another study, no improved AIS reclassification was achieved for individuals with complete SCI. The lack of efficacy in using FES alone over a shorter time course may relate to its inability to engage supraspinal and intraspinal neurocircuitry, despite its benefits of increasing aerobic capacity and preventing muscle atrophy (Griffin et al., 2009; Ragnarsson, 2008).

Continuous EES employs an electrical signal of constant amplitude and frequency, selected through experimentation (Harkema et al., 2011), to induce muscle contraction and facilitate generation of rhythmic movement. It can additionally engage the proprioceptive circuits via stimulation of the lumbosacral segment. By reactivating the CPG, EES enabled AIS-A people to stand, walk, or cycle with minimal assistance (Angeli et al., 2018; Harkema et al., 2011). EES has been combined with physical training of standing (Wenger et al., 2016), multimodal exercises (e.g., seated trunk balance, standing balance and stepping, range of motion; Wenger et al., 2016), and stepping or gait retraining with body weight support (Angeli et al., 2018; Wagner et al., 2018). However, continuous EES has been shown to disrupt afferent signals generated by mechanoreceptors of intact muscles, tendons, and joints (Formento et al., 2018), a type of input that is believed essential to activate and modulate the CPG (Lavrov et al., 2008). To solve this conflict, a computational model representing the salient anatomical and physiological features of lower limb muscles and CPG was developed

to modulate EES and minimize antidromic collisions along proprioceptive afferent pathways (Formento et al., 2018; Wagner et al., 2018). This spatiotemporal-modulated EES was applied to three participants (AIS-C/D) who subsequently regained voluntary control over paralyzed muscles and maintained it even in absence of stimulation (Wagner et al., 2018). Thus, refining appropriate timing, amplitude, frequency in stimulation delivery to facilitate segmental proprioceptive, and, perhaps, other deep sensory inputs appears necessary for recovering sustainable patterned motor neuron functions.

TES is a non-invasive alternative to EES, wherein electrodes are positioned on the surface of the skin, commonly between spinal processes (cathode) and on iliac crest (anode). TES modulates the excitability of the spinal cord, evoking the reconnection between supraspinal and intraspinal networks (Kakulas, 1999). In a prototype study, five participants with cervical or thoracic motor complete SCI (AIS-B) underwent an 18-week intervention. TES elicited a stepping-like motion when participants were concurrently asked to produce a voluntary effort. When participants were administered buspirone (7.5 mg, b.i.d., p.o.), the combined neuromodulation generated voluntary muscle contraction and additive motor effect (Gerasimenko et al., 2015). As described above, similar combinatorial TES and buspirone treatments also restored discernible degrees of hand function or locomotion in individuals with AIS-A/B injuries at different spinal cord levels (Freyvert et al., 2018; Gad et al., 2017).

Electrical stimulation of spinal cord or muscles is often paired with some form of mechanical assistance, delivered by a practitioner (e.g., limb mobilization during physical therapy) or via motorized devices. Mechanical assistance can help with limb movement when insufficient force is produced via volitional or electrically induced muscle contraction, but can also extend training session durations. However, to our best knowledge, no published study has shown that mechanical assistance alone, regardless of the delivery modality, resulted in reclassification of SCI from AIS A/B to AIS-C/D. Clearly, electrical stimulation, pharmacological neuromodulation, and mechanical assistance in combination can have additive or synergistic effects of improving motosensory function for chronic SCI.

(2) Brain-computer interfaces (BCIs): Concurrent engagement of efferent and afferent pathways during rehabilitation has been considered essential in the recovery of motosensory function. To this end, BCIs have been garnering interest as a vehicle to engage cerebral cortical networks during rehabilitation (Donati et al., 2016). BCIs translate acquired EEG signals into real world commands representing the intent of the user. The act of thinking

about performing a movement (i.e., motor imagery) is transformed into a command for the neuromechanical prosthesis and, consequently, into a real action. Importantly, BCI-enhanced neuromechanical prostheses have been engineered to bypass SCI lesions, enabling certain degrees of voluntary control of the affected limbs via electrical stimulation (Ethier et al., 2012). Current BCI paradigms, although only classifying a finite number of motor intentions (limited by the number of surface EEG channels and the signal quality), can detect gross motor initiatives (e.g., walking). This was made possible by advanced artificial intelligence and machine learning algorithms, which have been continuously improved by increasing the amount of EEG data decoded into coordinated motor instructions to directly operate the muscles (Lotte et al., 2018) or exoskeletons (Benabid et al., 2019).

Implanted electrode arrays to access cortical signals (Oxley et al., 2016; Musk and Neuralink, 2019) are leading to a generational leap in amount and quality of available EEG data. These innovations enabled an increased number of user-intentions that can be correctly classified by artificial intelligence algorithms but are unlikely to lead to physiological control of paralysed limbs. This is because neural information from the motor cortex cannot be directly mapped to movement, which has to work in consortium with the musculoskeletal system under the environmental context before a naturally synchronized limb movement can be generated. Similar challenges were met when trying to adapt artificial intelligence methods to correctly map EMG, signals mostly associated to muscle force production, to directly produce movement (Zhang et al., 2020; Kapelner et al., 2020). To overcome these hurdles, new systems that integrate computational models of the spinal cord, somatosensory apparatus, and musculoskeletal input (e.g., human digital twins; see below), can be designed in the future to generate better mapping between EEG and movement, to deliver truly embodied neuromechanical prostheses.

(3) The human digital twin: The digital twin is a high-fidelity multiscale computation model composed of numerous subsystems that are described and interconnected via mathematical relationships. This modern paradigm, initially devised and deployed in the aerospace industry, enables monitoring of physical quantities that cannot be observed via direct measurement (Glaessgen and Stargel, 2012). Following SCI, below lesion sensory input cannot be conveyed to the brain, causing a loss of body position, CPG function and movement awareness, and ensuing reorganization of the somatosensory cortex (Nardone et al., 2013). Neuromechanical prostheses can relay the disrupted proprioceptive information to the body sites with intact sensory inflow. This augmented sensory feedback, or artificially

generated biofeedback, has been utilized to enhance or substitute proprioception (Giggins et al., 2013; Shokur et al., 2016). Therefore, combinations of visual (e.g., virtual reality), haptic, or auditory monitors, and positional/orientation feedback (i.e., vestibular input) can be provided to people with SCI via the human digital twin that is designed to interpret heterogeneous biosensory data (e.g., EMG, ECG, EEG, forces and movement, respiration, hand gestures, etc.) to improve rehabilitation training outcomes (**Figure 2**). However, there is presently no consensus regarding the optimal delivery modality and amount of biofeedback, which type of afferent signals to augment, and how to best convert mechanoreceptor signals into interpretable feedback data. We and others postulated that providing biofeedback that is consistent and time-synchronized with native and intact proprioceptive signals might more effectively promote functional recovery in chronic stage of SCI (Jackson and Zimmermann, 2012; Pizzolato et al., 2019).

<Insert Figure 2 about here>

The importance of employing both efferent and afferent neuropathways during rehabilitation was indicated in a phase I study that combined BCI, biofeedback, and graded gait training (i.e., combination of exoskeleton and body weight supported walking; Donati et al., 2016). After a 12-month intervention, four participants (50%) improved from AIS-A/B to AIS-C. While AIS classification was unchanged in the other four (AIS-A/B), all participants showed some improvement in their measured EMG signals, with five showing improved ambulation ability compared to baseline. In a subsequent phase II study, all participants continued their training under the same protocol for an additional 14 months, resulting in AIS-C reclassification for all of participants (Shokur et al., 2018). Two of these, who were reclassified from AIS-A to AIS-C, also underwent additional gait retraining using a combination of BCI, haptic feedback, and FES, showing further increased lower extremity motor score after 22 training sessions (Selfslagh et al., 2019). The encouraging results from these studies, albeit with small sample sizes, suggested that reestablishment of neural function after severe SCI is possible with non-invasive multimodal therapeutic approaches.

Enabling coordination between assistive devices and the individual with SCI requires correct understanding of the complex and causal interplay between the different elements of the neuromusculoskeletal system, according to their roles in generating and sensing movement. In this context, a digital twin can be designed to interpret heterogeneous biosensory input and cybernetically coordinate the participant's conscious intention, appropriately augmented feedback, and proper control of the electromechanical multimodal

assistive devices (e.g., exoskeletons, electrical stimulation) (**Figure 2**). Moreover, digital twins implementing large-scale neuromusculoskeletal models, must entail mathematical descriptions of muscles, tendons, joints, and bones that closely match the individual's physiology, anatomy, and function capacity (Pizzolato et al., 2017), thereby allowing researchers to unveil the mechanistic causal relationships between neural motor command and final movement for different neuromuscular conditions (Sartori et al., 2017). Recent advancements in computational biomechanics have also produced real-time human digital twins, which are able to instantaneously predict the internal state of musculoskeletal tissues (e.g., stress/strain) and associated motor command (Pizzolato et al., 2020), and to interface with electromechanical assistive devices (Sartori et al., 2018). This will help pave the way to the application of the human digital twin in complex rehabilitation therapies, where the treatment is automatically optimized to match the neuromuscular capabilities of the individual.

Conclusions

Major progress made in the past 30 years has demonstrated that patterned motor activities can be rekindled below the spinal cord lesion without direct supraspinal innervation. The neurobiological mechanisms underlying such recoveries have been targeted for developing noninvasive therapies to improve function after SCI. Pharmacological approaches employing individual or combined drug treatments, which manipulated receptors of neuromodulators (e.g., 5HT_{1A} receptors) or monoamine neurotransmissions, exhibited efficacious clinical outcomes. The necessity for the intervention to simultaneously affect multiple components of the sensorimotor neural network was corroborated by data generated from cell therapy investigations for repairing the injured spinal cord (e.g., donor cell-exerted neurotrophic, anti-inflammatory, and homeostatic effects). Lastly, neuromechanical prostheses-based rehabilitation protocols were more effective for chronic SCI when they were performed in combination with drug administration (e.g., 5-HT_{1A} agonists). Furthermore, the potency of the treatments could be additively or synergistically enhanced by their combination with functional training therapies.

For qualitative therapy improvement, the human digital twin represents a powerful multimodal strategy to integrate innate volitional pathways with different forms of neuromechanical assistance to cybernetically tailor a particular rehabilitation therapy to meet an individual's clinical needs. If designed to also incorporate the effects of pharmacological

agents and TES/EES/FES, the system may constitute a novel drug-device therapy. Overall, the findings from the literature review suggested that to move the field forward, it is important for future research and development operations to focus on (i) formulating pharmacological and cellular treatment regimens that can concurrently modulate key neurotransmissions and homeostatic processes to mitigate secondary injury and optimize repair, (ii) developing BCI-integrated neuromechanical prostheses (e.g., human digital twins) that can coordinate the participant's conscious drive with biosensory inputs and outputs of FES, activity training, and drug administration devices, (iii) advancing the understanding of the neurobiological mechanisms of the CPG reactivation in the adult mammalian spinal cord after injury, and (iv) organizing phase II and III trials with adequate statistical power, proper control setting, and multicenter participation to standardize new SCI treatment.

Acknowledgements

C.P., D.P., G.G., S.H. and D.G.L. have been supported by funds from Motor Accident Insurance Commission, Queensland Government, Australia, Perpetual IMPACT and Griffith University. Teng Laboratory is currently supported by The US AMRMC (W81XWH-15-1-0621), The SCI Trust Fund of The Commonwealth of Massachusetts, The Gordon Project to Cure Clinical Paralysis, The Cele H. and William B. Rubin Family Fund, and The Roosevelt Warm Springs Foundation. With Dr. Teng's joint sponsorship, M.A.G. and J.W. are additionally supported by BAU Medical School, Turkey, and Wuhan Union Hospital, Tongji Medical College/HZUST, PRC, respectively. R.D., whose thesis research was promoted by Y.D.T., was a student at The University of Nottingham, UK.

References

- Aarabi, B., Harrop, J.S., Tator, C.H., et al., 2012. Predictors of pulmonary complications in blunt traumatic spinal cord injury. *J Neurosurg Spine* 17(1 Suppl), 38-45.
- Adams, M.M., Hicks, A.L., 2005. Spasticity after spinal cord injury. *Spinal Cord* 43, 577-586.
- Agrawal, S., Fehlings, M., 1996. Mechanisms of secondary injury to spinal cord axons in vitro: role of Na⁺, Na⁽⁺⁾-K⁽⁺⁾-ATPase, the Na⁽⁺⁾-H⁺ exchanger, and the Na⁽⁺⁾-Ca²⁺ exchanger. *J Neurosci* 16, 545-552.

- Ahuja, C.S., Wilson, J.R., Nori, S., et al. 2017. Traumatic spinal cord injury. *Nat Rev Dis Primers* 3, 17018.
- Al'joboori, Y.D., Edgerton, V.R., Ichiyama, R.M., 2020. Effects of Rehabilitation on Perineural Nets and Synaptic Plasticity Following Spinal Cord Transection. *Brain Sci* 10.
- Allen, A.R., 1911. Surgery of Experimental Lesion of Spinal Cord Equivalent to Crush Injury of Fracture Dislocation of Spinal Column. *Journal of the American Medical Association* LVII, 878-880.
- American Spinal Injury Association, 1982. Standards for neurological classification of spinal injured patients. ASIA, Chicago
- Andén, N.E., Jukes, M.G.M., Lundberg, A., 1966. The Effect of DOPA on the Spinal Cord. 2. A Pharmacological Analysis. *Acta Physiologica Scandinavica* 67, 387-397.
- Anderson, M.A., Burda, J.E., Ren, Y., Ao, Y., O'Shea, T.M., Kawaguchi, R., Coppola, G., Khakh, B.S., Deming, T.J., Sofroniew, M.V., 2016. Astrocyte scar formation aids central nervous system axon regeneration. *Nature* 532, 195-200.
- Angeli, C.A., Boakye, M., Morton, R.A., Vogt, J., Benton, K., Chen, Y., Ferreira, C.K., Harkema, S.J., 2018. Recovery of Over-Ground Walking after Chronic Motor Complete Spinal Cord Injury. *The New England journal of medicine* 379, 1244-1250.
- Arevalo-Martin, A., Garcia-Ovejero, D., Sierra-Palomares, Y., Paniagua-Torija, B., Gonzalez-Gil, I., Ortega-Gutierrez, S., Molina-Holgado, E., 2012. Early endogenous activation of CB1 and CB2 receptors after spinal cord injury is a protective response involved in spontaneous recovery. *PLoS One* 7, e49057.
- Baptiste, D.C., Tighe, A., Fehlings, M.G., 2009. Spinal cord injury and neural repair: focus on neuroregenerative approaches for spinal cord injury. *Expert Opin Investig Drugs* 18, 663-673.
- Barbeau, H., Rossignol, S., 1991. Initiation and modulation of the locomotor pattern in the adult chronic spinal cat by noradrenergic, serotonergic and dopaminergic drugs. *Brain Research* 546, 250-260.
- Barraud, Q., Obeid, I., Aubert, I., Barriere, G., Contamin, H., McGuire, S., Ravenscroft, P., Porras, G., Tison, F., Bezard, E., Ghorayeb, I., 2010. Neuroanatomical study of the A11 diencephalospinal pathway in the non-human primate. *PLoS One* 5, e13306.
- Behrman, A.L., Bowden, M.G., Nair, P.M., 2006. Neuroplasticity after spinal cord injury and training: an emerging paradigm shift in rehabilitation and walking recovery. *Phys Ther* 86, 1406-1425.

- Bell, J.E.S., Seifert, J.L., Shimizu, E.N., Sucato, D.J., Romero-Ortega, M.I., 2017. Atraumatic Spine Distraction Induces Metabolic Distress in Spinal Motor Neurons. *J Neurotrauma* 34, 2034-2044.
- Benabid, A.L., Costecalde, T., Eliseyev, A., et al. 2019. An exoskeleton controlled by an epidural wireless brain-machine interface in a tetraplegic patient: a proof-of-concept demonstration. *Lancet Neurol* 18, 1112-1122.
- Bloom, O., Herman, P.E., Spungen, A.M., 2020. Systemic inflammation in traumatic spinal cord injury. *Exp Neurol* 325, 113143.
- Bracken, M.B., Collins, W.F., Freeman, D.F., et al., 1984. Efficacy of methylprednisolone in acute spinal cord injury. *Jama* 251, 45-52.
- Bracken, M.B., Shepard, M.J., Collins, et al., 1990. A randomized, controlled trial of methylprednisolone or naloxone in the treatment of acute spinal-cord injury. Results of the Second National Acute Spinal Cord Injury Study. *N Engl J Med* 322, 1405-1411.
- Bracken, M.B., Shepard, M.J., Hellenbrand, et al., 1985. Methylprednisolone and neurological function 1 year after spinal cord injury. Results of the National Acute Spinal Cord Injury Study. *Journal of neurosurgery* 63, 704-713.
- Bracken, M.B., Shepard, M.J., Holford, T.R., et al. 1997. Administration of methylprednisolone for 24 or 48 hours or tirilazad mesylate for 48 hours in the treatment of acute spinal cord injury. Results of the Third National Acute Spinal Cord Injury Randomized Controlled Trial. National Acute Spinal Cord Injury Study. *Jama* 277, 1597-1604.
- Bradbury, E.J., Moon, L.D.F., Popat, R.J., King, V.R., Bennett, G.S., Patel, P.N., Fawcett, J.W., McMahon, S.B., 2002. Chondroitinase ABC promotes functional recovery after spinal cord injury. *Nature* 416, 636-640.
- Brown, A., Nabel, A., Oh, W., Etlinger, J.D., Zeman, R.J., 2014. Perfusion imaging of spinal cord contusion: injury-induced blockade and partial reversal by beta2-agonist treatment in rats. *J Neurosurg Spine* 20, 164-171.
- Busch, S.A., Horn, K.P., Silver, D.J., Silver, J., 2009. Overcoming macrophage-mediated axonal dieback following CNS injury. *J Neurosci* 29, 9967-9976.
- Cabaj, A.M., Majczynski, H., Couto, E., Gardiner, P.F., Stecina, K., Slawinska, U., Jordan, L.M., 2017. Serotonin controls initiation of locomotion and afferent modulation of coordination via 5-HT7 receptors in adult rats. *J Physiol* 595, 301-320.

- Cadotte, D.W., Bosma, R., Mikulis, D., Nugaeva, N., Smith, K., Pokrupa, R., Islam, O., Stroman, P.W., Fehlings, M.G., 2012. Plasticity of the injured human spinal cord: insights revealed by spinal cord functional MRI. *PLoS One* 7, e45560.
- Carlstedt, T., Cullheim, S., Risling, M., Ulfhake, B., 1989. Nerve fibre regeneration across the PNS-CNS interface at the root-spinal cord junction. *Brain Res Bull* 22, 93-102.
- Chan, L.W., Griebeling, T.L., Arnold, E.P., Chu, P.S., New, P.W., Wagg, A., 2018. Special considerations in the urological management of the older spinal cord injury patient. *World J Urol* 36, 1603-1611.
- Chariker, J.H., Ohri, S.S., Gomes, C., et al. 2019. Activity/exercise-induced changes in the liver transcriptome after chronic spinal cord injury. *Sci Data* 6, 88.
- Chen, B., Zhang, Y., Chen, L., Huang, S., Li, S., Yao, J., 2013. Dose-effects of aorta-infused clenbuterol on spinal cord ischemia-reperfusion injury in rabbits. *PLoS One* 8, e84095.
- Chikuda, H., Yasunaga, H., Takeshita, K., Horiguchi, H., Kawaguchi, H., Ohe, K., Fushimi, K., Tanaka, S., 2014. Mortality and morbidity after high-dose methylprednisolone treatment in patients with acute cervical spinal cord injury: a propensity-matched analysis using a nationwide administrative database. *Emerg Med J* 31, 201-206.
- Choi, H., Liao, W.L., Newton, K.M., Onario, R.C., King, A.M., Desilets, F.C., Woodard, E.J., Eichler, M.E., Frontera, W.R., Sabharwal, S., Teng, Y.D., 2005. Respiratory abnormalities resulting from midcervical spinal cord injury and their reversal by serotonin 1A agonists in conscious rats. *J Neurosci* 25, 4550-4559.
- Christie, S.D., Comeau, B., Myers, T., Sadi, D., Purdy, M., Mendez, I., 2008. Duration of lipid peroxidation after acute spinal cord injury in rats and the effect of methylprednisolone. *Neurosurg Focus* 25, E5
- Clark, J.M., Findlay, D.M., 2017. Musculoskeletal Health in the Context of Spinal Cord Injury. *Curr Osteoporos Rep* 15, 433-442.
- Côté, M.P., Murray, M., Lemay, M.A., 2017. Rehabilitation Strategies after Spinal Cord Injury: Inquiry into the Mechanisms of Success and Failure. *J Neurotrauma* 34, 1841-1857.
- Courtine, G., Gerasimenko, Y., van den Brand, R., Yew, A., Musienko, P., Zhong, H., Song, B., Ao, Y., Ichiyama, R.M., Lavrov, I., Roy, R.R., Sofroniew, M.V., Edgerton, V.R., 2009. Transformation of nonfunctional spinal circuits into functional states after the loss of brain input. *Nat Neurosci* 12, 1333-1342.

- Cyranoski, D., 2019a. Japan's approval of stem-cell treatment for spinal-cord injury concerns scientists. *Nature* 565, 544-545.
- Cyranoski, D., 2019b. The potent effects of Japan's stem-cell policies. *Nature* 573, 482-485.
- Darrow, J.J., Avorn, J., Kesselheim, A.S., 2020. FDA Approval and Regulation of Pharmaceuticals, 1983-2018. *Jama* 323, 164-176.
- de Rivero Vaccari, J.P., Brand, F., 3rd, Adamczak, S., Lee, S.W., Perez-Barcena, J., Wang, M.Y., Bullock, M.R., Dietrich, W.D., Keane, R.W., 2016. Exosome-mediated inflammasome signaling after central nervous system injury. *J Neurochem* 136 Suppl 1, 39-48.
- DeForge, D., Nymark, J., Lemaire, E., Gardner, S., Hunt, M., Martel, L., Curran, D., Barbeau, H., 2004. Effect of 4-aminopyridine on gait in ambulatory spinal cord injuries: a double-blind, placebo-controlled, crossover trial. *Spinal Cord* 42, 674-685.
- Dias, D.O., Kim, H., Holl, D., Werne Solnestam, B., Lundeborg, J., Carlen, M., Goritz, C., Frisen, J., 2018. Reducing Pericyte-Derived Scarring Promotes Recovery after Spinal Cord Injury. *Cell* 173, 153-165 e122.
- Domingo, A., Al-Yahya, A.A., Asiri, Y., Eng, J.J., Lam, T., 2012. A systematic review of the effects of pharmacological agents on walking function in people with spinal cord injury. *J Neurotrauma* 29, 865-879.
- Donati, A.R., Shokur, S., Morya, E., et al. 2016. Long-Term Training with a Brain-Machine Interface-Based Gait Protocol Induces Partial Neurological Recovery in Paraplegic Patients. *Sci Rep* 6, 30383.
- Dorton, M.C., Lucci, V.M., de Groot, S., Loughin, T.M., Cragg, J.J., Kramer, J.K., Post, M.W.M., Claydon, V.E., 2020. Evaluation of cardiovascular disease risk in individuals with chronic spinal cord injury. *Spinal Cord* doi: 10.1038/s41393-020-00566-5. Online ahead of print.
- Dunn, J., Blight, A., 2011. Dalfampridine: a brief review of its mechanism of action and efficacy as a treatment to improve walking in patients with multiple sclerosis. *Curr Med Res Opin* 27, 1415-1423.
- Eldahan, K.C., Williams, H.C., Cox, D.H., Gollihue, J.L., Patel, S.P., Rabchevsky, A.G., 2020. Paradoxical effects of continuous high dose gabapentin treatment on autonomic dysreflexia after complete spinal cord injury. *Exp Neurol* 323, 113083.
- Ethier, C., Oby, E.R., Bauman, M.J., Miller, L.E., 2012. Restoration of grasp following paralysis through brain-controlled stimulation of muscles. *Nature* 485, 368-371.

- Farooque, M., Hillered, L., Holtz, A., Olsson, Y., 1996. Changes of extracellular levels of amino acids after graded compression trauma to the spinal cord: an experimental study in the rat using microdialysis. *J Neurotrauma* 13, 537-548.
- Fawcett, J.W., 2020. The Struggle to Make CNS Axons Regenerate: Why Has It Been so Difficult? *Neurochem Res* 45, 144-158.
- Fehlings, M.G., Kim, K.D., Aarabi, B., Rizzo, M., Bond, L.M., McKerracher, L., Vaccaro, A.R., Okonkwo, D.O., 2018. Rho Inhibitor VX-210 in Acute Traumatic Subaxial Cervical Spinal Cord Injury: Design of the Spinal Cord Injury Rho Inhibition InvestiGation (SPRING) Clinical Trial. *J Neurotrauma* 35, 1049-1056.
- Fehlings, M.G., Tetreault, L.A., Wilson, J.R., Kwon, B.K., Burns, A.S., Martin, A.R., Hawryluk, G., Harrop, J.S., 2017. A Clinical Practice Guideline for the Management of Acute Spinal Cord Injury: Introduction, Rationale, and Scope. *Global Spine J* 7, 84S-94S.
- Fehlings, M.G., Theodore, N., Harrop, J., Maurais, G., Kuntz, C., Shaffrey, C.I., Kwon, B.K., Chapman, J., Yee, A., Tighe, A., McKerracher, L., 2011. A phase I/IIa clinical trial of a recombinant Rho protein antagonist in acute spinal cord injury. *J Neurotrauma* 28, 787-796.
- Fehlings, M.G., Vaccaro, A., Wilson, J.R., Singh, A., W. Cadotte, D., Harrop, J.S., Aarabi, B., Shaffrey, C., Dvorak, M., Fisher, C., Arnold, P., Massicotte, E.M., Lewis, S., Rampersaud, R., 2012. Early versus Delayed Decompression for Traumatic Cervical Spinal Cord Injury: Results of the Surgical Timing in Acute Spinal Cord Injury Study (STASCIS). *PLoS One* 7, e32037.
- Felleiter, P., Müller, N., Schumann, F., Felix, O., Lierz, P., 2012. Changes in the use of the methylprednisolone protocol for traumatic spinal cord injury in Switzerland. *Spine (Phila Pa 1976)* 37, 953-956.
- Fleming, J.C., Norenberg, M.D., Ramsay, D.A., Dekaban, G.A., Marcillo, A.E., Saenz, A.D., Pasquale-Styles, M., Dietrich, W.D., Weaver, L.C., 2006. The cellular inflammatory response in human spinal cords after injury. *Brain* 129, 3249-3269.
- Formento, E., Minassian, K., Wagner, F., Mignardot, J.B., Le Goff-Mignardot, C.G., Rowald, A., Bloch, J., Micera, S., Capogrosso, M., Courtine, G., 2018. Electrical spinal cord stimulation must preserve proprioception to enable locomotion in humans with spinal cord injury. *Nat Neurosci* 21, 1728-1741.

- Freund, P., Rothwell, J., Craggs, M., Thompson, A.J., Bestmann, S., 2011. Corticomotor representation to a human forearm muscle changes following cervical spinal cord injury. *Eur J Neurosci* 34, 1839-1846.
- Freund, P., Seif, M., Weiskopf, N., Friston, K., Fehlings, M.G., Thompson, A.J., Curt, A., 2019. MRI in traumatic spinal cord injury: from clinical assessment to neuroimaging biomarkers. *Lancet Neurol* 18, 1123-1135.
- Freund, P., Wannier, T., Schmidlin, E., Bloch, J., Mir, A., Schwab, M.E., Rouiller, E.M., 2007. Anti-Nogo-A antibody treatment enhances sprouting of corticospinal axons rostral to a unilateral cervical spinal cord lesion in adult macaque monkey. *J Comp Neurol* 502, 644-659.
- Freyvert, Y., Yong, N.A., Morikawa, E., Zdunowski, S., Sarino, M.E., Gerasimenko, Y., Edgerton, V.R., Lu, D.C., 2018. Engaging cervical spinal circuitry with non-invasive spinal stimulation and buspirone to restore hand function in chronic motor complete patients. *Sci Rep* 8, 15546.
- Friede, T., Posch, M., Zohar, S., et al. 2018. Recent advances in methodology for clinical trials in small populations: the InSPiRe project. *Orphanet J Rare Dis* 13, 186.
- Frigon, A., Johnson, M.D., Heckman, C.J., 2012. Differential modulation of crossed and uncrossed reflex pathways by clonidine in adult cats following complete spinal cord injury. *J Physiol* 590, 973-989.
- Frontera, W.R., Choi, H., Krishnan, G., Krivickas, L.S., Sabharwal, S., Teng, Y.D., 2006. Single muscle fiber size and contractility after spinal cord injury in rats. *Muscle Nerve* 34, 101-104.
- Fung, J., Stewart, J.E., Barbeau, H., 1990. The combined effects of clonidine and cyproheptadine with interactive training on the modulation of locomotion in spinal cord injured subjects. *Journal of the Neurological Sciences* 100, 85-93.
- Furlan, J.C., Fehlings, M.G., 2008. Cardiovascular complications after acute spinal cord injury: pathophysiology, diagnosis, and management. *Neurosurg Focus* 25, E13.
- Furlan, J.C., Noonan, V., Cadotte, D.W., Fehlings, M.G., 2011. Timing of decompressive surgery of spinal cord after traumatic spinal cord injury: an evidence-based examination of pre-clinical and clinical studies. *J Neurotrauma* 28, 1371-1399.
- Gad, P., Gerasimenko, Y., Zdunowski, S., Turner, A., Sayenko, D., Lu, D.C., Edgerton, V.R., 2017. Weight Bearing Over-ground Stepping in an Exoskeleton with Non-invasive Spinal Cord Neuromodulation after Motor Complete Paraplegia. *Front Neurosci* 11, 333.

- Gagnon, D.H., Escalona, M.J., Vermette, M., Carvalho, L.P., Karelis, A.D., Duclos, C., Aubertin-Leheudre, M., 2018. Locomotor training using an overground robotic exoskeleton in long-term manual wheelchair users with a chronic spinal cord injury living in the community: Lessons learned from a feasibility study in terms of recruitment, attendance, learnability, performance and safety. *J Neuroeng Rehabil* 15, 12.
- Ganzer, P.D., Beringer, C.R., Shumsky, J.S., Nwaobasi, C., Moxon, K.A., 2018. Serotonin receptor and dendritic plasticity in the spinal cord mediated by chronic serotonergic pharmacotherapy combined with exercise following complete SCI in the adult rat. *Exp Neurol* 304, 132-142.
- Gerasimenko, Y., Roy, R.R., Edgerton, V.R., 2008. Epidural stimulation: comparison of the spinal circuits that generate and control locomotion in rats, cats and humans. *Exp Neurol* 209, 417-425.
- Gerasimenko, Y.P., Lu, D.C., Modaber, M., Zdunowski, S., Gad, P., Sayenko, D.G., Morikawa, E., Haakana, P., Ferguson, A.R., Roy, R.R., Edgerton, V.R., 2015. Noninvasive Reactivation of Motor Descending Control after Paralysis. *J Neurotrauma* 32, 1968-1980.
- Giggins, O.M., Persson, U.M., Caulfield, B., 2013. Biofeedback in rehabilitation. *J Neuroeng Rehabil* 10, 60.
- Gill, M.L., Grahn, P.J., Calvert, J.S., et al. 2018. Neuromodulation of lumbosacral spinal networks enables independent stepping after complete paraplegia. *Nat Med* 24, 1677-1682.
- Giroux, N., Rossignol, S., Reader, T.A., 1999. Autoradiographic study of $\alpha 1$ - and $\alpha 2$ -noradrenergic and serotonin1A receptors in the spinal cord of normal and chronically transected cats. *J. Comp. Neurol* 406, 402-414.
- Glaessgen, E., Stargel, D., 2012. The digital twin paradigm for future NASA and US Air Force vehicles, 53rd AIAA/ASME/ASCE/AHS/ASC Structures, Structural Dynamics and Materials Conference 20th AIAA/ASME/AHS Adaptive Structures Conference 14th AIAA, p. 1818.
- Gozal, E.A., Neill, B.E., Sawchuk, M.A., Zhu, H., Halder, M., Chou, C.-C., Hochman, S., 2014. Anatomical and functional evidence for trace amines as unique modulators of locomotor function in the mammalian spinal cord. *Front. Neural Circuits* 8, 134.
- GrandPré, T., Li, S., Strittmatter, S.M., 2002. Nogo-66 receptor antagonist peptide promotes axonal regeneration. *Nature* 417, 547-551.

- Grau, J.W., Baine, R.E., Bean, P.A., Davis, J.A., Fauss, G.N., Henwood, M.K., Hudson, K.E., Johnston, D.T., Tarbet, M.M., Strain, M.M., 2020. Learning to promote recovery after spinal cord injury. *Exp Neurol* 330, 113334.
- Griffin, L., Decker, M.J., Hwang, J.Y., Wang, B., Kitchen, K., Ding, Z., Ivy, J.L., 2009. Functional electrical stimulation cycling improves body composition, metabolic and neural factors in persons with spinal cord injury. *J Electromyogr Kinesiol* 19, 614-622.
- Grijalva, I., Guízar-Sahagún, G., Castaneda-Hernández, G., Mino, D., Maldonado-Julián, H., Vidal-Cantú, G., Ibarra, A., Serra, O., Salgado-Ceballos, H., Arenas-Hernández, R., 2003. Efficacy and Safety of 4-Aminopyridine in Patients with Long-Term Spinal Cord Injury: A Randomized, Double-Blind, Placebo-Controlled Trial. *Pharmacotherapy* 23, 823-834.
- Grillner, S., Zangger, P., 1979. On the central generation of locomotion in the low spinal cat. *Exp. Brain Res.* 34, 241-261.
- Gruner, J.A., Yee, A.K., 1999. 4-Aminopyridine enhances motor evoked potentials following graded spinal cord compression injury in rats. *Brain Research* 816, 446-456.
- Guertin, P.A., Ung, R.V., Rouleau, P., 2010. Oral administration of a tri-therapy for central pattern generator activation in paraplegic mice: proof-of-concept of efficacy. *Biotechnol J* 5, 421-426.
- Guest, J.D., Hiester, E.D., Bunge, R.P., 2005. Demyelination and Schwann cell responses adjacent to injury epicenter cavities following chronic human spinal cord injury. *Exp Neurol* 192, 384-393.
- Gurden, G.G., Feringa, E.R., 1974. Effects of sympatholytic agents on CNS regeneration, cystic necrosis after spinal cord transection. *Neurology* 24, 187-191.
- Guth, L., Albuquerque, E.X., Deshpande, S.S., Barrett, C.P., Donati, E.J., Warnick, J.E., 1980. Ineffectiveness of enzyme therapy on regeneration in the transected spinal cord of the rat. 52, 73.
- Hall, E.D., Braughler, J.M., 1981. Acute effects of intravenous glucocorticoid pretreatment on the in vitro peroxidation of cat spinal cord tissue. *Exp Neurol* 73, 321-324.
- Hall, E.D., Wolf, D.L., Braughler, J.M., 1984. Effects of a single large dose of methylprednisolone sodium succinate on experimental posttraumatic spinal cord ischemia. Dose-response and time-action analysis. *J Neurosurg* 61, 124-130.

- Hall, O.T., McGrath, R.P., Peterson, M.D., Chadd, E.H., DeVivo, M.J., Heinemann, A.W., Kalpakjian, C.Z., 2019. The Burden of Traumatic Spinal Cord Injury in the United States: Disability-Adjusted Life Years. *Arch Phys Med* 100, 95-100.
- Harkema, S., Gerasimenko, Y., Hodes, J., Burdick, J., Angeli, C., Chen, Y., Ferreira, C., Willhite, A., Rejc, E., Grossman, R.G., Edgerton, V.R., 2011. Effect of epidural stimulation of the lumbosacral spinal cord on voluntary movement, standing, and assisted stepping after motor complete paraplegia: a case study. *Lancet* 377, 1938-1947.
- Harnie, J., Doelman, A., de Vette, E., Audet, J., Desrochers, E., Gaudreault, N., Frigon, A., 2019. The recovery of standing and locomotion after spinal cord injury does not require task-specific training. *Elife* 8, e50134.
- Hawthorne, A.L., Hu, H., Kundu, B., Steinmetz, M.P., Wylie, C.J., Deneris, E.S., Silver, J., 2011. The unusual response of serotonergic neurons after CNS Injury: lack of axonal dieback and enhanced sprouting within the inhibitory environment of the glial scar. *J Neurosci* 31, 5605-5616.
- Hayes, V.Y., Isackson, P.J., Fabrazzo, M., Follesa, P., Mocchetti, I., 1995. Induction of nerve growth factor and basic fibroblast growth factor mRNA following clenbuterol: contrasting anatomical and cellular localization. *Exp Neurol* 132, 33-41.
- Holanda, L.J., Silva, P.M.M., Amorim, T.C., Lacerda, M.O., Simão, C.R., Morya, E., 2017. Robotic assisted gait as a tool for rehabilitation of individuals with spinal cord injury: a systematic review. *J Neuroeng Rehabil* 14, 126.
- Holmes, G.M., Blanke, E.N., 2019. Gastrointestinal dysfunction after spinal cord injury. *Exp Neurol* 320, 113009.
- Hong, L.T.A., Kim, Y.M., Park, H.H., Hwang, D.H., Cui, Y., Lee, E.M., Yahn, S., Lee, J.K., Song, S.C., Kim, B.G., 2017. An injectable hydrogel enhances tissue repair after spinal cord injury by promoting extracellular matrix remodeling. *Nat Commun* 8, 533.
- Hostrup, M., Jacobson, G.A., Jessen, S., Lemminger, A.K., 2020. Anabolic and lipolytic actions of beta2 -agonists in humans and antidoping challenges. *Drug Test Anal* 12, 597-609.
- Hostrup, M., Reitelseder, S., Jessen, S., Kalsen, A., Nyberg, M., Egelund, J., Kreiberg, M., Kristensen, C.M., Thomassen, M., Pilegaard, H., Backer, V., Jacobson, G.A., Holm, L., Bangsbo, J., 2018. Beta2-adrenoceptor agonist salbutamol increases protein turnover rates and alters signalling in skeletal muscle after resistance exercise in young men. *J Physiol* 596, 4121-4139.

- Houle, J.D., Tom, V.J., Mayes, D., Wagoner, G., Phillips, N., Silver, J., 2006. Combining an autologous peripheral nervous system "bridge" and matrix modification by chondroitinase allows robust, functional regeneration beyond a hemisection lesion of the adult rat spinal cord. *J Neurosci* 26, 7405-7415.
- Huie, J.R., Morioka, K., Haefeli, J., Ferguson, A.R., 2017. What Is Being Trained? How Divergent Forms of Plasticity Compete To Shape Locomotor Recovery after Spinal Cord Injury. *J Neurotrauma* 34, 1831-1840.
- Hubscher, C.H., Herrity, A.N., Williams, C.S., Montgomery, L.R., Willhite, A.M., Angeli, C.A., Harkema, S.J., 2018. Improvements in bladder, bowel and sexual outcomes following task-specific locomotor training in human spinal cord injury. *PLoS One* 13, e0190998.
- Jackson, A., Zimmermann, J.B., 2012. Neural interfaces for the brain and spinal cord--restoring motor function. *Nat Rev Neurol* 8, 690-699.
- James, N.D., Bartus, K., Grist, J., Bennett, D.L.H., McMahon, S.B., Bradbury, E.J., 2011. Conduction Failure following Spinal Cord Injury: Functional and Anatomical Changes from Acute to Chronic Stages. *J. Neurosci* 31, 18543.
- Jeffrey-Gauthier, R., Josset, N., Bretzner, F., Leblond, H., 2018. Facilitation of Locomotor Spinal Networks Activity by Buspirone after a Complete Spinal Cord Lesion in Mice. *J Neurotrauma* 35, 2208-2221.
- Jogia, T., Ruitenberg, M.J., 2020. Traumatic Spinal Cord Injury and the Gut Microbiota: Current Insights and Future Challenges. *Front Immunol* 11, 704.
- Kabatas, S., Yu, D., He, X.D., Thatte, H.S., Benedict, D., Hepgul, K.T., Black, P.M., Sabharwal, S., Teng, Y.D., 2008. Neural and anatomical abnormalities of the gastrointestinal system resulting from contusion spinal cord injury. *Neuroscience* 154, 1627-1638.
- Kakulas, B.A., 1999. The applied neuropathology of human spinal cord injury. *Spinal Cord* 37, 79-88.
- Kanai, A., Zabbarova, I., Ikeda, Y., Yoshimura, N., Birder, L., Hanna-Mitchell, A., de Groat, W., 2011. Sophisticated models and methods for studying neurogenic bladder dysfunction. *Neurourol Urodyn* 30, 658-667.
- Kapelner, T., Sartori, M., Negro, F., Farina, D., 2020. Neuro-Musculoskeletal Mapping for Man-Machine Interfacing. *Sci Rep* 10, 5834.
- Kern, H., Hofer, C., Loeffler, S., Zampieri, S., Gargiulo, P., Baba, A., Marcante, A., Piccione, F., Pond, A., Carraro, U., 2017. Atrophy, ultra-structural disorders, severe atrophy and

- degeneration of denervated human muscle in SCI and Aging. Implications for their recovery by Functional Electrical Stimulation, updated 2017. *Neurol Res* 39, 660-666.
- Kim, M., Hong, S.K., Jeon, S.R., Roh, S.W., Lee, S., 2018. Early (≤ 48 Hours) versus Late (> 48 Hours) Surgery in Spinal Cord Injury: Treatment Outcomes and Risk Factors for Spinal Cord Injury. *World Neurosurg* 118, e513-e525.
- Kirshblum, S., Botticello, A., Benedetto, J., Donovan, J., Marino, R., Hsieh, S., Wagaman, N., 2020. A Comparison of Diagnostic Stability of the ASIA Impairment Scale Versus Frankel Classification Systems for Traumatic Spinal Cord Injury. *Arch Phys Med Rehabil* 101, 1556-1562.
- Knikou, M., A Guertin, P., Minassian, K., 2017. Probing the human spinal locomotor circuits by phasic step-induced feedback and by tonic electrical and pharmacological neuromodulation. *Curr. Pharm. Des* 23, 1805-1820.
- Koyanagi, I., Iwasaki, Y., Isu, T., Akino, M., Abe, H., Mitsumori, K., Nakagawa, T., Sakuragi, M., Saitoh, H., Nomura, M., 1989. [Myelotomy for acute cervical cord injury. Report of four cases]. *Neurol Med Chir (Tokyo)* 29, 302-306.
- Krebs, J., Koch, H.G., Hartmann, K., Frotzler, A., 2016. The characteristics of posttraumatic syringomyelia. *Spinal Cord* 54, 463-466.
- Krishna, V., Andrews, H., Varma, A., Mintzer, J., Kindy, M.S., Guest, J., 2014. Spinal cord injury: how can we improve the classification and quantification of its severity and prognosis? *J Neurotrauma* 31, 215-227.
- Krucoff, M.O., Rahimpour, S., Slutzky, M.W., Edgerton, V.R., Turner, D.A., 2016. Enhancing Nervous System Recovery through Neurobiologics, Neural Interface Training, and Neurorehabilitation. *FrontNeuroscience* 10, 584-584.
- Kucher, K., Johns, D., Maier, D., Abel, R., Badke, A., Baron, H., Thietje, R., Casha, S., Meindl, R., Gomez-Mancilla, B., Pfister, C., Rupp, R., Weidner, N., Mir, A., Schwab, M.E., Curt, A., 2018. First-in-Man Intrathecal Application of Neurite Growth-Promoting Anti-Nogo-A Antibodies in Acute Spinal Cord Injury. *Neurorehabil Neural Repair* 32, 578-589.
- Kuzhandaivel, A., Nistri, A., Mladinic, M., 2010. Kainate-mediated excitotoxicity induces neuronal death in the rat spinal cord in vitro via a PARP-1 dependent cell death pathway (Parthanatos). *Cell Mol Neurobiol* 30, 1001-1012.
- Landry, E.S., Lapointe, N.P., Rouillard, C., Levesque, D., Hedlund, P.B., Guertin, P.A., 2006. Contribution of spinal 5-HT1A and 5-HT7 receptors to locomotor-like

- movement induced by 8-OH-DPAT in spinal cord-transected mice. *Eur J Neurosci* 24, 535-546.
- Lavrov, I., Courtine, G., Dy, C.J., van den Brand, R., Fong, A.J., Gerasimenko, Y., Zhong, H., Roy, R.R., Edgerton, V.R., 2008. Facilitation of stepping with epidural stimulation in spinal rats: role of sensory input. *J Neurosci* 28, 7774-7780.
- Lee, B.B., Cripps, R.A., Fitzharris, M., Wing, P.C., 2014. The global map for traumatic spinal cord injury epidemiology: update 2011, global incidence rate. *Spinal Cord* 52, 110-116.
- Leemhuis, E., De Gennaro, L., Pazzaglia, A.M., 2019. Disconnected Body Representation: Neuroplasticity Following Spinal Cord Injury. *J Clin Med* 8.
- Li, Y., Lucas-Osma, A.M., Black, S., Bandet, M.V., Stephens, M.J., Vavrek, R., Sanelli, L., Fenrich, K.K., Di Narzo, A.F., Dracheva, S., Winship, I.R., Fouad, K., Bennett, D.J., 2017. Pericytes impair capillary blood flow and motor function after chronic spinal cord injury. *Nature Medicine* 23, 733-741.
- Li, Y., Ritzel, R.M., He, J., Cao, T., Sabirzhanov, B., Li, H., Liu, S., Wu, L.J., Wu, J., 2021. The voltage-gated proton channel Hv1 plays a detrimental role in contusion spinal cord injury via extracellular acidosis-mediated neuroinflammation. *Brain Behav Immun*. 91, 267-283
- Liu, D., Thangnipon, W., McAdoo, D.J., 1991. Excitatory amino acids rise to toxic levels upon impact injury to the rat spinal cord. *Brain Res* 547, 344-348.
- Liu, D., Xu, G.Y., Pan, E., McAdoo, D.J., 1999. Neurotoxicity of glutamate at the concentration released upon spinal cord injury. *Neuroscience* 93, 1383-1389.
- Liu, Z., Yao, X., Jiang, W., Li, W., Zhu, S., Liao, C., Zou, L., Ding, R., Chen, J., 2020. Advanced oxidation protein products induce microglia-mediated neuroinflammation via MAPKs-NF- κ B signaling pathway and pyroptosis after secondary spinal cord injury. *J Neuroinflammation* 17, 90.
- Loane, C., Politis, M., 2012. Buspirone: What is it all about? *Brain Res* 1461, 111-118.
- Losey, P., Young, C., Krimholtz, E., Bordet, R., Anthony, D.C., 2014. The role of hemorrhage following spinal-cord injury. *Brain Res* 20, 1569, 9-1518.
- Lotte, F., Bougrain, L., Cichocki, A., Clerc, M., Congedo, M., Rakotomamonjy, A., Yger, F., 2018. A review of classification algorithms for EEG-based brain-computer interfaces: a 10 year update. *J Neural Eng* 15, 031005.
- Maciejasz, P., Eschweiler, J., Gerlach-Hahn, K., Jansen-Troy, A., Leonhardt, S., 2014. A survey on robotic devices for upper limb rehabilitation. *J Neuroeng Rehabil* 11, 3.

- Mandadi, S., Leduc-Pessah, H., Hong, P., Ejdrygiewicz, J., Sharples, S.A., Trang, T., Whelan, P.J., 2016. Modulatory and plastic effects of kinins on spinal cord networks. *J Physiol* 594, 1017-1036.
- Maresh, S., Prowting, J.L., Vaughan, S., Kruppe, E., Alsabri, B., Yarandi, H., Badr, M.S., Sankari, A., 2020. Buspirone Decreases Susceptibility to Hypocapnic Central Sleep Apnea in Chronic SCI Patients. *J Appl Physiol* (1985).
- Martineau, L., Horan, M.A., Rothwell, N.J., Little, R.A., 1992. Salbutamol, a β_2 -adrenoceptor agonist, increases skeletal muscle strength in young men. *Clinical Science* 83, 615-621.
- Matsushita, T., Lankford, K.L., Arroyo, E.J., Sasaki, M., Neyazi, M., Radtke, C., Kocsis, J.D., 2015. Diffuse and persistent blood-spinal cord barrier disruption after contusive spinal cord injury rapidly recovers following intravenous infusion of bone marrow mesenchymal stem cells. *Exp Neurol* 267, 152-164.
- Mayne, R.J., van der Poel, C., Woods, M.G., Lynch, G.S., 2015. Skeletal effects of the alteration of masseter muscle function. *Aust Orthod J* 31, 184-194.
- Mautes, A.E., Weinzierl, M.R., Donovan, F., Noble, L.J., 2000. Vascular events after spinal cord injury: contribution to secondary pathogenesis. *Phys Ther* 80, 673-687.
- Maynard, F.M., Jr., Bracken, M.B., Creasey, G., Ditunno, J.F., Jr., Donovan, W.H., Ducker, T.B., Garber, S.L., Marino, R.J., Stover, S.L., Tator, C.H., Waters, R.L., Wilberger, J.E., Young, W., 1997. International Standards for Neurological and Functional Classification of Spinal Cord Injury. American Spinal Injury Association. *Spinal Cord* 35, 266-274.
- McDonald, J.W., Becker, D., Sadowsky, C.L., Jane, J.A., Sr., Conturo, T.E., Schultz, L.M., 2002. Late recovery following spinal cord injury. Case report and review of the literature. *J Neurosurgery* 97, 252-265.
- McKeon, R.J., Schreiber, R.C., Rudge, J.S., Silver, J., 1991. Reduction of neurite outgrowth in a model of glial scarring following CNS injury is correlated with the expression of inhibitory molecules on reactive astrocytes. *J Neurosci* 11, 3398-3411.
- McPherson, J.G., Miller, R.R., Perlmutter, S.I., 2015. Targeted, activity-dependent spinal stimulation produces long-lasting motor recovery in chronic cervical spinal cord injury. *Proc Natl Acad Sci U S A* 112, 12193-12198.
- Megía García, A., Serrano-Muñoz, D., Taylor, J., Avendaño-Coy, J., Gómez-Soriano, J., 2020. Transcutaneous Spinal Cord Stimulation and Motor Rehabilitation in Spinal Cord Injury: A Systematic Review. *Neurorehabil Neural Repair* 34, 3-12.

- Merkler, D., Metz, G.A., Raineteau, O., Dietz, V., Schwab, M.E., Fouad, K., 2001. Locomotor recovery in spinal cord-injured rats treated with an antibody neutralizing the myelin-associated neurite growth inhibitor Nogo-A. *J Neurosci* 21, 3665-3673.
- Middleton, J.W., Dayton, A., Walsh, J., Rutkowski, S.B., Leong, G., Duong, S., 2012. Life expectancy after spinal cord injury: a 50-year study. *Spinal Cord* 50, 803-811.
- Miyamoto, S., 2019. Japan responds: stem-cell therapy justified. *Nature* 569, 40.
- Miyao, Y., Sasaki, M., Taketsuna, S., Yokota, C., Umegaki, M., 2020. Early Development of Syringomyelia after Spinal Cord Injury: Case Report and Review of the Literature. *NMC Case Rep J* 7, 217-221.
- Monnier, P.P., Sierra, A., Schwab, J.M., Henke-Fahle, S., Mueller, B.K., 2003. The Rho/ROCK pathway mediates neurite growth-inhibitory activity associated with the chondroitin sulfate proteoglycans of the CNS glial scar. *Mol Cell Neurosci* 22, 319-330.
- Morse, L., Teng, Y.D., Pham, L., Newton, K., Yu, D., Liao, W.L., Kohler, T., Muller, R., Graves, D., Stashenko, P., Battaglino, R., 2008. Spinal cord injury causes rapid osteoclastic resorption and growth plate abnormalities in growing rats (SCI-induced bone loss in growing rats). *Osteoporos Int* 19, 645-652.
- Musienko, P., van den Brand, R., Marzendorfer, O., Roy, R.R., Gerasimenko, Y., Edgerton, V.R., Courtine, G., 2011. Controlling specific locomotor behaviors through multidimensional monoaminergic modulation of spinal circuitries. *J Neurosci* 31, 9264-9278.
- Musk, E., Neuralink, 2019. An Integrated Brain-Machine Interface Platform With Thousands of Channels. *J Med Internet Res* 21, e16194.
- Naftchi, N.E., 1982. Functional restoration of the traumatically injured spinal cord in cats by clonidine. *Science* 217, 1042-1044.
- Nardone, R., Holler, Y., Brigo, F., Seidl, M., Christova, M., Bergmann, J., Golaszewski, S., Trinka, E., 2013. Functional brain reorganization after spinal cord injury: systematic review of animal and human studies. *Brain Res* 1504, 58-73.
- Noga, B.R., Johnson, D.M.G., Riesgo, M.I., Pinzon, A., 2009. Locomotor-activated neurons of the cat. I. Serotonergic innervation and co-localization of 5-HT₇, 5-HT_{2A}, and 5-HT_{1A} receptors in the thoraco-lumbar spinal cord. *J Neurophysiol* 102, 1560-1576.
- O'Connor, G., Jeffrey, E., Madorma, D., Marcillo, A., Abreu, M.T., Deo, S.K., Dietrich, W.D., Daunert, S., 2018. Investigation of Microbiota Alterations and Intestinal Inflammation Post-Spinal Cord Injury in Rat Model. *J Neurotrauma* 35, 2159-2166.

- O'Shea, T.M., Burda, J.E., Sofroniew, M.V., 2017. Cell biology of spinal cord injury and repair. *J Clin Invest* 127, 3259-3270.
- Osaka, M., Honmou, O., Murakami, T., Nonaka, T., Houkin, K., Hamada, H., Kocsis, J.D., 2010. Intravenous administration of mesenchymal stem cells derived from bone marrow after contusive spinal cord injury improves functional outcome. *Brain Res* 1343, 226-235.
- Osterholm, J.L., 1974. Noradrenergic mediation of traumatic spinal cord autodestruction. *Life Sciences* 14, 1363-1384.
- Osterholm, J.L., Mathews, G.J., 1972. Altered norepinephrine metabolism, following experimental spinal cord injury. 2. Protection against traumatic spinal cord hemorrhagic necrosis by norepinephrine synthesis blockade with alpha methyl tyrosine. *J Neurosurgery* 36, 395-401.
- Otani, K., Abe, H., Kadoya, S., 1994. Beneficial effect of methylprednisolone sodium succinate in the treatment of acute spinal cord injury. *Sekitsui Sekizui* 7, 633-647
- Oxley, T.J., Opie, N.L., John, S.E., et al. 2016. Minimally invasive endovascular stent-electrode array for high-fidelity, chronic recordings of cortical neural activity. *Nat Biotechnol* 34, 320-327.
- Pearlstein, E., Mabrouk, F.B., Pflieger, J.F., Vinay, L., 2005. Serotonin refines the locomotor-related alternations in the in vitro neonatal rat spinal cord. *Eur J Neurosci* 21, 1338-1346.
- Perrier, J.F., Cotel, F., 2015. Serotonergic modulation of spinal motor control. *Curr Opin Neurobiol* 33, 1-7.
- Perrin, F.E., Gerber, Y.N., Teigell, M., Lonjon, N., Boniface, G., Bauchet, L., Rodriguez, J.J., Hugnot, J.P., Privat, A.M., 2011. Anatomical study of serotonergic innervation and 5-HT(1A) receptor in the human spinal cord. *Cell Death Dis* 2, e218.
- Pershhouse, K.J., Barker, R.N., Kendall, M.B., Buettner, P.G., Kuipers, P., Schuurs, S.B., Amsters, D.I., 2012. Investigating changes in quality of life and function along the lifespan for people with spinal cord injury. *Arch Phys Med Rehabil* 93, 413-419.
- Petitjean, M.E., Pointillart, V., Dixmerias, F., Wiart, L., Sztark, F., Lassié, P., Thicoïpé, M., Dabadie, P., 1998. [Medical treatment of spinal cord injury in the acute stage]. *Annales francaises d'anesthesie et de reanimation* 17, 114-122.
- Pizzolato, C., Lloyd, D.G., Barrett, R.S., Cook, J.L., Zheng, M.H., Besier, T.F., Saxby, D.J., 2017. Bioinspired Technologies to Connect Musculoskeletal Mechanobiology to the Person for Training and Rehabilitation. *Front Comput Neurosci* 11, 96.

- Pizzolato, C., Saxby, D.J., Palipana, D., Diamond, L.E., Barrett, R.S., Teng, Y.D., Lloyd, D.G., 2019. Neuromusculoskeletal Modeling-Based Prostheses for Recovery After Spinal Cord Injury. *Front Neurobot* 13, 97.
- Pizzolato, C., Shim, V.B., Lloyd, D.G., Devaprakash, D., Obst, S.J., Newsham-West, R., Graham, D.F., Besier, T.F., Zheng, M.H., Barrett, R.S., 2020. Targeted Achilles tendon training and rehabilitation using personalized and real-time multiscale models of the neuromusculoskeletal system. *Front Bioeng Biotechnol* 8, 878.
- Rabchevsky, A.G., Michael, F.M., Patel, S.P., 2020. Mitochondria focused neurotherapeutics for spinal cord injury. *Exp Neurol* 330, 113332.
- Radhakrishna, M., Steuer, I., Prince, F., Roberts, M., Mongeon, D., Kia, M., Dyck, S., Matte, G., Vaillancourt, M., Guertin, P.A., 2017. Double-Blind, Placebo-Controlled, Randomized Phase I/IIa Study (Safety and Efficacy) with Bupirone/Levodopa/Carbidopa (Spinalon™) in Subjects with Complete AIS A or Motor-Complete AIS B Spinal Cord Injury. *Curr Pharm Des* 23, 1789-1804.
- Ragnarsson, K.T., 2008. Functional electrical stimulation after spinal cord injury: current use, therapeutic effects and future directions. *Spinal Cord* 46, 255-274.
- Rodgers, H.M., Yow, J., Evans, E., Clemens, S., Brewer, K.L., 2019. Dopamine D1 and D3 receptor modulators restore morphine analgesia and prevent opioid preference in a model of neuropathic pain. *Neurosci* 406, 376-388.
- Rojas, P.S., Fiedler, J.L., 2016. What Do We Really Know About 5-HT1A Receptor Signaling in Neuronal Cells? *Front Cell Neurosci* 10, 272.
- Ropper, A.E., Thakor, D.K., Han, I., Yu, D., Zeng, X., Anderson, J.E., Aljuboory, Z., Kim, S.W., Wang, H., Sidman, R.L., Zafonte, R.D., Teng, Y.D., 2017. Defining recovery neurobiology of injured spinal cord by synthetic matrix-assisted hMSC implantation. *Proc Natl Acad Sci U S A* 114, E820-E829.
- Rosenberg, L.J., Teng, Y.D., Wrathall, J.R., 1999a. 2,3-Dihydroxy-6-nitro-7-sulfamoylbenzo(f)quinoxaline reduces glial loss and acute white matter pathology after experimental spinal cord contusion. *J Neurosci* 19, 464-475.
- Rosenberg, L.J., Teng, Y.D., Wrathall, J.R., 1999b. Effects of the sodium channel blocker tetrodotoxin on acute white matter pathology after experimental contusive spinal cord injury. *J Neurosci* 19, 6122-6133.
- Rosenblum, D., 1993. Clonidine-induced bradycardia in patients with spinal cord injury. *Arch Phys Med Rehabil* 74, 1206-1207.

- Rupp, R., Schließmann, D., Plewa, H., Schuld, C., Gerner, H.J., Weidner, N., Hofer, E.P., Knestel, M., 2015. Safety and efficacy of at-home robotic locomotion therapy in individuals with chronic incomplete spinal cord injury: a prospective, pre-post intervention, proof-of-concept study. *PLoS One* 10, e0119167.
- Saraswat Ohri, S., Bankston, A.N., Mullins, S.A., Liu, Y., Andres, K.R., Beare, J.E., Howard, R.M., Burke, D.A., Riegler, A.S., Smith, A.E., Hetman, M., Whittemore, S.R., 2018a. Blocking Autophagy in Oligodendrocytes Limits Functional Recovery after Spinal Cord Injury. *J Neurosci* 38, 5900-5912.
- Saraswat Ohri, S., Mullins, A., Hetman, M., Whittemore, S.R., 2018b. Activating Transcription Factor-6alpha Deletion Modulates the Endoplasmic Reticulum Stress Response after Spinal Cord Injury but Does Not Affect Locomotor Recovery. *J Neurotrauma* 35, 486-491.
- Sartori, M., Durandau, G., Dosen, S., Farina, D., 2018. Robust simultaneous myoelectric control of multiple degrees of freedom in wrist-hand prostheses by real-time neuromusculoskeletal modeling. *J Neural Eng* 15, 066026.
- Sartori, M., Yavuz, U.S., Farina, D., 2017. In Vivo Neuromechanics: Decoding Causal Motor Neuron Behavior with Resulting Musculoskeletal Function. *Sci Rep* 7, 13465.
- Sasada, S., Kato, K., Kadowaki, S., Groiss, S.J., Ugawa, Y., Komiyama, T., Nishimura, Y., 2014. Volitional walking via upper limb muscle-controlled stimulation of the lumbar locomotor center in man. *J Neurosci* 34, 11131-11142.
- Savic, G., DeVivo, M.J., Frankel, H.L., Jamous, M.A., Soni, B.M., Charlifue, S., 2017. Long-term survival after traumatic spinal cord injury: a 70-year British study. *Spinal Cord* 55, 651-658.
- Scivoletto, G., Tamburella, F., Laurenza, L., Torre, M., Molinari, M., 2014. Who is going to walk? A review of the factors influencing walking recovery after spinal cord injury. *Front Hum Neurosci* 8, 141-141.
- Segal, J.L., Brunnemann, S.R., 1998. 4-Aminopyridine Alters Gait Characteristics and Enhances Locomotion in Spinal Cord Injured Humans. *J Spinal Cord Med* 21, 200-204.
- Segal, J.L., Pathak, M.S., Hernandez, J.P., Himber, P.L., Brunnemann, S.R., Charter, R.S., 1999. Safety and Efficacy of 4-Aminopyridine in Humans with Spinal Cord Injury: A Long-Term, Controlled Trial. *Pharmacotherapy* 19, 713-723.
- Selfslagh, A., Shokur, S., Campos, D.S.F., Donati, A.R.C., Almeida, S., Yamauti, S.Y., Coelho, D.B., Bouri, M., Nicoletis, M.A.L., 2019. Non-invasive, Brain-controlled

- Functional Electrical Stimulation for Locomotion Rehabilitation in Individuals with Paraplegia. *Sci Rep* 9, 6782.
- Shank, C.D., Walters, B.C., Hadley, M.N., 2019. Current Topics in the Management of Acute Traumatic Spinal Cord Injury. *Neurocrit Care* 30, 261-271.
- Sharples, S.A., Koblinger, K., Humphreys, J.M., Whelan, P.J., 2014. Dopamine: a parallel pathway for the modulation of spinal locomotor networks. *Front Neural Circuits* 8, 55-55.
- Shokur, S., Donati, A.R.C., Campos, D.S.F., Gitti, C., Bao, G., Fischer, D., Almeida, S., Braga, V.A.S., Augusto, P., Petty, C., Alho, E.J.L., Lebedev, M., Song, A.W., Nicolelis, M.A.L., 2018. Training with brain-machine interfaces, visuo-tactile feedback and assisted locomotion improves sensorimotor, visceral, and psychological signs in chronic paraplegic patients. *PLoS One* 13, e0206464.
- Shokur, S., Gallo, S., Moiola, R.C., Donati, A.R.C., Morya, E., Bleuler, H., Nicolelis, M.A.L., 2016. Assimilation of virtual legs and perception of floor texture by complete paraplegic patients receiving artificial tactile feedback. *Sci Rep* 6, 32293.
- Signorile, J.F., Banovac, K., Gomez, M., Flipse, D., Caruso, J.F., Lowensteyn, I., 1995. Increased muscle strength in paralyzed patients after spinal cord injury: effect of beta-2 adrenergic agonist. *Arch Phys Med Rehabil* 76, 55-58.
- Springer, J.E., Azbill, R.D., Knapp, P.E., 1999. Activation of the caspase-3 apoptotic cascade in traumatic spinal cord injury. *Nature Medicine* 5, 943-946.
- Steward, O., Sharp, K., Yee, K.M., Hofstadter, M., 2008. A re-assessment of the effects of a Nogo-66 receptor antagonist on regenerative growth of axons and locomotor recovery after spinal cord injury in mice. *Exp Neurol* 209, 446-468.
- Stewart, J.E., Barbeau, H., Gauthier, S., 1991. Modulation of Locomotor Patterns and Spasticity with Clonidine in Spinal Cord Injured Patients. *Can J Neurol Sci* 18, 321-332.
- Stoffel, J.T., Van der Aa, F., Wittmann, D., Yande, S., Elliott, S., 2018. Fertility and sexuality in the spinal cord injury patient. *World J Urol* 36, 1577-1585.
- Strain, M.M., Hook, M.A., Reynolds, J.D., Huang, Y.J., Henwood, M.K., Grau, J.W., 2019. A brief period of moderate noxious stimulation induces hemorrhage and impairs locomotor recovery after spinal cord injury. *Physiol Behav* 212, 112695.
- Suberviola, B., González-Castro, A., Llorca, J., Ortiz-Melón, F., Miñambres E., 2008. Early complications of high-dose methylprednisolone in acute spinal cord injury patients. *Injury* 39, 748-752.

- Telemacque, D., Zhu, F.Z., Ren, Z.W., Chen, K.F., Drepaal, D., Yao, S., Yang, F., Qu, Y.Z., Sun, T.F., Guo, X.D., 2020. Effects of durotomy versus myelotomy in the repair of spinal cord injury. *Neural Regen Res* 15, 1814-1820.
- Teng, Y.D., 2019a. Functional Multipotency of Stem Cells and Recovery Neurobiology of Injured Spinal Cords. *Cell Transplant* 28, 451-459.
- Teng, Y.D., 2019b. Functional multipotency of stem cells: Biological traits gleaned from neural progeny studies. *Semin Cell Dev Biol* 95, 74-83.
- Teng, Y.D., Bingaman, M., DaSilva, A.M.T., Pace, P.P., Gillis, R.A., Wrathall, J.R., 2003. Serotonin 1A receptor agonists reverse respiratory abnormalities in spinal cord-injured rats. *J Neurosci* 23, 4182-4189.
- Teng, Y.D., Choi, H., Onario, R.C., Zhu, S., Desilets, F.C., Lan, S., Woodard, E.J., Snyder, E.Y., Eichler, M.E., Friedlander, R.M., 2004. Minocycline inhibits contusion-triggered mitochondrial cytochrome c release and mitigates functional deficits after spinal cord injury. *Proc Natl Acad Sci U S A* 101, 3071-3076.
- Teng, Y.D., Lavik, E.B., Qu, X., Park, K.I., Ourednik, J., Zurakowski, D., Langer, R., Snyder, E.Y., 2002. Functional recovery following traumatic spinal cord injury mediated by a unique polymer scaffold seeded with neural stem cells. *Proc Natl Acad Sci U S A* 99, 3024-3029.
- Teng, Y.D., Wrathall, J.R., 1997. Local blockade of sodium channels by tetrodotoxin ameliorates tissue loss and long-term functional deficits resulting from experimental spinal cord injury. *J Neurosci* 17, 4359-4366.
- Ung, R.-V.V., Rouleau, P., Guertin, P.A., 2012. Functional and physiological effects of treadmill training induced by buspirone, carbidopa, and L-DOPA in clenbuterol-treated paraplegic mice. *Neurorehabil Neural Repair* 26, 385-394.
- van De Meent, H., Hosman, A.J., Hendriks, J., Zwarts, M., Group, E.-S.S., Schubert, M., 2010. Severe degeneration of peripheral motor axons after spinal cord injury: a European multicenter study in 345 patients. *Neurorehabil Neural Repair* 24, 657-665.
- van Silfhout, L., Peters, A.E., Berlowitz, D.J., Schembri, R., Thijssen, D., Graco, M., 2016. Long-term change in respiratory function following spinal cord injury. *Spinal Cord* 54, 714-719.
- Vardjan, N., Horvat, A., Anderson, J.E., Yu, D., Croom, D., Zeng, X., Luznik, Z., Kreft, M., Teng, Y.D., Kirov, S.A., Zorec, R., 2016. Adrenergic activation attenuates astrocyte swelling induced by hypotonicity and neurotrauma. *Glia* 64, 1034-1049.

- Visavadiya, N.P., Patel, S.P., VanRooyen, J.L., Sullivan, P.G., Rabchevsky, A.G., 2016. Cellular and subcellular oxidative stress parameters following severe spinal cord injury. *Redox Biol* 8, 59-67.
- Vise, W.M., Yashon, D., Hunt, W.E., 1974. Mechanisms of norepinephrine accumulation within sites of spinal cord injury. *J Neurosurgery* 40, 76-82.
- Vismara, I., Papa, S., Rossi, F., Forloni, G., Veglianesi, P., 2017. Current Options for Cell Therapy in Spinal Cord Injury. *Trends Mol Med* 23, 831-849.
- Vivodtzev, I., Picard, G., O'Connor, K., Taylor, J.A., 2020. Serotonin 1A agonist and cardiopulmonary improvements with whole-body exercise in acute, high-level spinal cord injury: a retrospective analysis. *Eur J Appl Physiol* doi: 10.1007/s00421-020-04536-w. Online ahead of print.
- Wagner, F.B., Mignardot, J.B., Le Goff-Mignardot, C.G., et al. 2018. Targeted neurotechnology restores walking in humans with spinal cord injury. *Nature* 563, 65-71.
- Wenger, N., Moraud, E.M., Gandar, J., et al. 2016. Spatiotemporal neuromodulation therapies engaging muscle synergies improve motor control after spinal cord injury. *Nat Med* 22, 138-145.
- Wrathall, J.R., Teng, Y.D., Choiniere, D., Mundt, D.J., 1992. Evidence that local non-NMDA receptors contribute to functional deficits in contusive spinal cord injury. *Brain Res* 586, 140-143.
- Yoon, H., Moon, W.J., Nahm, S.S., Kim, J., Eom, K., 2018. Diffusion Tensor Imaging of Scarring, Necrosis, and Cavitation Based on Histopathological Findings in Dogs with Chronic Spinal Cord Injury: Evaluation of Multiple Diffusion Parameters and Their Correlations with Histopathological Findings. *J Neurotrauma* 35, 1387-1397.
- Young, W., Flamm, E.S., 1982. Effect of high-dose corticosteroid therapy on blood flow, evoked potentials, and extracellular calcium in experimental spinal injury. *J Neurosurgery* 57, 667-673.
- Yu, D., Neeley, W.L., Pritchard, C.D., Slotkin, J.R., Woodard, E.J., Langer, R., Teng, Y.D., 2009. Blockade of peroxynitrite-induced neural stem cell death in the acutely injured spinal cord by drug-releasing polymer. *Stem Cells* 27, 1212-1222.
- Zeman, R.J., Feng, Y., Peng, H., Etlinger, J.D., 1999. Clenbuterol, a beta(2)-adrenoceptor agonist, improves locomotor and histological outcomes after spinal cord contusion in rats. *Exp Neurol* 159, 267-273.

- Zeman, R.J., Zhang, Y., Etlinger, J.D., 1997. Clenbuterol, a beta2-adrenoceptor agonist, reduces scoliosis due to partial transection of rat spinal cord. *Am J Physiol* 272, E712-715.
- Zeman, R.J., Ludemann, R., Etlinger, J.D., 1987. Clenbuterol, a β 2-agonist, retards atrophy in denervated muscles. *Am J Physiol* 252, E152-E155.
- Zhang, X., Andren, P.E., Svenningsson, P., 2006. Repeated l-DOPA treatment increases c-fos and BDNF mRNAs in the subthalamic nucleus in the 6-OHDA rat model of Parkinson's disease. *Brain Res* 1095, 207-210.
- Zhang, L., Li, Z., Hu, Y., Smith, C., Farewik, E.M.G., Wang, R., 2020. Ankle Joint Torque Estimation Using an EMG-Driven Neuromusculoskeletal Model and an Artificial Neural Network Model. *IEEE Trans Autom Sci Eng*, 1-10.

Figure legends

Figure 1. An overview of pathophysiology and clinical outcome of traumatic spinal cord injury. Clinical spinal cord injury (SCI) produces acute (0-48h), subacute (2-14 days), intermediate (2 weeks-6 months) and chronic (> 6 months) phases of pathophysiological consequences. **(a)** In acute SCI, primary mechanical insult to neurons, glial cells, axons, and blood vessels sets a series of biochemical and pathophysiological events in motion, forming secondary injury processes. **(b)** Continued ischemia and inflammation cause delayed neural cell death, axon demyelination, neuron degeneration, glial malfunction, reactive gliosis, and suppression of endogenous repair responses in the subacute phase of SCI. The events jointly promote development of cystic microcavities (see details in **c**). The spinal cord, in response, strengthens its healing process consisting of astrocyte and pericyte proliferation, astrogliosis, inflammatory responses, etc., to close the wound despite major concession to neurological recovery. **(c)** Around the injury epicenter, post-traumatic syrinx formation is developing in more severely lesioned spinal cords in the intermediate phase and lasts into the chronic stage. Incessant in both phases are Wallerian degeneration, microglia activation, and production of pro-inflammatory cytokines. Inhibitory and excitatory neurotransmitters are imbalanced, paralyzing neural circuitry. Moreover, excessive accumulation of endogenous trace amines triggers capillary constriction and hypoxia. Serotonergic innervation is deprived below the lesion and catecholamine/glutamine/peptide neurotransmissions altered. The spinal cord, brain, and peripheral effectors (see specifics in **d**) endure maladaptive plasticity and chronic degeneration. **(d)** SCI results in systemic complications affecting multiple organs, especially

after high-level lesions. Commonly compromised are the motosensory, musculoskeletal, respiratory, cardiovascular, lower urinary tract, gastrointestinal (including gut microbiota), reproductive, autonomic, and immune systems, further aggravating disability. If not effectively treated, the complications may instigate SIRS and MODS.

Abbreviations: (1) 5-HT: 5-hydroxytryptamine (also called serotonin); (2) CSPGs: chondroitin sulfate proteoglycans; (3) Cyt *c*: cytochrome *c*; (4) MODS: multiple organ dysfunction syndrome; (5) RNS: reactive nitrogen species; (6) ROS: reactive oxygen species; and (7) SIRS: systemic inflammatory response syndrome.

Figure 2. Schematic representation of the interaction between the central and peripheral nervous systems with neuromechanical prostheses and pharmacological adjuncts. Disrupted descending neural tracts prevent motor commands to reach spinal circuits distal to the injury to activate muscles for movement generation. The interrupted motor pathway can be re-functionalized via a neuromechanical prosthesis, wherein electroencephalograms (EMG) associated to the motor intent of the individual are interpreted by a brain-computer interface and converted into a target motion and muscle activation patterns through a personalized digital twin of the individual. Muscle activation patterns and joint torques simulated by the digital twin are provided to assistive devices (e.g., functional electrical stimulation and rehabilitation robotics) that generate limb motion. As ascending afferent pathways are also damaged, sensory inflow from cutaneous mechanoreceptors, as well as that of the visual, auditory, and vestibular systems can be used to augment or substitute the afferent signal from proprioception mechanoreceptors in the lower limbs. In this regard, the digital twin can simulate proprioception mechanoreceptor data from muscles, tendons, plantar foot, and joints to redirect it to corresponding somatosensory areas via cross-reality interface (e.g., virtual reality, haptic feedback, and auditory feedback). Biosensory signals, derived from EMG, heart and respiratory rate, as well as patterned motion from the upper limbs (e.g., hand gestures), are interpreted by the digital twin to adapt and control the rehabilitation therapy. Finally, drugs such as buspirone can be co-administered to improve respiratory function in order to better aerobic exercise training intensity tolerance, promote neuroplasticity and neurorepair, and increase motor excitability (note: plotting of Figure 2 was assisted by BioRender.com).

Tables

Table 1. Summary of pharmacological neuromodulatory agents for subacute and chronic spinal cord injury

Drug	Mechanism of Action	Effect in Spinal Cord Injury	References
Clonidine	α_2 -adrenoceptor agonist	In animals: potentiated locomotor training and enhanced recovery of locomotion. In humans: variable effect on locomotion (e.g., reduced clonus, improved phasic EMG activity).	Barbeau and 2012; Naftch
Clenbuterol	β_2 -adrenoceptor agonist	In animals: reduced muscle atrophy and bone demineralization; enhanced functional recovery and tissue sparing at the site of injury; and stimulation of NGF expression.	Hayes et al., 1999
Salbutamol	β_2 -adrenoceptor agonist	In humans: increased voluntary muscle force.	Martineau et
Metaproterenol	β_2 -adrenoceptor agonist	In humans: increased voluntary muscle force and protein turnover rates in skeletal muscles.	Hostrup et al
Formoterol	β_2 -adrenoceptor agonist	In humans: induced anabolic response in skeletal muscles.	Mayne et al.,
Buspirone	Full agonist on presynaptic 5-HT _{1A} autoreceptors and partial agonist on postsynaptic 5-HT _{1A} receptors	In animals: reversed respiratory abnormalities and facilitated locomotor function by re-establishing symmetrical gait with long lasting effects. In humans: improved muscle coordination and upper limb strength when combined with appropriate rehabilitation; enhanced respiration, cardiovascular function, and thermoregulation; facilitated neurotransmission in dormant descending pathways; and reduced susceptibility to induced hypocapnic central sleep apnea.	Choi et al., 2018; Freyvert et al., 2018; Gerasimenko et al., 2018; Lo et al., 2020; Ter et al., 2020
DOPA	Increased dopamine and noradrenaline production and terminal release of neurotransmitters from the descending noradrenergic pathway	In spinal animals: generated rhythmic motor output in the absence of phasic peripheral feedback when co-administered with the monoamine oxidase inhibitor nialamide.	Andén et al.,

(Abbreviations: 5-HT: 5-hydroxytryptamine; DOPA: l-3,4-dihydroxyphenylalanine or levodopa; EMG: electromyogram; NGF: nerve growth factors)

Title: Non-Invasive Approaches to Functional Recovery after Spinal Cord Injury: Therapeutic Targets and Multimodal Device Interventions

Authors: Claudio Pizzolato^{1,2}, Mehmet A. Gunduz⁷, Dinesh Palipana^{2,3,4,5}, Jingnan Wu⁷, Gary Grant⁶, Susan Hall⁶, Rachel Dennison⁷, Ross D. Zafonte⁷, David G. Lloyd^{1,2}, Yang D. Teng^{7*}

¹ School of Allied Health Sciences, Griffith University, Gold Coast, QLD, Australia,

² Griffith Centre of Biomedical and Rehabilitation Engineering, Menzies Health Institute Queensland, and Advanced Design and Prototyping Technologies Institute (ADAPT), Griffith University, Gold Coast, QLD, Australia

³ The Hopkins Centre, Menzies Health Institute Queensland, Griffith University, Gold Coast, QLD, Australia

⁴ Gold Coast Hospital and Health Service, Gold Coast, QLD, Australia

⁵ School of Medicine, Griffith University, Gold Coast, QLD, Australia

⁶ School of Pharmacy and Pharmacology, Griffith University, Gold Coast, Australia

⁷ Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Mass General Brigham, and Harvard Medical School, Boston, MA, United States

a. Acute Phase

b. Subacute Phase

c. Intermediate & Chronic Phases

d. Systemic Complications

Figure 1

Figure 2