Postural control performance on the Functional Reach Test: validity of the Kids-Balance Evaluation Systems Test (Kids-BESTest) criteria.

Rosalee M Dewar ^{BPhy (Hons)¹}, Kylie Tucker ^{PhD²}, Andrew P Claus ^{PhD¹}, Robert S Ware ^{PhD³}, Leanne M Johnston ^{PhD¹}

¹ *The University of Queensland, School of Health and Rehabilitation Sciences, Brisbane, Australia,* ² *The University of Queensland, School of Biomedical Sciences, Brisbane, Australia,* ³ *Griffith University, Menzies Health Institute Queensland, Brisbane, Australia.*

Correspondence: Rosalee Dewar, School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane QLD 4072, Australia. Phone: +61 7 3365 2791.
Email: rosalee.sheather@uq.net.au

Publication data

Submitted for publication 1st June 2020

Acknowledgements: We would like to acknowledge the Children's Motor Control Research Collaboration and The University of Queensland Laboratory for Motor Control and Pain Research for support to conduct the study.

Competing interests: None

Source(s) of support: We acknowledge funding support awarded by the Research Foundation, Cerebral Palsy Alliance (PG4114) through the Children's Motor Control Research Collaboration.

ABSTRACT

Objective: Evaluate the validity of the Kids-BESTest clinical criteria for the Functional Reach Test (FRT) forward and lateral with laboratory measures of postural control in children with Cerebral Palsy (CP).

Design: Psychometric study of face, concurrent and content validity

Setting: Clinical laboratory

Participants: Fifty-eight children aged 7-18 years (ambulant CP n=17, typically developing [TD] n=41).

Intervention: Not applicable

Main outcome measures: Stability limits in standing were assessed using the Kids-BESTest FRT forwards [$FRT_{FORWARD}$], FRT lateral preferred [$FRT_{LATERAL(P)}$] and FRT lateral non-preferred [$FRT_{LATERAL(NP)}$]. Force platforms and kinematic markers were used to collect information on centre-of-pressure (CoP) and joint movement during reach. Analyses included: face validity (Kids-BESTest scores compared between CP and TD groups); concurrent validity (agreement between Kids-BESTest scores and kinematically-derived scores); and content validity (relationships between Kids-BESTest scores with kinematic and CoP data).

Results: Face validity was demonstrated with lower scores for CP compared to TD groups for $FRT_{FORWARD}$ ($p<0.001$) and $FRT_{LATERAL(NP)}$ ($p=0.03$) but not $FRT_{LATERAL(P)}$ ($p=0.12$). For concurrent validity, agreement between Kids-BESTest scores and kinematically-derived scores was good to excellent for $FRT_{LATERAL(P/NP)}$ (88-100%) and good for $FRT_{FORWARD}$ (86-88%) for both groups. For content validity, for the CP group, Kids-BESTest scores were correlated with $CoP\text{-}Range_{FORWARD}$ during $FRT_{FORWARD}$ ($r_s=0.68$) and $CoP\text{-}Range_{LATERAL(NP)}$ during $FRT_{LATERAL(NP)}$ ($r_s=0.57$) but not $FRT_{LATERAL(P)}$ ($r_s=0.07$). For kinematic data, correlations were moderate-high between Kids-BESTest scores and range of hip flexion ($r_s=0.51$) and ankle plantar flexion ($r_s=0.75$) during $FRT_{FORWARD}$, and trunk lateral flexion ($r_s=0.66$) during $FRT_{LATERAL(NP)}$.

Conclusion: The $FRT_{FORWARD}$ demonstrated face, concurrent and content validity. The $FRT_{LATERAL(P/NP)}$ demonstrated concurrent validity, but partial face and content validity. To improve validity of Kids-BESTest FRT criteria qualitative descriptors should be added to the quantitative measure of reach.

ABBREVIATIONS

Cerebral palsy (CP)

Centre-of-pressure (CoP)

Functional Reach Test (FRT)

Kids-Balance Evaluation Systems Test (Kids-BESTest)

Typically developing (TD)

Key words: Cerebral Palsy; Postural Balance; Child; Validity; Functional Reach Test

Introduction

The Functional Reach Test (FRT) examines the limits of stability in standing, or extent to which a person can reach, when they move their arms, trunk and head away from the neutral standing position without changing their base of support (i.e. their foot position) or losing balance [1]. This activity is frequently required in daily life for school-aged children for example when reaching forward or sideways to pick up something off the floor, reaching for a book on a shelf or an object in the refrigerator, or to drink from a drinking fountain. The FRT is used in children with cerebral palsy (CP) as a discriminative and evaluative measure of postural control [2]. In a systematic review of clinical tools to assess balance in children with CP, a version of the FRT forward was reported to have moderate evidence for reliability (inter and intra rater) and face validity but not concurrent or content validity [2, 3]. Further examination of validity will strengthen psychometric evidence for use of the FRT in clinical practice and research.

The Kids-Balance Evaluation Systems Test (Kids-BESTest), is the most comprehensive clinical test of postural control available for children [4], and can be used to identify and classify a broad range of postural control difficulties in children with and without CP [4, 5]. The FRT is a key item included in the Stability Limits/Verticality domain of the Kids-BESTest. While the Kids-BESTest Stability Limits/Verticality domain has good-excellent intra-rater, inter-rater and test-retest reproducibility for children with CP (ICC 0.70-0.92) [5], specific validity of the FRT criteria has not been examined.

Kids-BESTest item criteria, such as the FRT, require laboratory validation to determine how well they assess the domain of postural control they purport to evaluate. To establish the validity of the Kids-BESTest version of the FRT to measure limits of stability competence, the criteria needs to be compared to laboratory measures of both balance (i.e., controlling the centre-of-mass [CoM] in relation to the base of support [BoS]) and orientation (i.e., controlling the relationship between the body segments) [6]. Balance can be evaluated using force plates, which produce kinetic data representing Centre-of-Pressure (CoP) movement. Importantly, measurement of CoP outcomes, such as CoP excursion/range, has good reliability during reaching tasks in children with CP

[7-9]. Orientation can be evaluated using kinematic data, such as measurement of joint angle changes, representing body segment position and movement. In children with CP, trunk and limb kinematic data have been used to quantify position in reaching tasks [10, 11].

The aim of this study was to determine validity of the Kids-BESTest clinical criteria for the FRT forward (FRT_{FORWARD}) and lateral to the preferred (P) and non-preferred (NP) side ($FRT_{\text{LATERAL(P)/(NP)}}$) test items. Objectives were to determine: (i) face validity by comparing Kids-BESTest scores between children with and without CP; (ii) concurrent validity, by comparing agreement between Kids-BESTest clinical scores and corresponding laboratory data for reach distance (measurement of reach from digital image) and (iii) content validity, by examining possible relationships between clinical Kids-BESTest scores and laboratory measures for balance (CoP range) and orientation (kinematic - overall joint movement).

Method

Study Design

This study was a psychometric study of validity. Ethical approval was obtained from Human Research Ethics Committees in accordance with Australian National Health and Medical Research Council (NHMRC) guidelines.

Participants

Participants were children with and without CP. Typically developing (TD) children were recruited from the community and children with CP were recruited from a statewide CP service and CP Register. Children were included if they were: i) aged seven to 18 years; and ii) able to follow child-friendly test instructions. Children who were TD were also required to have a history of term birth (38 weeks or greater) and demonstrate motor skills within the typical range (percentile rank >15%) when assessed using the Bruininks-Oseretsky Test of Motor Proficiency, second edition, short form (BOT-2: SF) [12]. Children with CP were required to be able to stand independently for the FRT assessment (Gross Motor Function Classification System (GMFCS) Level I-III) [13]. Children were excluded if they had a history of: i) spasticity management,

such as Botulinum Toxin, in last three months; ii) orthopaedic or neurological surgery in previous 12 months; iii) intellectual, behavioural or medical difficulties that would limit full participation in assessment.

Information forms were provided to children and their parents/guardians along with a verbal explanation of the protocol and a chance to ask questions. Guardians signed a consent form and children signed an assent form prior to participation.

Procedure

All children participated in measurement of height and weight. Children with CP were also classified according to motor distribution (bilateral or unilateral) and severity (GMFCS level). The FRT was then administered as part of the Kids-BESTest, which was being administered as part of a larger study.

Outcome measures

Functional Reach Test – Clinical Scoring

The FRT was administered according to Kids-BESTest instructions published by Dewar and colleagues in 2017 [4]. The FRT is contained within the Kids-BESTest *Stability Limits/Verticality* domain as *Item 7* FRT_{FORWARD} and *Item 8* FRT_{LATERAL}. In each item, children stood independently with feet aligned to tape on the force plate i.e. 10 cm apart at the malleoli and the medial border of the feet at a 15-degree angle. Each item was performed as per the instructions documented in Appendix 1. Reach distance is recorded as the horizontal distance between two marks on paper.

Using Kids-BESTest clinical criteria, each trial was scored from 0 ‘no measurable lean or must be caught’ to 3 ‘maximum to limits’ (See Figure 1). The better of two trials is scored.

Item 7. FUNCTIONAL REACH FORWARD (Best of two trials)				
				Distance reached: ____ cm / in
0	1	2	3	
No measurable lean – or must be caught.	Poor < 16.5cm (6.5in).	Moderate 16.5cm -32cm (6.5–12.5in).	Maximum to limits: >32cm (12.5in).	

Item 8. FUNCTIONAL REACH LATERAL (Best of two trials)				
				Distance reached Left_ cm /in
				Distance reached Right_ cm/in
No measurable lean, or must be caught.	Poor: < 10cm (4in).	Moderate: 10- 25.5cm (4-10in).	Maximum to limit: > 25.5cm (10in).	Left
				Right

Figure 1 Kids-BESTest criteria for Item 7 Functional Reach Test Forward and Item 8 Functional Reach Test Lateral

ImageJ - Joint Kinematics

Joint kinematic analysis was used to assess concurrent and content validity and was performed using video of trials. Video recording using anatomical markers was used to assess (i) distance reached measured from change in wrist position and (ii) change in joint angles of the lower limb and trunk. Video was taken according to the Kids-BESTest protocol [4] with FRT_{FORWARD} collected from the sagittal view and FRT_{LATERAL} collected from the frontal view [4]. Anatomical markers were placed at the wrist (styloid process of the radius), shoulder (acromion), pelvis (anterior superior iliac spine), hip (greater trochanter), knee (lateral tibial condyle), ankle (lateral malleolus) and foot (the fifth metatarsal head). Screen shots were taken from video recordings of the target trials when the researcher said “start” and again when the researcher said “stop”. ImageJ software (v1.52a, National Institutes of Health, USA, <http://imagej.nih.gov/ij>) was used to measure (i) anterior or lateral distance the wrist moved between the start to the end position (See Figure 2A&B) and used to allocate a digital-derived score using the criteria from Figure 1 and (ii) change in joint angles

measured at the ankle, knee and hip between the start to the end position (See Figure 2C&D).

Figure 2 Images, taken from screen shots of video recordings, analysed by ImageJ for: A. Forward distance reached measured at the wrist; B. Lateral distance reached measured at the wrist; C. Changes in angles at the trunk-thigh, knee and ankle for Functional Reach Test Forward and D. Changes in trunk-thigh angle for Functional Reach Test Lateral.

Force platform (Bertec 4060 series) – Centre-of-pressure

While performing the FRT children stood with both feet on a single force plate for the FRT_{FORWARD} or with each foot on a separate plate for FRT_{LATERAL} (See Figure 2). Force plate data were collected during all trials. Data were digitized (16 bit) at a sampling rate

of 1000 samples/s using a Power 1401 data acquisition system with Spike 2 software (Cambridge Electronic Design Limited, Cambridge, UK).

To enable comparison of data obtained from the force plate with clinical data, for content validity, CoP range data was extracted from the same trial used to allocate the Kids-BESTest clinical score i.e. the longest reach of two trials. Start and finish time of trial was recorded on the CoP trace at the time of recording. This was checked visually at the time of data extraction and a cursor placed at start and end time. Start time was identified from the CoP trace, as a small deviation in ML direction, as this coincided with the paper being marked by the pen. End time was identified as the time in which the max AP CoP displacement occurred, just prior to a small deviation of the CoP in the ML direction, as this again coincided with the paper being marked by the pen. Spike 2 software was used to extract peak-to-peak CoP excursion in the antero-posterior (Range_{AP}) and medio-lateral (Range_{ML}) directions between these two time points.

Analysis

Summary statistics are presented as mean (standard deviation; SD) for continuous data and frequency (percentage) for categorical data, and were compared between groups using Student's t-tests or the Chi^2 tests. Face validity was assessed by comparing clinical Kids-BESTest FRT scores between groups (CP/TD) using Mann-Whitney U non-parametric tests. Concurrent validity was determined by calculating the percentage of exact agreement between FRT Kids-BESTest clinical scores as classified according to Kids-BESTest criteria versus score allocated using the distance reached measured from ImageJ. Percentage exact agreement was interpreted as: excellent if >90%, good if >80%, fair if >60% or poor if <60% [5]. Content validity was determined by examining (i) associations between clinical Kids-BESTest FRT scores and CoP data from the $\text{FRT}_{\text{FORWARD}}$ and $\text{FRT}_{\text{LATERAL}}$ for each group and (ii) between group differences between the clinical and laboratory data. Associations between clinical and laboratory data was examined using Spearman rank correlation coefficients. Correlations were interpreted as: very high +/-0.90-1.00, high +/-0.70 to 0.89, moderate +/-0.50 to 0.69, low +/- 0.30 to 0.49 and negligible +/- 0.00 to 0.29 [14]. Between group differences were examined by performing independent samples t-tests on (i) FRT distance reached

in cm and (ii) CoP Range_{AP} and Range_{ML}. Statistical analysis was performed using SPSS software (IBM, Armonk, NY, USA). Significance was set at $P < 0.05$.

RESULTS

Participants

A total of 62 families responded to the expression of interest. Four children with CP were excluded (one had received lower limb surgery in the previous 12 months, two had intellectual disability that limited their full participation in the test items, and one had ataxic type CP). Final participants included 58 children including 41 TD children and 17 children with CP (See Table 1). There were no statistically significant differences between groups for age, weight, height and foot length. Left Hand preference was identified in 2 (5%) of TD children and 4 (24%) of children with CP.

Table 1 Participant characteristics for children with cerebral palsy (CP) and typically developing (TD) children. Data is reported as Mean $\pm$ SD unless otherwise stated.

	CP Group n=17	TD Group n=41	P-value
Male [n (%)]	11 (65%)	21 (51%)	0.40
Age [years] (range)	11.7 $\pm$ 2.7 (7.9-16.9)	10.9 $\pm$ 2.3 (7.8-17.8)	0.27
Body mass index [kg/m ²]	18.1 $\pm$ 4.5	16.8 $\pm$ 3.8	0.29
Height [cm]	146.1 $\pm$ 14.4	150 $\pm$ 13.9	0.42
Weight [kg]	41.1 $\pm$ 15.8	38.9 $\pm$ 14.6	0.62
GMFCS			
I [n (%)]	11 (65%)	N/A	
II [n (%)]	6 (35%)		
MACS			
I [n (%)]	13 (76%)	N/A	
II [n (%)]	4 (24%)		
Motor Distribution			
Hemiplegia [n (%)]	13 (76%)	N/A	
Diplegia [n (%)]	4 (24%)		
Left Hand preference [n(%)]	4(24)	2(5)	
Kids-BESTest			
Total score /108	75 $\pm$ 14.9	99 $\pm$ 5.2	<0.01

GMFCS, Gross Motor Function Classification System; MACS, Manual Ability Classification System; Kids-BESTest, Kids Balance Evaluation Systems Test, N/A, not applicable.

Face Validity

Typically developing children performed well on the FRT_{FORWARD} with all children scoring 2 points (59%, n=24) or 3 points (41%, n=17). For the FRT_{LATERAL} TD children scored slightly lower with the more scoring 2 points (95%, preferred n=39, 90%, non-preferred n= 37) and fewer scoring 3 points (Preferred n=1, non-preferred n=3) or 1 point (preferred n=1, non-preferred n=1).

For the CP group, when performing the FRT_{FORWARD}, more scored 2 points (65%, n=11) than 1 point (23.5%, n=4) or 3 points (11.5%, n=2). For the FRT_{LATERAL} no child with CP scored 3 points and fewer children with CP scored 2 points than TD children (88% preferred n= 15, 82%, non-preferred n=14) and more children with CP scored 1 point (preferred n=2, non-preferred n= 3). For the CP group clinical scores were lower than the TD group for FRT_{FORWARD} ($p<0.01$) and FRT_{LATERAL(NP)} ($p=0.03$) but not FRT_{LATERAL(P)} ($p=0.12$) (See Table 2).

Table 2 Clinical measures and laboratory measures for children with CP (n=17) and typically developing children (n=41).

		Functional Reach Forward (FRT _{FORWARD})				Functional Reach Lateral –Preferred (FRT _{LATERAL(P)})				Functional Reach Lateral – Non-preferred (FRT _{LATERAL(NP)})			
		TD*	CP*	Mean diff& 95% CI	P- value	TD*	CP*	Mean diff& 95% CI	P- value	TD*	CP*	Mean diff& 95% CI	P- value
Clinical Measures													
FRT [cm]		30.6 (6.1)	22.6 (7.0)	8.0 4.3, 11.7	<0.01	18.7 (3.9)	14.7 (4.3)	4.0 1.7, 6.4	<0.01	18.6 (4.2)	15.6 (5.0)	3.0 0.4, 5.6	0.02
Kids-BESTest FRT score		2 [2-3]	2 [1-3]	N/A	<0.01	2 [1-3]	2 [1-2]	N/A	0.12	2 [1-3]	2 [1-2]	N/A	0.03
Laboratory Measures													
CoP [mm]	Range _{AP}	103.3 (24.6)	81.7 (31.6)	21.6 6.1, 37.1	<0.01	32.1 (16.1)	28.6 (9.8)	1.9 -4.9, 11.9	0.41	34.0 (14.8)	37.3 ^a (20.1)	-3.4 -13.1, 4.8	0.49
	Range _{ML}	27.6 (9.3)	37.3 (18.1)	-9.7 -16.9, -2.5	<0.01	104.6 (27.5)	90.9 (30.3)	8.4 -8.0, 24.7	0.31	104.8 (33.7)	108.1 ^a (41.2)	2.8 -24.5, 17.9	0.76
Joint Kinematics	Change in Angles:												
	Trunk-thigh	49.8 ^a (14.0)	33.8^b (14.9)	15.0 6.5, 23.5	<0.01	23.2 ^c [8.3]	16.8 [7.7]	6.4 1.7, 11.2	<0.01	22.0 ^c (7.3)	14.4 (7.0)	6.5 3.3, 11.8	<0.01
	Knee	3.6 ^a (4.0)	2.7 ^b (5.6)	0.9 -1.8, 3.6	0.50								
	Ankle	6.5 ^a (3.7)	5.3 ^b (5.3)	1.2 -1.3, 3.8	0.33								

*Summary statistics presented as mean (standard deviation) for all measures except Kids-BESTest FRT score, which is median (range) & TD and CP children compared using Student's t-test, except for Kids BESTest score (Mann Whitney U test). N/A = not applicable. Missing values due to video malfunction ^an=1, ^bn=2, ^cn=3.

Concurrent validity

Agreement between clinical Kids-BESTest scores and digital-derived scores for the $FRT_{FORWARD}$ was ‘good’ for both the CP (86%; 13/15) and the TD groups (88%; 35/40). Data was missing for two children with CP and one TD child due to incomplete videos. Agreement between clinical Kids-BESTest scores and digital-derived scores for the $FRT_{LATERAL}$ was ‘good to excellent’ for both the CP and TD groups with the agreement $FRT_{LATERAL(NP)}$ 88-100% (51/53; 5 missing values) and $FRT_{LATERAL(P)}$ 100% (51/51; 7 missing values).

Content Validity

Association between clinical and laboratory data

When comparing the clinical Kids-BESTest score with CoP range the TD group showed low significant correlations with $CoP\ Range_{AP}$ for $FRT_{FORWARD}$ and $CoP\ Range_{ML}$ for $FRT_{LATERAL(P/NP)}$ (See Table 3). When comparing clinical Kids-BESTest scores with kinematic data, the TD group showed a low significant correlation for $FRT_{FORWARD}$ with increase in trunk-thigh angle but not for change in the angle of the knee or ankle. For the $FRT_{LATERAL(P/NP)}$ no significant correlations between the clinical score and trunk-thigh angle were seen.

The CP group showed moderate correlations between the clinical Kids-BESTest score and $CoP\ Range_{AP}$ for the $FRT_{FORWARD}$ and $CoP\ Range_{ML}$ for $FRT_{LATERAL(NP)}$ but not $FRT_{LATERAL(P)}$. For kinematic data when reaching forward children with CP showed moderate correlation between clinical scores and increase in trunk-thigh angle, similar to TD children however they also showed high correlation with increased ankle angle (plantar flexion). The CP group demonstrated a significant moderate correlation between the clinical Kids-BESTest score and increase in trunk-thigh angle for the $FRT_{LATERAL(NP)}$ but not $FRT_{LATERAL(P)}$.

Table 3 Association between clinical Kids-BESTest scores and laboratory data including (i) CoP data and (ii) Kinematic data for three Functional Reach Conditions (forward, lateral left and right) for children with CP and TD children. Statistics presented are Spearman's correlation (r_s) and p-value.

Laboratory data	Functional Reach Forward		Functional Reach Lateral (Preferred)		Functional Reach Lateral (Non-preferred)	
	TD	CP	TD	CP	TD	CP
Range _{AP}	0.38	0.68	0.32	-0.19	0.39	-0.09
[p-value]	[0.01]	[<0.01]	[0.04]	[0.47]	[0.01]	[0.75]
Range _{ML}	0.24	0.44	0.36	0.07	0.31	0.57
[p-value]	[0.13]	[0.08]	[0.02]	[0.78]	[<0.05]	[<0.05]
Change in						
Thigh-Trunk Angle	0.43	0.51	0.30	0.37	0.27	0.66
[p-value]	[<0.01]	[0.05]	[0.06]	[0.14]	[0.1]	[0.03]
Change in Knee	0.10	0.49				
Angle [p-value]	[0.53]	[0.07]				
Change in Ankle	0.20	0.75				
Angle [p-value]	[0.20]	[<0.01]				

CP and TD groups differences in FRT: distance reached, CoP and joint position

To analyse between group differences it is helpful to consider the tasks of FRT_{FORWARD} and FRT_{LATERAL} separately. Analysis for the FRT_{FORWARD} indicated that children in the CP group achieved significantly smaller mean reach distance (cm) compared to TD children (See Table 2). The CP group performing the FRT_{FORWARD} demonstrated significantly smaller CoP Range_{AP} and larger Range_{ML} than the TD children. For change in joint position, from start to the end of reaching, the CP group showed smaller increase in hip flexion, knee extension and ankle plantar flexion. However only the smaller increase in hip flexion was significant.

In the FRT_{FORWARD} children with CP used different balance and orientation strategies when compared to TD children. Children who were TD were able to sustain the starting position; to reach, they used a strategy of flexion of the hips with arm extension.

Forward movement of the trunk was counterbalanced by backwards movement of the pelvis, achieved by extension of the knees and plantar flexion at the ankles (88%, n=36, See Figure 3). Less children with CP used this counterbalancing strategy (71%, n=12). Children with CP used a variety of other strategies including an asymmetrical strategy or symmetrical flexion at the knee and/or at the ankle (See Figure 3).

Analysis for the $FRT_{LATERAL}$ indicated that the CP group achieved significantly less mean $FRT_{LATERAL(P/NP)}$ reach distance (cm) compared to TD children (See Table 2) Clinical measures and laboratory measures for children with CP (n=17) and typically developing children (n=41). However, their CoP Range_{ML} was not significantly less than TD children. Video observation showed that for change in joint position, from start to the end of reaching, the CP group showed significantly smaller change in trunk-thigh angle than the TD children.

In the $FRT_{LATERAL}$ children with TD and CP used different balance and orientation strategies however the current clinical score cannot distinguish this, particularly when reaching to the preferred side. Children TD attained the starting position, were able to weight shift to the reaching side and laterally flex at the trunk. Children with CP had reduced or no weight shift to the reaching side and either increased their CoP range or constrained it (See Figure 3).

		Examples of Atypical strategies			
		Hemiplegia: Asymmetrical	Hemiplegia: Symmetrical	Diplegia: Crouch	Diplegia: Aligned
FRT_{FORWARD}
Performance					
CoP Range _{AP}	90 mm	80 mm	121 mm	132 mm	78 mm
Kids-BESTest Item Score	3	2	2	3	1
Attains starting position		<i>Non preferred</i>	<i>Preferred</i>		
Shoulders at 90					
Trunk and lower limb aligned					
End reach position					
Hip Flexed		↓	↓	↓	
Knee into Extension		↓			
Ankle into Plantar flexion		↓		↓	
FRT_{LATERAL}
Performance					
CoP Range _{ML}	120 mm	133 mm	207 mm	177 mm	114 mm
Kids-BESTest Item Score	3	2	2	3	2
Attains starting position		<i>Non preferred</i>	<i>Preferred</i>		
Shoulder 90, full elbow extension					
Trunk and pelvis aligned					
End reach position					
Trunk lateral flexion			↓	↓	SL↓
Heel up on non-reaching side					SL↓
Weight shift to the reaching side		↓	↓		

Figure 3 Examples of typical versus atypical strategies for forward and lateral reach.

FRT, Functional Reach Test; CoP, centre of pressure; AP, anterior-posterior; ML, medial-lateral; Kids BESTest, Kids Balance Evaluation Systems Test; SL, slight.

Discussion

This study has produced significant new psychometric data for the FRT. Face validity, concurrent and content validity was confirmed for the FRT_{FORWARD} and partially confirmed for FRT_{LATERAL}. Face validity for FRT_{FORWARD} was demonstrated with children with CP scoring lower than TD children, confirming the Kids-BEST criteria for the FRT can identify deficits in stability limits with forward reach. For the FRT_{LATERAL} both the TD and CP groups scored lower on the Kids-BEST than for the FRT_{FORWARD} indicating that the FRT_{LATERAL} is more difficult postural control task for children. Face validity was confirmed for FRT_{LATERAL(NP)} but not FRT_{LATERAL(P)} indicating the Kids-BEST criteria for the lateral reach may not distinguish different strategies used by children with CP, particularly when reaching to the preferred side. Concurrent validity of the clinical measurement of reach with digital-derived measures of reach was good to excellent for both FRT_{FORWARD} and FRT_{LATERAL(P/NP)} confirming that the clinical measurement is an accurate measurement of reach distance. Content validity was confirmed for the FRT_{FORWARD} and FRT_{LATERAL(NP)} conditions. This was not the case for FRT_{LATERAL(P)}. This confirms that the Kids-BEST clinical criteria is a valid representation of ability to balance during reach but content analysis of kinematic data reveal that clinical criteria do not distinguish orientation or balance strategies used by children with CP, and therefore do not distinguish if a child can perform a functional task when reaching.

When performing the FRT_{FORWARD} task TD children in our study demonstrated an counterbalancing strategy by flexing the trunk at the hip, increasing knee extension and relative plantar flexion at the ankle to keep the CoM well within the BoS. Fewer children with CP used this strategy. There are several possible reasons for this. Children with spastic CP have musculoskeletal impairments that prevent backward translation of the knee and pelvis, such as: calf spasticity and/or contracture, poor eccentric muscle control and poor coordination of selective motor control between agonist and antagonist muscles for the knee and ankle [15, 16]. These impairments make using the same strategies as TD children difficult or impossible. Therefore, to reach forward some children with CP use a hip strategy to achieve trunk flexion without counterbalancing,

i.e. flexion rather than extension at the knee and/or DF rather than PF at the ankle (Figure 3). Use of the hip strategy during the forward reach test was subjectively observed in another study on children with bilateral CP [3]. In our study, when using this strategy, children with CP appear to bring their CoM closer to the edge of the BoS (into the toes), demonstrated as an increase in the CoP Range_{AP}. If the CoM is close to the edge of the BoS, balance may be easily lost at the end of reach and therefore this may affect a child's ability to perform a functional task when reaching.

Few studies have investigated medio-lateral stability and the use of the FRT_{LATERAL} in children [17]. Our study showed that children with CP scored lower than TD children on the Kids-BESTest, however scores did not necessarily distinguish the different postural strategy used by children with different motor distribution of CP i.e. unilateral or bilateral. Typically developing children demonstrated a pattern of weight shift onto the leg of the reaching side and then flexing the trunk laterally to maximize their reach. When reaching to the non-preferred side, children with either unilateral or bilateral CP demonstrated decreased weight shift and decreased trunk lateral flexion. Difficulties with weight shift in standing have been demonstrated for children with CP in other research [18], in particular reaching to the non-preferred side by children with unilateral CP [19]. In addition, children with unilateral or bilateral CP have both been shown to have poor trunk lateral flexion when leaning or reaching in sitting, especially children with bilateral CP [20]. In our study, when children reached to the preferred side, children with and without unilateral CP achieved similar Kids-BESTest clinical scores, as these were based on reach distance (i.e. score 2 or 3). However it appeared that children with unilateral CP still had some difficulty with achieving maximum weight shift and trunk lateral flexion compared to TD children. As expected children with bilateral CP had difficulty reaching to their preferred side, showing a range of compensatory strategies similar to those demonstrated for their non-preferred side (Figure 3). This supports the value of both quantitative and qualitative descriptors of performance for the FRT.

Given the outcomes of this study, we recommended that qualitative descriptors be added to the Kids-BESTest clinical criteria to more effectively describe postural

strategies used by children when reaching outside their base of support (Appendix 1). These descriptors indicate; (i) start position postural orientation; (ii) end position postural orientation and; (iii) postural balance or where the CoM is in relation to the limits of stability. The qualitative descriptors may give insight into the child's ability to maintain postural control (orientation and balance) [21]. For example two children may be able to reach a similar distance however at the limit of stability have poor postural orientation, as we observed in children with CP, which limits their capacity to maintain or sustain their postural balance (CoM over BoS) to achieve a functional task e.g. picking up an object. These descriptors also give direction for possible interventions e.g. during functional reach forward a child with bilateral CP may benefit from a triceps surae lengthening program to improve ankle range of motion and a strengthening program to improve eccentric control.

Strengths, limitations and future directions for research

This study is the first to investigate the validity of the Kids-BESTest clinical criteria for the FRT forward and lateral in children with and without CP. Statistical differences were identified between children with and without CP, but future research could involve samples to differentiate postural strategies used by children with different CP motor distributions, motor types and ages. This paper involved examination of postural strategies used at the limits of stability in standing. Future research is needed to study postural strategies used to control limits of stability in sitting, as well as dynamic functional tasks such as sit-to-stand and interventions to improve limits of stability.

Conclusion

This study supports the face, concurrent and content validity of the forward FRT. Concurrent validity was confirmed for lateral FRT however face and content validity were only confirmed when reaching to the non-preferred side in children with CP. To improve the ability to reliably describe impairments in forward and lateral reach, we recommend the addition of qualitative descriptors to the Kids-BESTest FRT criteria in addition to quantitative reach distance.

Supporting information

Appendix 1 Kids-BESTest - Item 7 Funtional Reach Test Forward and Item 8 Functional Reach Test Lateral (i) instructions and (ii) score sheet with quantitative score (based on reach distance) and additional qualitative descriptors

II. STABILITY LIMITS	
Examiner Instructions	Child Instructions
7. FUNCTIONAL REACH FORWARD	
A large sheet of paper is taped to a wall. The child holds a pen in their closed hand closest to the wall. A mark is made on the paper in the starting position, by the examiner. The child reaches forward. A second mark is made by the examiner at the end position. The child may not lift heels, rotate trunk, or protract scapula excessively. The child must keep their arms parallel to the floor and may use less involved arm. The horizontal distance between these two marks is then measured after two trials. Record best reach. Record quality of movement observations on score sheet.	Stand normally. Please lift both arms straight in front of you, holding this pen in your hand closest to the wall. I will make a mark on the paper then you will reach forward as far as you can. Don't lift your heels. Don't touch the paper or the wall. Once you've reached, as far forward as you can, I will make a mark on the paper. Then return to a normal standing position. I will ask you to do this two times. Reach as far as you can.
8. FUNCTIONAL REACH LATERAL	
A large sheet of paper is taped to a wall. The child holds a pen in their closed hand of the reach arm. A mark is made on the paper in the starting position, by the examiner. The child reaches. A second mark is made by the examiner at the end position. The horizontal distance between these two marks is then measured after two trials. Record the best reach. Make sure the subject starts in neutral. The patient is allowed to lift one heel off the floor but not the entire foot. Record quality of movement observations on score sheet.	Stand normally with your feet shoulder width apart. Arms at your sides. Hold this pen in your hand and lift your arm out to the side. Your arm should not touch the wall behind you. Reach out as far as you can. Do not lift your toes off the floor. Reach as far as you can. (REPEAT other side).

Item 7. FUNCTIONAL REACH FORWARD (Best of two trials)

Distance reached: _____ cm / in

Quantitative Score (Reach distance)	☐ 0 points No measurable lean – or must be caught.	☐ 1 point Poor < 16.5 cm (6.5 in).	☐ 2 points Moderate 16.5 - 32 cm (6.5–12.5 in).	☐ 3 points Maximum to limits: >32 cm (12.5 in).
--	--	---	---	--

Qualitative Scores	Start position	Reach position			
Upper limb (R) (L)	☐ Sh. flexion 90° ☐ Sh. flexion 90°	☐ Sh. flexion <50° ☐ Sh. flexion <50°	☐ Sh. flexion <70° ☐ Sh. flexion <70°	☐ Sh. flexion 90° ☐ Sh. flexion 90°	
Hip (R) (L)	☐ Hip neutral F/E ☐ Hip neutral F/E	☐ Hip flexion <45° ☐ Hip flexion <45°	☐ Hip flexion 70-45° ☐ Hip flexion 70-45°	☐ Hip flexion >70° ☐ Hip flexion >70°	
Knee (R) (L)	☐ Knee neutral F/E ☐ Knee neutral F/E	☐ >5° flexion ☐ >5° flexion	☐ >10° hyperextension	☐ 5 to -5° flexion	
Ankle (R) (L)	☐ Ankle neutral DF/PF ☐ Ankle neutral DF/PF	☐ Ankle in DF ☐ Ankle in DF	☐ Ankle neutral DF/PF ☐ Ankle neutral DF/PF	☐ Ankle in PF ☐ Ankle in PF	
Weight bearing	☐ Weight symmetrical	☐ Weight in toes	☐ Weight asymmetrical	☐ Weight in heels	

Comments:

Item 8a. FUNCTIONAL REACH LATERAL – Preferred (Best of two trials)

 Indicate reaching arm: ☐ Left ☐ Right

Distance reached _____ cm / in

Quantitative Score (Reach distance)	☐ 0 points No measurable lean, or must be caught.	☐ 1 point Poor: < 10 cm (4 in).	☐ 2 points Moderate: 10-25.5 cm (4-10 in).	☐ 3 points Maximum to limit: > 25.5 cm (10 in).
--	---	---	--	--

Qualitative Scores	Start position	Reach position			
Shoulder	☐ Sh. flexion 90°	☐ Sh. flexion <50°	☐ Sh. flexion <70°	☐ Sh. flexion 90°	
Elbow	☐ Elb. extended	☐ Elb. flexion ≥90°	☐ Elb. flexion 45-90°	☐ Elb. extended	
Trunk	☐ Trunk vertical	☐ LF <10°	☐ LF 10-20°	☐ LF > 20°	
Pelvis	☐ Pelvis neutral	☐ Pelvis rotated	☐ Pelvis sl. rotated	☐ Pelvis neutral	
Weight bearing	☐ Weight symmetrical on both feet	☐ Reach side heel up; Non-reach heel down	☐ Both heels down	☐ Reach side heel down Non-reach side heel up	
Weight shift	☐ Symmetrical	☐ No weight shift to reaching side	☐ Some weight shift to reaching side	☐ Max weight shift to reaching side	

Comments:

Item 8b. FUNCTIONAL REACH LATERAL – Non-Preferred (Best of two trials)

 Indicate reaching arm: ☐ Left ☐ Right

Distance reached _____ cm / in

Quantitative Score (Reach distance)	☐ 0 points No measurable lean, or must be caught.	☐ 1 point Poor: < 10 cm (4 in).	☐ 2 points Moderate: 10-25.5cm(4-10 in).	☐ 3 points Maximum to limit: > 25.5 cm (10 in).
--	---	---	--	--

Qualitative Scores	Start position	Reach position			
Shoulder	☐ Sh. flexion 90°	☐ Sh. flexion <50°	☐ Sh. flexion <70°	☐ Sh. flexion 90°	
Elbow	☐ Elb. extended	☐ Elb. flexion ≥90°	☐ Elb. flexion 45-90°	☐ Elb. extended	
Trunk	☐ Trunk vertical	☐ LF <10°	☐ LF 10-20°	☐ LF > 20°	
Pelvis	☐ Pelvis neutral	☐ Pelvis rotated	☐ Pelvis sl. rotated	☐ Pelvis neutral	
Weight bearing	☐ Weight symmetrical on both feet	☐ Reach side heel up; Non-reach heel down	☐ Both heels down	☐ Reach side heel down Non-reach side heel up	
Weight shift	☐ Symmetrical	☐ No weight shift to reaching side	☐ Some weight shift to reaching side	☐ Max weight shift to reaching side	

Comments:

(R), Right; (L), Left; Sh, Shoulder; F, Flexion; E, extension; DF, Dorsiflexion; PF, Plantarflexion; Elb, Elbow; LF, Lateral Flexion; Sl, slight.

References

- [1] Horak FB, Wrisley DM, Frank J. The balance evaluation systems test (BESTest) to differentiate balance deficits. *Phys Ther*. 2009;89:484-98.
- [2] Saether R, Helbostad JL, Riphagen II, Vik T. Clinical tools to assess balance in children and adults with cerebral palsy: a systematic review. *Dev Med Child Neurol*. 2013;55:988-99.
- [3] Niznik TM, Turner D, Worrell TW. Functional reach as a measurement of balance for children with lower extremity spasticity. *Phys Occup Ther Pediatr*. 1996;15:1-16.
- [4] Dewar R, Claus AP, Tucker K, Ware R, Johnston LM. Reproducibility of the Balance Evaluation Systems Test (BESTest) and the Mini-BESTest in school-aged children. *Gait Posture*. 2017;55:68-74.
- [5] Dewar R, Claus AP, Tucker K, Ware RS, Johnston LM. Reproducibility of the Kids-BESTest and the Kids-Mini-BESTest for Children With Cerebral Palsy. *Arch Phys Med Rehabil*. 2019.
- [6] Dewar R, Love S, Johnston LM. Exercise interventions improve postural control in children with cerebral palsy: a systematic review. *Dev Med Child Neurol*. 2015;57:504-20.
- [7] Zaino CA, McCoy SW. Reliability and comparison of electromyographic and kinetic measurements during a standing reach task in children with and without cerebral palsy. *Gait Posture*. 2008;27:128-37.
- [8] Liu WY, Zaino CA, McCoy SW. Anticipatory postural adjustments in children with cerebral palsy and children with typical development. *Pediatr Phys Ther*. 2007;19:188-95.
- [9] Van Der Heide JC, Fock JM, Otten B, Stremmelaar E, Van Eykern LA, Hadders-Algra M. Postural control during reaching in preterm children with cerebral palsy. *Dev Med Child Neurol*. 2004;46:253-66.
- [10] Van Der Heide JC, Hadders-Algra M. Postural muscle dyscoordination in children with cerebral palsy. *Neural Plast*. 2005;12:197-203.
- [11] Hadders-Algra M, Van Der Fits I, Stremmelaar EF, Touwen BCL. Development of postural adjustments during reaching in infants with CP. *Dev Med Child Neurol*. 1999;41:766-76.
- [12] Deitz JC, Kartin D, Kopp K. Review of the Bruininks-Oseretsky test of motor proficiency, (BOT-2). *Phys Occup Ther Pediatr*. 2007;27:87-102.
- [13] Palisano RJ, Rosenbaum P, Bartlett D, Livingston MH. Content validity of the expanded and revised Gross Motor Function Classification System. *Dev Med Child Neurol*. 2008;50:744-50.
- [14] Mukaka M. A guide to appropriate use of Correlation coefficient in medical research. *Malawi Med J*. 2012;24:69-71.
- [15] Rose J. Selective motor control in spastic cerebral palsy. *Dev Med Child Neurol*. 2009;51:578-9.
- [16] Barrett RS, Lichtwark GA. Gross muscle morphology and structure in spastic cerebral palsy: a systematic review. *Dev Med Child Neurol*. 2010;52:794-804.
- [17] Bartlett D, Birmingham T. Validity and reliability of a pediatric reach test. *Pediatr Phys Ther*. 2003;15:84-92.
- [18] Liao HF, Mao PJ, Hwang AW. Test-retest reliability of balance tests in children with cerebral palsy. *Dev Med Child Neurol*. 2001;43:180-6.
- [19] Ledebt A, Becher J, Kapper J, Rozendaal R, Bakker R, Leenders IC, et al. Balance training with visual feedback in children with hemiplegic cerebral palsy: effect on stance and gait. *Motor Control*. 2005;9:459.
- [20] Heyrman L, Desloovere K, Molenaers G, Verheyden G, Klingels K, Monbaliu E, et al. Clinical characteristics of impaired trunk control in children with spastic cerebral palsy. *Res Dev Disabil*. 2013;34:327-34.
- [21] Dewar R, Claus AP, Tucker K, Johnston LM. Perspectives on Postural Control Dysfunction to Inform Future Research: A Delphi Study for Children With Cerebral Palsy. *Arch Phys Med Rehabil*. 2017;98:463-79.