Global Biogeochemical Cycles

RESEARCH ARTICLE

10.1029/2020GB006816

Key Points:

- We developed and evaluated a wetland biogeochemical model (BRTSim-BAMS4) to quantify global wetland CO₂, CH₄, and N₂O emissions and nutrient sequestration rates
- Modeling indicates that agricultural N fertilization and runoff increase N₂O emissions by 13% in wetlands in the Northern Hemisphere
- At equal amount of C and N sequestered, fertilization leads to greater C and N gaseous emissions, while no differences were found for S sequestration

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

C. Pasut,
c.pasut@sydney.edu.au

Citation:

Pasut, C., Tang, F. H. M., Hamilton, D., Riley, W. J., & Maggi, F. (2021). Spatiotemporal assessment of GHG emissions and nutrient sequestration linked to agronutrient runoff in global wetlands. *Global Biogeochemical Cycles*, 35, e2020GB006816. <https://doi.org/10.1029/2020GB006816>

Received 1 SEP 2020
Accepted 24 FEB 2021

Spatiotemporal Assessment of GHG Emissions and Nutrient Sequestration Linked to Agronutrient Runoff in Global Wetlands

Chiara Pasut¹ , Fiona H. M. Tang¹ , David Hamilton² , William J. Riley³ , and Federico Maggi¹

¹Laboratory for Advanced Environmental Engineering Research, School of Civil Engineering, The University of Sydney, Sydney, NSW, Australia, ²Australian Rivers Institute, Griffith University, Nathan Campus, Nathan, QLD, Australia,

³Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

Abstract Wetlands play a key role in regulating global greenhouse gas (GHG) emissions but anthropogenic impacts on nutrients may severely alter this balance. Recent assessments indicate that almost 22% of the global wetland area may be affected by agricultural runoff. In this work, we developed and applied a dynamic mechanistic reaction network model of soil organic matter linking the carbon (C), nitrogen (N), and sulfur (S) cycles at 0.5° × 0.5° spatial resolution across the globe. The model was used to estimate GHG emissions and nutrient sequestration rates in wetlands, driven by environmental stressors including N, P, and S fertilization. Wetland annual GHG emissions are estimated to be 136 ± 12.5 Tg C-CH₄, 589 ± 45.8 Tg C-CO₂, and 0.3 ± 0.04 Tg N-N₂O; in contrast, C, N, and S annual sequestration rates are estimated to be 576 ± 88.1 Tg C, 20 ± 4.4 Tg N, and 7.4 ± 0.8 Tg S, between 2000 and 2017. N fertilization inputs were responsible for 13% N₂O emissions in wetlands in the Northern Hemisphere, while tropical wetlands were major reservoirs for C, N, and S. Temperature, net primary productivity, and methanogenic microorganisms exert the major control on GHG emissions. Wetland CH₄ and CO₂ emissions were found to have a hysteretic relationship with seasonal soil temperature, but not N₂O. A global-scale assessment is pivotal for best nutrient management practices, reducing nutrient losses, and controlling gas emissions.

1. Introduction

Significant and rapid changes in nutrient turnover and budget have occurred over the past century (Li & Bowman, 2001; Vitousek et al., 1997). Temperature, precipitation, soil weathering, microbial activity, and nutrient runoff control carbon (C), nitrogen (N), and sulfur (S) availability and their turnover rates in wetlands (Matias et al., 2011; Whitehead, 2000). However, these dynamics have largely been overlooked in global-scale assessments, even though wetlands contribute globally between 25% and 30% of CH₄ emissions (Saunio et al., 2020), store almost 30% of the organic C (Lal, 2008), and have higher C soil content than forests and grasslands (Kayranli et al., 2010), while occupying only 7% of the Earth's surface (Lehner & Döll, 2004).

Unprecedented high temperatures in the northern latitudes have accelerated permafrost melting (Treat et al., 2016), leaving an unfrozen, active peat layer that remobilizes large amounts of C and other nutrients into wetlands, and change their hydrological and biogeochemical functioning (Malone et al., 2013) resulting in higher N₂O, CH₄, and CO₂ emissions (Abbott & Jones, 2015; Loiko et al., 2017). At lower latitudes, significant cropland expansion to meet the high global food and fiber demand has increased nonpoint nutrient pollution linked to fertilizer applications (Wagner et al., 2008). In addition, intense hydroclimatic fluctuations (i.e., prolonged droughts followed by sudden and severe floods) have affected watershed runoff resulting in nutrient-rich stormflow pulses into wetlands (Kaushal et al., 2014; Solomon, 2007) usually located at low elevation within the catchment (Were et al., 2019).

Changes in nutrient loads, whether anthropogenic or not, can interfere with the preexisting biogeochemical cycles within wetlands and significantly affect the gas emission quality and soil nutrient stocks. Anomalous concentrations of N or S can have inhibitory effects on the C cycle (Kaushal et al., 2010, 2011), such as S fertilization inhibiting CH₄ emissions due to energetic preference of SO₄²⁻ reduction over CH₄ genesis

(Bronson et al., 1997; Gauci et al., 2004). A high rate of N fertilization increases N_2O emissions (Bonnnett et al., 2013; Burgin & Groffman, 2012). However, a different response may occur in N-poor soil where an increase in N input may increase net primary productivity (NPP) and bioavailable substrates, in turn increasing CH_4 emissions (Dise & Verry, 2001). Over the long term, increases in NPP may also increase plant nutrient uptake, hence less N may be available to soil denitrifying bacteria (Mosier et al., 2002). Changes in nutrient inputs also affect microbial ecology, hence the nutrient turnover and storage. Excess nutrient inputs can increase rates of soil organic matter (SOM) decomposition and cause a shift toward wetlands as a net C source rather than a sink, and warming can also synergistically amplify this effect (Kaushal et al., 2014).

Many recent studies (Arora et al., 2018; Bloom et al., 2017; Hayman et al., 2014; L. Liu et al., 2020; Melton et al., 2013; Riley et al., 2011; Zhu et al., 2015) have investigated wetlands biogeochemistry coupled to hydroclimatic forcing, vegetation dynamics, soil properties, and soil biology to improve the CH_4 budget assessment at regional and global scales. The Wetland CH_4 Intercomparison of Model Projects (WETCHIMP; Melton et al., 2013; Wania et al., 2013) analyzed and compared the most advanced models available for global CH_4 emissions using a common protocol for hydroclimatic variables next improved in Saunio et al. (2020). These projects can be considered the state-of-the-art in wetland modeling and highlight the strengths and weaknesses of current models. Besides the level of complexity used to describe the C cycle and climate feedbacks in each model, only a relatively small number of models incorporate multiple nutrient cycles, although they usually limit their descriptions to C and N (Saunio et al., 2020). Among those, Community Land Model 4 (CLM4Me; Riley et al., 2011), Dynamic land Ecosystem Model (DLEM; Tian et al., 2015; Tian, Melillo, et al., 2011; Tian, Xu, et al., 2011; Xu et al., 2010), *ecosys* (Chang et al., 2019; Grant, Mekonnen, & Riley, 2019; Grant, Mekonnen, Riley, Arora, et al., 2019), and Sheffield Dynamic Global Vegetation Model (SDGVM; Beerling et al., 2011; Woodward et al., 1995) combine a global dynamic vegetation module, soil hydroclimatic changes, permafrost dynamics, land cover change, and agriculture management (e.g., fertilization, irrigation, and harvesting). However, each model solves each geographic grid cell independently; hence, no water runoff is explicitly accounted for (Wania et al., 2013), and only a very simplified version of the N cycle is represented. Therefore, the combined effects of global land use linked to hydroclimatic forces on global greenhouse gas (GHG) emissions and nutrient stocks are not fully described, resulting in model uncertainties. Addressing grid cell connectivity and feedback and the interrelationships between biologically mediated processes governing not only C and N but also S and P cycling will improve the understanding of watershed features.

The aim of this study was to investigate the relationship between multiple nutrient interactions (i.e., C, N, S, and P) and environmental forcing factors for wetland GHG emissions. We used a mechanistic nutrient biogeochemical model, Biotic and Abiotic Model for SOM-version 4 (BAMS4), in a dynamic solver, BRT-Sim, to describe in space and time the coupled ecohydrological and biogeochemical cycles over a global grid of $0.5^\circ \times 0.5^\circ$ resolution during the period 2000–2017. The mechanistic description of the wetland was coupled with a simplified runoff model that estimates agricultural runoff from the surrounding catchment to account for N, P, and S fertilizer inputs. Using BRTSim-BAMS4 we (1) report new estimates of wetland CH_4 , N_2O , and CO_2 emissions globally; (2) investigate and quantify nutrient sequestration rates in wetlands at different latitudes; (3) quantify the environmental factors that influence GHG emissions and nutrient sequestration; and (4) examine fertilization effects on GHG emission dynamics.

2. Materials and Methods

2.1. C–N–S Reaction Network (BAMS4)

The BAMS4 reaction network is a simplification of the SOM cycle described in Pasut et al. (2020) and elaborated from Riley et al. (2014), Tang et al. (2019), and Ceriotti et al. (2020). It consists of two organic polymer pools of PolyC (lignin and cellulose) and PolyCN (peptidoglycan), three organic monomer pools of MonoC, MonoCN, and MonoCS, representing the organic carbon, nitrogen, and sulfur, respectively, and two inorganic carbon molecules (CH_4 and CO_2). The C cycle includes four microbial functional groups, three aerobic (fungi F_{DEP} , heterotrophic bacteria B_{AER} , and methanotrophs B_{MOB}) and one anaerobic (methanogenic microbes B_{MGB}), which control the depolymerization, mineralization, CH_4 oxidation, and CH_4 genesis, respectively. The C cycle was coupled with the N and S cycles, which include nitrification, denitrification, nitrogen

fixation, S pools reduction, S pools oxidation, and thiosulfate and sulfur trioxide disproportionation ($\text{NH}_3 \rightarrow \text{NH}_4^+ \rightarrow \text{NO}_2^- \rightarrow \text{NO}_3^- \rightarrow \text{NO} \rightarrow \text{N}_2\text{O} \rightarrow \text{N}_2 \rightarrow \text{NH}_4^+$, and $\text{SO}_4^{2-} \rightarrow \text{SO}_3^{2-} \rightarrow \text{S}_2\text{O}_4^{2-} \rightarrow \text{S}_0 \rightarrow \text{HS}^-$, respectively). NH_3 gas emissions can be large in agricultural system but typically very small in wetlands because the equilibrium reaction between NH_4^+ and $\text{NH}_3(\text{g})$ is strongly directed toward NH_4^+ , resulting in very small wetland $\text{NH}_3(\text{g})$ exsolution and emission to the atmosphere, especially for low NH_4^+ concentration usually found in wetland (e.g., 10^{-5} or 10^{-6}). Each C, N, and S pool can be present in aqueous, protected (e.g., on the mineral surface binding), or gaseous phase. Each reaction includes specific microbial functional groups (ammonia-oxidizing bacteria B_{AOB} , nitrite-oxidizing bacteria B_{NOB} , denitrifying bacteria B_{DEN} , sulfur reducing bacteria B_{SRB} , thiosulfate- and sulfide-reducing bacteria B_{ThSRB} , sulfate-reducing bacteria B_{SRB} , thiosulfate and sulfide disproportioning bacteria B_{SDB} , and photolithoautotroph oxidizing bacteria B_{SOB}) described in detail in Pasut et al. (2020) and represented in Figure 1. The first two functional groups are aerobic, while the others are anaerobic. BAMS4 also represents a simplified P cycle that includes PO_4^{3-} production from MonoCN mineralization, PO_4^{3-} protection, and PO_4^{3-} plant uptake (R4, R55, and R32).

The microbial dynamics include C and N immobilization, growth, mortality, and necromass decomposition described by Michaelis–Menten–Monod kinetics. Less than 3% of microbial biomass is composed by P and S (Paul, 2014); therefore, their immobilization was neglected. The biological response is controlled by temperature, pH, water stress (Figure S1), O_2 availability, and other stressors related to the competition for electron donors and acceptors, and inhibitory factors. SOM protection on the soil matrix is described as a kinetic reaction after Riley et al. (2014), while the protection of inorganic species is described at equilibrium. Plant nutrient uptake (NH_4^+ , NO_3^- , SO_4^{2-} , and PO_4^{3-}) is accounted for by MM kinetics, and CH_4 is transported to the atmosphere by diffusion and plant uptake.

We used the NPP from the Moderate Resolution Imaging Spectroradiometer (MODIS) on NASA's Terra satellite (Imagery produced by the NASA Earth Observations team based on data provided by the MODIS Land Science Team) as a proxy of SOM input to the soil, and we used the C:N and C:S ratios of above and belowground litter quality to partition C, N, and S in soil (Table S7).

Tables S1–S6 summarize all the reactions and corresponding parameters for both biotic and abiotic processes used in BAMS4, which were derived from earlier estimation and validation (Maggi et al., 2008; Pasut et al., 2020; Riley et al., 2014) and are not expanded here.

2.2. Data Description

This study used georeferenced databases of dynamic and constant environmental variables at $0.5^\circ \times 0.5^\circ$ resolution globally. The dynamic quantities consist of monthly fractional wetland area (SWAMPS v3.2; Poulter et al., 2017; Schroeder et al., 2015), NPP (MODIS Land Science Team, 2019), evapotranspiration (Zhang et al., 2016), runoff, the 6-h Climatic Research Unit (CRU-TS 3.21) time series of precipitation, longwave and shortwave solar radiation, and atmospheric temperature, which were used to construct daily input data from 2000 to 2017. The constant quantities consist of the soil physical properties (i.e., porosity, bulk density, and soil texture; Hengl et al., 2017), soil hydrothermal properties (i.e., Brooks and Corey model parameters [Brooks & Corey, 1964], permeability, heat capacity, and conductivity; Dai et al., 2013), equilibrium soil organic C (Hengl et al., 2017), land cover (MODIS – LC; Cover & Change, 1999; Friedl & Sulla-Menashe, 2015; Sulla-Menashe et al., 2019), N and P fertilization (Potter et al., 2011a, 2011b), and biogeochemical parameters. Kinetic parameters were not geographically distributed. Additionally, we created a global data set of S fertilization and four land cover-specific (grassland, forest, wetland, and tundra) maximum and average plant root density profiles (based on Canadell et al., 1996; Paul, 2014), methane plant efficiency emissions (based on Walter et al., 2001), nitrogen fixation rate (based on Paul, 2014), and C:N ratio of aboveground litter quality (based on Bréchet et al., 2017; Hättenschwiler & Jørgensen, 2010; Paul, 2014; Pei et al., 2019; Rouifed et al., 2010; Snowdon et al., 2005), which are distributed as described in the Data Availability Statement and listed in Table S7.

The agricultural area surrounding the wetland was derived from the MODIS-IGBP land cover product (MODIS – LC; Cover & Change, 1999; Friedl & Sulla-Menashe, 2015; Sulla-Menashe et al., 2019). We calculated the long-term average area of both wetland and cropland from 2000 to 2017 and overlapped them

Figure 1. BAMS4 reaction network. RX identifies the reactions, which are expanded in Tables S1–S6 and are characterized by parameters earlier estimated and validated in Pasut et al. (2020). BAMS4, Biotic and Abiotic Model for SOM-version 4.

to identify the grid cells affected by agriculture, which were tagged across the entire simulation. The watershed discharge Q was estimated following the Runoff Curve Number model (NRCS-CN) as Hong and Adler (2008):

$$Q = \frac{(P - I_a)^2}{P - I_a + S}, \quad (1)$$

$$S = \frac{25,400}{CN} - 254, \quad (2)$$

where P is the precipitation, I_a is the initial abstraction ($I_a = 0.2S$), S is the potential retention, and CN is the curve number estimated from hydrologic soil group characteristics (Figure S9), the class of land use, and

the antecedent soil moisture condition (see supporting information and summary in Table S8). The flow direction was derived from the steepest down-slope gradient of a digital elevation map (Data Announcement 88-MGG-02, Digital relief of the Surface of the Earth. NOAA, National Geophysical Data Center, Boulder, CO, 1988). The nutrient concentration in the runoff was calculated as 30% of the total fertilizer inputs to account for crop nutrient uptake. We did not explicitly model crop dynamics in this work.

The details and reference for each database used in our modeling are summarized in Table S9.

2.3. Computational Domain

The computational domain involves about 25,000 grid cells describing wetlands, and neglects lakes, rivers, reservoirs, rice paddies (due to the lack of globally distributed data on the agricultural management), and saline estuaries and salt marshes (due to the high salt concentration that inhibits microbial metabolism, Poulter et al., 2017). From the original SWAMPS database (Poulter et al., 2017), we selected the grid cells where the maximum average wetland area fraction is greater than 5% to reduce computational effort and eliminate potential ephemeral wetlands (Figures S2a–S2c), and where the permafrost is less than 80% of the ground area (Brown et al., 2001) due to the lack of permafrost thawing modeling. The selected grid cells cover a wetland area of almost 5 million km²; which almost 22% is affected by agricultural runoff (Figure S2d). Each grid cell included a 2-m vertical profile divided over three soil layers of rootzone (30, 30, and 40 cm thickness), a soil layer of 100 cm below the rootzone, and four atmospheric layers (30, 30, 40, and 100 cm thickness) to allow for water ponding, and heat and gas exchange with the atmosphere.

Each grid cell was initialized with SOM concentrations estimated in the SoilGrids (Menne et al., 2012) and run for 250 years, relooping the same dynamic quantities of the 2000–2017 period to reach a statistical steady state in the biogeochemical reactions. The last 18 years were used for our analyses.

2.4. BRTSim Solver

BAMS4 was integrated in a general-purpose multiphase and multispecies bioreactive transport simulator, BRTSim-v4.0e solver, for the aqueous species advection and diffusion, gas diffusion, SOM protection in the soil matrix, and gas exsolution/dissolution, together with variably saturated soil and heat flow (Maggi, 2019). The vertical profile of each chemical and biological species in the soil was not imposed, neither over time nor over space, but emerged from multiple biologically mediated reactions in BAMS4. In this way, the system can respond to sudden environmental changes, particularly important for pH and O₂ availability.

2.5. Methods of Validation

The simplified BAMS4 reaction network proposed here was first tested against the full-size BAMS3 network applied to a peatland in Michigan (Shannon & White, 1994) to ensure equivalent outputs. Next, the monthly and long-term annual average land surface temperature of the topsoil (top 30 cm) was assessed against the land surface temperature database of NOAA (top 5 cm; Menne et al., 2012). The long-term average of modeled pH and SOC in the topsoil was assessed against the values in SoilGrids (Hengl et al., 2017). Modeled CH₄ fluxes were benchmarked against the annual average field observations from 12 sites globally distributed (Table S10 data from FLUXNET2015 data set Pastorello et al. [2020]) and data for the Amazon catchment reported in Pangala et al. (2017). The global average annual CH₄ emission was also compared to global modeling assessments (L. Liu et al., 2020) and top-down and bottom-up estimation from the Global Methane Budget 2000–2017 project related to wetland modeling in Saunio et al. (2020) (Table S10 and Figure S10). The N₂O global emission was benchmarked against the DLEM global N₂O estimation from terrestrial ecosystems in Tian et al. (2015). Average carbon sequestration rates were benchmarked against literature values in Mitsch et al. (2013), Villa and Bernal (2018), and Cheng et al. (2020).

2.6. Analyses

Among all the outputs of BRTSim-BAMS4, the C, N, and S sequestration, and CH₄, CO₂, and N₂O gas emissions were selected as target state variables in the analyses between 2000 and 2017. Emission rates were calculated as the variation of CH₄, CO₂, and N₂O gaseous concentration across the four atmospheric nodes within a specific time interval. CO₂ emissions include the CO₂ produced during aerobic and anaerobic respiration, autotrophic CH₄ oxidation, denitrification, S pool reductions, and CH₄ oxidation during aerenchyma transport. C, N, and S sequestration rates were calculated as the total input (the above and belowground SOM input, N fixation, and N and S fertilizations) minus all the losses (gas emission, plant uptake of N and S, and leaching below the root zone). Data are presented as average value $\pm$ the standard deviation σ ($\sigma = \sqrt{\sum(x_i - \mu)^2 / n}$, where x_i is a state variable, μ is its average, and n is the sample size). Linear correlation coefficients were used to evaluate potential relationships between each GHG emission rate and wetland surface area. The lag time between the spike in gas emissions and the maximum area was calculated using cross-correlation analysis.

We grouped the wetlands according to the Köppen–Geiger (KG) climate classification (Köppen & Geiger, 1930) to carry out a statistical analysis of the controlling stressors on GHG emissions and C, N, and S sequestration rates, and a dynamic analysis of the effect of fertilizers on GHG emissions and several controlling factors (temperature, SOM input, B_{MGB} , and B_{DEN}). For the first analysis, the Pearson coefficient was used to study the correlation of the forcing boundary conditions (S, N, and P fertilization, N fixation, the difference between precipitation and evapotranspiration, and SOM input), microbial functional group concentrations, soil variables (SOC, temperature, and pH), and soil properties (porosity, hydraulic conductivity, root depth, soil bulk density, and CH₄ plant uptake efficiency) against each target output for each climatic class. In this analysis, we accounted for the annual average of each variable between 2000 and 2017. The most relevant parameters resulting from this first stage were then analyzed seasonally by, comparing results with and without fertilizer input, and using the monthly average within each KG climatic class.

3. Results

3.1. Model Validation

Figure S3 shows that the simplified BAMS4 accurately capture the response of the full BAMS3 network for total C (TC), total N (TN), total S (TS) in soil, and the annual gas emissions of CH₄, N₂O, CO₂, and H₂S ($R = 0.995$, $p \leq 0.001$). In addition, BAMS4 captured the monthly and long-term annual average land surface temperature data in Menne et al. (2012) (Figure S8a, $R = 0.98$). The latter database reports only the first 5 cm of the topsoil, while our model refers to the first 30 cm, which explains part of the mismatch. The benchmarking of pH and SOC in the topsoil against SoilGrids was satisfactory (Figures S8b and S8c, $R = 0.72$ and $R = 0.99$, respectively). Even though a background H⁺ recovery reaction was included in BAMS4 to act as a buffer (R57–R58 in Table S6), the pH was overestimated in some range of expected values. The reason can be ascribed to nonlinear effects on H⁺ production and consumption between the aerobic and anaerobic transition phases, and specific geochemistry of the soil, which are challenging to capture.

Benchmarking of CH₄ emissions against field observations and the simulated global annual emission resulted in a relatively good match (scatter-plot in Figure 2a, Tables S10 and S11, and Figure S10, respectively). The simulated value of the Amazon catchment matched the top-down regional estimate (32.0 ± 4.2 Tg C-CH₄ observed against the 29.3 ± 1.6 Tg C-CH₄ simulated). Field measurements may include uncertainty, such as temporal gaps in the recording and limited numbers of years of data, hence hindering a direct comparison with the model output. Regional-scale measurements are not physically possible (Mekonnen et al., 2016), thus other global assessments from process-based and top-down models provide support to benchmark model outcomes. Our N₂O global annual emission (0.3 ± 0.04 Tg N-N₂O) is lower than the value of Tian et al. (2015) (0.95 ± 0.04 Tg N-N₂O). This discrepancy may result from differences in the modeled N cycle; DLEM in Tian et al. (2015) represents a simplified version of the N cycles using an empirical equation to partition N gases into N₂O, NO, and N₂ as a function of porosity and volumetric water content, whereas BRTSim-BAMS4 attempts to account for other environmental factors, such as an explicit microbial dynamics, O₂, and pH.

Figure 2. Global averages and isolatitude sums emission (at 0.5° latitude) of (a and b) CH_4 , (c and d) CO_2 , and (e and f) N_2O emissions. In (a, c, and d), the average emission is normalized by the average wetland area. Light-gray shaded areas in (b, d, and f) represent the standard deviations of the isolatitude sums between 2000 and 2017. Panel (f) compares the assessments with and without the effect of fertilizer runoff (no significant effect on CH_4 and CO_2 emissions—data not shown). Inset in (a) shows the simulated CH_4 emissions against 12 field observations.

Figure 3. Global monthly average wetland area (black line in a–c), (a) CH_4 , (b) CO_2 , and (c) N_2O emissions between 2008 and 2017. Panel (c) compares the assessments with and without the effect of fertilizer input (blue line).

The reader can find the benchmark for temperature, pH, and SOC (Figures S8a–S8c), the corresponding anomaly maps (Figures S4, S6, and S7), as well as benchmarking samples for the seasonal temperature in 20 random grid cells over time (Figure S5).

3.2. CH_4 , CO_2 , and N_2O Emissions

The estimated annual global average CH_4 , CO_2 , and N_2O emissions were 135.6 ± 12.5 Tg C- CH_4 , 589.1 ± 45.8 Tg C- CO_2 , and 0.3 ± 0.04 Tg N- N_2O , respectively, between 2000 and 2017. A significantly high correlation exists between global monthly wetland CH_4 and CO_2 emissions ($R = 0.98$ and $p \leq 0.01$), while no correlation links CH_4 and N_2O emissions ($R = -0.08$ and $p \geq 0.01$), indicating that different environmental factors influence the N_2O emissions (Figure 3). In the long term, all GHG emissions and the wetland area show a seasonal pattern (Figures 3a–3c) that follows the temperature change in the Northern Hemisphere, which causes the most significant seasonal changes in wetland area due to snow melting and runoff (Figures S2a–S2c). CH_4 and CO_2 peak between June and August, approximately 4–6 weeks after the peak in wetland area, suggesting a lag time between the optimal biomass response and the rise in water table and temperature. In contrast, the spike in N_2O emissions occurs about 6 weeks before the peak in area, due to a more prompt response of denitrifying bacteria to temperature (optimum between 13°C and 37°C rather than 20°C and 37°C of B_{MGB}), a faster reaction rate, and less O_2 inhibition for B_{DEN} .

Spatially, the tropical areas in the Amazon, central Africa, and southeast Asia were the principal hot-spots of CH_4 and CO_2 emissions annually releasing between 100 and 150 g C- CH_4 m^{-2} of wetland (Figures 2a and 2c) and 350 and 750 g C- CO_2 m^{-2} (Figures 2b and 2d), respectively. The northern latitude regions (e.g., Canada, Alaska, and north Russia) show the second highest emissions due to the prevalence of summer wetlands (Figures S2a–S2c) and the high SOM input from the peatland and bog vegetation, despite the low winter temperature. In contrast, southern latitudes (e.g., south Australia, Patagonia), Sub-Sahara, and

central Europe show significantly lower emissions, on average 1–40 g C-CH₄ m⁻² year⁻¹, and 150–350 g C-CO₂ m⁻² year⁻¹, respectively.

N₂O emissions follow a spatial distribution directly linked to N fertilization; hence, N₂O gases are greater for those areas where wetlands and croplands are in proximity, mostly in the north temperate regions and north-east India (yellow grid cells in Figure S1d). Here, the annual average emission is between 0.2 and 0.5 g N-N₂O m⁻² year⁻¹, which contributes to the spike in emissions for a given latitude range. Tropical areas have relatively high emissions as well, on average between 0.12 and 0.2 g N-N₂O m⁻² year⁻¹, and there is a close relationship between SOM input and N₂O emissions for areas not affected by N fertilization. In contrast, lower emissions occur in wetlands in Australia and Sub-Saharan regions, which are characterized by sparse vegetation and negligible runoff from agriculture, releasing ≤0.05 g N-N₂O m⁻² year⁻¹.

A simulation conducted without fertilizer input (blue line, Figure 2f) shows that N₂O emissions in the Southern Hemisphere are not significantly affected by N fertilization, while the overall emission is almost 13% higher than with fertilization runoff between 60°N and 10°N. Comparing the two simulations (Figure 3c), N₂O emissions without fertilization follow the same seasonality as those with fertilization; however, the peak is significantly lower.

3.3. Nutrient Sequestration Rates

Globally, the average C, N, and S sequestration rates are 120.5 ± 122.5 g C m⁻² of wetland year⁻¹, 3.75 ± 5.6 g N m⁻² year⁻¹, and 0.98 ± 2.05 g S m⁻² year⁻¹. The tropics have the highest sequestration rates of C, N, and S; on average, 273.6 g C m⁻² year⁻¹, 11.4 g N m⁻² year⁻¹, and 4.2 g S m⁻² year⁻¹ (Figures 4a, 4d, and 4g and Table 1). Carbon sequestration was also very high in peatland regions in northern latitudes (continental class [Dfc]), ranging between 200 and 400 g C m⁻² of wetland year⁻¹. Negative C sequestration rates (losses from soil) are observed mainly in the Sub-Sahara Desert, while negative N sequestration rates are found in northern Tibet. In contrast, negative S sequestration rates are globally distributed without a clear pattern. We further compared each nutrient sequestration rate with the corresponding total gas emissions, where each square in Figures 4c, 4f, and 4i represents the bin average of 10 g C m⁻² year⁻¹, 1 g N m⁻² year⁻¹, and 1 g S m⁻² year⁻¹ increments. We found that an increase in C sequestration corresponds to an increase in C emission (the sum of CH₄ and CO₂), up to a maximum of 350 g C m⁻² year⁻¹, after which the emissions slightly decrease and stabilize between 700 and 400 g C m⁻² year⁻¹. Wetlands with high sequestration to emission ratios are located in the Southern Hemisphere (Amazon catchment, Malaysia, Indonesia, east Australia, and New Zealand) and are characterized by high SOM input and pH ≤ 6, which slows anaerobic reaction rates. We found the same patterns between N sequestration and total N emissions (corresponding to the sum of NO, N₂O, and N₂), but the N emissions decrease once reached the maximum. No clear trend between S sequestration and emissions (H₂S) has been found. Comparing the wetlands affected by agricultural inputs and those without, we found that the emission of C and N is greater in those with fertilization for the same amount of C and N sequestered; in contrast, the two values almost coincide for S sequestration.

The total long-term average of sequestered C, N, and S is 575 Tg C year⁻¹, 20 Tg N year⁻¹, and 7.4 Tg S year⁻¹, respectively. In terms of sum per degree latitude (Figures 4b, 4e, and 4h), there are two peaks: the first across 60°N and the second in the tropical area. The total value for carbon sequestration is lower than the one found in Mitsch et al. (2013), however, that value is an estimation from 21 wetlands distributed globally.

3.4. Controlling Variables

We analyzed the dominant variables that affect each GHG emission and nutrient sequestration rate, and we investigated whether these effects are associated with the KG climatic class. Figures 5 and 6 report the Pearson correlation coefficient between pairs of variables sorted from the positive (red) to the negative (blue) correlation.

The SOM input in the system (label SOM input) has the highest correlation with all the three GHG emissions and the forcing variables. This correlation also helps to explain why the N₂O emissions are high in northeast and southeast of the USA, despite N fertilization ranges between 0.1 and 5 kg N ha⁻¹. Analyzing the aggregated KG classes, the net recharge (label P-ETA) has a positive but weak correlation with GHG.

Figure 4. (a) Average annual carbon (C) sequestration, (b) average C total per 0.5° latitude sequestration rate, and (c) average C sequestration against total C emissions (CH_4 and CO_2); (d) average annual nitrogen (N) sequestration, (e) average N total per 0.5° latitude sequestration rate, and (f) average annual N sequestration against total N emissions (NO , N_2O , and N_2); and (g) average annual sulfur (S) sequestration, (h) average S total per latitude 0.5° sequestration rate, and (i) average annual S sequestration S emissions (H_2S). Each point represents a grid cell, while each square represents the bin average of a step of $10 \text{ g C m}^{-2} \text{ year}^{-1}$, $1 \text{ g N m}^{-2} \text{ year}^{-1}$, and $\text{g S m}^{-2} \text{ year}^{-1}$.

However, a strong correlation was found in the continental regions (Dwa/Dwb/Dwc/Dwd/Dfa/Dfb classes), which tends to have a variable precipitation regime. N, P, and S fertilizations (labeled as N, P, and S Fert.) have weak correlations with each GHG emissions, either positive or negative depending on the target variable. N and S fertilizations have a weak and negative correlation with CH_4 , likely because the amended N and S increase the substrate competition and inhibition, except in the arid (Bsk and Bwh) and continental (Dfa) wetlands, where we found a positive correlation. In general, these are nutrient-poor environments, meaning that additional N input helps to sustain the microbial activity. N_2O emission and N and S fertilization are generally positively correlated but are weakly negatively correlated in the temperate regions (Cs/Cwa/Cwb/Cwc/Cfa/Cfb/Cfc).

Wetland GHG emissions are significantly correlated with microbial diversity. We found that B_{MGB} is positively correlated not only with CH_4 and CO_2 emissions, which are the direct products of the reaction R6

Table 1
Annual Average Nutrient Sequestration Rates for Different Climatic Classes (Value $\pm \sigma$ of g C/N/S m⁻² Year⁻¹)

Variable	Tropical	Arid	Temperate	Continental	Polar	Global	Reference
C seq	273.6 $\pm$ 169.0	31.7 $\pm$ 26.1	132.4 $\pm$ 112.0	113.1 $\pm$ 63.6	26.9 $\pm$ 24.2	120.37 $\pm$ 122.5	This work
C seq ^a	55.8–168.5	39–85.3	55.2–120	–	15.6–137.1	–	Villa and Bernal (2018)
C seq ^b	–	–	12–1,180	–	2.96–407	185.2	Cheng et al. (2020)
N seq	11.4 $\pm$ 7.8	0.3 $\pm$ 0.9	4.0 $\pm$ 6.1	2.9 $\pm$ 2.1	0.5 $\pm$ 0.9	3.75 $\pm$ 5.6	This work
S seq	4.19 $\pm$ 2.99	0.04 $\pm$ 0.15	0.65 $\pm$ 1.35	0.37 $\pm$ 0.45	0.05 $\pm$ 0.14	0.98 $\pm$ 2.05	This work
Total C seq ^c	563	–	160	–	111	830	Mitsch et al. (2013)
Total C seq ^c	368	–	152	–	55	575	This work

^aUpper and lower edges are the 25th and 75th quantiles. ^bMinimum and maximum values of the range. ^cTotal carbon sequestration (Tg C year⁻¹).

(methanogenesis) and R7 (methane oxidation), but also with N₂O emissions. The presence of B_{MGB} guarantees anaerobic conditions necessary for denitrification. The same trade-off explains the high correlation between B_{DEN} and CH₄ and CO₂ emissions.

Interestingly, we found that among the soil variables, the simulated soil organic carbon (label SOC) has the highest correlation with each GHG, while temperature did not have a significant effect, in particular on CO₂ emissions (see also Section 3.5). Finally, pH has a negative correlation with all the target variables except

Figure 5. Pearson coefficient of forcing for carbon input (label SOM input), net recharge (label PRE-ETA), N₂ fixation (N fixation), and P, N, and S fertilization (label P, N, S Fert.), microbial functional groups concentration, that is, B_{MGB} , B_{MOB} , B_{AOB} , B_{DEN} , F_{DEP} , B_{AER} , B_{SRB} , and B_{NOB} , soil variables, that is, soil organic carbon (label SOC), temperature (label T), and pH (label pH), and soil properties, that is, porosity (label PHI), hydraulic saturation (label KS), plant CH₄ emission efficiency (label PLEM), root depth (label ERTD), and bulk density (label BLD), against long-term average annual CH₄, CO₂, and N₂O emissions during 2000–2017. Results in each group are sorted from the highest positive (red) to the lowest negative (blue) correlation coefficient values. SOM, soil organic matter.

Figure 6. Pearson coefficient of forcing for carbon input (label SOM input), net recharge (label PRE-ETA), N_2 fixation (N fixation), and P, N, and S fertilization (label P, N, S Fert.), microbial functional groups concentration, that is, B_{MGB} , B_{MOB} , B_{AOB} , B_{DEN} , F_{DEP} , B_{AER} , B_{SRB} , and B_{NOB} , soil variables, that is, soil organic carbon (label SOC), temperature (label T), and pH (label pH), and soil properties, that is, porosity (label PHI), hydraulic saturation (label KS), plant CH_4 emission efficiency (label PLEM), root depth (label ERTD), and bulk density (label BLD), against long-term average annual carbon (C), nitrogen (N), and sulfur (S) sequestration rate. Results in each group are sorted from the highest positive (red) to the lowest negative (blue) correlation coefficient values. SOM, soil organic matter.

in the arid and tropical climates (Af/Am/As/Aw), where it positively affects CH_4 , N_2O , and CO_2 . Of all the soil properties, none seems relevant to GHG emissions except the porosity (label PHI), which is positively correlated with CH_4 in some temperate and continental regions (Cs/Cwa and Dwa/Dwb/Dwc/Dwd), and bulk density (label BDL), which is negatively correlated with CH_4 , CO_2 , and N_2O in some continental regions (Dwa/Dwb/Dwc/Dwd).

The carbon input (label SOM input) has the highest and most uniform correlation among all the forcing conditions against each nutrient sequestration. By contrast, N, P, and S fertilizations (labeled as N, P, and S Fert.) have a weak negative correlation because the fertilizer increases the input of rapidly degradable substrates. However, in the arid (Bsk and Bwh) and continental (Dfa) wetlands, we found a positive correlation with fertilizer input. As mentioned above, additional nutrient inputs to these soils helps to sustain microbial activity, but because of the extreme temperature and precipitation regimes, only a part of the fertilizer nutrients are sequestered in the soil matrix. Increases in microbial activity decrease nutrient sequestration. In particular, heterotrophic respiration is 10 times faster than the anaerobic one. In fact, we found that B_{MGB} is highly positively correlated with sequestration of all nutrients, while B_{AER} is negatively correlated in most of the KG classes.

Among the soil variables, the simulated soil organic carbon (label SOC) is the most highly correlated with sequestration of C, N, and S, while pH is strongly negatively correlated, except in arid regions (Bsk/Bwh/Bwk). As per GHG emissions, the porosity (label PHI) is positively correlated with each KG class, and the bulk density (label BDL) is negatively correlated.

Figure 7. Dynamic analysis: CH_4 with changing temperature (a), SOM input (b), and B_{MGB} concentration (c) during 1 year; CO_2 with changing temperature (d), SOM input (e), and B_{MGB} concentration (f) during 1 year; N_2O with changing temperature (g), SOM input (h), and B_{DEN} concentration (i) during 1 year; January is the initial point, represented with a dot, then the arrow shows the direction. Data refer only the grid cells affected by agricultural runoff. Gray dashed line represents each KG class without fertilization inputs. SOM, soil organic matter; KG, Köppen–Geiger.

3.5. Dynamic Analysis

We examined if a specific subseasonal relationship exists between GHG emissions and temperature, SOM input, B_{MGB} , and B_{DEN} , and whether fertilizer inputs affect this relationship. The KG climate classification groups wetlands within the same climate and the same or similar vegetation cover (Rohli et al., 2015). The starting point (January) is represented in Figure 7 with a dot while the arrow shows the time direction. Figure 7 reports only the KG classes (and therefore only the grid cells) affected by fertilization inputs, while Figure S11, shows the results for all KG classes.

Fertilization inputs slightly decrease CH_4 emissions (Figures 7a–7c) due to electron competition except in arid regions (Bsh, Bsk, and Bwh), where the emissions remain almost constant. In contrast, CO_2 emissions increase slightly (Figures 7d–7f) because of a decrease in CH_4 production and oxidation, and a significant increase in N_2O production (Figures 7g–7i). We found hysteresis in the relationship between CH_4 and CO_2 emissions and temperature, with higher emissions during autumn than in spring at the same temperature (Chang et al., 2020), while SOM input and B_{MGB} do not.

Dry climates (Bsh, Bsk, and Bwh) have the lowest CH_4 emissions and do not show a wide hysteresis against temperature (Figure 7a). These areas have extreme temperatures that affect the vegetation (the major biome is desert, Rohli et al., 2015), with very low SOM input (Figure 7b). The tropical area is divided into two groups: rain forest (Af and Am in Figure S11) has high CH_4 emissions, SOM input, and B_{MGB} concentrations and is not affected by agricultural runoff; and the savanna (As/Aw), characterized mainly by grasslands (Rohli et al., 2015) with low SOM input, emissions, and B_{MGB} concentrations. Continental and temperate regions show the most significant change in CH_4 emissions as a function of temperature (Figure 7a). Despite the highest emissions in Dfc (subarctic), Dfb (warm summer continental), and Cfb/Cfc (humid subtropical climate and oceanic climate), the effect of fertilizer is minor. Dfc and Dfb are characterized by grassland, peatland, and temperate forest (Rohli et al., 2015); hence, the annual SOM input is very high (Figure 7b), while Cfb/Cfc is characterized by tundra, grassland, and temperate forest, with high SOM input during summertime. Surprisingly, the B_{MGB} concentration is quite stable over the year in each KG climatic group, with no significant changes, even accounting for fertilization (Figure 7c).

CO_2 emissions show dynamics similar to CH_4 , highlighting again the strong correlation between these two gases (Figures 7d–7f).

Figures 7g–7i show very different dynamics for N_2O emissions, with no hysteresis against temperature and a much wider cycle due to fertilization. Moreover, higher average emissions occur around April in Cfb/Cfc, Cfa, Dfb, Dfc, Dfa, and Cwb/Cwc climatic regions (Figure 7g). Emissions in tropical areas are lower than in other areas because of fewer anthropogenic pressures, including agricultural fertilization (Figures 7g–7i and S11g–S11i). In addition, tropical areas consist of rainforest (Rohli et al., 2015), therefore the C:N litter aboveground is very high ($\text{C:N}_{\text{NPPAB}} = 64.30$), releasing less N to the soil. Additional moderate inputs of N fertilizer have little effect on emissions. B_{DEN} dynamics are very similar to CO_2 or CH_4 emissions and B_{MGB} , with little change over the year (Figure 7i); however, here, fertilizers have a significant effect on biomass, in particular in the Cfb/Cfc and Dfa classes.

4. Discussion

Current global estimates of wetland CH_4 emissions range between 95 and 213 Tg C- CH_4 year^{−1} (Kirschke et al., 2013; L. Liu et al., 2020; Melton et al., 2013; Poulter et al., 2017; Saunio et al., 2020; Tian et al., 2015; Zhu et al., 2015) across different historical periods, different estimation approaches, different climatic forcing, and wetland area fractions. Here, we estimated an average CH_4 global emission of 135 Tg C- CH_4 year^{−1} between 2000 and 2017. On average, our estimates are slightly greater than other bottom-up estimates in Saunio et al. (2020) (Figure S10b) but still in the range of top-down estimated emissions. Most of the current models linked CH_4 production to NPP, temperature, pH, and redox conditions using multiplication factors, but without accounting for microbial growth and carbon sequestration dynamics, the latter being very important in C-rich northern peatlands. Rising temperature and permafrost thawing make part of prestocked C in these areas available to microorganisms, thus increasing GHG emissions significantly (Baldwin et al., 2014; Davidson & Janssens, 2006; Kuhn et al., 2018). The tropical area is also the primary contributor to intra-annual variations given its favorable temperature and persistently large wetland fractional area. The northern latitudes contribute most of the high intra-annual variability (Figure 3).

Agronutrient inorganic fertilization had minor effects on wetland CH_4 and CO_2 emissions but significantly affected the N_2O emissions (Barnard et al., 2005; Kroeze & Seitzinger, 1998). Neglecting runoff of N from fertilization would underestimate the available concentration of NH_4^+ and NO_3^- in wetland soils (Morse et al., 2012), resulting in 13% reduction in N_2O emissions in the Northern Hemisphere. The overall emission of N_2O from wetlands is not particularly high compared to agriculture's contribution. However, N_2O

is almost 300 times more powerful as a GHG than CO₂, which could represent a potential problem, both in terms of global warming and biodiversity conservation. Nutrient-rich inputs to wetlands also increase plant productivity and may change the dominant plant composition (Egertson et al., 2004; Hagerthey et al., 2008; Johnson & Rejmánková, 2005), thus directly affecting nutrient decomposition, retention, and turnover rates (Verhoeven et al., 2006). Increased nutrient inputs, together with an increase in temperature, may increase N₂O emissions synergistically after considering the positive correlation between the two variables (Figure 5). The relationship between emissions and either temperature or SOM input shows a high degree of complexity (Figure 7) as it strongly depends on the type of major biomes present. A deeper understanding of these dynamics in each climatic region may give a better description of SOM input in terms of quality (C:N and C:S litter ratio) and quantity (amount of litter entering the soil) and could be used to improve descriptions of biogeochemical cycles in wetlands. Moreover, N fertilization was positively correlated with CH₄ emissions in arid climates, where additional N helped to sustain microbial activity in a nutrient-poor environment.

The addition of fertilizers interferes not only with the biogeochemical cycles of a wetland but also with the ecological system. Reis et al. (2017) estimated the impact of human pressure on inland wetlands and found that tropical areas, although generally pristine, have the highest “human influential factor” due to forest conversion into agricultural land and for population expansion. Both contributions may result in a high nutrient load reaching the wetland, thus altering the emissions, and the ecological equilibrium, that is, increasing eutrophication and threatening the local fauna. The actual impact of agriculture is more complex, and it involves consideration of essential ecosystem services (McBratney et al., 2014). A better framework that combines complex biogeochemical and ecosystem service assessment is essential to set up a more effective conservation plan.

Our results demonstrated the importance of accounting for coupled protection, dissolved, and gaseous phases. We found that the tropical and temperate climates showed the highest rates of nutrient sequestration, followed by continental, polar, and arid climates, in agreement with previous literature (Table 1). Our modeled global average C sequestration rate is 120.4 g C m⁻² year⁻¹ regardless of the type of wetland or climatic class and is lower than in Cheng et al. (2020), who reported 185.2 g C m⁻² year⁻¹. We also found that high C and N sequestration rates (e.g., 350 g C m⁻² year⁻¹ and 22 g N m⁻² year⁻¹) coincide with high gas emission rates, similar to field experiments in Nahlik and Mitsch (2010) and Bernal and Mitsch (2013). This result indicates that high SOM input rates positively affect emissions, supplying substrate to soil microorganisms, and concurrently leading to a high potential sequestration rate. It also highlights the strong link between factors that affect microbial dynamics and soil sequestration and the need for these processes to be described together (Villa & Bernal, 2018). BRTSim-BAMS4 can allocate the carbon input in different pools, that is, slow and fast biodegradable constituents. Having highly mobile species and highly protected species that dynamically change according to local conditions provides a better nutrient budget balance at different time scales.

We found that fertilization inputs do not promote C, N, and S sequestration, and the two variables are weakly negatively correlated in contrast to results from Cheng et al. (2020). However, additional N inputs help sustain plant growth particularly in nutrient-poor environments such as in arid areas, thus resulting in greater emissions and also sequestration (Hobbie, 2005). Another important factor controlling the emission is the temperature, which indirectly affects the plant yield, hence the SOM input, and the biological reaction rates, even if only the latter feature was included in our model formulation.

As with all models, uncertainties exist in our model projections. BRTSim-BAMS4 includes a wide range of complex biogeochemical reactions and feedback responses from the environment. Here, we did not investigate the errors in parameters that may affect the level of uncertainty of this study, although a sensitivity analysis was developed in previous works with the parent reaction network (Ceriotti et al., 2020; Pasut et al., 2020), highlighting the main biogeochemical processes that controls GHG emissions and C and N stocks in different ecosystems and hydroclimatic conditions. Whether the 13% decrease in N₂O emissions is within the order of magnitude of the model error is uncertain. More long-term observations of N₂O emission from flux towers in wetlands are required, but these are still a great challenge. In addition, both the reaction networks and simulation setup may be improved. For instance, both wetland and cropland areas have likely changed in these last 18 years, while we only account for changes in wetland area. A better

description should also include cropland expansion and therefore variable quality and quantity of runoff. Additionally, the biochemical kinetic parameters in the model are all spatially homogeneous, except for a few specific parameters that are vegetation dependent (e.g., C:N ratio of NPP_{AG} , N_2 fixation, root depth, and CH_4 plant emission efficiency). Neglecting the spatial distribution of these parameters results in a simplified description of microbial adaptation, which may introduce uncertainties in tropical areas, where pH is, on average, very low (Menne et al., 2012) and the temperature is mostly continuously high, and in northern latitudes, where temperatures are very low (Meng et al., 2012; Zhu et al., 2015). The effect of pH on each biologically mediated reaction was taken into account using a Michaelis–Menten term describing the microbial growth as neutrophil (Paul, 2014), hence with optima between pH 5 and 8. However, it has been found that the reduction of N_2O to N_2 is slowed down, sometimes even prevented, in acid soils, thus resulting in high N_2O emissions (B. Liu et al., 2014). This may significantly affect the estimated emissions in peat soil, where the average pH is lower than 6. Despite the complexity of the reaction network described here, some biogeochemical processes are still missing in BAMS4. For instance, we do not account for iron, manganese, and copper cycles, and their relationship with sulfur precipitation, C cycle mineralization rate (Luke et al., 2015), and toxicity to plants and bacteria, because these processes have not been fully understood yet. For the simulation setup, we used a dynamic wetland area to postprocess the data, thus potentially introducing uncertainties in our estimates (Petrescu et al., 2010). Neglecting a proper description of the expansion and contraction of the wetland area may only partially describe the O_2 diffusion in soil, hence the redox potential control on microbial processes. In addition, the effect of catchment drainage can be improved by accounting for point sources such as urban drainage, wastewater, and livestock discharges, which have recently been addressed as major sources of N input in watersheds (Bao et al., 2019). Finally, the runoff model does not explicitly account for river and stream flow. It has been found that N_2O emission rates strongly depend on the flow rate, river size, and the order of the stream of the river (Marzadri et al., 2017; Turner et al., 2015). Hence, similar pattern may occur to wetlands: depending on their position in the catchment, they can be affected differently by dilution and previous degradation, affecting the nutrients turnover.

5. Conclusions

We found that fertilizer inputs mainly affect N_2O emissions in the Northern Hemisphere, and neglecting them may underestimate by 13% the N_2O emissions. SOM input, SOC, temperature, and methanogenic bacteria are the major variables affecting GHG emissions and C, N, and S sequestration. We also found hysteresis in the relationship of temperature with CH_4 and CO_2 emissions over a year, but not with N_2O . Fertilization does not modify this relationship; however, CH_4 emissions slightly decrease and CO_2 emissions slightly increase with fertilization. Nutrient sequestration is one of the most important features of wetlands, and we found that for an equal C and N stock, wetlands affected by agriculture have higher GHG emissions. Our results show that coupling SOM with N and S cycles may support a more comprehensive biogeochemical model assessment of a wetland, both in terms of emissions and sequestration, including ecological feedbacks regulating microbial responses. Our results point out the need to focus future research on linking nutrient cycles and their complex dynamics in different phases to improve past, present, and future assessment of global nutrient budgets.

Data Availability Statement

We distribute the model output of CH_4 , CO_2 , and N_2O emissions, soil temperature, soil moisture, long-term average soil pH, carbon input, long-term average annual carbon, long-term average soil carbon, nitrogen, and sulfur sequestration rate. We include also CH_4 plant emission efficiency for aerenchyma transport, C:N and C:S ratio of litter in grassland, forest, and shrubland, N_2 fixation rate in grassland, forest, shrubland, and wetland, and the average root density for forest, grassland, shrublands, and wetland. All the data are at $0.5^\circ \times 0.5^\circ$ resolution, released in .NC format and available at <https://figshare.com/s/e50db94ddff-baa0f8304>. DOI: 10.6084/m9.figshare.12904385.

Acknowledgments

This work was supported by the SREI2020 EnviroSphere research program and the SREI Voucher of the University of Sydney. The authors acknowledge the Sydney Informatics Hub and the University of Sydney's high performance computing cluster Artemis for providing the high performance computing resources that have contributed to the results reported within this work. The authors acknowledge the use of the National Computational Infrastructure (NCI) which is supported by the Australian Government, and accessed through the Sydney Informatics Hub HPC Allocation Scheme, which is supported by the Deputy Vice-Chancellor (Research), University of Sydney and the ARC LIEF, 2019: Smith, Muller, Thornber, et al., Sustaining and strengthening merit-based access to National Computational Infrastructure (LE190100021). William J. Riley was supported by U.S. Department of Energy, Office of Science, Biological and Environmental Research, Regional and Global Climatic Modeling Program through the RUBISCO Scientific Focus Area under contract DE-AC02-05CH11231 to Lawrence Berkeley National Laboratory.

References

- Abbott, B. W., & Jones, J. B. (2015). Permafrost collapse alters soil carbon stocks, respiration, CH₄, and N₂O in upland tundra. *Global Change Biology*, 21(12), 4570–4587.
- Arora, V. K., Melton, J. R., & Plummer, D. (2018). An assessment of natural methane fluxes simulated by the CLASS-CTEM model. *Biogeosciences*, 15(15), 4683–4709.
- Baldwin, A. H., Jensen, K., & Schönfeldt, M. (2014). Warming increases plant biomass and reduces diversity across continents, latitudes, and species migration scenarios in experimental wetland communities. *Global Change Biology*, 20(3), 835–850.
- Bao, W., Yang, Y., Fu, T., & Xie, G. H. (2019). Estimation of livestock excrement and its biogas production potential in china. *Journal of Cleaner Production*, 229, 1158–1166.
- Barnard, R., Leadley, P. W., & Hungate, B. A. (2005). Global change, nitrification, and denitrification: A review. *Global Biogeochemical Cycles*, 19, GB1007. <https://doi.org/10.1029/2004GB002282>
- Beerling, D. J., Fox, A., Stevenson, D. S., & Valdes, P. J. (2011). Enhanced chemistry–climate feedbacks in past greenhouse worlds. *Proceedings of the National Academy of Sciences*, 108(24), 9770–9775.
- Bernal, B., & Mitsch, W. J. (2013). Carbon sequestration in two created riverine wetlands in the midwestern United States. *Journal of Environmental Quality*, 42(4), 1236–1244.
- Bloom, A. A., Bowman, K. W., Lee, M., Turner, A. J., Schroeder, R., Worden, J. R., & Jacob, D. J. (2017). A global wetland methane emissions and uncertainty dataset for atmospheric chemical transport models (WetCHARTs version 1.0). *Geoscientific Model Development*, 10(6), 2141–2156.
- Bonnett, S. A. F., Blackwell, M. S. A., Leah, R., Cook, V., O'Connor, M., & Maltby, E. (2013). Temperature response of denitrification rate and greenhouse gas production in agricultural river marginal wetland soils. *Geobiology*, 11(3), 252–267.
- Bréchet, L., Le Dantec, V., Ponton, S., Goret, J.-Y., Sayer, E., Bonal, D., et al. (2017). Short- and long-term influence of litter quality and quantity on simulated heterotrophic soil respiration in a lowland tropical forest. *Ecosystems*, 20(6), 1190–1204.
- Bronson, K. F., Neue, H.-U., Abao, E. B., & Singh, U. (1997). Automated chamber measurements of methane and nitrous oxide flux in a flooded rice soil: I. Residue, nitrogen, and water management. *Soil Science Society of America Journal*, 61, 981–987.
- Brooks, R. H., & Corey, A. T. (1964). Hydraulic properties of porous media. In *Hydrology papers* (No. 3). Colorado State University.
- Brown, J., Ferrians, O., Jr., Heginbottom, J., & Melnikov, E. (2001). *Circum-arctic map of permafrost and ground-ice conditions* (National Snow and Ice Data Center/World Data Center for Glaciology, Boulder, CO). Digital Media.
- Burgin, A. J., & Groffman, P. M. (2012). Soil O₂ controls denitrification rates and N₂O yield in a riparian wetland. *Journal of Geophysical Research*, 117, G01010. <https://doi.org/10.1029/2011JG001799>
- Canadell, J., Jackson, R. B., Ehleringer, J. B., Mooney, H. A., Sala, O. E., & Schulze, E.-D. (1996). Maximum rooting depth of vegetation types at the global scale. *Oecologia*, 108(4), 583–595. <https://doi.org/10.1007/bf00329030>
- Cerriotti, G., Tang, F. H., & Maggi, F. (2020). Similarities and differences in the sensitivity of soil organic matter (SOM) dynamics to biogeochemical parameters for different vegetation inputs and climates. *Stochastic Environmental Research and Risk Assessment*, 34, 2229–2244.
- Chang, K.-Y., Riley, W. J., Brodie, E. L., McCalley, C. K., Crill, P. M., & Grant, R. F. (2019). Methane production pathway regulated proximally by substrate availability and distally by temperature in a high-latitude mire complex. *Journal of Geophysical Research: Biogeosciences*, 124, 3057–3074. <https://doi.org/10.1029/2019JG005355>
- Chang, K.-Y., Riley, W. J., Crill, P. M., Grant, R. F., & Saleska, S. R. (2020). Hysteretic temperature sensitivity of wetland CH₄ fluxes explained by substrate availability and microbial activity. *Biogeosciences Discussions*, 17, 5849–5860.
- Cheng, C., Li, M., Xue, Z., Zhang, Z., Lyu, X., Jiang, M., & Zhang, H. (2020). Impacts of climate and nutrients on carbon sequestration rate by wetlands: A meta-analysis. *Chinese Geographical Science*, 30(3), 483–492. <https://doi.org/10.1007/s11769-020-1122-3>
- Cover, M. L., & Change, L.-C. (1999). *MODIS Land Cover Product Algorithm Theoretical Basis Document (Atbd) Version 5.0; MODIS documentation* (vol. 23, pp. 42–47). Maryland, USA: MODIS documentation; University of Maryland.
- Dai, Y., Shangquan, W., Duan, Q., Liu, B., Fu, S., & Niu, G. (2013). Development of a china dataset of soil hydraulic parameters using pedotransfer functions for land surface modeling. *Journal of Hydrometeorology*, 14(3), 869–887. <https://doi.org/10.1175/jhm-d-12-0149.1>
- Davidson, E. A., & Janssens, I. A. (2006). Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*, 440(7081), 165–173. <https://doi.org/10.1038/nature04514>
- Dise, N. B., & Verry, E. S. (2001). Suppression of peatland methane emission by cumulative sulfate deposition in simulated acid rain. *Biogeochemistry*, 53(2), 143–160. <https://doi.org/10.1023/a:1010774610050>
- Egertson, C. J., Kopaska, J. A., & Downing, J. A. (2004). A century of change in macrophyte abundance and composition in response to agricultural eutrophication. *Hydrobiologia*, 524(1), 145–156. <https://doi.org/10.1023/b:hydr.0000036129.40386.ce>
- Friedl, M., & Sulla-Menashe, D. (2015). MCD12Q1: MODIS/Terra + Aqua land cover type yearly L3 global 500m SIN grid v006. <https://doi.org/10.5067/MODIS/MCD12Q1.006>
- Gauci, V., Matthews, E., Dise, N., Walter, B., Koch, D., Granberg, G., & Vile, M. (2004). Sulfur pollution suppression of the wetland methane source in the 20th and 21st centuries. *Proceedings of the National Academy of Sciences*, 101(34), 12583–12587. <https://doi.org/10.1073/pnas.0404412101>
- Grant, R. F., Mekonnen, Z. A., & Riley, W. J. (2019). Modeling climate change impacts on an arctic polygonal tundra: 1. Rates of permafrost thaw depend on changes in vegetation and drainage. *Journal of Geophysical Research: Biogeosciences*, 124, 1308–1322. <https://doi.org/10.1029/2018JG004644>
- Grant, R. F., Mekonnen, Z. A., Riley, W. J., Arora, B., & Torn, M. S. (2019). Modeling climate change impacts on an Arctic Polygonal Tundra: 2. Changes in CO₂ and CH₄ exchange depend on rates of permafrost thaw as affected by changes in vegetation and drainage. *Journal of Geophysical Research: Biogeosciences*, 124, 1323–1341. <https://doi.org/10.1029/2018JG004645>
- Hagerthey, S. E., Newman, S., Rutchev, K., Smith, E. P., & Godin, J. (2008). Multiple regime shifts in a subtropical peatland: Community-specific thresholds to eutrophication. *Ecological Monographs*, 78(4), 547–565. <https://doi.org/10.1890/07-0538.1>
- Hättenschwiler, S., & Jørgensen, H. B. (2010). Carbon quality rather than stoichiometry controls litter decomposition in a tropical rain forest. *Journal of Ecology*, 98(4), 754–763. <https://doi.org/10.1111/j.1365-2745.2010.01671.x>
- Hayman, G. D., O'Connor, F. M., Dalvi, M., Clark, D. B., Gedney, N., Huntingford, C., et al. (2014). Comparison of the HadGEM2 climate–chemistry model against in situ SCIAMACHY atmospheric methane data. *Atmospheric Chemistry and Physics*, 14(23), 13257–13280. <https://doi.org/10.5194/acp-14-13257-2014>
- Hengl, T., de Jesus, J. M., Heuvelink, G. B., Gonzalez, M. R., Kilibarda, M., Blagotić, A., et al. (2017). SoilGrids250m: Global gridded soil information based on machine learning. *PLoS One*, 12(2), e0169748. <https://doi.org/10.1371/journal.pone.0169748>

- Hobbie, S. E. (2005). Contrasting effects of substrate and fertilizer nitrogen on the early stages of litter decomposition. *Ecosystems*, 8(6), 644–656. <https://doi.org/10.1007/s10021-003-0110-7>
- Hong, Y., & Adler, R. F. (2008). Estimation of global SCS curve numbers using satellite remote sensing and geospatial data. *International Journal of Remote Sensing*, 29(2), 471–477. <https://doi.org/10.1080/01431160701264292>
- Johnson, S., & Rejmánková, E. (2005). Impacts of land use on nutrient distribution and vegetation composition of freshwater wetlands in northern Belize. *Wetlands*, 25(1), 89–100. [https://doi.org/10.1672/0277-5212\(2005\)025\[0089:ioluo\]2.0.co;2](https://doi.org/10.1672/0277-5212(2005)025[0089:ioluo]2.0.co;2)
- Kaushal, S. S., Groffman, P. M., Band, L. E., Elliott, E. M., Shields, C. A., & Kendall, C. (2011). Tracking nonpoint source nitrogen pollution in human-impacted watersheds. *Environmental Science & Technology*, 45(19), 8225–8232. <https://doi.org/10.1021/es200779e>
- Kaushal, S. S., Mayer, P. M., Vidon, P. G., Smith, R. M., Pennino, M. J., Newcomer, T. A., et al. (2014). Land use and climate variability amplify carbon, nutrient, and contaminant pulses: A review with management implications. *Journal of the American Water Resources Association*, 50(3), 585–614. <https://doi.org/10.1111/jawr.12204>
- Kaushal, S. S., Pace, M. L., Groffman, P. M., Band, L. E., Belt, K. T., Meyer, P. M., & Welty, C. (2010). Land use and climate variability amplify contaminant pulses. *Eos, Transactions American Geophysical Union*, 91(25), 221–222. <https://doi.org/10.1029/2010eo250001>
- Kayranli, B., Scholz, M., Mustafa, A., & Hedmark, Å. (2010). Carbon storage and fluxes within freshwater wetlands: A critical review. *Wetlands*, 30(1), 111–124. <https://doi.org/10.1007/s13157-009-0003-4>
- Kirschke, S., Bousquet, P., Ciais, P., Saunio, M., Canadell, J. G., Dlugokencky, E. J., et al. (2013). Three decades of global methane sources and sinks. *Nature Geoscience*, 6(10), 813–823. <https://doi.org/10.1038/ngeo1955>
- Köppen, W., & Geiger, G. (1930). *Handbuch der klimatologie-gebrüder borntraeger*. (Vol. 1). Berlin: Gebrüder Borntraeger.
- Kroeze, C., & Seitzinger, S. P. (1998). The impact of land use on N₂O emissions from watersheds draining into the northeastern Atlantic Ocean and European seas. *Environmental Pollution*, 102(1), 149–158. [https://doi.org/10.1016/s0269-7491\(98\)80027-4](https://doi.org/10.1016/s0269-7491(98)80027-4)
- Kuhn, M., Lundin, E. J., Giesler, R., Johansson, M., & Karlsson, J. (2018). Emissions from thaw ponds largely offset the carbon sink of northern permafrost wetlands. *Scientific Reports*, 8(1), 1–7. <https://doi.org/10.1038/s41598-018-27770-x>
- Lal, R. (2008). Carbon sequestration. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1492), 815–830. <https://doi.org/10.1098/rstb.2007.2185>
- Lehner, B., & Döll, P. (2004). Development and validation of a global database of lakes, reservoirs and wetlands. *Journal of Hydrology*, 296(1–4), 1–22. <https://doi.org/10.1016/j.jhydrol.2004.03.028>
- Li, Z., & Bowman, R. S. (2001). Retention of inorganic oxyanions by organo-kaolinite. *Water Research*, 35(16), 3771–3776. [https://doi.org/10.1016/s0043-1354\(01\)00120-8](https://doi.org/10.1016/s0043-1354(01)00120-8)
- Liu, B., Frostegård, Å., & Bakken, L. R. (2014). Impaired reduction of N₂O to N₂ in acid soils is due to a posttranscriptional interference with the expression of nosZ. *mBio*, 5(3). <https://doi.org/10.1128/mbio.01383-14>
- Liu, L., Zhuang, Q., Oh, Y., Shurpali, N. J., Kim, S., & Poulter, B. (2020). Uncertainty quantification of global net methane emissions from terrestrial ecosystems using a mechanistically based biogeochemistry model. *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005428. <https://doi.org/10.1029/2019JG005428>
- Loiko, S. V., Pokrovsky, O. S., Raudina, T. V., Lim, A., Kolesnichenko, L. G., Shirokova, L. S., et al. (2017). Abrupt permafrost collapse enhances organic carbon, CO₂, nutrient and metal release into surface waters. *Chemical Geology*, 471, 153–165. <https://doi.org/10.1016/j.chemgeo.2017.10.002>
- Luke, S., Preston, M. D., Basiliko, N., & Watmough, S. A. (2015). Microbial communities, biomass, and carbon mineralization in acidic, nutrient-poor peatlands impacted by metal and acid deposition. *Water, Air, & Soil Pollution*, 226(2), 19. <https://doi.org/10.1007/s11270-014-2265-6>
- Maggi, F. (2019). *BRTSim, a general-purpose computational solver for hydrological, biogeochemical, and ecosystem dynamics*. arXiv preprint arXiv:1903.07015.
- Maggi, F., Gu, C., Riley, W., Hornberger, G., Venterea, R., Xu, T., & Oldenburg, C. (2008). A mechanistic treatment of the dominant soil nitrogen cycling processes: Model development, testing, and application. *Journal of Geophysical Research: Biogeosciences*, 113, G02016. <https://doi.org/10.1029/2007JG000578>
- Malone, L., Lacelle, D., Kokelj, S., & Clark, I. D. (2013). Impacts of hillslope thaw slumps on the geochemistry of permafrost catchments (Stony Creek Watershed, NWT, Canada). *Chemical Geology*, 356, 38–49. <https://doi.org/10.1016/j.chemgeo.2013.07.010>
- Marzadri, A., Dee, M. M., Tonina, D., Bellin, A., & Tank, J. L. (2017). Role of surface and subsurface processes in scaling N₂O emissions along riverine networks. *Proceedings of the National Academy of Sciences of the United States of America*, 114(17), 4330–4335. <https://doi.org/10.1073/pnas.1617454114>
- Matías, L., Castro, J., & Zamora, R. (2011). Soil-nutrient availability under a global-change scenario in a Mediterranean mountain ecosystem. *Global Change Biology*, 17(4), 1646–1657. <https://doi.org/10.1111/j.1365-2486.2010.02338.x>
- McBratney, A., Field, D. J., & Koch, A. (2014). The dimensions of soil security. *Geoderma*, 213, 203–213. <https://doi.org/10.1016/j.geoderma.2013.08.013>
- Mekonnen, Z. A., Grant, R. F., & Schwalm, C. (2016). Sensitivity of modeled NEP to climate forcing and soil at site and regional scales: Implications for upscaling ecosystem models. *Ecological Modelling*, 320, 241–257. <https://doi.org/10.1016/j.ecolmodel.2015.10.004>
- Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., et al. (2013). Present state of global wetland extent and wetland methane modelling: Conclusions from a model inter-comparison project (WETCHIMP). *Biogeosciences*, 10, 753–788. <https://doi.org/10.5194/bg-10-753-2013>
- Meng, L., Hess, P. G. M., Mahowald, N. M., Yavitt, J. B., Riley, W. J., Subin, Z. M., et al. (2012). Sensitivity of wetland methane emissions to model assumptions: Application and model testing against site observations. *Biogeosciences*, 9(7), 2793–2819. <https://doi.org/10.5194/bg-9-2793-2012>
- Menne, M. J., Durre, I., Vose, R. S., Gleason, B. E., & Houston, T. G. (2012). An overview of the global historical climatology network-daily database. *Journal of Atmospheric and Oceanic Technology*, 29(7), 897–910. <https://doi.org/10.1175/jtech-d-11-00103.1>
- Mitsch, W. J., Bernal, B., Nahlik, A. M., Mander, Ü., Zhang, L., Anderson, C. J., et al. (2013). Wetlands, carbon, and climate change. *Land-scape Ecology*, 28(4), 583–597. <https://doi.org/10.1007/s10980-012-9758-8>
- Mosier, A. R., Morgan, J. A., King, J. Y., Lecain, D., & Milchunas, D. G. (2002). Soil-atmosphere exchange of CH₄, CO₂, NO_x, and N₂O in the Colorado shortgrass steppe under elevated CO₂. *Plant and Soil*, 240(2), 201–211. <https://doi.org/10.1023/a:1015783801324>
- Nahlik, A. M., & Mitsch, W. J. (2010). Methane emissions from created riverine wetlands. *Wetlands*, 30(4), 783–793. <https://doi.org/10.1007/s13157-010-0038-6>
- Pangala, S. R., Enrich-Prast, A., Basso, L. S., Peixoto, R. B., Bastviken, D., Hornibrook, E. R. C., et al. (2017). Large emissions from flood-plain trees close the amazon methane budget. *Nature*, 552(7684), 230–234. <https://doi.org/10.1038/nature24639>

- Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., et al. (2020). The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data. *Scientific Data*, 7(1), 1–27.
- Pasut, C., Tang, F. H., & Maggi, F. (2020). A mechanistic analysis of wetland biogeochemistry in response to temperature, vegetation, and nutrient input changes. *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005437. <https://doi.org/10.1029/2019JG005437>
- Paul, E. A. (2014). *Soil microbiology, ecology and biochemistry*. Academic Press.
- Pei, G., Liu, J., Peng, B., Gao, D., Wang, C., Dai, W., et al. (2019). Nitrogen, lignin, C/N as important regulators of gross nitrogen release and immobilization during litter decomposition in a temperate forest ecosystem. *Forest Ecology and Management*, 440, 61–69. <https://doi.org/10.1016/j.foreco.2019.03.001>
- Petrescu, A., Van Beek, L., Van Huissteden, J., Prigent, C., Sachs, T., Corradi, C., & Dolman, A. (2010). Modeling regional to global CH₄ emissions of boreal and arctic wetlands. *Global Biogeochemical Cycles*, 24, GB4009. <https://doi.org/10.1029/2009GB003610>
- Potter, P., Ramankutty, N., Bennett, E., & Donner, S. (2011a). *Global fertilizer and manure, version 1: Nitrogen fertilizer application*. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC).
- Potter, P., Ramankutty, N., Bennett, E., & Donner, S. (2011b). *Global fertilizer and manure, version 1: Phosphorus fertilizer application*. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC).
- Poulter, B., Bousquet, P., Canadell, J. G., Ciais, P., Peregon, A., Saunio, M., et al. (2017). Global wetland contribution to 2000–2012 atmospheric methane growth rate dynamics. *Environmental Research Letters*, 12(9), 094013. <https://doi.org/10.1088/1748-9326/aa8391>
- Reis, V., Hermoso, V., Hamilton, S. K., Ward, D., Fluet-Chouinard, E., Lehner, B., & Linke, S. (2017). A global assessment of inland wetland conservation status. *BioScience*, 67(6), 523–533. <https://doi.org/10.1093/biosci/bix045>
- Riley, W. J., Maggi, F., Kleber, M., Torn, M. S., Tang, J. Y., Dwivedi, D., & Guerry, N. (2014). Long residence times of rapidly decomposable soil organic matter: Application of a multi-phase, multi-component, and vertically resolved model (BAMS1) to soil carbon dynamics. *Geoscientific Model Development*, 7(4), 1335–1355. <https://doi.org/10.5194/gmd-7-1335-2014>
- Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., et al. (2011). Barriers to predicting changes in global terrestrial methane fluxes: Analyses using CLM4Me, a methane biogeochemistry model integrated in CESM. *Biogeosciences*, 8(7), 1925–1953. <https://doi.org/10.5194/bg-8-1925-2011>
- Rohli, R. V., Joyner, T. A., Reynolds, S. J., & Ballinger, T. J. (2015). Overlap of global Köppen–Geiger climates, biomes, and soil orders. *Physical Geography*, 36(2), 158–175. <https://doi.org/10.1080/02723646.2015.1016384>
- Rouified, S., Handa, I. T., David, J.-F., & Hättenschwiler, S. (2010). The importance of biotic factors in predicting global change effects on decomposition of temperate forest leaf litter. *Oecologia*, 163(1), 247–256. <https://doi.org/10.1007/s00442-009-1528-1>
- Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., & others (2020). The Global Methane Budget 2000–2017. *Earth System Science Data*, 12(3), 1561–1623.
- Schroeder, R., McDonald, K., Chapman, B., Jensen, K., Podest, E., Tessler, Z., et al. (2015). Development and evaluation of a multi-year fractional surface water data set derived from active/passive microwave remote sensing data. *Remote Sensing*, 7(12), 16688–16732. <https://doi.org/10.3390/rs71215843>
- Shannon, R. D., & White, J. R. (1994). A three-year study of controls on methane emissions from two Michigan peatlands. *Biogeochemistry*, 27(1), 35–60. <https://doi.org/10.1007/bf00002570>
- Snowdon, P., Ryan, P., & Raison, R. J. (2005). *Review of C:N ratios in vegetation, litter and soil under Australian native forests and plantations*. CSIRO Research Publications Repository.
- Solomon, S. (2007). The physical science basis: Contribution of Working Group I to the fourth assessment report of the Intergovernmental Panel on Climate Change. Intergovernmental Panel on Climate Change (IPCC). *Climate Change*, 2007, 996.
- Sulla-Menashe, D., Gray, J. M., Abercrombie, S. P., & Friedl, M. A. (2019). Hierarchical mapping of annual global land cover 2001 to present: The MODIS collection 6 land cover product. *Remote Sensing of Environment*, 222, 183–194. <https://doi.org/10.1016/j.rse.2018.12.013>
- Tang, F. H. M., Riley, W. J., & Maggi, F. (2019). Hourly and daily rainfall intensification causes opposing effects on C and N emissions, storage, and leaching in dry and wet grasslands. *Biogeochemistry*, 144(2), 197–214. <https://doi.org/10.1007/s10533-019-00580-7>
- Tian, H., Chen, G., Lu, C., Xu, X., Ren, W., Zhang, B., et al. (2015). Global methane and nitrous oxide emissions from terrestrial ecosystems due to multiple environmental changes. *Ecosystem Health and Sustainability*, 1(1), 1–20. <https://doi.org/10.1890/ehs14-0015.1>
- Tian, H., Melillo, J., Lu, C., Kicklighter, D., Liu, M., Ren, W., et al. (2011). China's terrestrial carbon balance: Contributions from multiple global change factors. *Global Biogeochemical Cycles*, 25, GB1007. <https://doi.org/10.1029/2010GB003838>
- Tian, H., Xu, X., Lu, C., Liu, M., Ren, W., Chen, G., & Liu, J. (2011). Net exchanges of CO₂, CH₄, and N₂O between China's terrestrial ecosystems and the atmosphere and their contributions to global climate warming. *Journal of Geophysical Research: Biogeosciences*, 116, G02011. <https://doi.org/10.1029/2010JG001393>
- Treat, C. C., Wollheim, W. M., Varner, R. K., & Bowden, W. B. (2016). Longer thaw seasons increase nitrogen availability for leaching during fall in tundra soils. *Environmental Research Letters*, 11(6), 064013. <https://doi.org/10.1088/1748-9326/11/6/064013>
- Turner, P. A., Griffis, T. J., Lee, X., Baker, J. M., Venterea, R. T., & Wood, J. D. (2015). Indirect nitrous oxide emissions from streams within the US corn belt scale with stream order. *Proceedings of the National Academy of Sciences of the United States of America*, 112(32), 9839–9843. <https://doi.org/10.1073/pnas.1503598112>
- Verhoeven, J., Arheimer, B., Yin, C., & Hefting, M. (2006). Regional and global concerns over wetlands and water quality. *Trends in Ecology & Evolution*, 21(2), 96–103. <https://doi.org/10.1016/j.tree.2005.11.015>
- Villa, J. A., & Bernal, B. (2018). Carbon sequestration in wetlands, from science to practice: An overview of the biogeochemical process, measurement methods, and policy framework. *Ecological Engineering*, 114, 115–128. <https://doi.org/10.1016/j.ecoleng.2017.06.037>
- Vitousek, P. M., Aber, J. D., Howarth, R. W., Likens, G. E., Matson, P. A., Schindler, D. W., et al. (1997). Human alteration of the global nitrogen cycle: Sources and consequences. *Ecological Applications*, 7(3), 737–750. [https://doi.org/10.1890/1051-0761\(1997\)007\[0737:haotgn\]2.0.co;2](https://doi.org/10.1890/1051-0761(1997)007[0737:haotgn]2.0.co;2)
- Wagner, L., Vidon, P., Tedesco, L., & Gray, M. (2008). Stream nitrate and doc dynamics during three spring storms across land uses in glaciated landscapes of the midwest. *Journal of Hydrology*, 362(3–4), 177–190. <https://doi.org/10.1016/j.jhydrol.2008.08.013>
- Walter, B. P., Heimann, M., & Matthews, E. (2001). Modeling modern methane emissions from natural wetlands: 1. Model description and results. *Journal of Geophysical Research*, 106(D24), 34189–34206. <https://doi.org/10.1029/2001jd900165>
- Wania, R., Melton, J. R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., et al. (2013). Present state of global wetland extent and wetland methane modeling: Methodology of a model inter-comparison project (WETCHIMP). *Geoscientific Model Development*, 6(3), 617–641. <https://doi.org/10.5194/gmd-6-617-2013>
- Were, D., Kansime, F., Fetahi, T., Cooper, A., & Jjuuko, C. (2019). Carbon sequestration by wetlands: A critical review of enhancement measures for climate change mitigation. *Earth Systems and Environment*, 3, 327–340.
- Whitehead, D. (2000). *Nutrient elements in grassland: Soil–plant–animal relationships*. CABI Publishing.

- Woodward, F. I., Smith, T. M., & Emanuel, W. R. (1995). A global land primary productivity and phytogeography model. *Global Biogeochemical Cycles*, 9(4), 471–490. <https://doi.org/10.1029/95gb02432>
- Xu, X., Tian, H., Zhang, C., Liu, M., Ren, W., Chen, G., & Bruhwiler, L. (2010). Attribution of spatial and temporal variations in terrestrial methane flux over North America. *Biogeosciences Discussions*, 7(4). <https://doi.org/10.5194/bg-7-3637-2010>
- Zhang, Y., Pena Arancibia, J., McVicar, T., Chiew, F., Vaze, J., Zheng, H., & Wang, Y. (2016). Monthly global observation-driven Penman–Monteith–Leuning (PML) evapotranspiration and components (v2). CSIRO. Data Collection. <https://doi.org/10.4225/08/5719A5C48DB85>
- Zhu, Q., Peng, C., Chen, H., Fang, X., Liu, J., Jiang, H., et al. (2015). Estimating global natural wetland methane emissions using process modelling: Spatio-temporal patterns and contributions to atmospheric methane fluctuations. *Global Ecology and Biogeography*, 24(8), 959–972. <https://doi.org/10.1111/geb.12307>