Growth inhibitory properties of extracts prepared from selected *Leptospermum* and *Melaleuca* species against a panel of pathogenic bacteria

Lindiwe Nomathemba Mpala¹, Getmore Rumbudzai Chikowe¹, Ian Edwin Cock^{1,2*}

¹School of Natural Sciences, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland 4111, AUSTRALIA.

²Environmental Futures Research Institute, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland 4111, AUSTRALIA.

ABSTRACT

Introduction: *Leptospermum longifolium* (C.T. White & W.D. Francis) S.T. Blake, *Leptospermum petersonii* Bailey and *Melaleuca alternifolia* (Maiden & Betche) Cheel are aromatic native Australian trees with uses as traditional medicines. Essential oils produced from leaves of these species have reputed antiseptic properties against many bacteria. Despite this, *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf solvent extractions have not been rigorously examined for antibacterial properties against many pathogens. **Methods:** The antimicrobial activity of methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts was investigated by disc diffusion and growth time course assays against a panel of pathogenic bacteria. The growth inhibitory activity was quantified by MIC determination. Toxicity was determined using the *Artemia franciscana* nauplii bioassay. **Results:** The methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts inhibited the growth of a wide range of bacterial species. Growth of both gram positive and gram negative bacteria was inhibited by all extracts. The *L. longifolium* and *L. petersonii* extracts were generally more potent inhibitors of bacterial growth than was the *M. alternifolia* extract against most bacterial species. *A. hydrophilia*, *C. freundii*, *P. mirabilis* and *B. cereus* growth was particularly susceptible to the extracts, with MIC values as low as 147 µg/mL (inhibition of *A. hydrophilia* growth by the *L. longifolium* extract). The antibacterial activity of the *L. longifolium*, *L. petersonii* and

M. alternifolia extracts were further investigated by growth time course assays, with significant growth inhibition recorded in all cultures within 1 h of exposure. All extracts were determined to be nontoxic in the *Artemia franciscana* nauplii bioassay, indicating their safety for therapeutic uses.

Conclusions: The lack of toxicity of the methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts and their growth inhibitory bioactivity against a panel of pathogenic bacteria partially validate Australian Aboriginal usage of these species as antiseptic agents and indicate their potential in the development of antiseptic agents.

Key words: Myrtaceae, *Leptospermum longifolium*, *Leptospermum petersonii*, *Melaleuca alternifolia*, Tea-tree, Australian plants, Antibacterial activity, Medicinal plants.

Correspondence:

Ian Edwin Cock, School of Natural Sciences, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland 4111, AUSTRALIA.
Phone no: +61 7 37357637; Fax: +61 7 37355282

E-mail: I.Cock@griffith.edu.au (I. E. Cock)

DOI : 10.5530/pc.2016.4.4

INTRODUCTION

Plants produce a wide variety of secondary metabolites which provide characteristic pigment, odour and flavour characteristics. In addition, these compounds may also provide the plants with protection against microbial challenge.¹ Traditional plant derived medicines have been used for thousands of years in most parts of the world and with the increase in microbial antibiotic resistance, their use in fighting bacterial pathogens is becoming the focus of intense study.^{2,3} Whilst much of the research into traditional medicinal plant use has focused on Asian,⁴ African⁵ and South American⁶ plants, the therapeutic potential of the flora of Australia has also been recognised for thousands of years. The first Australians had well developed ethnopharmacological systems and understood the therapeutic properties of a wide variety of aromatic Australian plants.⁷ Despite this, relatively few studies have rigorously examined the antibacterial activity of Australian native plants. However, recently there has been increased study in this field.

The healing properties of Australian plants of the family Myrtaceae have been understood by Australian Aborigines for many thousands of years. The genera *Leptospermum* and *Melaleuca* are particularly prevalent in traditional Aboriginal healing systems.⁷ More recently, Sir Joseph Banks (the botanist aboard Captain James Cook's voyage of the Endeavour) coined the phrase 'tea-tree' for the endemic Australian plant *Melaleuca alternifolia* due to similarities in use with *Camellia sinensis*, and some of its perceived therapeutic properties.⁷ However, this common name creates some taxonomic confusion, with several *Leptospermum* and

Kunzea species also collectively known as 'tea-tree'. Nowadays, a thriving trade exists, with 'tea tree' essential oils marketed globally as an antiseptic. The bacterial growth inhibitory properties of many genera within the family Myrtaceae have been documented. In particular, *Callistemon* spp.,⁸ *Eugenia* spp.,⁷ *Kunzea* spp.,^{7,9} *Leptospermum* spp.^{7,10,11} and *Syzygium* spp.¹²⁻¹⁴ have been reported to inhibit the growth of a wide panel of bacteria, including many medicinally important pathogens. It is noteworthy that many of these studies have screened essential oils for antibacterial activity and the growth inhibitory properties of several Myrtaceae extracts are yet to be rigorously examined.

Leptospermum longifolium (C.T. White & W.D. Francis) S.T. Blake (Figure 1a) and *Leptospermum petersonii* Bailey (Figure 1b) are large shrubs/small trees which are native to sclerophyll forests and subtropical rainforest areas of eastern Australia. *Melaleuca alternifolia* (Maiden & Betche) Cheel (commonly known as narrow leaved paperbark, narrow leaved tea-tree; Figure 1c) is a related species which has a similar but limited occurrence in eastern Australian subtropical sclerophyll forests, particularly besides streams and in swampy areas.¹ The leaves of all 3 species are distilled commercially to produce essential oils rich in geraniol (Figure 1d), neral (Figure 1e), citronellal (Figure 1f), pinene (Figure 1g), sabinene (Figure 1h), α-terpinene (Figure 1i), limonene (Figure 1j) and 1,8-cineole (Figure 1k).⁷

Most studies into the antibacterial potential of the family Myrtaceae focus on the essential oil of the leaves.^{7,15,16} In most *Leptospermum* and

Melaleuca species, the levels and composition of the essential oil terpenoid components receives the most interest. In particular, the monoterpenoid compositions of several species have been extensively reported.⁷ *Leptospermum* spp. essential oils contain especially high levels of citral, which comprises a mixture of geranial (β -citral; Figure 1d) and neral (α -citral; Figure 1e).^{7,15,16} Both neral and geranial have been previously reported to have potent antibacterial activity against a variety of bacteria.¹⁷⁻¹⁹ Similarly, a high composition of other monoterpenoids in various *Myrtaceae* essential oils has been reported, with citronellal (Figure 1f), pinene (Figure 1g), sabinene (Figure 1h), α -terpinene (Figure 1i), limonene (Figure 1j) and 1,8-cineole (Figure 1k) often reported as major components. The antiseptic properties of these compounds is well established, with potent bacteriostatic and bactericidal activities reported.²²

Most of the studies reporting antibacterial properties for *Leptospermum* spp. and *Melaleuca* spp. have examined essential oils. However, the use of essential oils for the testing of antimicrobial activity can be problematic. The relative insolubility of many of the oil components retards their diffusion through agar gels in agar dilution or disc diffusion studies. Many studies have utilised solubilising agents (e.g. Tween 80) to aid oil component diffusion, resulting in variable results.^{20,21} Solubilising agents appear to increase the susceptibility of some bacteria to antimicrobial agents, decrease the susceptibility of others, whilst having no effect on yet other bacteria. The current study was undertaken to examine 3 taxonomically related *Myrtaceae* spp. (*L. longifolium*, *L. petersonii* and *M. alternifolia*) for growth inhibitory properties against a panel of pathogenic bacteria.

MATERIALS AND METHODS

Plant collection and extraction

Leptospermum longifolium (C.T. White & W.D. Francis) S.T. Blake, *Leptospermum petersonii* Bailey and *Melaleuca alternifolia* (Maiden & Betche) Chee leaves were obtained from and identified by Philip Cameron, senior botanic officer, Mt Cootha Botanical Gardens, Brisbane, Australia. Leaf samples were dried in a Sunbeam food dehydrator and stored at -30°C. Prior to use, the leaves were freshly ground to a coarse powder and 1 g quantities were weighed into separate tubes. A volume of 50 mL methanol (Ajax, Australia; AR grade) was added to individual tubes and extracted for 24 hrs at 4°C with gentle shaking. The extracts were filtered through filter paper (Whatman No. 54) under vacuum, followed by drying by rotary evaporation in an Eppendorf concentrator 5301. The resultant pellets were dissolved in 10 mL sterile deionised water (containing 1% DMSO). The extracts were passed through a 0.22 μ m filter (Sarstedt) and stored at 4°C until use.

Qualitative phytochemical studies

Phytochemical analysis of the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts for the presence of saponins, phenolic compounds, flavonoids, phytosteroids, triterpenoids, cardiac glycosides, anthraquinones, tannins and alkaloids was conducted by previously described assays.²³⁻²⁴

Antibacterial screening

Test microorganisms

All media was supplied by Oxoid Ltd., Australia. Clinical isolate microbial strains of *Aeromonas hydrophila*, *Alcaligenes faecalis*, *Bacillus cereus*, *Citrobacter freundii*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas fluorescens*, *Salmonella newport*, *Serratia marcescens*, *Shigella sonnei*, *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Streptococcus pyogenes* were obtained from Ms Michelle Mendell and Ms Jane Gifkins, Griffith University. All stock cultures were subcultured and maintained in nutrient broth at 4°C.

Evaluation of antimicrobial activity

Antimicrobial activity of all plant extracts was determined using a modified disc diffusion assay.²⁵⁻²⁷ Briefly, 100 μ L of each bacterial culture was grown in 10 mL of fresh nutrient broth until they reached a count of $\sim 10^8$ cells/mL. A volume of 100 μ L of the bacterial suspension was spread onto nutrient agar plates and extracts were tested for antibacterial activity using 5 mm sterilised filter paper discs. Discs were infused with 10 μ L of the plant extracts, allowed to dry and placed onto the inoculated plates. The plates were allowed to stand at 4°C for 2 h before incubation at 30°C for 24 h. The diameters of the inhibition zones were measured to the closest whole millimetre. Each assay was performed in at least triplicate. Mean values ($\pm$ SEM) are reported in this study. Standard discs of ampicillin (10 μ g) were obtained from Oxoid, Australia and were used as positive controls to compare antibacterial activity. Filter discs infused with 10 μ L of distilled water were used as a negative control.

Minimum inhibitory concentration (MIC) determination

The minimum inhibitory concentration (MIC) of each extract against susceptible bacteria was determined as previously described.^{28,29} Briefly, the plant extracts were diluted in deionised water and tested across a range of concentrations. Discs were infused with 10 μ L of the test dilutions, allowed to dry and placed onto inoculated plates. The assay was completed as outlined above and graphs of the zone of inhibition versus concentration were plotted for each extract. Linear regression was used to determine the MIC values of each extract.

Bacterial growth time course assay

Bacterial growth time course studies were performed as previously described.³⁰ Briefly, 3 mL of the and *A. hydrophila*, *C. freundii* and *P. mirabilis* bacterial cultures in nutrient broth were added individually to 27 mL nutrient broth containing 3 mL of 10 mg/mL methanolic plant extract to give a final concentration of 1000 μ g/mL in the assay. The tubes were incubated at 30°C with gentle shaking. The optical density was measured hourly at 550 nm for a 6 h incubation period. Control tubes were incubated under the same conditions but without the extract. All assays were performed in triplicate.

Toxicity screening

Reference toxin for toxicity screening

Potassium dichromate ($K_2Cr_2O_7$) (AR grade, Chem-Supply, Australia) was prepared as a 4 mg/mL solution in distilled water and was serially diluted in artificial seawater for use in the *Artemia franciscana* nauplii bioassay.

Artemia franciscana nauplii toxicity screening

Toxicity was tested using an adapted *Artemia franciscana* nauplii lethality assay.³¹⁻³³ Briefly, 400 μ L of seawater containing approximately 54 (mean 54.2, n=75, SD 11.5) *A. franciscana* nauplii were added to wells of a 48 well plate and immediately used for bioassay. A volume of 400 μ L of diluted plant extracts or the reference toxin were transferred to the wells and incubated at $25 \pm 1^\circ$ C under artificial light (1000 Lux). A 400 μ L seawater negative control was run in triplicate for each plate. All treatments were performed in at least triplicate. The wells were checked at regular intervals and the number of dead counted. The nauplii were considered dead if no movement of the appendages was detected within 10 seconds. After 24 h, all nauplii were sacrificed and counted to determine the total % mortality per well. The LC_{50} with 95% confidence limits for each treatment was determined using probit analysis.

Statistical analysis

Data are expressed as the mean $\pm$ SEM of at least three independent experiments. One way ANOVA was used to calculate statistical significance between control and treated groups with a P value < 0.01 considered to be statistically significant.

RESULTS

Liquid extraction yields and qualitative phytochemical screening

Extraction of 1 g of *L. longifolium*, *L. petersonnii* and *M. alternifolia* leaves with methanol yielded dried extracts ranging from 93 mg (methanolic *M. alternifolia* extract) to 130 mg (methanolic *L. longifolium* extract) (Table 1). The dried extracts were resuspended in 10 mL of deionised water (containing 1% DMSO), resulting in the extract concentrations shown in Table 1. Qualitative phytochemical studies showed that all extracts contained similar classes of phytochemicals. All had high levels of phenolics and tannins. Generally, all extracts also contained moderate levels of flavonoids and moderate to high levels of saponins. Low to moderate levels of triterpenoids, phytosterols and alkaloids were also present in all extracts. The *L. longifolium*, *L. petersonnii* and *M. alternifolia* extracts were devoid of all other classes of phytochemicals.

Antimicrobial activity

To determine the growth inhibitory activity of the *L. longifolium*, *L. petersonnii* and *M. alternifolia* extracts against the panel of pathogenic bacteria, aliquots (10 μ L) of each extract were screened in the disc diffusion assay. All extracts inhibited a broad spectrum of the gram negative bacterial species (Figure 2). Indeed, all extracts inhibited the growth of 5 of the 10 gram negative bacteria screened (50%). The methanolic *L. longifolium* extract was a potent growth inhibitor against most bacterial species (as assessed by the sizes of the zones of inhibition), with zones of inhibition > 10 mm against *A. hydrophilia*, *C. freundii* and *P. mirabilis*. This inhibition was particularly noteworthy compared to the inhibition by the ampicillin (10 μ g; inhibition zones of approximately 8.6, 8.3 and 10.3 mm respectively against the same bacterial species) and chloramphenicol controls (2 μ g; inhibition zones of approximately 9.7, 9.3 and 8.6 mm respectively against the same bacterial species). The *L. longifolium* extract was also a good inhibitor of *A. faecalis* and *K. pneumoniae* growth, albeit with smaller zones of inhibition (< 10 mm). Similar bacterial growth inhibitory trends were noted for the *L. petersonnii* and *M. alternifolia* extracts against the gram negative bacterial species (Figure 2). As reported for the *L. longifolium* extract, the growth of *A. faecalis*, *A. hydrophilia*, *C. freundii*, *K. pneumoniae* and *P. mirabilis* was strongly inhibited by the *L. petersonnii* and *M. alternifolia* extracts.

The growth of gram positive bacteria was also potently inhibited by the *L. longifolium*, *L. petersonnii* and *M. alternifolia* extracts (Figure 3). All of the gram positive bacterial species tested were inhibited by each of the extracts. *B. cereus* and *S. aureus* were the most susceptible gram positive bacteria (as judged by the zones of inhibition), with zones of inhibition as large as 10.6 mm (*L. longifolium* and *L. petersonnii* inhibition of *S. aureus*). Notably, the *Leptospermum* spp. extracts were substantially more potent inhibitors of gram positive bacterial growth than was the *M. alternifolia* extract.

The antimicrobial efficacy was further quantified by determining the MIC values for each extract against the microbial species which were determined to be susceptible. All extracts were potent growth inhibitors of several bacterial species (as judged by MIC; Table 2). *P. mirabilis* was highly susceptible to all of the extracts, with MIC values generally < 500 μ g/mL (< 5 μ g infused into the disc). As *P. mirabilis* infection is a common cause of urinary tract infections and has also been identified

as a trigger of rheumatoid arthritis,^{38,39} the *L. longifolium*, *L. petersonnii* and *M. alternifolia* leaf extracts have potential for the prevention and treatment of these diseases in genetically susceptible individuals.

The *L. longifolium* and *L. petersonnii* extracts, but not the *M. alternifolia* extract, were also potent *K. pneumoniae* growth inhibitors (MIC's of 320 and 687 μ g/mL for the *L. longifolium* and *L. petersonnii* extracts respectively). As *K. pneumoniae* can trigger ankylosing spondylitis in genetically susceptible individuals,^{40,41} these extracts may also be useful in the prevention and treatment of this disease. Furthermore, all extracts were moderate inhibitors of *S. pyogenes* growth, with MIC values 1000-1500 μ g/mL. *S. pyogenes* may cause a myriad of diseases including streptococcal pharyngitis, impetigo and rheumatic heart disease, depending on which tissue it infects.^{42,43} Thus, the methanolic *L. longifolium*, *L. petersonnii* and *M. alternifolia* leaf extracts may also be useful in the prevention and treatment of these diseases.

L. longifolium, *L. petersonnii* and *M. alternifolia* extracts were also good inhibitors of *A. hydrophilia*, with MICs of 147, 279 and 705 μ g/mL respectively. The *L. longifolium* and *L. petersonnii* extracts, but not the *M. alternifolia* extract, were also potent inhibitors of *C. freundii*, *B. cereus*, *S. aureus* and *S. epidermidis* growth, with MIC values substantially < 1000 μ g/mL. The *M. alternifolia* extract also inhibited the growth of these bacteria, albeit with higher MICs indicative of only moderate growth inhibition.

Bacterial growth time course assay

The antibacterial activity of the *L. longifolium*, *L. petersonnii* and *M. alternifolia* extracts was further investigated in the reference bacterial strains by bacterial growth time course assays in the presence and absence of the extract. As *A. hydrophilia*, *C. freundii*, and *P. mirabilis* were the most susceptible to the *L. longifolium*, *L. petersonnii* and *M. alternifolia* extracts, these species were selected for growth time course studies. The starting concentration of the extract used in these assays was 1000 μ g/mL. All extracts significantly inhibited *A. hydrophilia* (Figure 4a), *C. freundii* (Figure 4b) and *P. mirabilis* (Figure 4c) growth within 1 h, indicating a rapid antimicrobial action. Furthermore, growth inhibition of all bacterial species by the *L. longifolium*, *L. petersonnii* and *M. alternifolia* extracts was still significantly inhibited by the end of the 6 h time course study. This may indicate that these extracts have bactericidal activity against these bacteria at the doses tested.

Quantification of toxicity

The toxicity of the *L. longifolium*, *L. petersonnii* and *M. alternifolia* extracts was initially tested in the *Artemia franciscana* nauplii bioassay at a concentration of 2000 μ g/mL (Figure 5). All extracts induced low levels of mortality at 24 and 48 h, similar to the % mortality seen for the seawater control. As none of the extracts induced $> 50\%$ mortality, all were deemed to be nontoxic. Extracts with an LC_{50} of greater than 1000 μ g/mL towards *Artemia* nauplii have previously been defined as being nontoxic.³² In contrast, the potassium dichromate positive controls induced mortality within 4 h (results not shown), with 100% mortality induction seen by 24 h.

DISCUSSION

Plant derived remedies are becoming increasingly sought after in the treatment of a myriad of diseases and disorders due both to their perception of greater safety than synthetic drugs, and the failure of current drug regimens to effectively treat many diseases. Our study examined the ability of methanolic *L. longifolium*, *L. petersonnii* and *M. alternifolia* leaf extracts to inhibit the growth of a panel of medically important bacterial pathogens, and on their toxicity. The gram positive and gram negative bacteria tested in this study were susceptible to all of

Table 1: The mass of dried extracted material, the concentration after resuspension in deionised water and qualitative phytochemical screenings of the methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* extracts

Plant Species	Mass of Dried Extract (mg)	Resuspended Extract (mg/mL)	Phytochemical Analysis Results																
			Phenols			Cardiac Glycosides	Saponins	Triterpenes	Phytosteroids	Alkaloids	Flavanoids	Tannins	Anthraquinones						
			Total Phenolics	Water Soluble	Water Insoluble														
			Keller-Kiliani Test	Froth Persistence	Emulsion test	Salkowski Test	Acetic Anhydride Test	Meyers Test	Wagners Test	Draggendorffs Test	Shinoda Test	Kumar test	Ferric Chloride Test	Lead Acetate Test	Free	Combined			
<i>L. longifolium</i>	130	13	+++	+++	+++	-	+++	+	-	+	+	+	+	+	+	+	-		
<i>L. petersonii</i>	117	11.7	+++	+++	+++	-	+	+	+	+	-	-	-	++	++	+++	-		
<i>M. alternifolia</i>	93	9.3	+++	+++	+++	-	+++	+	+	++	++	+	+	+	++	+++	-		

+++ indicates a large response; ++ indicates a moderate response; + indicates a minor response; - indicates no response in the assay.

Table 2: Minimum bacterial growth inhibitory concentration ($\mu\text{g/mL}$) of the *L. longifolium*, *L. petersonii* and *M. alternifolia* extracts against susceptible bacterial species

Bacterial species	<i>L. longifolium</i>	<i>L. petersonii</i>	<i>M. alternifolia</i>
Gram negative			
<i>A. faecalis</i>	1438	849	2600
<i>A. hydrophilia</i>	147	279	705
<i>C. freundii</i>	619	720	1350
<i>E. coli</i>	-	-	815
<i>K. pneumoniae</i>	320	687	-
<i>P. mirabilis</i>	485	648	449
<i>P. fluorescens</i>	-	-	-
<i>S. newport</i>	-	-	-
<i>S. marcescens</i>	-	-	-
<i>S. sonnei</i>	-	-	-
Gram positive			
<i>B. cereus</i>	755	284	1500
<i>S. aureus</i>	428	389	2842
<i>S. epidermidis</i>	978	846	2978
<i>S. pyogenes</i>	1230	1084	1486

Numbers indicate the mean MIC values of triplicate determinations. - indicates no inhibition.

the extracts, although a greater susceptibility was noted for gram positive bacterial species. Many previous studies with other plant species have also reported a greater susceptibility of gram positive bacteria towards solvent extracts for South American,³⁴ African³⁵ and Australian plant extracts.^{36,37}

The *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts were particularly potent inhibitors of *P. mirabilis* (MIC values $<500 \mu\text{g/mL}$). As *P. mirabilis* infection is a common cause of urinary tract infections and has also been identified as a trigger of rheumatoid arthritis,^{38,39} the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts have potential for the prevention of these diseases in genetically susceptible individuals. The *L. longifolium* and *L. petersonii* extracts were also potent *K. pneumoniae* growth inhibitors. As *K. pneumoniae* can trigger

ankylosing spondylitis in genetically susceptible individuals^{40,41} this extract may also be useful in the prevention and treatment of this autoimmune disease. Furthermore, all extracts were moderate inhibitors of *S. pyogenes* growth with MIC values 1000-1500 $\mu\text{g/mL}$. *S. pyogenes* may cause a myriad of diseases including streptococcal pharyngitis, impetigo and rheumatic heart disease, depending on which tissue it infects.^{42,43} Thus, the methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts may also be useful in the prevention and treatment of these diseases.

Aside from inhibition of the growth of the bacterial triggers of the autoimmune disease discussed above, the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts were also moderate to good inhibitors of several other bacterial pathogens. All extracts inhibited *A. faecalis*, *A. hydrophilia*, *C. freundii* and *B. cereus* growth, with MICs generally $<1000 \mu\text{g/mL}$. Thus, these extracts have potential in the treatment diseases caused by these pathogens (e.g. food poisoning, diarrhoea and dysentery). The *L. longifolium* and *L. petersonii* extracts were also good inhibitors of *S. aureus* and *S. epidermidis* growth (MICs 400-1000 $\mu\text{g/mL}$). The *M. alternifolia* extract was a moderate inhibitor of the growth of these bacteria, with MICs $<3000 \mu\text{g/mL}$. As both of these bacteria are skin disease pathogens, these extracts also may have applications as topical treatments of these diseases.

Whilst an investigation of the phytochemistry of the methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts was beyond the scope of our study, previous studies have reported high terpenoid contents for essential oils of these species.^{7,15} Similarly, other plants of the family *Myrtaceae* are also well known for their high terpenoid contents.⁷ Monoterpenes have been reported to exert a wide variety of biological effects including antibacterial, antifungal, anti-inflammatory and anti-tumour activities²² and therefore may contribute to the bacterial growth inhibitory activity of the *L. longifolium*, *L. petersonii* and *M. alternifolia* extracts reported in our study. A wide variety of monoterpenoids including camphor, carvone, cineole, borneol, menthone, pinene, terpinene, as well as their derivatives, inhibit the growth of an extensive panel of pathogenic and food spoilage bacteria.⁴⁴ Interestingly, several of these monoterpenoids have also been reported to suppress NF- κ B signaling (the major regulator of inflammatory diseases).⁴⁵⁻⁴⁸ This may be particularly relevant for the extracts which inhibited *P. mirabilis* (a bacterial trigger of rheumatoid arthritis)^{38, 39} and *K. pneumoniae* (a trigger of ankylosing spondylitis).^{40,41} The terpene components in

Figure 1: (a) *L. longifolium*, (b) *L. petersonii*, (c) *M. alternifolia*, (d) geraniol, (e) neral, (f) citronellal, (g) α-pinene, (h) sabinene, (i) α-terpinene, (j) limonene, (k) 1,8-cineole.

Figure 2: Growth inhibitory activity of the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts against gram negative bacterial species and the ampicillin (10 µg) and chloramphenicol (2 µg) controls. All determinations were performed in at least triplicate and the results are expressed as mean zones of inhibition (mm) ± SEM.

these extracts may therefore have a pleuripotent mechanism in blocking the autoimmune inflammatory diseases and relieving its symptoms by acting on both the initiator and downstream inflammatory stages of the disease. Further phytochemical evaluation studies and bioactivity driven isolation of active components is required to further evaluate the mechanism(s) of bacterial growth inhibition.

Another commonality between the inhibitory methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts was that all contained high levels of tannins (Table 1). Many studies have reported potent growth inhibitory activities for a number of tannin compounds. Gallotannins have been reported to inhibit the growth of a broad spectrum of bacterial species⁴⁹ through a variety of mechanisms including binding cell surface molecules including lipoteichoic acid and proline-rich cell surface proteins,^{50,51} and by inhibiting glucosyltransferase enzymes.⁵² Ellagitannins are also highly potent inhibitors of bacterial growth, with MIC values as low as 62.5 µg/mL.^{49,51} Ellagitannins have also been reported to function via several antibiotic mechanisms including interaction with cytoplasmic oxidoreductases and by disrupting bacterial cell walls.^{49,51} Thus, it is likely that multiple compounds within the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts may contribute to the inhibition of bacterial growth.

The findings reported here also demonstrate that the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts were nontoxic towards *Artemia franciscana* nauplii, with LC₅₀ values substantially > 1000 µg/mL. Extracts with LC₅₀ values > 1000 µg/mL towards *Artemia* nauplii are defined as being nontoxic.³² Whilst our preliminary toxicity studies indicate that these extracts may be safe for therapeutic use, studies using human cell lines are required to further evaluate the safety of these extracts. Furthermore, whilst these studies have demonstrated the potential of the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf

extracts in the development of future antibiotic chemotherapeutics for the prevention and treatment of urinary tract infections, autoimmune diseases (particularly rheumatoid arthritis and ankylosing spondylitis) and some skin diseases, more work is required to isolate the inhibitory components and determine the mechanism(s) of inhibition.

CONCLUSION

The results of this study demonstrate the potential of the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts as inhibitors of pathogenic bacteria growth. Furthermore, their lack of toxicity indicates that they may be safe for internal as well as topical treatment in the doses tested. Further studies aimed at the purification and identification of bioactive components are needed to examine the mechanisms of action of these agents.

ACKNOWLEDGEMENTS

The authors are grateful to Philip Cameron for providing the *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf leaves used in this study, and to Michelle Mendell and Jane Gifkins for the gift of the bacterial strains. Financial support for this work was provided by the Environmental Futures Research Institute and the School of Natural Sciences, Griffith University, Australia.

CONFLICTS OF INTEREST

The authors report no conflicts of interest.

ABBREVIATIONS USED

DMSO: Dimethyl sulfoxide; **LC₅₀:** The concentration required to achieve 50 % mortality; **MIC:** minimum inhibitory concentration.

Figure 4: Bacterial growth curves for (a) *A. hydrophilia*, (b) *C. freundii*, (c) *P. mirabilis*. All bioassays were performed in at least triplicate and are expressed as mean $\pm$ SEM. * = growth results in the presence of *L. longifolium* extract that are significantly different to the untreated control growth ($p < 0.01$); # = growth results in the presence of *L. petersonii* extract that are significantly different to the untreated control growth ($p < 0.01$); ^ = growth results in the presence of *M. alternifolia* extract that are significantly different to the untreated control growth ($p < 0.01$).

Figure 5: The lethality of the aqueous and methanolic *L. longifolium*, *L. petersonii* and *M. alternifolia* leaf extracts (2000 µg/mL), potassium dichromate (1000 µg/mL) and a seawater control. Blue bars represent the % mortality following 24 h exposure to the extract/toxin. Green bars represent the % mortality following 48 h exposure to the extract/toxin. NC=negative (seawater) control; PC=positive control (1000 µg/mL potassium dichromate). All bioassays were performed in at least triplicate and are expressed as mean ± SEM.

REFERENCES

- Cowan MM. Plant products as antibacterial agents. *Clinical Microbiology Reviews*. 1999;12(4):564-82.
- Bhavnani SM, Ballow CH. New agents for Gram-positive bacteria. *Current Opinion in Microbiology*. 2000;3(5):528-34.
- Chiariandy CM, Seaforth CE, Phelps RH. Screening of medicinal plants from Trinidad and Tobago for antimicrobial and insecticidal properties. *Journal of Ethnopharmacology*. 1999;64(3):265-70.
- Patwardhan B, Warude D, Pushpangadan P. Ayurveda and traditional Chinese medicine: a comparative overview. Evidence-based Complimentary and Alternative Medicine. 2005;2(4):465-73.
- Hostettmann K, Marston A, Ndjoko K. The potential of African plants as a source of drugs. *Current Organic Chemistry*. 2000;4(10):973-1010.
- Paz EA, Cerdeiras MP, Fernandez J. Screening of Uruguayan medicinal plants for antimicrobial activity. *Journal of Ethnopharmacology*. 1995;45(1):67-70.
- Cock IE. Medicinal and aromatic plants—Australia. In *Ethnopharmacology, Encyclopedia of Life Support Systems (EOLSS)*, 2011. Developed under the auspices of UNESCO. Oxford, UK: EOLSS Publishers; 2011. Available from: <http://www.eolss.net>. Accessed 1 April 2013.
- Cock IE. Antimicrobial activity of *Callistemon citrinus* and *Callistemon salignus* methanolic extracts. 2012 *Pharmacognosy Communications*. 2012;2(3):50-7. DOI: 10.5530/pc.2012.3.11
- Wright MH, Matthews B, Arnold MSJ *et al*. The prevention of fish spoilage by high antioxidant Australian culinary plants: *Shewanella putrefaciens* growth inhibition. *International Journal of Food Science and Technology* 2016; DOI: 10.1111/ijfs.13026
- Cock IE. Antimicrobial activity of *Leptospermum bracteata* and *Leptospermum juniperium* methanolic extracts. *Pharmacognosy Communications*. 2013; 3(3):45-52. DOI: 10.5530/pc.2013.3.9
- Cock IE. Antibacterial activity of selected Australian plant species. *The Internet Journal of Microbiology*. 2008;6:2.
- Sautron C, Cock IE. Antimicrobial activity and toxicity of *Syzygium australe* and *Syzygium leuhmannii* fruit extracts. *Pharmacognosy Communications*. 2014;4(1):53-60. DOI: 10.5530/pc.2014.1.8
- Chikowe G, Mpala L, Cock IE. Antibacterial activity of selected Australian *Syzygium* species. *Pharmacognosy Communications*. 2013;3(4):77-83. DOI: 10.5530/pc.2013.4.11
- Cock IE. Antimicrobial activity of *Syzygium australe* and *Syzygium leuhmannii* methanolic extracts. *Pharmacognosy Communications*. 2012;2(2):71-7. DOI: 10.5530/pc.2012.2.11
- Webb MA. *Bush Sense. Australian Essential oils and aromatic compounds*. Griffin Press, 2000, Adelaide Australia.
- Opdyke DLJ. *Citral. Monographs on fragrance raw materials*. Food and Cosmetics Technology. 1976;14:615.
- Wilkinson JM, Hipwell M, Ryan T. Bioactivity of *Backhousia citriodora*: antibacterial and antifungal activity. *Journal of Agriculture and Food Chemistry*. 2003;51(1):76-81.
- Inouye S, Takizawa T, Yamaguchi H. Antimicrobial activity of essential oils and their major constituents against respiratory tract pathogens by gaseous contact. *Journal of Antimicrobial Chemotherapy*. 2001;47(5):565-73.
- Kim JM, Marshall MR, Cornell JA. Antibacterial activity of carvacrol, citral and geraniol against *Salmonella typhimurium* in culture media and in fish cubes. *Journal of Food Science*. 1995;60(6):1364-8.
- Griffin SG, Markham JL, Leach DN. An agar dilution method for the determination of the minimum inhibitory concentration of essential oils. *Journal of Essential Oil Research*. 2000;12(2):249-55.
- Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. *Journal of Applied Microbiology*. 1999;86(6):985-90.
- Cock IE. The phytochemistry and chemotherapeutic potential of *Tasmannia lanceolata* (Tasmanian pepper): A review. *Pharmacognosy Communications*. 2013;3(4):1-13. DOI: 10.5530/pc.2013.4.3
- Kalt FR, Cock IE. Gas chromatography-mass spectroscopy analysis of bioactive *Petalostigma* extracts: Toxicity, antibacterial and antiviral activities. *Pharmacognosy Magazine*. 2014;10(37 Suppl):S37-S49. DOI: 10.4103/0973-1296.127338
- Vesoul J, Cock IE. The potential of Bunya nut extracts as antibacterial functional foods. *Pharmacognosy Communications*. 2012;2(1):72-9. DOI: 10.5530/pc.2012.1.3
- Hart C, Ilanko P, Sirdaarta J, *et al*. *Tasmannia stipitata* as a functional food/natural preservative: Antimicrobial activity and toxicity. *Pharmacognosy Communications*. 2014;4(4):33-47. DOI: 10.5530/pc.2014.4.4
- Winnett V, Boyer H, Sirdaarta J, *et al*. The potential of *Tasmannia lanceolata* as a natural preservative and medicinal agent: Antimicrobial activity and toxicity. *Pharmacognosy Communications*. 2014;4(1):42-52. DOI: 10.5530/pc.2014.1.7
- Arkhipov A, Sirdaarta J, Rayan P, *et al*. An examination of the antibacterial, antifungal, anti-Giardial and anticancer properties of *Kigelia africana* fruit extracts. *Pharmacognosy Communications*. 2014;4(3):62-76. DOI: 10.5530/pc.2014.3.7

28. Courtney R, Sirdaarta J, Matthews B, *et al.* Tannin components and inhibitory activity of Kakadu plum leaf extracts against microbial triggers of autoimmune inflammatory diseases. *Pharmacognosy Journal*. 2015;7(1):18-31. DOI: 10.5530/pj.2015.72
29. Sirdaarta J, Matthews B, Cock IE. Kakadu plum fruit extracts inhibit growth of the bacterial triggers of rheumatoid arthritis: Identification of stilbene and tannin components. *Journal of Functional Foods*. 2015;17:610-20. DOI: 10.1016/j.jff.2015.06.019
30. Cock IE. Antimicrobial activity of *Acacia aulacocarpa* and *Acacia complanta* methanolic extracts. *Pharmacognosy Communications*. 2012;2(1):66-71. DOI: 10.5530/pc.2012.1.12
31. Ruebhart DR, Wikramasinghe WA, Cock IE. Protective efficacy of the antioxidants vitamin E and Trolox against *Microcystis aeruginosa* and microcystin-LR in *Artemia franciscana* nauplii. *Journal of Toxicology and Environmental Health Part A*. 2009;72(24):1567-75.
32. Cock IE, Ruebhart DR. Comparison of the brine shrimp nauplii bioassay and the ToxScreen-II test for the detection of toxicity associated with *Aloe vera* (*Aloe barbadensis* Miller) leaf extract. *Pharmacognosy Research*. 2009;1(2):98-101.
33. Cock IE. Assessment of the toxicity of selected Australian native plant extracts using the *Artemia franciscana* nauplii bioassay. *The Internet Journal of Toxicology*. 2008;5:2.
34. Mohanty S, Cock IE. Evaluation of the antibacterial activity of *Mycaria cauliflora* methanolic leaf and fruit extracts. *The Internet Journal of Microbiology*. 2009;7(2).
35. Cock IE, van Vuuren SF. South African food and medicinal plant extracts as potential antimicrobial food agents. *Journal of Food Science and Technology*. 2015;52(11):6879-99. DOI: 10.1007/s13197-015-1806-3
36. Cock IE, Kukkonen L. An examination of the medicinal potential of *Scaevola spinescens*: Toxicity, antibacterial, and antiviral activities. *Pharmacognosy Research*. 2011;3(2):85-94. DOI: 10.4103/0974-8490.81955
37. Vesoul J, Cock IE. An examination of the medicinal potential of *Pittosporum phylliraeoides*: Toxicity, antibacterial, and antifungal activities. *Pharmacognosy Communications*. 2011;1(2):8-17. DOI: 10.5530/pc.2011.2.1
38. Cock IE, van Vuuren SF. Anti-*Proteus* activity of some South African plants: Their potential for the prevention of rheumatoid arthritis. *Inflammopharmacology*. 2014;22(1):23-36. DOI: 10.1007/s10787-013-0179-3
39. Ebringer A, Rashid T. Rheumatoid arthritis is an autoimmune disease triggered by *Proteus* urinary tract infection. *Clinical and Developmental Immunology*. 2006;13(1):41-8.
40. Cock IE, van Vuuren SF. The potential of selected South African plants with anti-*Klebsiella* activity for the treatment and prevention of ankylosing spondylitis. *Inflammopharmacology*. 2015;23(1):21-35. DOI: 10.1007/s10787-014-0222-z
41. Ebringer A, Rashid T, Wilson C. Ankylosing spondylitis as an auto-immune disease linked to intestinal *Klebsiella* infection: prospects for a new therapeutic approach. *Current Rheumatology Reviews*. 2006;2:55-68.
42. Gerber MA, Baltimore RS, Eaton CB. Prevention of rheumatic fever and diagnosis and treatment of acute Streptococcal pharyngitis. A scientific statement from the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee of the Council on Cardiovascular Disease in the Young, the Interdisciplinary Council on Functional Genomics and Translational Biology, and the Interdisciplinary Council on Quality of Care and Outcomes Research: Endorsed by the American Academy of Pediatrics. *Circulation*. 2009;119(11):1541-51.
43. Wright MH, Arnold MSJ, Lee CJ, *et al.* Qualitative phytochemical analysis and antibacterial activity evaluation of Indian *Terminalia* spp. against the pharyngitis causing pathogen *Streptococcus pyogenes*. *Pharmacognosy Communications*. 2016;6(2):85-92. DOI: 10.5530/pc.2016.2.6
44. Bakkali F, Averbeck S, Averbeck D. Biological effects of essential oils – A review. *Food and Chemical Toxicology*. 2008;46(2):446-75.
45. Salminen A, Lehtonen M, Suuronen T. Terpenoids: Natural inhibitors of NF- κ B signalling with anti-inflammatory and anticancer potential. *Cell and Molecular Life Sciences*. 2008;65(19):2979-99.
46. Lu XG, Zhan LB, Feng BA. Inhibition of growth and metastasis of human gastric cancer implanted in nude mice by d-limonene. *World Journal of Gastroenterology*. 2004;10(14):2140-4.
47. Crowell PL. Prevention and therapy of cancer by dietary monoterpenes. *Journal of Nutrition*. 1999;129(3):775S-8S.
48. Zhou JY, Tang FD, Mao GG. Effect of α -pinene on nuclear translocation of NF- κ B in THP-1 cells. *Acta Pharmacologica Sinica*. 2004;25(4):480-4.
49. Buzzini P, Arapitsas P, Goretti M. Antimicrobial activity of hydrolysable tannins. *Mini-Reviews in Medicinal Chemistry*. 2008;8:1179-87.
50. Wolinsky LE, Sote EO. Isolation of natural plaque-inhibiting substances from 'Nigerian chewing sticks'. *Caries Research*. 1984;18(3):216-25.
51. Hogg SD, Embury G. Blood-group-reactive glycoprotein from human saliva interacts with lipoteichoic acid on the surface of *Streptococcus sanguis* cells. *Archives in Oral Biology*. 1982;27(3):261-8.
52. Wu-Yuan CD, Chen CY, Wu RT. Gallotannins inhibit growth, water-soluble glucan synthesis, and aggregation of *Streptococci mutans*. *Journal of Dental Research*. 1988;67(1):51-5.

PICTORIAL ABSTRACT

SUMMARY

- *L. longifolium*, *L. petersonii* and *M. alternifolia* extracts displayed broad spectrum antibacterial activity against gram positive and gram negative bacteria.
- All extracts were potent inhibitors of *P. mirabilis* growth, with MICs generally <500 µg/mL.
- *A. hydrophila*, *C. freundii*, and *B. cereus* were also particularly susceptible (MICs as low as 147 µg/mL)
- All *L. longifolium*, *L. petersonii* and *M. alternifolia* extracts were nontoxic.

ABOUT AUTHORS

Ms Lindiwe Mpala completed at BSc at Griffith University in life sciences. Following graduation, she undertook a research project in Dr Ian Cock's laboratory in the School of Natural Sciences at Griffith University. The project examined the growth inhibitory properties of a variety of Australian native plants against an extensive panel of bacterial pathogens.

Ms Getmore Chikowe completed at BSc at Griffith University in life sciences. Following graduation, she undertook a research project in Dr Ian Cock's laboratory in the School of Natural Sciences at Griffith University. The project examined the growth inhibitory properties of a variety of Australian native plants against an extensive panel of bacterial pathogens.

Dr Ian Cock leads a research team in the Environmental Futures Research Institute and the School of Natural Sciences at Griffith University, Australia. His research involves bioactivity and phytochemical studies into a variety of plant species of both Australian and international origin, including *Aloe vera*, South Asian and South American tropical fruits, as well as Australia plants including *Scaevola spinescens*, *Pittosporum phylliraeoides*, *Terminalia ferdinandiana* (Kakadu plum), Australian *Acacias*, *Syzygiums*, *Petalostigmas* and *Xanthorrhoea johnsonii* (grass trees). This range of projects has resulted in nearly 200 publications in a variety of peer reviewed journals.