The limit to distribution of a rainforest marsupial folivore is consistent with thermal intolerance

Andrew K. Krockenberger^{1*}

William Edwards¹

John Kanowski²

¹Centre for Tropical Biodiversity and Climate Change Research,

School of Marine and Tropical Biology

James Cook University

P.O. Box 6811 Cairns, QLD 4870

² Australian Wildlife Conservancy, North-east Australia, and

Environmental Futures Centre, Griffith University, Nathan, Qld 4111, Australia

*Author for correspondence:

A.K.Krockenberger

Ph: +61 7 4042 1238

Fax: +61 7 4042 1319

Email: Andrew.Krockenberger@jcu.edu.au

Running head: Thermal intolerance in a marsupial folivore

Abstract

Correlative models of impacts of climate change on species are generally based on correlations between current distributions and climatic parameters, rather than on a detailed understanding of the ecological factors and processes that actually limit distribution. Many of the vertebrates endemic to the rainforests of north-east Australia are restricted to upland forests and considered threatened by climate change. However, for most of these species, the factors controlling distribution are unknown. We examined the role of thermal intolerance as a possible limit to distribution of *Pseudochirops archeri* (green ringtail possum), a specialist arboreal folivore restricted to rainforests above 300 m altitude in Australia's Wet Tropics. We measured short-term metabolic responses to a range of ambient temperatures and found that *P. archeri* stores heat when ambient temperatures exceed 30°C, reducing water requirements for evaporative cooling. Due to the rate at which body temperature increases with ambient temperatures >30 °C, this strategy is not effective over periods longer than around 6 h. We hypothesise that the distribution of *P. archeri* is limited by interactions between (i) the duration and severity of extreme temperatures (over 30°C), (ii) a scarcity of free water in the rainforest canopy in the dry season, and (iii) constraints on water intake from foliage imposed by plant secondary metabolites and fibre. We predict that dehydration becomes biologically limiting for *P. archeri* where extreme temperatures (>30°C) persist for more than 6 h per day over 4-6 days or more. We also predict this mechanism to be most pronounced in lactating females, who have the highest intake requirements of any individuals in the population. Consistent with our hypothesis, the abundance of *P. archeri* in the field is correlated with the occurrence of extreme temperatures, declining markedly at sites where the

average maximum temperature of the warmest week of the year is above 30°C.

Assuming the mechanism of limitation is based on extreme temperatures, we expect impacts of climate change on *P. archeri*, to occur in discrete, rapid events rather than as a slow contraction in range.

Keywords: climate change, limits to distribution, metabolic rate, plant secondary metabolites, temperature extremes, water availability

Introduction

Worldwide, animals and plants restricted to cool upland habitats are considered particularly vulnerable to the effects of climate change (Thomas et al. 2004; Pounds et al. 2006). In north-east Australia, for example, a 1°C increase in average temperature is predicted to reduce the extent of environments presently supporting high elevation rainforests by 40-50% (Hilbert et al. 2001), while a 3.5°C increase is predicted to eliminate the core climatic habitat of 30 of 65 vertebrates endemic to the region and reduce the core climatic habitat of the remaining species by nearly 90% (Williams et al. 2003). However, most of these predictions are based on correlations between current distributions and climatic parameters, rather than on a detailed understanding of the ecological factors and processes that actually limit distribution (Berteaux et al. 2006).

The vertebrates endemic to the rainforests of north-eastern Australia include four species of ringtail possum (Pseudocheiridae). All are restricted to upland forests, have small geographical ranges and patterns of distribution that reflect alternate contraction and expansion of rainforest habitat through the Pleistocene (Winter 1997). A plausible explanation for the upland distribution of these possums is that they are adapted to a cool climate (Nix 1991; Winter 1997). Several alternative hypotheses have recently been dismissed, including altitudinal variation in the floristic composition of forests, foliar chemistry and predation (Kanowski et al. 2003; Kanowski 2004). Briefly, many of the plant taxa known to be important in the diet of the ringtail possums occur in lowland forests; none of the known or likely predators of the ringtail possums are restricted to lowland forests; and foliar chemistry (defined in terms of nutrients such as nitrogen, or tannins) does not explain altitudinal variation in possum abundance.

In this study, we examine the hypothesis that the upland distribution of rainforest ringtail possums reflects their intolerance of hot temperatures experienced in the lowlands. Specifically, we examine (i) the metabolic responses of individuals of *Pseudocheirus archeri* Collett (green ringtail possum) to a range of temperatures, particularly to short term periods of high temperature; and (ii) the relationship between the abundance of *P. archeri* and the occurrence of high temperatures throughout its range. *P. archeri* is a specialist arboreal folivore restricted to rainforests above 300 m elevation in the Australian Wet Tropics (Winter 1997; Jones et al. 2006). Unlike other Australian rainforest ringtail possums, *P. archeri* does not den in hollow trees or dreys (nests), but spends daylight hours curled up on a branch in the canopy (Winter et al. 2008), and hence is exposed to temperatures prevailing in the forest canopy. Having determined the metabolic responses of green ringtail possums to variation in temperature and the relationship between their distribution and occurrence of extreme temperature, we proceed to develop a hypothesis for the mechanism of intolerance of temperature extremes, linking limits on intake imposed by plant secondary metabolites with demands for evaporative cooling during extreme temperature events.

Materials and methods

METABOLIC RESPONSES TO TEMPERATURE

Seven individuals of *P. archeri* were collected during the dry season (July to October) from two sites in complex notophyll vine forest at 740-780 m elevation on the Atherton Tableland, north Queensland, Australia (17°15'S, 145°30'E). Possums were located at night (19:00-24:00) using spotlights from the edge of the forest, and captured using a gas-powered tranquiliser dart-gun (Montech Black Wolf Tranquil

Arms Company, Vic. Australia) and 0.5 ml darts (Minidarts; Tranquil Arms Company, Vic. Australia) containing 30 mg Zoletil (Zolazepam and Tiletamine 50:50; Virbac Pty Ltd). Tranquillised possums were weighed and transported 5–10 km to the laboratory where they were held overnight in a small cage (500 X 400 X 400 mm) before measuring metabolism as oxygen consumption. In our previous work with this species we have found that possums are fully alert and active around 2 h after being tranquilised in this way, and, although there have been no pharmacokinetic studies of these drugs in a herbivore or any marsupial, plasma half-lives for clearance of these drugs of between 1 and 5 hours have been reported across a range of mammals (cat, dog, monkey, rat, Lin et al. 1993; polar bear, Semple et al. 2000). Consequently at least 2, and up to 10 plasma half-lives elapsed between capture and energetic measurements, giving us confidence that there was little, if any effect of the tranquilisation on the following measurements. To allow for the possibility that there was some residual effect of tranquiliser on the results, time since tranquilisation was included as a factor in the statistical analysis. The possums were released after dark at the point of capture and no more than 24 h after capture.

Oxygen consumption was measured by positive pressure, flow-through respirometry starting at 07:00 the morning after capture and finishing by 17:00 that day. This is the normal resting phase for these possums. Each possum was held in a metabolism chamber (300 X 300 X 300 mm) containing a branch for a perch over wire mesh and a layer of vegetable oil to prevent evaporation from faeces or urine produced during the measurement. The metabolism chamber was placed inside a temperature-controlled cabinet. The possum's body temperature was measured using a fine (32 gauge) thermocouple (Type T) inserted approximately 50 mm into the rectum and held in place by taping to the tail. Chamber temperature was measured using another

(8 gauge) thermocouple protruding approximately 50 mm into the chamber. These thermocouples exited the metabolism chamber through a gas-tight cable gland to a thermocouple reader (TC-1000; Sable Systems International NV, USA). Possums were allowed to adjust to the chamber for one hour, then exposed for 90 minutes to each of six temperatures sequentially (5, 12, 18, 24, 30 and 35°C). After allowing 60 minutes for metabolism to reach a steady state at each temperature, measurements were taken for the final 30 minutes of exposure to determine the response of possums to that temperature. At the highest temperature (35°C), possums were monitored constantly and removed from the chamber when their body temperature reached 39°C or when they became agitated, to avoid excessive heat stress. In this case, measurements have been averaged from the final 30 minutes of exposure. This never included substantial activity, because possums were either inactive, or became agitated, in which case they were removed from the chamber, with little intermediate activity.

Air was pumped (3-4 l min⁻¹) from outside the laboratory, through a desiccant (Drierite, anhydrous Na₂SO₄, Hammond Drierite Co. OH, USA) then a mass-flow controller/meter (Sidetrak 840L-2-0V1-SV1-E-V1-S1 Sierra Instruments Inc., CA, USA with a Sable Systems Mass Flow Controller v 1.0, Sable Systems International NV, USA), a copper coil within the cabinet to equilibrate its temperature with that of the cabinet, then into the chamber. Excurrent air from the chamber was subsampled (100 ml min⁻¹) via a manifold and passed through a humidity sensor (HMP-45A, Viasala Oy, Finland), a desiccant scrubber (Drierite), then a carbon dioxide scrubber (Ascarite; sodium hydroxide coated silica, Sigma Chemicals USA, followed by desiccant, Drierite, to redry the air) and the oxygen analyser (Sable Systems FC-2 Oxzilla, Sable Systems International NV, USA). The oxygen analyser was set up for

differential gas analysis, continually comparing the excurrent air with incurrent air subsampled after the dessicant but before the flow meter. At the start of each measurement, and periodically (at least every 60 min) a baseline was recorded by manually switching from the excurrent air to a subsample of the incurrent air. Output from all sensors were acquired every 5 s onto a personal computer using a 12-bit analog to digital converter (Lab-NB; National Instruments Corp. TX, USA) and Labhelper software (Warthog Systems; www.warthog.ucr.edu). Oxygen consumption was estimated according to Withers (1977) using LabAnalyst software (Warthog Systems; www.warthog.ucr.edu) and converted to energy equivalents using the value provided by Gessaman and Nagy (1988) for the metabolism of lipid ($19.8 \text{ kJ.l O}_2^{-1}$). The oxygen analyser was standardised with dry, carbon dioxide free air before commencing each day's measurements. The respirometry system was calibrated using the iron-burn method of Young et al. (1984).

Respiration rate and tidal volume were measured by whole body plethysmography according to the method described by Dawson et al. (2000b). Pressure fluctuations caused by warming and wetting of inspired air were measured using a pressure transducer (Sable Systems PT-100, Sable Systems International NV, USA) and acquired at 20-60 hz using the system described above. Volume change equivalents of the pressure fluctuations were calibrated by recording pressure fluctuations associated with injecting a known volume of air (25 ml) into the chamber after each recording of ventilation. The injection procedure was repeated 10 times at rates such that deflection kinetics were similar to those recorded from the possums at that temperature and the mean deflection of those repeats was used in calculations. Ventilatory frequency was determined directly from the average period of pressure

fluctuations recorded and tidal volume (V_T) was estimated using the method of Malan (1973; Equation 6), assuming that lung temperature was equal to body temperature. To determine the influence of temperature on metabolic rate, body temperature, evaporative water loss, respiratory rate, and tidal volume and minute volume we fitted a repeated measures non-linear mixed effects model with temperature, temperature², $\log_e(\text{mass})$, time since tranquilisation, time in chamber and time of day fitted as fixed effects and animal ID treated as a random factor. Temperature² was included based on theoretical expectations that response to varying temperature would not be linear across the measurement range (Withers 1992), which were confirmed via the existence of a parabolic scatter of all response variables against temperature in initial diagnostic plots. For all response variables an autoregressive correlation structure (AR1) was implemented using a dummy time variable to account for potential temporal non-independence. Autoregressive residual correlation was implemented at the lowest level (i.e. to allow for possibility that there were differences in correlation structure between individuals) (Zurr et al. 2009). The order in which the explanatory variables were entered was chosen to reflect our expectation of their biological importance. There are strong theoretical reasons to suggest that temperature, temperature² (because the effect of temperature is curvilinear) and body mass will be influential in determining all response variables (Withers 1992). Thus, these factors were fitted first and in this order. The three remaining explanatory variables were fitted to the model in the following order: time since tranquilisation, time spent in chamber and time of day. For each response variable we began by fitting the full model (including all main effects), and examined the significance of all terms in the overall fit. When main effects were not significant ($P > 0.05$), we removed the term with least explanatory power (the least significant in its contribution to describing the

data) and refitted the model. We repeated this process until all remaining terms were significant and no further terms could be removed (Zurr et al. 2009). We deemed the final result of the sequential term deletion process to represent the minimum adequate model required to describe the data. Model fitting was performed in NMLE package in R (R Development Core Team, 2010). In the case of respiration rate, residuals from the final linear mixed model (dependent on temperature and temperature²) were not normally distributed and we were unable to transform the data to find an appropriate model. Consequently we have analysed the data using a repeated measures analysis of variance with temperature as a repeated measure, followed post-hoc by paired t-tests with Bonferroni corrections. This approach assumes that time itself was not a significant factor, which we feel is reasonable given that it was not significant (Table 1) for any other variable considered.

ABUNDANCE OF POSSUMS IN RELATION TO TEMPERATURE

The abundance of *P. archeri* was surveyed by spotlight at 50 sites in rainforests on the Atherton Tablelands and adjacent lowlands, north Queensland (Kanowski et al. 2001). For each site, an index of the occurrence of extreme temperatures, the average temperature of the hottest week of the year, was estimated from interpolation of climatic data using ANUCLIM software (see Acknowledgements).

Results

METABOLIC RESPONSES TO TEMPERATURE

The seven individuals of *P. archeri* measured in this experiment ranged from 1.04 to 1.55 kg in mass, with a mean mass of 1.17 ± 0.07 (SE) kg. Resting metabolic rate was strongly related to ambient temperature (Table 1; Fig. 1a). The minimum of the

modelled rate of metabolism was 231 kJ d^{-1} at an ambient temperature of 27.1°C and the average mass of 1.17 kg . Below this thermoneutral zone metabolic rate rose (Fig. 1a), indicating thermoregulatory heat production. Body temperature was also dependent on ambient temperature (Table 1; Fig. 1b) and increased significantly from a modelled minimum of 35.8°C at $T_a = 16.6^\circ\text{C}$ to 38°C at $T_a = 35^\circ\text{C}$ (Fig. 1b). Body temperature rose monotonically as a function of the time (up to 145 minutes) the possum was exposed to temperatures greater than 30°C (Fig. 2; Body temperature $[\text{C}] = 36.5 + 0.01439 * \text{Time} [\text{mins}]$; $F_{1,13}=11657$, $P < 0.001$, $r^2= 0.999$). It should be noted that responses after 145 minutes were not measured, and the relationship could change after that time. The sample size represented in this relationship also declined after 85 minutes of exposure as possums either reached body temperatures of 39°C (which was set as a conservative limit for body temperature due to animal welfare considerations) or became agitated and dislodged the thermocouple, necessitating the termination of the experiment.

Evaporative water loss accelerated with increasing ambient temperature above 10°C (Table 1; Fig. 1c). Respiration rate tended to decline slightly as ambient temperature rose, but then rose sharply at higher temperatures (Table 1; Fig. 1d) and tidal volume fell slowly at first, then increasingly rapidly as ambient temperature rose (Table 1; Fig. 1e), and at 35°C was only 37% of the tidal volume at 5°C .

ABUNDANCE OF POSSUMS IN RELATION TO TEMPERATURE

The abundance of *P. archeri* was negatively correlated with the mean maximum temperature of the hottest week (Spearman rank correlation, $r_s = -0.45$, $P = 0.001$: Fig. 3). The relationship was not linear. Possums were on average an order of magnitude more abundant at sites where the mean maximum temperature of the

warmest week was less than 30°C (0.38 ± 0.05 (SE) individuals ha⁻¹) than at sites which experienced higher temperatures (0.03 ± 0.02 (SE) individuals ha⁻¹) (t-test on $\ln(x+1)$ transformed data, P (determined from randomisation) = 0.001).

Discussion

METABOLIC RESPONSES OF *P. ARCHERI* TO TEMPERATURE

At low ambient temperatures, *P. archeri* displayed a predictable mammalian pattern of metabolism (Fig. 1a). The minimum resting metabolic rate in the thermoneutral zone at 27°C was 100–110% of the level expected for a marsupial of the same mass (Dawson and Hulbert 1970; Withers et al. 2000; Withers et al. 2006) and rose rapidly as temperatures dropped below 20°C, indicating that possums were increasing metabolism to maintain their body temperature. This was accompanied by slow deep breathing at low temperatures, with ventilation rate and tidal volume changing relatively little across the range of lower temperatures. At temperatures above 30°C the body temperature of possums rose rapidly. Over the range of ambient temperatures examined, their body temperature rose as a linear function of the time they were exposed to temperatures above 30°C (up to 145 minutes of exposure). This was initially evident as an increase in body temperature at the ambient temperature of 30°C (Fig. 1b). Body temperatures rises of around 4–6°C are sufficient to impair function in most birds and mammals (Cossins and Bowler 1987). Critical thermal maxima vary substantially from species to species, ranging from an LD₅₀ body temperature of 41.7°C in dogs (Adolph 1947) to tolerance of 43°C in Angolan free-tailed bats (Maloney et al. 1999). Using a mid-range estimate of lethal body temperature rise of 5°C (Cossins and Bowler 1987), to 41.5°C, matches the LD₅₀ body temperature of dogs quite closely (Adolph 1947) as well as a single record of lethal

body temperature of 40.9°C in the quokka (Robinson and Morrison 1957), giving us some confidence in adopting 41.5°C as a putative limit in *P. archeri*. Extrapolating from the rate of increase in body temperature (Fig. 2), the possums would reach lethal body temperatures after 6 h of exposure to temperatures over 30°C, unless they substantially increased their evaporative heat loss at those high temperatures.

The response of *P. archeri* to high temperature (a 2.1 °C increase in body temperature with an increase in ambient temperatures from 18–35 °C) is very similar to that reported for another small Pseudocheirid, *Petauroides volans* (greater glider) (a 2.3 °C increase in body temperature over the same range of ambient temperatures: Rubsamen et al. 1984). Withers (1992) suggests that this heat storage is a relatively common characteristic of mammals, which is taken to its extreme in the strategy of ‘adaptive heterothermy’ exhibited by certain large bodied arid-zone herbivores (Schmidt-Nielsen et al. 1957; Bartholomew and Rainy 1971; Mitchell et al. 2002; Ostrowski et al. 2003; Ostrowski and Williams 2006). A number of other small marsupials show a similar rise in body temperature when exposed to high ambient temperatures (although most reach a new stable temperature after about an hour (Robinson and Morrison 1957). Adaptive heterothermy is most strongly exhibited in dehydrated individuals, and the extent of temperature changes seen in *P. archeri* are similar to those found in much larger, dehydrated grey kangaroos (Dawson et al. 2007). The alternative is to maintain body temperature at high ambient temperatures by an increase in evaporative cooling, largely through panting, increased ventilation frequency and reduced tidal volume, and in some species by licking well-vascularised portions of the skin (Needham et al. 1974; Degabriele and Dawson 1979; Dawson et al. 2000a,b; Jessen 2001). Consequently this heterothermy serves to reduce evaporative water loss for cooling as well as heat gain from the environment

(Schmidt-Nielsen et al. 1957; Bartholomew and Rainy 1971; Withers 1992; Mitchell et al. 2002; Ostrowski et al. 2003; Ostrowski and Williams 2006). At the highest ambient temperature measured in this study, individuals of *P. archeri* did increase ventilation frequency and reduce tidal volume, significantly increasing evaporative water loss, but this was insufficient, within the time of the measurements, to maintain a stable body temperature. Individuals were not observed licking their skin in this study, but may have done so if their body temperatures continued to rise. In contrast to possums in this study, Robinson and Morrison (1957) noted that “possums” (*Trichosurus*, *Petaurus* and *Petauroides*) were the most active “coat-licking” marsupials and Rubsamen et al. (1984) reported that *P. volans* intensively licked its hindlimbs and forearms, but only when body temperature exceeded 38°C.

An important caveat to this study is that responses of captive animals may be different to animals under free-ranging conditions. Nychthemeral body temperature cycles of bandicoots vary in timing and intensity between free-ranging and captive individuals, possibly due to differences in the timing of activity (Warnecke et al. 2007). Captive sugar gliders also change their thermal biology, becoming less likely to enter torpor (Geiser et al. 2007). However, in both these cases, the captive animals had been held in captivity for substantial periods and displayed less variability in body temperature than free-ranging animals, making it less likely that the body temperature variability observed in this study were exaggerated due to captivity.

HETEROTHERMY AS A WATER CONSERVATION STRATEGY IN *P.*

ARCHERI

We hypothesize that heterothermy in *P. archeri* is a strategy for minimising water intake requirements during the hot, dry conditions experienced seasonally in the

rainforest canopy, and makes an important contribution to their ability to tolerate temperature extremes. This hypothesis is based on the following premises: (i) free water is seasonally scarce in the canopy of Australian tropical rainforests; (ii) rainforest ringtail possums are likely to have difficulty satisfying their water requirements from foliage alone, particularly if those requirements are elevated by high temperatures, and (iii) the water savings provided by heterothermy are likely to be ecologically significant to *P. archeri*.

Seasonal scarcity of free water in the rainforest canopy

Australia's tropical rainforests are subject to marked seasonality of rainfall. For example, median monthly precipitation in the driest quarter (August-October) is less than 25 mm at Atherton, the closest weather station to the study area (Bureau of Meteorology 2007). For extended periods during the dry season, there may be no free water in the rainforest canopy, apart from dew where atmospheric conditions are favourable, and from mist in forests within the cloud layer (McJannet et al. 2008). Extreme temperatures in Australia's wet tropical rainforests tend to occur during the late dry season before wet season cloud cover increases, particularly during El Nino years with a late onset of the wet season (Tracey 1982). Maximum temperatures also occur mid-afternoon on a diurnal scale and rainforest ringtail possums do not feed during daylight hours (Winter et al. 2008) meaning that possums experience the hottest conditions during the time when water is least readily available to them.

Water intake requirements of rainforest ringtail possums

Observations of water flux in wild Pseudocheirids indicate that all rely on free water to meet part of their water requirements (Foley et al. 1990; Goudberg 1990; Munks and Green 1995). We argue this is also likely to be true for *P. archeri*. An average

size individual (1170 g) of *P. archeri* is predicted to require 172 ml d⁻¹ of water, based on the equation for non-arid zone marsupials presented by Nagy and Bradshaw (2000). (This equation closely predicts water use by two sympatric rainforest ringtails, *Hemibelideus lemuroides* Collett and *Pseudochirulus herbertensis* Collett, within 3 – 12% of documented rates (Goudberg 1990)). To satisfy predicted water requirements from foliage alone, and allowing for the production of metabolic water (around 20 ml d⁻¹ for other Pseudocheirids: Munks and Green 1995, Foley et al. 1990), an average-sized individual of *P. archeri* would have to consume 65–82 g d⁻¹ DW foliage (calculations based on foliar water content of 65–70% for tree species preferred by *P. archeri*: Goudberg 1990; Jones and Krockenberger unpublished data). This predicted food intake rate is higher than published food intake rates for other Pseudocheirids: for example, food intake rates for *P. volans* and *Pseudocheirus peregrinus* (common ringtail possum) are mostly 45–55 g DW kg^{0.75} d⁻¹ (Foley et al. 1990; Marsh et al. 2003b), equivalent to 49–59 g d⁻¹ DW for an average-sized *P. archeri*. However, *P. archeri* has a relatively long distal colon and may have a more efficient water economy than other Pseudocheirids (Crowe and Hume 1997).

Constraints on foliage intake

While the water requirements of *P. archeri* may be able to be satisfied by foliage under ‘normal’ conditions (or at least leave only a small deficit to be obtained from free water), we argue it is unlikely that *P. archeri* could meet elevated water requirements such as those imposed by active thermoregulation or lactation (Green 1997) by increasing foliage intake. Again there are no specific data for *P. archeri*, but studies of other small folivorous mammals have shown that rates of food intake are strongly constrained by the gut-filling effects of fibre (Cork and Foley 1991) and by

various physiological effects of plant secondary metabolites (PSMs) (Torregrossa and Dearing 2009), including toxicity (McLean et al. 2004), acidosis (Foley et al. 1995), the depletion of energy and nutrients used to eliminate PSMs from the body (Foley 1992; Sorensen et al. 2005) and diuresis (Dearing et al. 2001, 2002). Although folivores have evolved sophisticated mechanisms for detoxifying and excreting PSMs, these processes are rate-limited (Foley et al. 1995; Marsh et al. 2006a,b). There is considerable evidence that folivores regulate the intake of PSMs so as not to exceed physiological thresholds (Lawler et al. 2000; Marsh et al. 2003a; Scrivener et al. 2004; Foley and Moore 2005; Moore et al. 2005; Iason and Villaba 2006), and that these constraints on intake can be sufficiently strong to override elevated demands for nutrients and energy (Torregrossa and Dearing 2009), including those imposed by active thermoregulation (Stapley et al. 2000). In this scenario, the most vulnerable individuals in the population are likely to be lactating females, whose food intake would already be the highest of any individuals in the population (Gittleman and Thompson 1988; Krockenberger 2003; Krockenberger and Hume 2007). This conclusion assumes that most, or all, of the extra water intake by lactating females is exported in the milk and thus is not available for evaporative cooling. In common ringtail possums, *Pseudocheirus peregrinus*, this assumption is robust, as most (~86%) of the increased water intake at peak lactation is exported in the milk (Munks 1990; Munks and Green 1995). However, if the assumption does not apply to *P. archeri*, and lactating females increase foliage water intake for the energetic requirements of milk production more than by the amount of the water they export in milk, then lactating females could have an excess of water intake and thus not be as vulnerable to the impacts of heatwave events.

There is no reason to believe that rainforest ringtail possums can escape the constraints on intake discussed above. Correlative studies of diet selection have shown that the rainforest ringtail possums all tend to avoid foliage with high concentrations of condensed tannins (Goudberg 1990); while *P. archeri* in particular avoids individuals of a favoured tree species, *Aleurites rockinghamensis*, with high concentrations of cyanogenic glycosides (Jones and Krockenberger, unpublished data). Compared with sympatric ringtail possums, *P. archeri* selects foliage with a relatively high fibre content (Goudberg 1990), but this does not mean that overall levels of food intake by *P. archeri* are unconstrained by fibre (Cork 1994). Herbivores can attempt to circumvent constraints on fibre intake by selectively feeding on young leaves, which have a higher water content and lower fibre content than mature foliage (Coley and Barone 1996), but young leaves are often strongly defended by low-molecular weight PSMs such as cyanogenic glycosides (Coley and Barone 1996; Miller et al. 2006). For example, *P. archeri* does not eat young leaves of the favoured tree species *A. rockinghamensis* (Procter-Gray 1985; Jones et al. 2006), which have concentrations of cyanogenic glycosides an order of magnitude higher than mature foliage (Jones and Krockenberger unpublished data).

Water conservation provided by heterothermy

We argue that the water savings due to heterothermy over relatively short periods could be significant to water conservation by *P. archeri*, as follows. If an average-sized possum (1.17 kg) stored metabolic heat equivalent to a 5°C rise in body temperature over a 6 h period while exposed to an ambient temperature of 35°C, i.e., close to body temperature and therefore with very slow dry dissipation of heat, 19.9 kJ of heat energy would be stored that otherwise would have taken 8.3 g water to

dissipate by evaporation (assuming heat capacity of tissue = $3.4 \text{ J g}^{-1} \text{ }^{\circ}\text{C}^{-1}$; latent heat of vaporisation of water at 40°C is 2400 J/g Withers 1992). That is, a strategy of maintaining a constant body temperature at an ambient temperature of 35°C would raise evaporative water loss across that 6 hours by 70%, compared with the strategy adopted by *P. archeri* (32 mg/min evaporative water loss at 35°C , see Fig. 1c, while storing heat at 0.86°C/h , see Fig.2, which is a total of 11.69 g water loss over 6 h). In the scenario presented above, maintaining constant body temperature results in an extra loss of 8.3 g of water, or 1 % of the body water content of *P. archeri* (average mass 1170 g and 73.5% body water content; Krockenberger, unpublished data). Most mammals are relatively intolerant to dehydration, with alterations in physiological function noted as low as 4-6 % dehydration (Thibodeau and Patton 2006), so the water conservation represented by heat storage is likely to be significant over an extended period of extreme temperature such as a heat wave of 4-6 days duration. It should be noted that another consequence of elevated temperature is that metabolic heat production rises due to the Q_{10} effect, so body temperature would rise progressively more quickly if evaporative water loss did not increase, although the rise in dry heat loss rate as the temperature gradient between the possum and its environment increased could partially compensate. If a Q_{10} of 2.3 (Dawson et al. 2007) is applied, a possum at a body temperature of 41.5°C would have to dissipate 49% more metabolic heat than at its resting metabolism. This could ultimately shorten the time taken to reach lethal limits unless evaporative heat loss increased to partially compensate. It also means that the greatest water savings occur if the ambient temperature declines before an upper body temperature limit is reached, so that heat can again be shed by dry means. The 49% increase in metabolism due to a rise in body temperature substantially reduces or negates the water savings of the

strategy if the individual is eventually forced to evaporate water to control body temperature (Dawson et al. 2007). Similarly, if our hypothesis of heat storage breaks down at body temperatures above 39°C and *P. archeri* defend a lower body temperature than 41.5°C, the limit due to either dehydration or intoxication that we propose still applies. However, under these circumstances the mechanism would operate at a shorter time threshold, because possums would be forced to substantially increase evaporative water loss to defend that temperature, particularly as metabolic heat production rises (Dawson et al. 2007). The smaller body to ambient temperature gradient would also limit increases in dry heat loss.

In summary, use of heat storage means that *P. archeri* would be tolerant of extreme temperatures that were around 30-35°C for up to 6 h a day without having to markedly increase water intake. If temperatures remained above 30°C for over 6 h a day, however, it is likely that possums could not continue to store heat, but would be forced to increase evaporative cooling. In situations where free water is not available, we argue that *P. archeri* is likely to have difficulty meeting elevated water requirements from foliage alone. In these circumstances, dehydration would become biologically limiting where extreme temperatures (>30°C) persisted for more than 6 h per day over 4-6 days or more. This is concordant with the observed pattern of distribution of *P. archeri* in relation to the occurrence of extreme temperatures (Fig. 3). Given that lactating females have the highest intake of any individuals in the population (Gittleman and Thompson 1988; Krockenberger 2003; Krockenberger and Hume 2007), we expect this effect to be most marked in lactating females (although subject to the caveat discussed above) and may act to reduce reproductive success in marginal habitat.

Limitations on water intake may also account for the upland distribution of other rainforest ringtail possums in north-east Australia, which show similar threshold responses to extreme temperatures (Kanowski 2004), and for the montane distribution of most species of rainforest ringtail possums in New Guinea (Flannery 1994). While folivorous marsupials do occur in the New Guinea lowlands, most are Phalangerids (cuscus), with numerous differences in gut morphology, physiology and breadth of diet from ringtail possums (Crowe and Hume 1997). In particular, phalangerids are more generalised herbivores than ringtail possums, having a greater reliance on fruit and higher rates of food intake, attributes which would allow increased an intake of water in their diet. Folivorous primates, which inhabit lowland rainforests elsewhere in the tropics, also tend to be moderately frugivorous (Montgomery 1978).

Predicted impacts of climate change

If our hypothesis is correct, then predictions of the impacts of climate change on the range of *P. archeri* and perhaps the other rainforest ringtail possums (Williams et al. 2003; Kanowski 2004) are likely to be underestimates, as the correlative models used in these predictions are based largely on climatic averages, rather than extremes. Extreme thermal stresses (heat and cold), along with average summer temperature provided the best descriptors of local extinctions of pikas, *Ochotona princeps*, in America's Great Basin over the last 60 years (Beever et al. 2010). The frequency and severity of extreme temperature events are expected to increase markedly under enhanced greenhouse conditions (IPCC 2007). In Cairns in north-east Australia, for example, the frequency of hot days ($>35^{\circ}\text{C}$) over summer is predicted to increase by 47-150% by 2030 (CSIRO and Bureau of Meteorology 2007). In addition, current predictions do not consider synergistic effects of climate change, such as the lifting of

the cloud base (Still et al. 1999), or possible increases in the production of carbon-based defences of rainforest species under enhanced carbon dioxide concentrations (Kanowski 2001). Because the proposed mechanism limiting the distribution of *P. archeri* involves interactions with plant defenses, elevated levels of PSMs may further constrain foliage intake and exacerbate the effects of high temperatures on water requirements.

Our model provides a testable mechanism linking strongly supported predictions of changes in the physical climate (IPCC 2007) to often poorly understood impacts on fauna (Berteaux et al. 2010; Parmesan 2006). We predict that range contractions of *P. archeri* are likely to follow long dry seasons with periods of extreme temperatures where daily temperatures exceed 30°C for six or more hours over a period of four to six ,or more, consecutive days. Under these conditions we expect populations of *P. archeri* at sites with, or close to, threshold maximum temperatures to decline or disappear. Lastly, because the mechanism is based on extremes rather than slow accumulation of warming we can expect impacts of climate change on *P. archeri* to occur in discrete events rather than as a slow contraction in range. Given the increasing frequency of extremely hot years, we may soon be accorded the opportunity to test these predictions in the field.

Acknowledgements

We would like to thank the Nasser family for permission to study the possums on their property and our volunteers, especially Steve McKenna, for assistance with fieldwork. Thanks to Dr John Winter for discussion and stimulating this work, Dr David Westcott, Dr Chris Margules and CSIRO TFRI Atherton for providing laboratory space for the respirometry and reviewers for comments that helped us to

improve the manuscript. The laboratory measurements were funded by the James Cook University Merit Research Grant Scheme and Rainforest CRC, and ongoing analysis by the Marine and Tropical Sciences Research Facility. The climate data was provided by the Spatial Modelling and Data Exchange project of the Rainforest CRC using the ANUCLIM software and data developed by the Centre for Resource and Environmental Studies at the Australian National University. This study complied with the legal and ethical requirements of the state of Queensland and was conducted under approvals from the JCU Animal Ethics Committee (A856) and Queensland EPA (WISP01557603).

References

- Adolph EF (1947) Tolerance to heat and dehydration in several species of mammals. *American Journal of Physiology* 151: 564-575.
- Bartholomew GA, Rainy M (1971) Regulation of body temperature in the rock hyrax, *Heterohyrax brucei*. *Journal of Mammalogy*, 52: 81-95.
- Beever EA, Ray C, Mote PW, Wilkening JL (2010) Testing alternative models of climate-mediated extirpations. *Ecological Applications* 20: 164-178.
- Berteaux D, Humphries MM, Krebs CJ, Lima M, McAdams AG, Pettorelli N, Réale D, Saitoh T, Tilkadlec E, Weladji RB, Stenseth NChr (2006) Constraints to projecting the effects of climate change on mammals. *Climate Research* 32: 151-158.
- Bureau of Meteorology (2007) Climate averages: Atherton Post Office.
www.bom.gov.au/climate/averages/tables/cw_031002.shtml
- Coley PD, Barone JA (1996) Herbivory and plant defences in tropical forests. *Annual Review of Ecology and Systematics* 27: 305-335.

- Cork SJ (1994) Digestive constraints on dietary scope in small and moderately-small mammals: how much do we really understand? In: Chivers DJ, Langer P (eds) *The Digestive System in Mammals: Food, Form and Function*. Cambridge University Press, Cambridge, pp. 337-369.
- Cork SJ, Foley WJ (1991) Digestive and metabolic strategies of arboreal mammalian folivores in relation to chemical defences in temperate and tropical forests. In: Palo RT, Robbins CT (eds) *Plant Defences Against Mammalian Herbivory*. CRC Press, Boca Raton, Florida, pp. 133-166.
- Cossins AR, K Bowler (1987) *Temperature Biology of Animals*. Chapman and Hall, London
- Crowe O, Hume ID (1997) Morphology and function of the gastrointestinal tract of Australian folivorous possums. *Australian Journal of Zoology* 45: 357-368.
- CSIRO, Bureau of Meteorology (2007) *Climate Change in Australia*. Technical Report. www.climatechangeinaustralia.gov.au
- Dawson TJ, Hulbert AJ (1970) Standard metabolism, body temperature, and surface area of Australian marsupials. *American Journal of Physiology* 218: 1233–1238.
- Dawson TJ, Blaney CE, McCarron HCK, Maloney SK (2007) Dehydration, with and without heat, in kangaroos from mesic and arid habitats: different thermal responses including varying patterns in heterothermy in the field and laboratory. *Journal of Comparative Physiology B* 177: 797-807.
- Dawson T, Blaney C, Munn A, Krockenberger AK, Maloney S (2000a) Thermoregulation by kangaroos from mesic and arid habitats: Influence of temperature on routes of heat loss in eastern grey kangaroos (*Macropus*

- giganteus*) and red kangaroos (*Macropus rufus*). Physiological and Biochemical Zoology 73: 374-381
- Dawson T, Munn A, Blaney C, Krockenberger AK, Maloney S (2000b) Ventilatory accomodation of oxygen demand and respiratory water loss in kangaroos from mesic and arid environments, the eastern grey kangaroo (*Macropus giganteus*) and red kangaroo (*Macropus rufus*). Physiological and Biochemical Zoology 73: 382-388.
- Dearing MD, Mangione AM, Karasov WH (2001) Plant secondary compounds as diuretics: An overlooked consequence. American Zoologist 41: 890–901.
- Dearing MD, Mangione AM, Karasov WH (2002) Ingestion of plant secondary compounds causes diuresis in desert herbivores. Oecologia 130: 576-584.
- Degabriele R, Dawson TJ (1979) Metabolism and heat balance in an arboreal marsupial, the Koala (*Phascolarctos cinereus*). Journal of Comparative Physiology B 134: 293-301.
- Foley WJ (1992) Nitrogen and energy retention and acid-base status in the common ringtail possum (*Pseudocheirus peregrinus*): evidence of the effects of absorbed allelochemicals. Physiological Zoology 65: 403-421.
- Foley WJ, Moore BD (2005) Plant secondary metabolites and vertebrate herbivores – from physiological regulation to ecosystem function. Current Opinion in Plant Biology 8: 430-435.
- Foley WJ, Kehl JC, Nagy KA, Kaplan IR, Borsboom AC (1990) Energy and water metabolism in free-living greater gliders, *Petauroides volans*. Australian Journal of Zoology 38: 1-9.

- Foley WJ, McLean S, Cork SJ (1995) Consequences of biotransformation of plant secondary metabolites on acid-base metabolism in mammals - a final common pathway? *Journal of Chemical Ecology* 21: 721-743.
- Geiser F, Holloway JC, Kortner G (2007) Thermal biology, torpor and behaviour in sugar gliders: a laboratory-field comparison. *Journal of Comparative Physiology B* 177: 495-501.
- Gessaman JA, Nagy KA (1988) Energy metabolism: errors in gas exchange conversion factors. *Physiological Zoology* 61: 507–513.
- Gittleman JL, Thompson SD (1988) Energy allocation in mammalian reproduction. *American Zoologist* 28: 863 – 875.
- Goudberg N (1990) The feeding ecology of three species of north Queensland upland rainforest possums, *Hemibelideus lemuroides*, *Pseudocheirus herbertensis* and *Pseudocheirus archeri* (Marsupialia: Petauridae). PhD thesis, James Cook University, Townsville.
- Green B (1997) Field energetics and water fluxes in marsupials. In: Saunders NR, LA Hinds LA (eds) *Marsupial Biology: Recent Research, New Perspectives*. UNSW Press, Sydney, pp. 143-162.
- Hilbert DW, Ostendorf B, Hopkins MS (2001) Sensitivity of tropical forests to climate change in the humid tropics of north Queensland. *Austral Ecology* 26: 590-603.
- Iason GR, Villalba JJ (2006) Behavioral strategies of mammal herbivores against plant secondary metabolites: The Avoidance–Tolerance Continuum. *Journal of Chemical Ecology* 32: 1115–1132.
- IPCC (2007) *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental

- Panel on Climate Change. Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds). Cambridge University Press, Cambridge.
- Jessen C (2001) Temperature Regulation in Humans and Other Mammals. Springer Verlag, Berlin.
- Jones KM, MacLagan SJ, Krockenberger AK (2006) Diet selection in the green ringtail possum (*Pseudochirops archeri*): a specialist folivore in a diverse forest. *Austral Ecology* 31: 799-807.
- Kanowski J (2001) Effects of elevated CO₂ on the foliar chemistry of seedlings of two rainforest trees from north-east Australia: Implications for folivorous marsupials. *Austral Ecology* 26: 165-172.
- Kanowski J (2004) Ecological determinants of the distribution and abundance of folivorous possums inhabiting rainforests of the Atherton Tablelands, north-east Queensland. In: Goldingay R, Jackson S (eds) *The Biology of Australian Possums and Gliders*. Surrey Beatty and Sons, Chipping Norton NSW, pp. 539-548.
- Kanowski J, Hopkins MS, Marsh H, Winter JW (2001) Ecological correlates of folivore abundance in north Queensland rainforests. *Wildlife Research* 28: 1-8.
- Kanowski J, Irvine AK, Winter JW (2003) The relationship between the floristic composition of rain forests and the abundance of folivorous marsupials in north-east Queensland. *Journal of Animal Ecology* 72: 627-632.
- Krockenberger AK (2003) Meeting the energy demands of reproduction in female koalas, *Phascolarctos cinereus*: evidence for energetic compensation. *Journal of Comparative Physiology B* 173: 531–540

- Krockenberger AK, Hume ID (2007) A flexible digestive strategy accommodates the nutritional demands of reproduction in a free-living folivore, the Koala (*Phascolarctos cinereus*). *Functional Ecology* 21: 748 – 756.
- Lawler IR, Foley WJ, Eschler BM (2000) Foliar concentration of a single toxin creates habitat patchiness for a marsupial folivore. *Ecology* 81: 1327–1338.
- Lin HC, Thurmon JC, Benson GJ, Tranquilli WJ (1993) Telazol- a review of its pharmacology and use in veterinary medicine. *Journal of Veterinary Pharmacology and Therapeutics* 16: 383-418.
- Malan A (1973) Ventilation measured by body plethysmography in hibernating mammals and in poikilotherms. *Respiration Physiology* 17: 32–44.
- Maloney SK, Bronner GN, Buffenstein R (1999) Thermoregulation in the Angolan Free-Tailed Bat *Mops condylurus*: A Small Mammal That Uses Hot Roosts. *Physiological and Biochemical Zoology* 72: 385-396
- Marsh KJ, Foley WJ, Cowling A, Wallis IR (2003a) Differential susceptibility to *Eucalyptus* secondary compounds explains feeding by the common ringtail (*Pseudocheirus peregrinus*) and common brushtail possum (*Trichosurus vulpecula*). *Journal of Comparative Physiology B* 173: 69-78.
- Marsh KJ, Wallis IR, Foley WJ (2003b) The effect of inactivating tannins on the intake of *Eucalyptus* foliage by a specialist *Eucalyptus* folivore (*Pseudocheirus peregrinus*) and a generalist herbivore (*Trichosurus vulpecula*). *Australian Journal of Zoology* 51: 31-42.
- Marsh KJ, Wallis IR, Andrew RL, Foley WJ (2006a) The detoxification limitation hypothesis: where did it come from and where is it going? *Journal of Chemical Ecology* 32: 1247-1266.

- Marsh KJ, Wallis IR, McLean S, Sorenson JS, Foley WJ (2006b) Conflicting demands on detoxification pathways influence how common brushtail possums choose their diets. *Ecology* 87: 2103-2112.
- McJannet D, Wallace J, Fitch P, Disher M, Reddell P (2008) Hydrological processes in the tropical rainforests of Australia. In: Stork NE, Turton SM (eds) *Living in a Dynamic Tropical Forest Landscape*. Blackwell, Oxford, pp. 197-209.
- McLean S, Brandon S, Davies NW, Foley WJ, Muller HK (2004) Jensenone: biological reactivity of a marsupial antifeedant from *Eucalyptus*. *Journal of Chemical Ecology* 30: 19-36.
- Miller RE, Jensen R, Woodrow IE (2006) Frequency of cyanogenesis in tropical rainforests of far north Queensland, Australia. *Annals of Botany* 97: 1017-1044.
- Mitchell D, Maloney SK, Jessen J, Laburn HP, Kamerman PR, Mitchell G, Fuller A (2002) Adaptive heterothermy and selective brain cooling in arid-zone mammals. *Comparative Biochemistry and Physiology B* 131: 571-585.
- Montgomery GG (1978) *The Ecology of Arboreal Folivores*. Smithsonian Institution Press, Washington DC.
- Moore BD, Foley WJ, Wallis IR, Cowling A, Handasyde KA (2005) *Eucalyptus* foliar chemistry explains selective feeding by koalas. *Biology Letters* 1: 64-67.
- Munks, SA (1990). Ecological energetics and reproduction in the common ringtail possum, *Pseudocheirus peregrinus*. Ph.D. Thesis, University of Tasmania, Hobart
- Munks SA, Green B (1995) Energy allocation for reproduction in a marsupial arboreal folivore, the common ringtail possum (*Pseudocheirus peregrinus*). *Oecologia* 101: 94-104.

- Nagy KA, Bradshaw SD (2000) Scaling of energy and water fluxes in free-living arid-zone Australian marsupials. *Journal of Mammalogy* 81: 962-970.
- Needham AD, Dawson TJ, Hales JRS (1974) Forelimb blood flow and saliva spreading in the thermoregulation of the red kangaroo, *Megaleia rufa*. *Comparative Biochemistry and Physiology* 49A: 555–565.
- Nix HA (1991) Biogeography: pattern and process. In: Nix HA, Switzer MA (eds) *Rainforest Animals: Atlas of Vertebrates Endemic to Australia's Wet Tropics*. Australian National Parks and Wildlife Service, Canberra, pp. 11-39.
- Ostrowski S, Williams JB (2006) Heterothermy of free-living Arabian sand gazelles (*Gazella subgutturosa marica*) in a desert environment. *The Journal of Experimental Biology* 209: 1421-1429.
- Ostrowski S, Williams JB, Ismael K (2003) Heterothermy and the water economy of free-living Arabian oryx (*Oryx leucoryx*). *The Journal of Experimental Biology* 206: 1471-1478.
- Parnesan C (2006) Ecological and Evolutionary Responses to Recent Climate Change. *Annual Review of Ecology, Evolution and Systematics* 37: 637-669.
- Pounds JA, Bustamante MR, Coloma LA, Consuegra JA, Fogden MPL, Foster P N, La Marca E, Masters KL, Merino-Viteri A, Puschendorf R, Ron SR, Sánchez-Azofeifa GA, Still CJ, Young BE (2006) Widespread amphibian extinctions from epidemic disease driven by global warming. *Nature* 439: 161-167.
- Procter-Gray E (1985) The behaviour and ecology of Lumholtz's tree-kangaroo, *Dendrolagus lumholtzi* (Marsupialia: Macropodidae). PhD thesis, Harvard University, Cambridge, Massachusetts.

- R Development Core Team (2010) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- Robinson KW, Morrison PR (1957) The reaction to hot atmospheres of various species of Australian marsupial and placental animals. *Journal of Cellular and Comparative Physiology* 49: 455-478.
- Rubsamen K, Hume ID, Foley WJ, Rubsamen U (1984) Implications of the large surface area to body mass ratio on the heat balance of the greater glider (*Petauroides volans*: Marsupialia). *Journal of Comparative Physiology B* 154: 105-111.
- Schmidt-Nielsen K, Schmidt-Nielsen B, Jørgensen SA, Haupt TR (1957) Body temperature of the camel and its relation to water economy. *American Journal of Physiology* 188: 103–112.
- Scrivener NJ, Johnson CN, Wallis IR, Takasaki M, Krockenberger AK (2004) Which trees do wild common brushtail possums (*Trichosurus vulpecula*) prefer? Problems and solutions in scaling laboratory findings to diet selection in the field. *Evolutionary Ecology Research* 6: 77–87.
- Semple HA, Gorecki DK, Farley SD, Ramsay MA (2000) Pharmacokinetics and tissue residues of Telazol in free-ranging polar bears. *Journal of Wildlife Diseases* 36: 653-662.
- Sorensen JS, McLister JD, Dearing MS (2005) Plant secondary metabolites compromise the energy budgets of specialist and generalist mammalian herbivores. *Ecology* 86: 125-139.

- Stapley J, Foley WJ, Cunningham R, Eschler B (2000) How well can common brushtail possums regulate their intake of *Eucalyptus* toxins? Journal of Comparative Physiology B 170: 211-218.
- Still C J, Foster N F, Schneider S H (1999) Simulating the effects of climate change on tropical montane cloud forests. Nature 398: 608–610.
- Thibodeau GA, Patton KT (2006) Anatomy and Physiology 6th edition. Elsevier, Philadelphia.
- Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont L, Collingham YC, Erasmus BFN, Ferreira de Siqueira M, Grainger A, Hannah L, Hughes L, Huntley B, van Jaarsveld AS, Midgley GF, Miles L, Ortega-Huerta MA, Townsend Peterson A, Williams SE (2004) Extinction risk from climate change. Nature 427: 145-148.
- Torregrossa A-M, Dearing MD (2009) Nutritional toxicology of mammals: regulated intake of plant secondary compounds. Functional Ecology 23: 48-56.
- Tracey JG. (1982) The vegetation of the humid tropical region of north Queensland. CSIRO, Melbourne 124pp.
- Warnecke L, Withers PC, Schleucher E, Maloney SK (2007) Body temperature variation of free-ranging and captive southern brown bandicoots *Isodon obesulus* (Marsupialia: Peramelidae). Journal of Thermal Biology 32: 72-77.
- Warthog Systems: data acquisition and analysis software for the Macintosh. Mark Chappell and the Regents of the University of California.
www.warthog.ucr.edu
- Williams SE, Bolitho E, Fox S (2003) Climate change in Australian tropical rainforests: an impending environmental catastrophe. Proceedings of the Royal Society London 270: 1887-1892.

- Winter JW (1997) Responses of non-volant mammals to late Quaternary climatic changes in the Wet Tropics region of north-eastern Australia. *Wildlife Research* 24: 493-511.
- Winter JW, Krockenberger AK, Moore NJ (2008) Green Ringtail Possum, *Pseudochirops archeri*. In: Van Dyke S and Strahan R (eds) *The Mammals of Australia* (), Reed New Holland Pty Ltd, Sydney Pp 245-247.
- Withers PC (1977) Measurement of VO_2 , VCO_2 and evaporative water loss with a flow through mask. *Journal of Applied Physiology* 42: 120–123.
- Withers PC (1992) *Comparative Animal Physiology*. WB Saunders, Sydney.
- Withers PC, Thompson GG, Seymour RS (2000) Metabolic physiology of the north-western marsupial mole, *Notoryctes caurinus* (Marsupialia: Notorycidae). *Australian Journal of Zoology* 48: 241–258.
- Withers PC, Cooper CE, Larcombe AN (2006) Environmental correlates of physiological variables in marsupials. *Physiological and Biochemical Zoology* 79: 437-453.
- Young BA, Fenton TW, McLean JA (1984) Calibration methods in respiratory calorimetry. *Journal of Applied Physiology* 56: 1120–1125.
- Zurr AF, Ieno EN, Walker NJ, Savelier AA, Smith GM (2009). *Mixed Effects Models and Extension in Ecology with R*. Springer Science and Business Media, New York.

Table 1. Significant factors in final non-linear mixed effects model analyses of the effect of temperature on the metabolic physiology of green ringtail possums, *Pseudochirops archeri*. Results are from model fit for seven individual possums measured at 6 temperatures, using an autoregressive correlation structure (CorAR1) with individual treated as a random factor.

Factor	F _{1,33}	P	Estimate	SE	$\hat{d}^a$	ϕ^b
Metabolic rate (kJ/day)						
Intercept	276.38	<0.0001	486.3	24.2	0.0213	0.899
Temperature ²	23.01	<0.0001	0.3461	0.0421		
Temperature	114.36	<0.0001	-18.79	1.76		
Oxygen consumption (ml/min)						
Intercept	276.38	<0.0001	17.06	0.85	0.000395	0.899
Temperature ²	23.01	<0.0001	0.01214	0.00148		
Temperature	114.36	<0.0001	-0.6589	0.0616		
Body temperature (°C)						
Intercept	44421.39	<0.0001	37.50	0.28	0.00004	0.783
Temperature ²	53.47	<0.0001	0.00623	0.00064		
Temperature	60.54	<0.0001	-0.2067	0.0266		
Evaporative water loss (mg/min)						
Intercept	199.94	0.00011	15.07	2.29	2.798	0.665
Temperature ²	132.39	<0.0001	0.03122	0.0051		
Temperature	7.98	0.0079	-0.5958	0.2109		
Tidal volume (ml)						
Intercept	127.56 ^c	<0.0001	9.396	0.732	0.000255	0.755
Temperature ²	40.04 ^c	<0.0001	-0.004715	0.000745		
Minute volume (ml)						
Intercept	322.56	<0.0001	444.9	36.5	31.84	-0.016
Temperature ²	9.27	0.0046	0.3679	0.0943		
Temperature	21.65	0.0001	-18.18	3.91		

^a Estimate of standard deviation of random effect intercept

^b Correlation between sequential measures in an individual

^c DF = 1,34

7 **Figure captions**

8 Fig. 1. The effect of ambient temperature on the metabolic physiology of the green
 9 ringtail possum, *Pseudochirops archeri*. The curves indicate the final mixed effects
 10 models fitted to the data. a) resting metabolic rate (kJ/day), b) body temperature (°C),
 11 c) evaporative water loss (mg/min), d) respiration rate (breaths/min) (means $\pm$ SE, n =
 12 7). Letters indicate statistically similar means.e) respiratory tidal volume (ml).

13 Fig. 2. Body temperatures of green ringtail possums, *Pseudochirops archeri*, at 10
 14 minute intervals from 5 to 145 minutes after they were exposed to ambient
 15 temperatures greater than 30°C, and increasing over time to 35°C (means $\pm$ SE, n = 7
 16 for times from 5 to 84 minutes after temperature of 30 °C was reached; thereafter
 17 sample size is marked above the data point). Prior to the period represented here they
 18 had undergone an acclimatisation period of 90 minutes at 30 °C during measurement
 19 of resting metabolism at that temperature. The linear least-squares line of best fit to
 20 those means is displayed (Body temperature [°C] = 36.5 + 0.01439 * Time [mins];
 21 $F_{1,13}=11657$, $P < 0.001$, $r^2= 0.999$).

22 Fig. 3. Abundance of the green ringtail possum *Pseudochirops archeri* in relation to
 23 the mean maximum temperature of the warmest week, an index of extreme
 24 temperatures, at 50 sites in rainforest on the Atherton Tablelands and adjacent
 25 lowlands.

26

27 Figure 1a

28

29

30 Figure 1b

31

32

33 Figure 1c

34

35

36 Figure 1d

37

38

39 Figure 1e

40

41

Figure 2

46