Title page:

Title: Paediatric ICU nurses' knowledge and practices to prevent central venous access device infections: implications for clinicians

Authors: Amanda J Ullman^{a,b}; Debbie A Long^{a,b}, Claire M Rickard^{a,c,d}

^a NHMRC Centre of Research Excellence in Nursing Interventions for Hospitalised Patients, Centre for Health Practice Innovation Griffith Health Institute, Griffith University, Nathan, QLD, Australia

^b Paediatric Intensive Care Unit, Royal Children's Hospital, Herston, QLD, Australia

^c Research and Development Unit, Centre for Clinical Nursing, Royal Brisbane and Women's Hospital, Herston, QLD, Australia

^d Princess Alexandra Hospital, Woolloongabba, QLD, Australia

author email: a.ullman@griffith.edu.au

Conflict of interest statement: CR has received research grant funding and educational speaker fees that are unrelated to this project from a intravenous device manufacturer, Becton Dickinson and Company (BD). All other authors have no competing interest, financial or otherwise, in any of the products and their associated distributors, discussed within this manuscript.

The full report of this research was originally published in Nurse Education Today 34 (2014), Ullman A, Long D, Rickard "Prevention of central venous catheter infections: A survey of paediatric ICU nurse' knowledge and practice", pp. 202-207 Copyright 2014, with permission from Elsevier

Background of the study:

Paediatric intensive care unit (ICU) nurses are responsible for the management of children vulnerable to developing healthcare-associated infections, specifically central venous access device (CVAD)-associated infections. CVAD-associated infections have been demonstrated to reduce to zero with the adherence to best practice guidelines and interventions (Pronovost, 2008, Han et al., 2010, Berenholtz et al., 2004).

Why this study was conducted (aims):

This study aimed to examine paediatric ICU nurses knowledge and paediatric ICU practices, to prevent CVAD-associated infections, in comparison to best practice guidelines.

Study methods:

A cross-sectional, descriptive study was undertaken between 2010 and 2011 throughout all tertiary-level paediatric ICUs Australia and New Zealand. A questionnaire of CVAD-associated infection prevention knowledge was completed by paediatric ICU nurses, and a survey of practice was completed by representative nursing managers and researchers. The questionnaire was developed using a previously validated adult ten-item questionnaire (Labeau et al., 2009, Labeau et al., 2008), with modifications to ensure applicability to paediatrics, using the Centers for Disease Control and Prevention (CDC) Guidelines for the Prevention of Intravascular Catheter-Related Infections (O'Grady et al., 2002, O'Grady et al., 2011). The survey of practice was developed using the CDC Guidelines (O'Grady et al., 2002).

Study results:

All Australian and New Zealand paediatric ICUs participated in the survey of practice ($n=8$) and 253 paediatric ICU nurses completed the questionnaire (response rate 34%).

The knowledge questionnaire mean total score was 5.5 out of a possible ten ($SD = 1.4$; range: 1-9). There was significant variation of total scores between paediatric ICU sites (range: 4.8–6.0; 7 df; $p = 0.01$). Questions of knowledge which were frequently incorrectly answered were surrounding maximum sterile barrier precautions for the insertion of CVADs and the correct concentration of chlorhexidine gluconate for the disinfection of the insertion site and routine dressing changes.

There was mixed application of CDC guidelines in the survey of practice of paediatric ICUs. Positive applications included the practice of prompt removal of CVADs when no longer essential and 2% chlorhexidine gluconate for antiseptic of the insertion site. Some paediatric ICUs reported 'sometimes' or 'mostly' routinely changing CVAD even if there is no suspicion of CVAD-associated infection and never using sutureless securement devices; which is contrary to CDC guidelines for best practice.

Implications for clinicians:

This study has demonstrated that a gap remains between best practice guidelines and nurses' knowledge and practice in the prevention of CVAD-associated infection. The prevention of CVAD-associated infection via the implementation of readily available guidelines should be an important goal for healthcare professionals.

While the prevention of CVAD-associated infection in the paediatric ICU depends on many factors, some are associated with discretionary decisions made by healthcare professionals (Ramritu et al., 2008). Personal education and professional development is the responsibility of all clinicians.

Process-improvement strategies are available within critical care to facilitate the translation of evidence-based guidelines to the bedside. These knowledge translation strategies include the use of catheter-care bundles, checklists and standardised insertion trolleys (Institute for Healthcare Improvement, 2011, McPeake et al., 2012). These strategies have been demonstrated to significantly reduce the rate of CVAD-associated infections, resulting in improved health outcomes for patients and significant cost savings (Helder et al., 2013, Sacks et al., 2014, Kim et al., 2011).

References:

- BERENHOLTZ, S. M., PRONOVOST, P., LIPSETT, P. A., HOBSON, D., EARSING, K., FARLEY, J. E., MILANOVICH, S., GARRETT-MAYER, E., WINTERS, B. D., RUBIN, H. R., DORMAN, T. & PERL, T. 2004. Eliminating catheter-related bloodstream infections in the intensive care unit. *Critical Care Medicine*, 32, 2014-20.
- HAN, Z., LIANG, S. & MARSCHALL, J. 2010. Current strategies for the prevention and management of central line-associated bloodstream infections. *Infection and Drug Resistance*, 10, 147-63.
- HELDER, O., KORNELISSE, R., VAN DER STARRE, C., TIBBOEL, D., LOOMAN, C., WIJNEN, R., POLEY, M. & ISTA, E. 2013. Implementation of a children's hospital-wide central venous catheter insertion and maintenance bundle. *BMC Health Serv Res*, 13, 417.
- INSTITUTE FOR HEALTHCARE IMPROVEMENT. 2011. *Implement the IHI Central Line Bundle* [Online]. Available: <http://www.ihl.org/knowledge/Pages/Changes/ImplementtheCentralLineBundle.aspx> [Accessed 12/11/2012].
- KIM, J. S., HOLTOM, P. & VIGEN, C. 2011. Reduction of catheter-related bloodstream infections through the use of a central venous line bundle: epidemiologic and economic consequences. *Am J Infect Control*, 39, 640-6.
- LABEAU, S., VANDIJCK, D., RELLO, J., ADAM, S., ROSA, A., WENISCH, C., AGBAHT, K., CSOMOS, A., SEHA, M., DIMOPOULOS, G., VANDEWOUDE, K., BLOT, S. & FOR THE EVIDENCE STUDY INVESTIGATORS 2009. Centers for Disease Control and Prevention guidelines for preventing central venous catheter-related infection: Results of a knowledge test among 3405 European intensive care nurses. *Critical Care Medicine*, 37, 320-323.
- LABEAU, S., VEREECKE, A., VANDIJCK, D., CLAES, B. & BLOT, S. 2008. Critical care nurses' knowledge of evidence based guidelines for preventing infections associated with central venous catheters: An evaluation questionnaire. *American Journal of Critical Care*, 17, 65-71.
- MCPEAKE, J., CANTWELL, S., BOOTH, M. G. & DANIEL, M. 2012. Central line insertion bundle: experiences and challenges in an adult ICU. *Nurs Crit Care*, 17, 123-9.
- O'GRADY, N. P., ALEXANDER, M., BURNS, L. A., DELLINGER, E. P., GARLAND, J., O'HEARD, S., LIPSETT, P. A., MASUR, H., MERMEL, L. A., PEARSON, M. L., RAAD, I. I., RANDOLPH, A., RUPP, M. E., SAINT, S. & HEALTHCARE INFECTION CONTROL PRACTICES ADVISORY COMMITTEE (HICPAC) 2011. Guidelines for the prevention of intravascular catheter-related infections. *Clinical Infectious Diseases*, 52, e162-93.
- O'GRADY, N. P., ALEXANDER, M., DELLINGER, E. P., GERBERDING, J. L., HEARD, S. O., MAKI, D. G., MASUR, H., MCCORMICK, R. D., MERMEL, L. A., PEARSON, M. L., RAAD, I. I., RANDOLPH, A. & WEINSTEIN, R. A. 2002. Guidelines for the prevention of intravascular catheter-related infections. *MMWR Morbidity and Mortality Weekly Report*, 51, 1-29.
- PRONOVOST, P. 2008. Interventions to decrease catheter-related bloodstream infections in the ICU: The Keystone Intensive Care Unit Project. *Am J Infect Control*, 36, S171 e1-5.

- RAMRITU, P., HALTON, K., COOK, D., WHITBY, M. & GRAVES, N. 2008. Catheter-related bloodstream infections in intensive care units: a systematic review with meta-analysis. *Journal of Advanced Nursing*, 62, 3-21.
- SACKS, G. D., DIGGS, B. S., HADJIZACHARIA, P., GREEN, D., SALIM, A. & MALINOSKI, D. J. 2014. Reducing the rate of catheter-associated bloodstream infections in a surgical intensive care unit using the Institute for Healthcare Improvement central line bundle. *Am J Surg*.