Pre-print version

Please cite as:

Ballantyne, M., Treby, D.L., Quarmby, J. and Pickering, C.M., (2016) Comparing the impacts of different types of recreational trails on Grey Box grassy-woodland vegetation: lessons for conservation and management. *Australian Journal of Botany* **64**, 246-259.

Comparing the impacts of different types of recreational trails on Grey Box grassy-woodland vegetation: lessons for conservation and management

Mark Ballantyne^{a*}, Donna Louise Treby, Joseph Quarmby^b and Catherine Marina Pickering^a

^a Environmental Futures Research Institute, Griffith University, Gold Coast, Queensland 4222, Australia

^b South Australian Department of Environment, Water and Natural Resources, GPO Box 1047, Adelaide, South Australia 5000, Australia

*corresponding author email: mark.runkowski@griffithuni.edu.au, phone: +61(0)405783604

Abstract

Tourism and recreation are popular in natural areas but can damage plant communities including those of high conservation value in protected areas. This includes impacts from recreational trails, but what type of trail has the most impact and why? We compared the impacts of five different trails (narrow, intermediate and wide bare earth trails, intermediate gravel trails and wide tarmac trails) on the endangered Grey Box grassy-woodland (*Eucalyptus microcarpa*) in Belair National Park near Adelaide in South Australia. First, the extent, width and area of recreational trails in the remnant woodland were mapped. Then, vegetation parameters were recorded in quadrats at three distances from the edge of trails in the woodland, with 10 replicate sites per trail type and single quadrats at 10 control sites (i.e. total 60 sites, 160 quadrats). All trails resulted in vegetation loss on the trail surface and along the edges of the trails, as well as changes in vegetation composition, including reductions in shrubs and bulbs close to the trail. The most common types of trail were bare earth trails with an average width of 2.5 m (50% of trails) which resulted in the greatest soil loss ($> 88,000 \text{ m}^3$) and vegetation loss ($33,899 \text{ m}^2$ or 3.4 ha) in the 167 ha woodland remnant overall. Wider (5.4 m) hardened tarmac trails, however, were associated with low species richness, high cover of exotic grasses and few herbs, shrubs and bulbs compared to vegetation away from trails and closer to other trails. Therefore a mixed approach to the provision of trails may be most appropriate, with hardened trails used in areas of highest use, but in some circumstances leaving trails unhardened may be more appropriate where they are likely to remain narrow and where there is less likely to be erosion and/or safety issues.

Key words: trail impacts, recreational infrastructure, remnant vegetation, edge effects, fragmentation, compositional changes

Introduction

Globally, and in Australia, nature-based tourism and recreation is increasing including trail-based activities such as hiking and mountain-biking (Kuenzi and McNeely 2008; Balmford et al. 2009; Monz et al. 2010; Newsome et al. 2013; Rossi et al. 2015). As a result, the creation of recreational trails in natural areas is becoming more common, especially in countries such as the USA and Australia (Monz et al. 2010; US National Park Service 2014; Ballantyne et al. 2014). While providing access to natural areas is important for human health and wellbeing (Swanwick et al. 2003; Lee and Maheswaran 2011) and fosters support for nature conservation (Spenceley et al. 2015), the infrastructure required for nature-based tourism and recreation has a range of environmental impacts including on plant communities of high conservation value (Matlack 1993; Stenhouse 2004; Godefroid and Koedam 2004; Hill and Pickering 2006; Olive and Marion 2009; Marion and Leung 2011; Ballantyne et al. 2014).

Recreational trails vary in their design, construction, maintenance and use. They can have hardened or unhardened surfaces, they can be narrow arteries or wide road-like developments, they can be formally created by land managers or informally created by users, and they can be single trails or a proliferation of dense networks (Marion and Leung 2001). Associated with the diversity of trails is variation in the severity and types of their impacts on vegetation (Godefroid and Koedam 2004; Hill and Pickering 2006; Monz et al. 2010; Ballantyne and Pickering 2015a). They can damage vegetation: (1) during their construction and maintenance, (2) as a result of changes in abiotic conditions caused by the presence of the trail, and/or (3) from the use of the trail for different types of recreational activities (Figure 1). These impacts can occur at the direct local scale (along the trail corridor itself) and at the indirect local scale (as edge effects away from the trail), as well as cumulatively across the landscape (Brooks and Lair 2005) (Figure 1).

[INSERT FIGURE 1]

Impacts of trails can vary depending on the methods used in their construction and maintenance, but clearing and disturbing vegetation along the trail corridor and adjacent to the trail is common, especially where heavy machinery is used (Buckley 2004; Ledoux 2004). There are often changes in soil conditions including compaction, soil removal and introduction of new soil or other substrates to create the trail (Marion and Leung 2004; Müllerová et al. 2011). Impacts from the presence of the trail include changes in light regimes, precipitation, wind exposure and temperature as well as altered water flow and

reduced organic matter along edge gradients away from the trail (Laurance et al. 1998; Potito and Beatty 2005; Pohlman et al. 2007). Finally, the use of the trail has additional impacts such as soil erosion and compaction often on the trail surface, along with damage to vegetation on the trail edge from trampling, changes in hydrology and soil movement as a result of soil erosion on the trail (Hammit and Cole 1998; Cole 2004; Marion and Leung 2004; Pescott and Stewart 2014) with more intensively-used trails often experiencing more damage than trails with limited use (Kutiel et al. 1999; Potito and Beatty 2005; Hamberg et al. 2008; Kim and Daigle 2012). The use of trails by plant collectors, arsonists, as well as the spread of weeds, pathogens and feral animals is also of concern (Benninger-Truax et al. 1992; Matlack 1993; Buckley et al. 2004; Baret and Strasberg 2005; Pickering and Mount 2010).

There is an increasing body of literature comparing the impacts of different trail-based activities such as hiking and mountain biking (Cole and Bayfield 1993; Kutiel et al. 1999; Talbot et al. 2003; Cole 2004; Hill and Pickering 2009; Monz et al. 2010; Pickering et al. 2010a). These studies have found that the timing, type and intensity of use and the tolerance of ecosystems affect the severity and types of impacts on vegetation (Hammit and Cole 1998; Cole 2004; Marion and Leung 2004; Pickering 2010). Fewer studies have assessed impacts of the trail infrastructure itself (Ballantyne and Pickering 2015a), and only a handful compare impacts among different types of trails (Godefroid and Koedam 2004; Müllerová et al. 2011) with just two from Australia (Hill and Pickering 2006; Ballantyne and Pickering 2015b). This is despite the hundreds of kilometres of formal trails (e.g. visible maintenance, signage and/or access infrastructure and/or mapping by land-owners for public use), and informal trails (those created and maintained by visitors for recreation that are outside of the formally-managed trail system) in Australia many of which traverse plant communities of high conservation value.

The results from the few studies that have compared among trails have found that the severity of impacts can vary among trails. Vegetation along the edge of formal hardened trails, for example, can be affected by disturbance to vegetation and soils during construction and maintenance resulting in the colonisation of trail edges by weeds and ruderal species (Hill and Pickering 2006), decreases in native plant cover, especially disturbance-sensitive species (Hill and Pickering 2006), changes in soil nutrients (Müllerová et al. 2011) and ultimately longer-term changes in plant composition and structure (Hill and Pickering 2006; Malmivaara-Lämsä et al. 2008; Ballantyne and Pickering 2015b). Informal, unhardened and narrow trails, in contrast, may have fewer and less severe impacts along the trail edge as there

is often less disturbance to existing vegetation during their creation, but there can be continuing impacts on the trail itself (Marion and Leung 2004; Olive and Marion 2009; Marion and Leung 2011; Wimpey and Marion, 2011). This includes soil erosion and trail widening when the trail surface is not stabilised by hardening which can affect vegetation on the trail edge (Olive and Marion 2009; Wimpey and Marion 2010).

The aim of the current research was to assess how recreational trails differ in the severity and types of impacts they have on plant communities of high conservation value. This involved comparing impacts of different recreational trails in a large remnant of Grey Box grassy-woodland (*Eucalyptus microcarpa*) in Belair National Park. The woodland is listed as an endangered plant community under Australian legalisation (Australian Government 2010) and as critically endangered within the World Wildlife Fund's terrestrial ecoregions database (World Wildlife Fund 2014). Specifically we: (1) mapped the extent and types of recreational trails in the woodland, (2) compared on-trail impacts between trail types (direct local scale), and (3) compared the impacts of different trail types on the composition of vegetation along the edges of the trails (indirect local scale). Based on the results of this study, we then provide recommendations for the management of different types of recreational trails in this and other plant communities that are of high conservation value, and popular sites for trail based recreation.

Study Ecosystem

Grey Box grassy-woodland (*Eucalyptus microcarpa*) is restricted to a series of scattered remnants across south-eastern Australia from central New South Wales through north and central Victoria and into South Australia (Australian Government 2010, 2012). Before European settlement the ecosystem covered around 3.4 million ha, but today occupies just over 500,000 ha; a loss of 85% (Australian Government 2010, 2012). The primary causes of this decline have been land clearing for agriculture, development and resource extraction with ongoing pressures from weed invasion, grazing and altered fire regimes (Yates and Hobbs 1997; Spooner et al. 2002; Australian Government 2010).

In South Australia, near the capital city of Adelaide (population 1.25 million people), the loss of Grey Box grassy-woodland has been particularly severe declining from 20,000 ha to just 2,000 ha (90% loss) (Kraehenbuehl 1996; Paton 2008). The woodland typically occurs on undulating plains and slopes with productive soils derived from alluvial or colluvial deposits

with rocky outcrops and clay pans (Australian Government 2010). Canopy cover is slightly denser than elsewhere in its range, likely as a response to higher rainfall in the region which averages 375-700 mm/year and Mediterranean temperatures (mean summer maximum 27°C; mean winter maximum 12°C) (Paton 2008; Australian Government 2012). The canopy is dominated by around 50% cover of the dominant tree *Eucalyptus microcarpa* often intermixed with stands of *E. camaldulensis* and *E. leucoxylon* (Australian Government 2010). The shrub layer provides around 30-40% cover and is dominated by species such as *Acacia pycnantha* and *Dodonaea viscosa*. The understory is highly diverse and dominated by grasses such as *Rytidosperma caespitosum* and *Poa labillardieri* var. *labillardieri* with numerous herb, bulb, chenopod and cryptophyte species. There is also high cover of invasive European grasses such as *Avena barbata*, *Brachypodium distachyon* and *Briza major* (Australian Government 2010). Total plant species richness often exceeds 200 even in small remnants (Quarmby pers. obs.) and includes numerous nationally-threatened species such as the Pink-lipped Spider-orchid (*Caladenia behrii*) and Plum Leek-orchid (*Prasophyllum pruinsum*). The ecosystem is also a stronghold for some International Union for the Conservation of Nature (IUCN) Red-listed species such as the Swift Parrot (*Lathamus discolor*) and Flinders Ranges Worm-lizard (*Aprasia pseudopulchella*).

Study Site

The Adelaide Grey Box grassy-woodland is highly fragmented and consists of numerous small remnants, the largest of which is 190 ha, with most <100 ha. Most remnants are within public reserves and private land between Adelaide and the southern Flinders Ranges (Australian Government 2010). These remnants are increasingly used for a range of recreational trail-based activities including hiking and mountain biking, including the large 167 ha of Grey Box grassy-woodland within Belair National Park (835 ha) (Figure 2). This peri-urban, IUCN Category II Park is 13 km south-east of Adelaide (Figure 2). The Park was proclaimed in 1891 and underwent a period of use as a semi-formal arboretum and government residence as well as being used for recreation with the creation of tennis courts, grass ovals and picnic areas (South Australian Government 2012). Today it is managed for conservation and recreation with around 250,000 visitors per year (Department of Environment, Water and Natural Resources 2003; Mitcham City Council 2015).

[INSERT FIGURE 2]

Data Collection

All recreational trails within the 167 ha of Grey Box grassy-woodland in Belair National Park were mapped including formal and informal trails. Trails were mapped using a Trimble Yuma GPS Tablet with 1-2 m real time accuracy and a recording interval of 1 m.

The formal and informal trails were further categorised based on their average width and surface using a visual survey method similar to that used in trail condition class assessments (e.g. Farrell and Marion 2001; Ballantyne et al. 2014). Average width of the trail tread, defined as the most heavily trafficked area of the trail (Wimpey and Marion 2010), was measured at the start of each new section of trail and at 100 m intervals along the trail. Average widths were calculated for each section and then all trails grouped into narrow (< 1 m), intermediate (1 – 3 m) and wide (> 3 m) trails. Surfaces were visually assessed, with bare soil, gravel and tarmac surfaced trails all found within the woodland. As a result five trail types were identified in total: (1) narrow bare soil, (2) intermediate bare soil, (3) wide bare soil, (4) intermediate gravel and (5) wide tarmac trails. While bare soil and gravel trails are predominantly used for hiking, dog-walking and running, wide tarmac trails are used for recreational vehicles, tourist drives and access to leisure facilities. For each of the five types of trails, 10 stratified random sites were located across the combined length of each type using ET GeoWizards 10.2 in ArcMap Version 10.1 (Ballantyne et al. 2014) for more detailed sampling of the trail and vegetation.

To assess impacts on the trail surface, soil loss (cross-sectional area), trail slope alignment angle and surface compaction were measured at each of the 10 sites per trail type. Cross-sectional area and trail-slope alignment angle were used to estimate soil loss using the methodology in Olive and Marion (2009). Soil compaction was measured using a penetrometer with a maximum capacity of 4.5 kg/cm² at five equally-spaced points across the span of the trail tread at each site.

To assess the impacts of the five types of trails on vegetation adjacent to the trail, we surveyed all plants (excluding trees) in 1 x 3 m quadrats located immediately adjacent to the edge of the trail, at 5 m and at 10 m away from the trail edge at each site (total of 150 quadrats, e.g. three distances, 10 replicate sites for each of five trail types). In addition, vegetation was assessed in 1 x 3 m quadrats at each of 10 randomly-selected control sites that

were a minimum of 50 m from the edge of trails, roads, mown areas, burnt areas or the woodland edge. Field work was carried out from 20 - 31 October 2014.

In each of the 160 quadrats, total vegetation cover, cover of separate growth forms, species richness and canopy cover were recorded along with abiotic variables (see below). Vegetation cover was measured as the top-cover (vegetation, rock, bare ground, dead wood or moss) and the total overlapping cover of all plant species intercepted by a thin metal wire placed at 100 randomly-located points within the quadrat. The number of 'hits' per species was converted into percentage cover values and used to calculate both overlapping vegetation cover and overlapping cover of separate growth forms. Total species per quadrat was used to calculate total species richness and weed richness. Canopy cover was analysed as a percentage by photographing the quadrangle of airspace directly above each zone and then calculating the number of pixel cross-hair points occupied by canopy, sky or other (Monz et al. 2010; Ballantyne and Pickering 2015b).

At each site (50 trail sites + 10 controls) abiotic covariates that are likely to affect vegetation were also recorded. These included slope, altitude, aspect, soil type, distance to water and time since last fire. Slope was measured using the Hunter Research and Technology LLC Theodolite HD Version 3.2 app for iPad (available online <http://hunter.pairsite.com/theodolite/>) by aligning the top of two poles of equal height 5 m either side of the survey point and measuring the difference in angle. Altitude was measured using a 1 second SRTM-derived 3-second smoothed digital elevation model provided by the Australian Government for the Adelaide region (<http://www.ga.gov.au/data-pubs>). Aspect was measured using an electronic GPS-corrected compass bearing against true north. Distance to water was the Euclidean distance to nearest water body (stream, river, lake) using ArcMap Version 10.1. Soil type was measured using the soil landscape map units of southern South Australia map (<http://www.naturemaps.sa.gov.au/>). Time since last fire was taken from a fire management dataset recording major fires in protected areas (<http://www.naturemaps.sa.gov.au/>).

Data Analyses

The total area of each trail type, and therefore the total area of vegetation lost to the trail surface, was calculated by multiplying the total length (measured in ArcMap Version 10.1) by the average width of each type of trail. We were then able to calculate total soil loss for

217 each trail type by multiplying average soil loss per site (cm^2) by the total length (cm^3) of that
218 type of trail.

219 To analyse differences among the trail variables: average soil loss (log10 transformed), trail
220 slope alignment angle (log10 transformed) and surface compaction, a series of One-way
221 ANOVAs were conducted in SPSS Version 21 when the assumptions of normality and
222 variance were satisfied. Least significant differences were used as post-hoc tests to determine
223 which pairs of trails differed.

224 To analyse whether vegetation 10 m from the trails differed from the controls, data for these
225 quadrats was compared with that from control quadrats using a series of One-way ANOVAs.
226 As no significant differences were found, the control quadrats were removed from further
227 analyses, with the 10 m quadrats treated as the 'natural' condition of the vegetation in
228 subsequent analyses.

229 To assess differences in the impact of the five trail types on understory vegetation, and how
230 far from the trail impacts could be detected, linear mixed models were used to analyse species
231 richness, relative vegetation cover, relative weed cover (arcsine square-root transformed) and
232 weed species richness (log transformed) in SPSS Version 21. Trail type, distance and
233 distance*trail type were set as fixed factors, and sites (blocks) used as a random factor.
234 Covariates (abiotic and biotic) were added to the list of fixed factors with aspect and altitude
235 log10-transformed to satisfy assumptions of normality and homoscedasticity. Covariates were
236 added and removed sequentially to determine the most significant models according to
237 Akaike's Information Criterion (AIC). Finally, to compare any significant effects using the
238 most powerful model, pairwise comparisons using estimated marginal means were computed
239 for trail type and distance using Syntax commands. If an interaction was significant, separate
240 One-way ANOVAs were performed for each trail type and distance to understand the nature
241 of the interaction. A p value of ≤ 0.03 was used to account for the increased chance of falsely
242 assuming a significant difference, i.e. an increased chance of Type II errors.

243 To assess overall similarity of vegetation depending on how far the quadrat was from the trail
244 and the type of trail, values for Sørensen's index were calculated using presence/absence data
245 for all living vascular species per quadrat. A Bray-Curtis similarity matrix was used to
246 compare composition between quadrats [site(distance*trail type)] for all trails with data being
247 extracted and analysed using a One-way ANOVA. Tukey post-hoc tests were used to
248 determine the nature of any significant differences.

To analyse differences in species composition and combined relative cover of growth forms between distances and trail types, a permutational multivariate ANOVA (PERMANOVA) was used with trail type, distance and distance*trail type as fixed factors, and site(trail type) as a random factor in PRIMER Version 6 using a Bray-Curtis dissimilarity matrix on square-root transformed cover data for individual species (divided by total cover) with the cover of litter, bare ground, rock, dead wood and moss removed. To determine if composition between 10 m and control sites was similar, we used a One-way ANOSIM. We found that composition was similar between these two groups of sites and so 10 m was used as the control. To show compositional differences among the five trail types and three distances, multi-dimensional scaling (MDS) plots were used to show the clustering of samples in two-dimensional space with goodness of fit shown as a stress value. This method is useful as it (1) visually portrays patterns by clustering and ordinating samples and (2) using the PERMANOVA function, statistically analyses them with a non-parametric permutation-based test (9,999 permutations), similar to that of an ANOVA, but not requiring a normalised data distribution (Clarke 1993). Finally, a SIMPER analysis was used to determine which species contributed to any observed differences between groups.

Additionally, we used separated Linear Mixed Models on total cover of each growth form (arcsine square-root transformed) per quadrat as an independent variable for graminoids, herbs and bulbs to determine the overall strength and direction of any observed effects. Shrubs were removed as a variable as they were absent from many quadrats.

To assess whether the trails had an effect on the cover of any dominant species [occurring in at least 50% of quadrats (75 out of 150)], cover data (arcsine square-root transformed) for that species was included in separate Linear Mixed Models, using post-hoc pairwise comparisons to determine the nature of any significant differences. These species included: the native shrub *Dodonaea viscosa*, and the two exotic grasses *Briza major* and *Brachypodium distachyon*.

Results

On-trail impacts

Within the 60 sample sites in the 167 ha of Grey Box grassy-woodland in Belair National Park, there was a total of 109 plant species including 4 species of trees, 11 shrubs, 20 bulbs, 49 herbs, 19 grasses, 5 sedges/rushes and 1 fern. There was also 20.1 km of trails in these

woodlands, with 16.3 km formal and 3.8 km informal trails. There was approximately 2.2 km of narrow bare, 10.1 km of intermediate bare, 2 km of wide bare, 1.5 km of intermediate gravel and 4.4 km of wide tarmac trails (Table 1). The narrow bare earth trails were all created by users (informal), while the other four trail types were either entirely, or predominantly, trails provided and maintained by the Park management.

The five types of trails differed in average width ($F = 52.075$, $p < 0.001$), apart from intermediate bare and intermediate gravel trails ($p = 0.320$). Because of their hardened surface, the wide tarmac trails had no soil loss from the surface. Narrow bare trails had less soil loss ($1.4\text{ m}^3\text{ m}^{-1}$) than the other three types ($F = 12.991$, $p < 0.001$) which all lost around 60.1 m^3 of soil per metre of trail. Total soil loss was very high for intermediate bare trails, estimated at $68,218.9\text{ m}^3$ in total, due to the greater length of this type of trail. Surprisingly, total soil loss for intermediate gravel trails was high at $4,095.2\text{ m}^3$ (28.1 m^3 of soil per metre of trail); around 40% greater than narrow bare trails which were longer than the total length of intermediate gravel trails. Compaction ($F = 0.879$, $p = 0.484$) and trail slope alignment angle ($F = 0.807$, $p = 0.527$), did not differ among the five types of trails (Table 1).

Because the five types of trails varied in length and width, they differed in the amount of vegetation lost. Intermediate bare trails were associated with the greatest loss ($25,132.3\text{ m}^2$ or 2.5 m^2 lost per metre of trail) of vegetation as they were the most extensive trail type, while wide tarmac trails, which were much shorter accounted for similar levels of vegetation loss ($23,751.9\text{ m}^2$ or 5.4 m^2 lost per metre of trail) because they were wider. Narrow bare trails caused the least loss of vegetation ($1,726.1\text{ m}^2$ or 0.8 m^2 lost per metre of trail), in part because they were narrower. Total vegetation loss due to all five trails was $60,712.8\text{ m}^2$, which is around 2.1% (3.4 ha) of the total remnant of Grey Box woodland (167 ha).

Table 1: Means and standard deviations for different trail variables for each of the five trail types assessed in Grey Box grassy-woodland in Belair National Park. Significant differences indicated in bold ($p < 0.05$) according to One-way ANOVA with lettering showing the nature of the difference according to least significant difference post-hoc comparisons (e.g. ‘a’ is significantly different to ‘b’ but the same as ‘a’). TSAA = trail slope alignment angle.

	Narrow bare	Intermediate bare	Wide bare	Intermediate gravel	Wide tarmac	Totals
Length trails (m)	2,157	10,052	2,011	1,458	4,399	20,079
Area trails (m ²)	1,726	25,132	7,041	3,062	23,752	60,713
Av. veg. loss (m ²) / m	0.8	2.5	3.5	2.1	5.4	
Surfacing	Unsurfaced	Unsurfaced	Unsurfaced	Dolomite gravel	Bitumen	
% Formal	0	90	70	100	100	
% Informal	100	10	30	0	0	
Av. width (m)	0.8 ± 0.3^a	2.5 ± 0.7^b	3.5 ± 1.2^c	2.1 ± 0.6^b	5.4 ± 0.6^d	
Av. soil loss (m ³) / m	1.35 ± 15.5^a	67.9 ± 58.5^b	84.5 ± 83.9^b	28.1 ± 18.4^b	0	
Total soil loss (m ³)	2,908.4	68,218.9	17,000.9	4,095.2	0	92,223.4
Av. compaction (kg/cm ²)	4.2 ± 0.5	4.3 ± 0.5	4.4 ± 0.3	4.4 ± 0.2	> 4.5	
TSAA (°)	44.5 ± 23.1	37.5 ± 27.8	55.5 ± 29.5	34.5 ± 36.1	39.5 ± 26.8	
Predominant use	Hiking	Hiking, mountain biking	Hiking, mountain biking	Hiking, mountain biking	Driving, hiking	

Was there an effect of the trails on vegetation?

There were no significant differences in any of the dependent variables between the control quadrats (50 m from trails) and those 10 m from the trails except for cover of herbs, indicating that most effects of the trails were limited to vegetation < 10 m from the trails. Vegetation in the control and 10 m quadrats was characterised by high cover of the native sub-shrubs *Hibbertia exutiacies* and *Hibbertia australis* (23% cover) and graminoids composed largely of the exotic grasses *Briza major* (18%) and *Brachypodium distachyon* (11%) along with the native sedge *Lepidosperma curtisiae* (18%) and tussock grass *Rytidosperma caespitosa* (10%). The native bulbs *Arthropodium strictum* (5%) and *Lomandra multiflora* (5%) and the exotic herb *Plantago lanceolata* (11%) also had comparatively high cover.

Vegetation was affected by the trails, with significant differences in several variables between quadrats on the edge of trails and those 5 m and 10 m away from the trails. For all five trail types, there was less vegetation in quadrats on the edge of trails ($43\% \pm 28$; $F = 15.534$, $p = < 0.001$) compared to quadrats 5 m ($62\% \pm 25$, $p = 0.001$) and 10 m ($63\% \pm 23$, $p = < 0.001$) away, but no differences between the 5 m and 10 m quadrats ($p = 0.864$) (Tables 2 and 3). Bulb cover was also affected by trails ($F = 11.843$, $p < 0.001$), with reduced cover near the trail (Tables 2 and 3). There were no overall effect of trails on the cover of dominant species including the native shrub *Dodonaea viscosa*, and the two exotic grasses *Briza major* and *Brachypodium distachyon*.

Overall composition (based on the ANOSIM) differed with distance from trails, but was also affected by an interaction with trail type (see below). The cover of different growth forms was affected by the trails (Global Rho = 0.26, $p < 0.001$, Stress = 0.18) with an increase in the cover of shrubs and bulbs in quadrats 5 m and 10 m (18.4% and 20.9%) from the trail compared to those on the trail edge (11.8%). There was a slight decrease in the cover of graminoids for the 5 m and 10 m quadrats (49.7% and 47.6%) compared to the trail edge (55.6%). These patterns were in part due to decreasing cover of the exotic grasses *B. major* and *B. distachyon* further from the trail and a rise in the cover of native grasses such as *Rytidosperma caespitosa* and *Microlaena stipoides* var. *stipoides*. Vegetation on the edge of the trails was typically dominated by grasses (average cover 71%) with lower cover of sedges (7%), shrubs (8%) and bulbs (<1%). Vegetation in quadrats 5 m from the trails was also dominated by grasses (60%) but with higher cover of shrubs (13%), sedges (9%) and mostly

336 exotic bulbs (5%). Vegetation in quadrats 10 m and 50m from the trails was dominated by
 337 grasses (53%) but > 20% of them were native, and there was a higher cover of shrubs (15%),
 338 sedges (10%) and native bulbs (7%).

339 Although there were 39 species of weeds (1 shrub, 8 bulbs, 11 grasses, 19 herbs) recorded
 340 across the 160 quadrats surveyed, with average weed cover of 29% per quadrat, there was no
 341 effect of the trails on weed cover *per se* (Tables 2 and 3).

342

Table 2: Results from Linear Mixed Models (LMM) and PERMANOVA analyses of vegetation in quadrats at different distances from five types of trails. Significant differences indicated in bold ($p < 0.05$). Neg. = negative.

Dependent Variable	Trail type	Distance from trail	Interaction	Significant covariates
Species richness (LMM)				
All	F = 1.264 p = 0.297	F = 10.746 p = <0.001	F = 2.590 p = 0.013	
Weeds	F = 1.187 p = 0.329	F = 0.881 p = 0.418	F = 0.655 p = 0.729	Altitude neg. interaction F = 5.954, p = 0.019
Total % overlapping cover (LMM)				
Vegetation	F = 1.140 p = 0.350	F = 15.675 p = <0.001	F = 1.195 p = 0.311	
Weeds	F = 1.204 p = 0.323	F = 0.561 p = 0.573	F = 1.030 p = 0.422	Altitude neg. interaction F = 5.469, p = 0.023
Graminoid	F = 0.611 p = 0.656	F = 1.475 p = 0.234	F = 0.454 p = 0.885	
Herb	F = 2.558 p = 0.049	F = 2.902 p = 0.060	F = 0.715 p = 0.678	
Bulb	F = 0.529 p = 0.715	F = 11.843 p = <0.001	F = 1.341 p = 0.233	
Overlapping cover of dominant species				
<i>Dodonaea viscosa</i>	F = 0.062 p = 0.993	F = 3.007 p = 0.054	F = 0.614 p = 0.764	
<i>Briza major</i>	F = 2.654 p = 0.045	F = 2.062 p = 0.133	F = 2.071 p = 0.050	
<i>Brachypodium distachyon</i>	F = 0.863 p = 0.493	F = 0.311 p = 0.733	F = 0.753 p = 0.644	
Composition (PERMANOVA)				
All species	Pseudo-F = 1.111 p = 0.230	Pseudo-F = 2.363 p = 0.002	Pseudo-F = 1.274 p = 0.040	
Growth forms	Pseudo-F = 1.054 p = 0.398	Pseudo-F = 5.824 p = <0.001	Pseudo-F = 1.112 P = 0.314	
Sørensen Similarity Index	n/a n/a	n/a n/a	F = 4.564 p = 0.004	

Table 3: Characteristics of vegetation in quadrats for controls vs 10 m, and for quadrats on the edge of the five trail types. Significant differences indicated in bold ($p < 0.05$) according to linear mixed models with lettering showing the nature of the difference according to least significant difference post-hoc comparisons (e.g. 'a' is significantly different to 'b' but the same as 'a'). Inter. = interaction term.

	Controls + 10m	Narrow bare edge	Intermediate bare edge	Wide bare edge	Intermediate gravel edge	Wide tarmac edge	Interaction or Main Effect?
Species richness	9.6 ± 2.7 a	7.6 ± 2.9 b	8.0 ± 3.3 b	6.3 ± 2.4 c	6.8 ± 3.1 b	4.1 ± 2.3 c	Interaction
Total % overlapping cover							
Vegetation	80.9 ± 21.3 a	39.3 ± 23.3 b	42.4 ± 27.2 b	34.7 ± 14.9 b	45.4 ± 30.2 b	25.0 ± 14.2 b	Main
Graminoids	36.3 ± 20.9	26.7 ± 20.9	30.5 ± 30.1	17.9 ± 10.4	35.1 ± 33.0	20.1 ± 11.4	None
Herbs	33.2 ± 16.9 a	18.6 ± 17.4 a	9.0 ± 10.1 a	12.5 ± 8.7 a	12.2 ± 12.2 a	3.6 ± 5.0 b	Main
Bulbs	7.1 ± 6.2 a	4.8 ± 5.6 b	6.1 ± 6.4 b	1.9 ± 1.4 b	2.3 ± 2.8 b	1.1 ± 2.2 b	Main
Shrubs	2.7 ± 2.2 a	3.4 ± 3.6 a	3.9 ± 5.3 a	2.5 ± 4.1 a	4.5 ± 12.1 a	0.2 ± 0.4 b	Interaction
Overlapping cover of dominant species							
<i>Dodonaea viscosa</i>	0.6 ± 1.3	2.8 ± 3.6	3.3 ± 5.3	2.0 ± 4.1	1.7 ± 4.1	0.2 ± 0.4	None
<i>Brachypodium distachyon</i>	5.5 ± 8.3	4.1 ± 5.6	8.4 ± 19.7	3.9 ± 7.3	6.1 ± 9.7	11.8 ± 9.8	None
<i>Briza major</i>	Controls	Narrow bare edge	Intermediate bare edge	Wide bare edge	Intermediate gravel edge	Wide tarmac edge	Inter.
	16.9 ± 17.5 a	10.6 ± 16.6 a	9.4 ± 15.1 a	4.1 ± 5.3 b	5.4 ± 7.7 b	0.8 ± 1.5 c	
	Narrow bare 5m	Narrow bare 5m	Intermediate bare 5m	Wide bare 5m	Intermediate gravel 5m	Wide tarmac 5m	
	11.5 ± 18.1 ab	14.4 ± 15.8 a	14.4 ± 15.8 a	2.7 ± 3.8 c	5.4 ± 7.7 b	2.5 ± 4.9 c	
	Narrow bare 10m	Narrow bare 10m	Intermediate bare 10m	Wide bare 10m	Intermediate gravel 10m	Wide tarmac 10m	
	15.7 ± 19.1 a	7.9 ± 10.0 ab	7.9 ± 10.0 ab	0.4 ± 0.9 b	15.9 ± 18.2 ac	5.5 ± 9.4 abc	
Sørensen Similarity Index (%)	n/a	Narrow bare	Intermediate bare	Wide bare	Intermediate gravel	Wide tarmac	
Edge vs. 5m composition		58.2 ± 16.3	44.9 ± 11.5	43.7 ± 22.5	50.2 ± 25.9	29.4 ± 25.7	Inter.
Edge vs. 10m composition		49.9 ± 25.4 a	37.9 ± 12.7 ab	39.4 ± 17.8 ab	49.9 ± 11.6 a	20.7 ± 17.6 b	
5m vs. 10m composition		55.8 ± 13	54.3 ± 15.1	47.1 ± 13.8	57.7 ± 11.2	52.9 ± 12.5	

Was there a difference in impacts among trail types?

Some trail impacts were not simply a result of the presence of any type of trail, but varied among the trails. This included impacts on species richness and composition (Tables 2 and 3; Figure 3). Species richness was affected by an interaction between trail type and distance (Linear Mixed Model, $F = 2.207$, $p = 0.034$), as was composition (PERMANOVA, Pseudo- $F = 1.27$, $p = 0.04$, Stress = 0.23).

For the wider trails, species richness and composition were affected by distance from the trail, but there was no effect of distance for the narrow and intermediate bare earth trails (Table 2, Figure 3). For the wide tarmac trail, for example, quadrats on the edge of the trail had less than half the number of species as those further away (edge = 4.1 ± 2.3 ; 5 m = 6.8 ± 1.5 ; 10 m = 9.1 ± 2.6 ; $p < 0.021$). The vegetation in quadrats on the edge of the wide tarmac trail also differed in composition to those further away with more exotic grasses such as *B. distachyon*, *B. major* and *Avena barbata* close to the trail, but fewer native grasses such as *M. stipoides* var. *stipoides*, *Anthosachne scabra* and *Poa labillardieri* which were mainly found in quadrats 5 m and 10 m away from the trail.

For the wide bare trail there were also fewer species in quadrats on the edge of the trail (6.3 ± 2.4), than in quadrats 5 m and 10 m away (8.9 ± 3.1 ; 8.1 ± 2.1 , respectively; $p < 0.001$). Composition, however, was more similar (Figure 3) with a greater abundance of native sedges and graminoids (*L. curtisiae* and *R. caespitosa*) and moderate abundance of exotic grasses (*B. distachyon* and *B. major*) across all distances, but with more herb species further from the trails. For intermediate gravel trails, composition steadily changed from quadrats on the trail edge to those 10 m away with a steady decrease in exotic grasses (*B. distachyon* and *B. major*) and an increase in sub-shrubs (e.g. *H. australis*) and bulbs (e.g. *Arthropodium strictum*).

Average similarity in composition based on Sørensen's indices between pairs of quadrats was affected by an interaction between trail type and distance with less similarity between quadrats on the edge of trails and those 10 m away ($39.6\% \pm 20.1$ similarity; $F = 4.564$, $p = 0.004$). The greatest differences in composition were along wide tarmac trails where quadrats on the edge of the trail were very different to those 10 m away ($20.7\% \pm 17.6$ similarity). Along intermediate and wide bare trails, similarity was greater between edge and 10 m distances ($38.6\% \pm 15.2$) and along narrow bare and intermediate gravel trails, similarity was even higher ($49.5\% \pm 18.5$). Along all trails, composition 5 m from the trail was similar to

edge quadrats (av. $45.3\% \pm 22.4$ similarity; $F = 2.490$, $p = 0.056$) and even more similar to 10 m quadrats (av. $53.6\% \pm 13.1$ similarity; $F = 0.925$, $p = 0.458$), and therefore acted as a transition zone.

Herb cover was lower along wide tarmac trails, but this effect occurred irrespective of distance from the trail ($F = 2.558$, $p = 0.049$). Also, the 10 m quadrats for the wide tarmac trails differed from the control quadrats indicating either site-specific covariation or that the impact of this trail type on herbs extends beyond 10 m from the edge of the trail.

The cover of one dominant, *B. major*, was affected by an interaction between trail type and distance, with a slight reduction in cover for quadrats along the edge of wide bare, intermediate gravel and wide tarmac trails ($F = 2.071$, $p = 0.05$) compared to other trails and controls ($16.9\% \pm 17.5$). Along the edge of the wide bare trails, this species had low cover ($4.1\% \pm 5.3$), but it was even lower in quadrats further from the trail (5 m = $2.7\% \pm 3.8$; 10 m = $0.4\% \pm 0.9$). In quadrats along intermediate gravel and wide tarmac trails, cover of *B. major* increased with increasing distance from the trail (edge = $3.1\% \pm 4.6$; 5 m = $3.9\% \pm 6.3$; 10 m = $10.7\% \pm 13.8$; Table 3).

Abiotic covariates had limited effect on vegetation with altitude having a negative linear interaction with weed species richness ($F = 5.954$, $p = 0.019$) and with weed cover ($F = 5.469$, $p = 0.023$). Slope, aspect, distance to water, soil type, time since last fire, and canopy cover had no significant effect on vegetation.

[INSERT FIGURE 3]

Discussion

The five types of recreational trails differed in their impacts on vegetation in the endangered Grey Box grassy-woodland in Belair National Park. Remnants such as this one are important for recreational opportunities, but are also of high conservation value. A common problem for those managing this and other Parks, is which type of trails are best to minimise environmental impacts while also providing appropriate recreational opportunities. Part of this challenge has been the limited research comparing environmental impacts among different types of trails.

There is an extensive network of recreational trails in Grey Box woodland

There were over 20 km of recreational trails in the Grey Box grassy-woodland of which nearly 90% were formal, management-created trails. The most common of these were intermediate bare earth trails (60%), which collectively resulted in the loss of > 25,000 m² of vegetation. This was partly due to the total length of this type of trail in the woodland, but also due to the tendency of unsurfaced trails to widen over time (Olive and Marion 2009; Monz et al. 2010; Wimpey and Marion 2010). Other trails such as the wide tarmac trails, although less common (25%) in the woodland, also resulted in the loss of > 23,000 m² of vegetation as they were wider (often > 7 m) to accommodate recreational vehicles. The width of these trails exposes vegetation on the trail edge to a range of changes in abiotic conditions such as increased wind-blow which may further reduce tree numbers and canopy cover.

Interestingly, in Belair National Park, only 11% of trails were informal, user-created trails. This may be due in part to the closure of several informal trails in the Park in the past, as part of upgrades to the trail network (Taylor Cullity Lethlean 2009). In other remnants of the same woodland in the region, there were more narrow informal trails (Ballantyne and Treby 2015). Narrow informal trails can be important components of trail systems included in protected areas (Leung et al. 2011; Ballantyne et al. 2014; Ballantyne and Treby 2015). They can result in extensive loss of habitat (up to 20%), habitat fragmentation and complex internal networks of disturbance disrupting the once-contiguous ecosystems (Ballantyne et al. 2014). The resulting impacts from uncontrolled trail networks are complex and hard to manage (Leung et al. 2011; Leung et al. 2012; Pickering et al. 2012) highlighting the importance of limiting the spread of informal trail networks particularly in areas of high conservation value (Leung et al. 2012; Wimpey and Marion 2011).

All five trails affected vegetation adjacent to the trails

All five trails had some impacts on vegetation in the Grey Box grassy-woodland. This included reduced vegetation cover along the edge of the trails and up to 5 m away, but also the reduced cover of bulbs and shrubs and increased cover of graminoids. These changes are likely to be the result of: (1) trampling damage along the edge of the trail mostly by hikers and mountain-bikers, (2) trail maintenance activities (e.g. herbicide spraying, slashing and digging), and (3) changes in abiotic conditions caused by the presence of the trail itself. Abiotic changes could include drier soils compared to vegetated areas, greater convection and conduction of heat, and reductions in local atmospheric moisture (Delgado et al., 2007). Additionally, the open nature of the trail corridors could lead to increased wind-blow damaging vegetation along the edge of the trail, especially trees and shrubs (Laurance et al. 1998; Harper and MacDonald 2002; Delgado et al. 2007; Ballantyne and Pickering 2015b).

For the Grey Box woodland, it appears there are already many species with traits able to deal with conditions common along the edges of trails, particularly exotic grasses that are native to the European-Mediterranean region. These species often accumulate dead matter with age, have vertical, recurved leaves and late-spring abscisic acid-accumulation which increases their capacity to deal with exposed hot conditions (Diemer 1996; Körner 2003; Volaire et al. 2009). Species native to the Grey Box grassy-woodland, such as sclerophyllous shrubs and sub-shrubs (e.g. *Hibbertia* spp.) may also be well adapted to hotter, drier and windier conditions, but like many other sclerophyllous species, are less tolerant of trampling damage (Rickard et al. 1994; Bernhardt-Römermann et al. 2011; Pescott and Stewart 2014).

Trails differ in their impacts with more on-trail issues from unhardened informal trails

When impacts were compared among trails, we found strong contrasts. Conditions on the trail were generally poorest for bare trails while hardened trails were in good condition, as would be expected. Bare trails varied in width ranging from 0.8 - 4 m, and for the wider trails, soil loss was high. Cumulatively, bare earth trails accounted for 95.6% of total soil loss from trails in the woodland ($> 88,128.2 \text{ m}^3$ or 6.2 m^3 lost per metre of trail). This again illustrates how unhardened trails can be prone to on-trail degradation (e.g. soil erosion) (Olive and Marion 2009; Wimpey and Marion 2010; Pickering et al. 2010b) including over time (Marion and Leung 2004).

In contrast, bare trails, were narrow (e.g. informal narrow bare earth trails < 1 m), with limited soil loss and on-trail conditions were as good as, or even better than, some hardened trails. Indeed, the average soil loss from intermediate gravel trails was 52% higher than from narrow bare earth trails despite there being more narrow bare trails in the woodland. Maintaining such narrow trails through use of education, low gradients, grade reversals and drainage features, (e.g. water bars, drainage dips), strategically placed pass-points could be an efficient and cost-effective way of maintaining on-trail conditions. These methods, however, have had limited success in the past (Olive and Marion 2009) and the apparently ‘good’ condition of the narrow bare earth trails in the Grey Box woodland may be because these trails are relatively new (Taylor Cullity Lethlean 2009) or they receive very low levels of traffic.

For vegetation adjacent to trails, there were more impacts from wider hardened trails

When comparing impacts on vegetation adjacent to the trails, we found that wider and hardened trails had more severe impacts that extended further from the trails than narrower bare trails, although narrower bare trails did have poor surface conditions and these proliferate out as trails widen over time. While trail-hardening can be a popular management option to improve on-trail conditions (Wimpey and Marion 2010; Pickering et al. 2010b; Leung et al. 2011), and can maintain sustainable trail networks (Hill and Pickering 2006; Cahill et al. 2008; Tomczyk 2011; Zhang et al. 2012), they often dramatically alter vegetation composition along the trail edge (Godefroid and Koedam 2004; Hill and Pickering 2006; Müllerová et al. 2011; Ballantyne and Pickering 2015b) and, in some cases, negatively affect visitor experiences (Cahill et al. 2008). One factor we were unable to measure accurately was the age of the trails, which is likely to have had some effect on the composition of trail-side vegetation. It is likely that some of the earlier hardened trails will have exerted greater abiotic changes and have been maintained rigorously for a long time, thus leading to more marked edge effects (Godefroid and Koedam 2004).

Vegetation along the edge of wide tarmac trails contained far fewer species, was dominated by exotic grasses (*Brachypodium distachyon* and *Avena barbata*) and had less herb cover than control quadrats in the woodland. The construction and maintenance of these trails often damages vegetation along the edge of the trail, including during the use of equipment, in the introduction of materials, grading, levelling and laying of heated tarmac and the use of multiple chemical additives and finishers (Buckley 2004; Delgado et al. 2007). This can

damage shrubs and trees in, and close to, the work zone which often exceeds 20 m either side of the actual trail tread (Ballantyne pers. obs.). Once laid, wide tarmac trails often have low albedo and so heat up rapidly, creating localized wind vortices resulting in drier conditions near the trails (Delgado et al. 2007; Pohlman et al. 2007). These effects may account for why much of the vegetation close to the tarmac trails in the Grey Box woodland consisted of disturbance-tolerant species close (< 5 m) to the trail.

For intermediate gravel trails, there was a more gradual decline in the cover of exotic grasses and increases in native bulbs and shrubs away from the trail. This wider, but less severe, edge effect may be due to the substrates on the trails. The base-rich dolomite gravel used for these types of gravel trails can spread into adjacent vegetation and may leach alkalis into an otherwise base-poor environment through surface-flow (Quarmby pers. obs.). In other regions with base-poor soils, the use of foreign alkali gravel has had a dramatic effect on species richness and composition that extends beyond 10 m from the edge of the trail, particularly down-slope of the trail due to changes in soil nutrients (Godefroid and Koedam 2004; Müllerová et al. 2011). Reduced disturbance during installation, but gradual leaching from gravel trails may explain why edge effect from these trails on composition was wider, but less severe in the Grey Box woodland.

Wide bare trails (average width 3.5 m) also resulted in fewer species close to the trail in the Grey Box woodland. Wider trails that require the removal of canopy vegetation expose the ground surface along and adjacent to the trail to greater insolation, convection and conduction resulting in drier conditions (Pohlman et al. 2007). The composition of vegetation along the edge of the wide bare trails in the Grey Box woodland, however, was generally similar to controls with a more even composition of exotic vs. native graminoids and more herbs. In particular, the cover of *Briza major* was low on the edge of wide bare trails and continued to decrease with distance in contrast to tarmac and gravel trails where it increased positively with distance from those trails. There may have been less disturbance in the creation of the wide bare trails (largely just clearing) compared to the wider hardened trails, allowing the retention of complex native edge structure, nutrient levels and soil condition, reducing the dominance of weeds close to the trail edge (Ballantyne and Pickering 2015b).

Management recommendations

This research demonstrates that recreational trails differ in their severity and types of impacts caused to vegetation, and so future management of natural areas must account for this variation when developing sustainable trail networks. The results support the use of hardened trails along the most intensely-used areas, on steep slopes or in areas inappropriate for unhardened trails (e.g. riparian areas) where there is a risk of deteriorating on-trail conditions and trail widening. Hardening may also help restrict the creation of informal trails networks (Marion and Leung 2011; Wimpey and Marion 2011; Leung et al. 2012). Hardened trails, however, do have local impacts on edge vegetation and so their use should be limited, where possible, in areas of high conservation value, such as in Grey Box grassy-woodlands. The use of wide tarmac trails for vehicle access and base-rich substrates such as dolomite gravel should be avoided because they appear to have had the greatest effects on plant composition appearing to favour novel disturbance-tolerant species $\geq 10\text{m}$ from the trail edges. Minimising damage during construction, using locally-sourced surfacing substrates and actively rehabilitating edge vegetation may help mitigate some of these impacts (Ballantyne and Pickering 2015b).

The results indicate that, where unhardened trails are strategically-located and maintained, they can both sustain good on-trail conditions and have limited effect on edge vegetation. This is especially important in areas of high conservation value containing disturbance-sensitive species. As long as unhardened trails are not developed in topographically-sensitive areas such as over soils prone to water-logging, deep or Aeolian soils, steep slopes and riparian zones, trail gradients are kept to a minimum and trails are placed parallel to contour lines with adequate drainage systems, they can remain in good condition over time (Marion and Leung 2004). Maintaining the trail in good condition can be difficult with high use, if it results in vegetation loss, soil erosion and trail widening. These type of issues with unhardened trails can be minimised by using natural anchor points (rocks and logs), strategically-placed pass points and by actively managing for different types and intensities of use. Finally, monitoring how trails affect vegetation and soils, as well as fauna and aquatic systems, and how their use changes over time is important for long-term trail sustainability.

Conclusion

We compared the types and severity of impacts by five types of recreational trails on vegetation in an endangered woodland. These types of ecosystems are undergoing increasing

loss and degradation globally due to urbanisation and clearing with the development of recreational trails adding to these threats. We found trails varied in their impacts. Wider unhardened bare earth trails caused the most soil and vegetation loss, while hardened trails were less common and so resulted in less overall damage. However, hardened trails had the greatest effect on vegetation along the edge of the trail including decreased species richness, altered composition and decreases in sensitive plant species. It is likely that these changes in vegetation are a result of disturbance during the construction and maintenance of the trails. Based on the results of the current study, and the few others comparing different types of trails, we suggest a mixed management approach may be best, that involves hardening trails in popular areas, but with minimal disturbance during construction, the use of local materials where possible and vegetation rehabilitation once built. In less popular areas and/or those of particularly high biodiversity and conservation value such as Grey Box grassy-woodland, leaving trails unhardened but investing effort into more strategic trail placement and sustainable trail design may help maintain on-trail conditions and limit the spread of informal trails. These broad recommendations will help assist natural area managers to more effectively account for both recreation and conservation demands, which is of particular importance in already threatened ecosystems such as Grey Box grassy-woodland.

Acknowledgements

The authors wish to thank the Adelaide and Mount Lofty Ranges Natural Resources Management Board for contributing funding towards this project as well as the South Australian Department of Environment, Water and Natural Resources for providing field equipment, spatial data and access to the research sites. Griffith University also provided funding for this project. Thanks also to Mr James McGregor for his assistance in the field, in species identification and for providing spatial data for Belair National Park and to James Michael Arthur for his assistance with statistical analyses.

List of Figure Legends

Fig 1 Recreational trails impact natural vegetation in three main ways: through the construction and maintenance of the initial trail corridor: (1), through maintained presence of the exposed trail corridor leading to changes in local abiotic conditions (2) and finally, through the use of the corridor by recreational activities (3). These processes cause impacts at

various scales from direct local effects on the trail (A), to indirect local effects at the trail edge (B) and to dispersed landscape scale effects that occur cumulatively across the trail network (C).

Fig 2 Map of Belair National Park (thick white boundary line), south-eastern Adelaide, South Australia (35°01'22.70" S, 138°04'84.80" E) showing the location of Grey Box grassy-woodland remnants (highlighted patches to north-west of Belair National Park) and recreational trail networks within these remnants. All spatial data obtained from the Department of Environment, Water and Natural Resources, South Australia (DEWNR). Map created using ESRI ArcMap Version 10.1.

Fig 3 Effect of five different types of trails on species richness and vegetation composition in Grey Box grassy-woodland. Lines signify changes in species richness (large dashed line) and species composition (small dotted line) adjacent the edge, 5 m and 10 m from the edge of the trail and among the five trail types. Species richness values for each distance are given above each graph while the line for species composition symbolises the nature of the change without respect to denominative changes in variables.

References

- Australian Bureau of Statistics (2013) Regional population growth for South Australia 2011-2012.
<http://www.abs.gov.au/ausstats/abs@.nsf/Previousproducts/3218.0Main%20Features72011-12?opendocument&tabname=Summary&prodno=3218.0&issue=2011-12&num=&view=>. Accessed 26 March 2015.
- Australian Government (2010) Commonwealth listing advice on Grey Box (*Eucalyptus microcarpa*) grassy-woodlands and derived native grasslands of south-eastern Australia. Department of the Environment, Water, Heritage and the Arts.
<http://www.environment.gov.au/cgi-bin/sprat/public/publicshowcommunity.pl?id=86>. Accessed 11 March 2015.
- Australian Government (2012) Grey Box (*Eucalyptus microcarpa*) grassy-woodlands and derived native grasslands of south-eastern Australia: a guide to the identification, assessment and management of a nationally threatened ecological community (EPBC Act, 1999).

623 [http://www.environment.gov.au/system/files/resources/e6041636-388e-40cc-9bd4-](http://www.environment.gov.au/system/files/resources/e6041636-388e-40cc-9bd4-8c8b2dbe6419/files/grey-box-booklet.pdf)
624 [8c8b2dbe6419/files/grey-box-booklet.pdf](http://www.environment.gov.au/system/files/resources/e6041636-388e-40cc-9bd4-8c8b2dbe6419/files/grey-box-booklet.pdf). Accessed 21 January 2015.

625 Ballantyne M, Gudes O, Pickering CM (2014) Recreational trails are an important cause of
626 fragmentation in endangered urban forests: a case-study from Australia. *Landsc Urban Plan*
627 130:112-124.

628 Ballantyne M, Pickering CM (2015a) The impacts of trail infrastructure on vegetation and
629 soils: current literature and future directions. *J Env Manage* 164:53-64.

630 Ballantyne M, Pickering CM (2015b) Differences in the impacts of formal and informal
631 recreational trails on urban forest loss and tree structure. *J Env Manag* 159:94-105.

632 Ballantyne M, Treby DL (2015) The impacts of recreational trails in Grey Box grassy-
633 woodland in the Adelaide region. Department of Environment, Water and Natural Resources,
634 South Australia.

635 Balmford A, Beresford J, Green J, Naidoo R, Walpole M, Manica A (2009) A global
636 perspective of trends in nature-based tourism. *PLoS Biol* 7:1-6.

637 Baret S, Strasberg D (2005) The effects of opening trails on exotic plant invasion in protected
638 areas on La Réunion island (Mascarene Archipelago, Indian Ocean). *Revue d'Écologie*
639 60:325-332.

640 Benninger-Truax M, Vankat JL, Schaefer RL (1992) Trail corridors as habitat and conduits
641 for movement of plant species in Rocky Mountain National Park, Colorado, USA. *Landsc*
642 *Ecol* 6:269-278.

643 Bernhardt-Römermann M, Gray A, Vanbergen AJ, Bergés L, Bohner A, Brooker RW, De
644 Bruyn L, De Cinti B, Dirnböck T, Grandin U, Hester AJ, Kanka R, Klotz S, Loucougaray G,
645 Lundin L, Matteucci G, Mészáros I, Oláh V, Preda E, Prévosto B, Pykälä J, Schmidt W,
646 Taylor ME, Vadineanu A, Waldmann T, Stadler J (2011) Functional traits and local
647 environment predict vegetation responses to disturbance: a pan-European multi-site
648 experiment. *J Ecol* 99:777-787.

649 Brooks ML, Lair B (2005) Ecological effects of vehicular routes in a desert ecosystem. US
650 Department of the Interior, Geological Survey, Western Ecological Research Center, Las
651 Vegas, USA.

652 Buckley R (2004) Environmental impacts of ecotourism. CABI Publishing, Wallingford, UK.

653 Buckley R, King N, Zubrinich T (2004) The role of tourism in spreading dieback disease in
654 Australian vegetation. In: Buckley R (ed) Environmental impacts of ecotourism. CABI
655 Publishing, Wallingford, UK, pp. 317-324.

656 Buckley R (2009) Ecotourism: principles and practises. CABI Publishing, Wallingford, UK.

657 Cahill KL, Marion JL, Lawson SR (2008) Exploring visitor acceptability for hardening trails
658 to sustain visitation and minimise impacts. *J Sustain Tourism* 16:232-245.

659 Cole DN, Bayfield NG (1993) Recreational trampling of vegetation: standard experimental
660 procedures. *Biol Conserv* 63:209-215.

661 Cole DN (2004) Impacts of hiking and camping on soils and vegetation: a review. In:
662 Buckley R (ed) Environmental impacts of ecotourism. CABI Publishing, Wallingford, UK,
663 pp. 41-60.

664 Delgado JD, Arroyo NL, Arévalo JR, Fernández-Palacios JM (2007) Edge effects of roads on
665 temperature, light, canopy cover and canopy height in laurel and pine forests (Tenerife,
666 Canary Islands). *Landsc Urban Plan* 81:328-240.

667 Department of Environment, Water and Natural Resources, (2003) Belair National Park
668 management plan, Mount Lofty Ranges, South Australia.
669 file:///C:/Users/s2731206/Downloads/PARKS_PDFS_BELAIR_MP.pdf. Accessed 21
670 January 2015.

671 Farrell TA, Marion JL (2001) Identifying and assessing ecotourism visitor impacts at eight
672 protected areas in Costa Rica and Belize. *Environ Conserv* 28:215-225.

673 Godefroid S, Koedam N (2004) The impact of forest paths upon adjacent vegetation: effects
674 of the path surfacing material on the species composition and soil compaction. *Biol Conserv*
675 119:405-419.

676 Hamberg L, Lehvävirta S, Minna ML, Rita H, Kotze DJ (2008) The effects of habitat edges
677 and trampling on understorey vegetation in urban forests in Helsinki, Finland. *Applied Veg*
678 *Sci* 11:83-98.

679 Hill W, Pickering CM (2006) Vegetation associated with different walking track types in the
680 Kosciuszko alpine area, Australia. *J Environ Manage* 78:24-34.

681 Hill W, Pickering CM (2009) Differences in resistance of three sub-tropical vegetation types
682 to experimental trampling. *J Environ Manage* 90:1305-1312.

683 Kim MK, Daigle JJ (2012) Monitoring of vegetation impact due to trampling on Cadillac
684 Mountain summit using high spatial resolution remote sensing data sets. *Environ Manage*
685 50:956-968.

686 Kraehenbuehl DN (1996) Pre-European vegetation of Adelaide: a survey from the Gawler
687 River to Hallett Cove. Nature Conservation Society of South Australia Publishing, Adelaide,
688 Australia.

689 Kuenzi, C. and McNeely, J., (2008) Nature-based tourism. In O Renn and KD Walker (eds.),
690 *Global Risk Governance*. IUCN Publishing, Gland, Switzerland. pp. 155.178.

691 Kutiel P, Zhevelev H, Harrison R (1999) The effect of recreational impacts on soil and
692 vegetation of stabilised Coastal Dunes in the Sharon Park, Israel. *Ocean Coast Manage*
693 42:1041-1060.

694 Laurance WF, Ferreira LV, Rankin-de Merona JM, Laurance SG, Hutchings RW, Lovejoy
695 TE (1998) Effects of forest fragmentation on recruitment patterns in Amazonian tree
696 communities. *Conserv Biol* 12:460-464.

697 Ledoux CB (2004) Determining safe clearing limits for skid road/trail construction. In:
698 Yaussy DA, Hix DM, Long RP, Goebel PC (eds), *Proceedings of the 14th Central Hardwood*
699 *Forest Conference*, Newtown Square, USA, pp. 148-153.

700 Lee ACK, Maheswaran R (2011) The health benefits of urban green spaces: a review of the
701 evidence. *J Public Health* 33:212-222.

702 Leung YF, Shaw N, Johnson K, Duhaime R (2002) More than a database: integrating GIS
703 data with the Boston Harbor Islands carrying capacity study. *George Wright Forum* 19:69-78.

704 Leung YF, Newburger T, Jones M, Kuhn B, Woiderski B (2011) Developing a monitoring
705 protocol for visitor-created informal trails in Yosemite National Park, USA. *Environ Manage*
706 47:93-106.

707 Leung YF, Pickering CM, Cole DN (2012) Informal trails and fragmentation effects: a
708 conceptual and research overview. In: Fredman P, Stenseke M, Liljendahl H, Mossing A,
709 Laven D (eds), *Sixth international conference on monitoring and management of visitors in*
710 *recreational and protected areas*, Stockholm, Sweden. pp. 358-359.

711 Malmivaara-Lämsä M, Hamberg L, Haapamäki E, Liski J, Kotze DJ, Lehvävirta S, Fritze H
712 (2008) Edge effects and trampling in boreal urban forest fragments - impacts on the soil
713 microbial community. *Soil Biol Biochem* 40:1612-1621.

714 Marion JL, Leung YF (2001) Trail resource impacts and an examination of alternative
715 assessment techniques. *J Park Recreat Admi* 19:17-37.

716 Marion JL, Leung YF (2004) Environmentally sustainable trail management. In: Buckley R
717 (ed), *Environmental impacts of ecotourism*, CABI Publishing, Wallingford, UK. pp. 229-244.

718 Marion JL, Leung YF (2011) Indicators and protocols for monitoring impacts of formal and
719 informal trails in protected areas. *J Tourism Leisure Studies* 17:215-236.

720 Matlack GR (1993) Sociological edge effects: spatial distribution of human impact in
721 suburban forest fragments. *Environ Manage* 17:829-835.

722 Mitcham City Council (2015) Belair National Park information sheet.
723 <http://www.mitchamcouncil.sa.gov.au/page.aspx?u=1698&t=uList&ulistId=0&c=9494>.
724 Accessed 26 March 2015.

725 Monz CA, Cole DN, Leung YF, Marion JL (2010) Sustaining visitor use in protected areas:
726 future opportunities in recreation ecology research based on the USA experience. *Environ*
727 *Manage* 45:551-562.

728 Müllerová J, Vítková M, Vitek O, (2011) The impacts of road and walking trails upon
729 adjacent vegetation: effects of road building materials on species composition in a nutrient
730 poor environment. *Sci Total Environ* 409:3839-3849.

731 Newsome D, Moore SA, Dowling RK (2013) *Natural area tourism: ecology, impacts and*
732 *management – second edition*. Channel View Publications, Bristol, UK.

733 Olive ND, Marion JL (2009) The influence of use-related, environmental and managerial
734 factors on soil loss from recreational trails. *J Environ Manage* 90:1483-1493.

735 Paton P (2008) Field testing – condition thresholds – Inland Grey Box (*Eucalyptus*
736 *microcarpa*) woodlands, South Australia. Department of Environment, Water, Heritage, and
737 the Arts, Canberra, Australia.

738 Pescott OL, Stewart GB (2014) Assessing the impact of human trampling on vegetation: a
739 systematic review and meta-analysis of experimental evidence. *PeerJ* DOI 10.7717/peerj.360.

740 Pickering CM (2010) Ten factors that affect the severity of environmental impacts of visitors
741 in protected areas. *AMBIO* 39:70-77.

742 Pickering CM, Mount A (2010) Do tourists disperse weed seed? A global review of
743 unintentional human-mediated terrestrial seed dispersal on clothing, vehicles and horses. *J*
744 *Sustain Tourism* 18:239-256.

745 Pickering CM, Hill W, Newsome D, Leung YF (2010a) Comparing hiking, mountain biking
746 and horse riding impacts vegetation and soils in Australia and the United States of America. *J*
747 *Environ Manage* 91:551-562.

748 Pickering CM, Castley JG, Hill W, Newsome D (2010b) Environmental, safety and
749 management issues of unauthorised trail technical features for mountain bicycling. *Landsc*
750 *Urban Plan* 97:58-67.

751 Pickering CM, Castley JG, Richardt K (2012) Informal trails fragmenting endangered
752 remnant vegetation in Australia. In: Fredman P, Stenseke M, Liljendahl H, Mossing A, Laven
753 D (eds), *Sixth international conference on monitoring and management of visitors in*
754 *recreational and protected areas*, Stockholm, Sweden. pp. 362-363.

755 Pohlman CL, Turton SM, Goosem M (2007) Edge effects of linear canopy openings on
756 tropical rain forest understory microclimate. *Biotropica* 39:62-71.

757 Potito AP, Beatty SW (2005) Impacts of recreation trails on exotic and ruderals species
758 distribution in grassland areas along the Colorado Front Range. *Environ Manage* 36:230-236.

759 Rickard CA, McLachlan A, Kerley GIH (1994) The effects of vehicular and pedestrian traffic
760 on dune vegetation in South Africa. *Ocean Coast Manage* 23:225-247.

761 Rossi SD, Byrne JA, Pickering CM (2015) The role of distance in peri-urban national park
762 use: Who visits them and how far do they travel? *App Geog* 63:77-88.

763 South Australian Government (2012) History of Belair National Park.
764 [http://www.environment.sa.gov.au/parks/Find_a_Park/Browse_by_region/Adelaide_Hills/Bel](http://www.environment.sa.gov.au/parks/Find_a_Park/Browse_by_region/Adelaide_Hills/Belair_National_Park/History)
765 [air_National_Park/History](http://www.environment.sa.gov.au/parks/Find_a_Park/Browse_by_region/Adelaide_Hills/Belair_National_Park/History). Accessed 16th April 2015.

766 Spenceley A, Kohl J, McArthur S, Myles P, Notarianni M, Paleczny D, Pickering C,
767 Worboys GL (2015) Visitor management. In: Worboys GL, Lockwood M, Kothari A, Feary
768 S, Pulsford I (eds), *Protected area governance and management*, ANU Press, Canberra,
769 Australia. pp. 715-750.

770 Spooner P, Lunt I, Robinson W (2002) Is fencing enough? The short-term effects of stock
 771 exclusion in remnant grassy-woodlands in southern NSW. *Ecol Manage Restor* 3:117-126.

772 Stenhouse RN (2004) Fragmentation and internal disturbance of native vegetation resources
 773 in the Perth metropolitan area, Western Australia. *Landsc Urban Plan* 68:389-401.

774 Swanwick C, Dunnett N, Woolley H (2003) Nature, role and value of green space in towns
 775 and cities: an overview. *Built Environ* 29:94-106.

776 Talbot LM, Turton SM, Graham AW (2003) Trampling resistance of tropical rainforest soils
 777 and vegetation in the wet tropics of north east Australia. *J Environ Manage* 69:63-69.

778 Taylor Cullity Lethlean (2009) Belair National Park trails master plan. Prepared for the
 779 Department of Environment, Water and Natural Resources, South Australia.
 780 <http://www.tcl.net.au/projects>. Accessed 20 May 2015.

781 US National Park Service (2014) National trails system database. Available on request from
 782 the US National Park Service. Accessed 9 December 2014.

783 Volaire F, Seddaiu G, Ledda L, Lelievre F (2009) Water deficit and induction of summer
 784 dormancy in perennial Mediterranean grasses. *Ann Bot* 103:1337-1346.

785 Wimpey J, Marion JL (2010) The influence of use, environmental and managerial factors on
 786 the width of recreational trails. *J Environ Manage* 91:2028-2037.

787 Wimpey J, Marion JL (2011) A spatial exploration of informal trail networks within Great
 788 Falls Park, VA. *J Environ Manage* 92:1012-1022.

789 World Wildlife Fund (2014) Mediterranean forests, woodlands and scrubs – terrestrial
 790 ecosystem information. <http://www.worldwildlife.org/biomes/mediterranean-forests-woodlands-and-scrubs>. Accessed 21 December 2014.

791
 792 Yates CJ and Hobbs RJ (1997) Temperate eucalypt woodlands: a review of their status,
 793 processes threatening their persistence and techniques for restoration. *Aust J Bot* 45:949-973.
 794