

# GENETIC LINKAGE MAPS OF CHICKEN CHROMOSOMES 6, 7, 8, 11 AND 13 FROM A BRAZILIAN RESOURCE POPULATION

Marcel Ambo<sup>1</sup>; Raquel de Lello Rocha Campos<sup>1</sup>; Ana Sílvia Meira Tavares Moura<sup>4</sup>; Clarissa Boschiero<sup>4</sup>; Millor Fernandes do Rosário<sup>2</sup>; Mônica Corrêa Ledur<sup>5</sup>; Kátia Nones<sup>6</sup>; Luiz Lehmann Coutinho<sup>3\*</sup>

<sup>1</sup>USP/ESALQ - Programa de Pós-Graduação em Ciência Animal e Pastagens.

<sup>2</sup>USP/ESALQ - Programa de Pós-Graduação em Genética e Melhoramento de Plantas.

<sup>3</sup>USP/ESALQ - Depto. de Zootecnia, C.P. 09 - 13418-900 - Piracicaba, SP - Brasil.

<sup>4</sup>UNESP/FMVZ - Depto. de Produção Animal - 18618-000 - Botucatu, SP - Brasil.

<sup>5</sup>Embrapa Suínos e Aves, C.P. 21 - 89700 - 000 - Concórdia, SC - Brasil.

<sup>6</sup>Crop & Food Research, Private Bag 4704 - Christchurch 8140 - New Zealand.

\*Corresponding author <llcoutin@esalq.usp.br>

**ABSTRACT:** A linkage map is essential not only for quantitative trait loci (QTL) mapping, but also for the organization and location of genes along the chromosomes. The present study is part of a project whose major objective is, besides from construction the linkage maps, the whole genome scan for mapping QTL for performance traits in the Brazilian experimental chicken population. Linkage maps of chicken chromosomes 6 to 8, 11 and 13 were constructed based on this population. The population was developed from two generations of crossbreeding between a broiler and a layer line. Fifty-one microsatellite markers were tested, from which 28 were informative: 4, 8, 7, 4 and 5 for chromosomes 6, 7, 8, 11 and 13, respectively. A SNP located in the leptin receptor gene was included for chromosome 8. Ten parental, 8 F<sub>1</sub> and 459 F<sub>2</sub> chickens from five full-sib families were genotyped with these markers. The number of total informative meioses per locus varied from 232 to 862, and the number of phase-known informative meioses from 0 to 764. Marker orders in the chromosomes coincided with those of the chicken consensus map, except for markers ADL0147 and MCW0213, on chromosome 13, which were inverted. The reduced number of phase-known informative meioses for ADL0147 (150) may be pointed out as a possible cause for this inversion, apart from the relative short distance between the two markers involved in the inversion (10.5 cM).

**Key words:** *Gallus gallus*, quantitative trait loci (QTL), genetic maps, microsatellite markers, animal breeding

## MAPAS DE LIGAÇÃO DOS CROMOSSOMOS 6, 7, 8, 11 E 13 DE UMA POPULAÇÃO BRASILEIRA DE GALINHA

**RESUMO:** O mapa de ligação além de ser fundamental no mapeamento de locos de características quantitativas (QTLs) é importante na organização e localização de genes distribuídos ao longo dos cromossomos. O presente estudo é parte de um trabalho cujo objetivo maior, é a análise de mapeamento de QTLs para características de desempenho no genoma de uma população experimental desenvolvida no Brasil. Com base nesta população foram construídos os mapas de ligação dos cromossomos 6 a 8, 11 e 13 da galinha. A população foi desenvolvida a partir de duas gerações de cruzamentos entre uma linhagem de corte e uma de postura. Foram testados 51 marcadores microssatélites, dos quais 28 foram informativos: 4, 8, 7, 4 e 5 dos cromossomos 6, 7, 8, 11 e 13, respectivamente. Um SNP localizado no gene do receptor da *leptina* foi incluído no cromossomo 8. Os 10 parentais, 8 F<sub>1</sub> e um total de 459 aves F<sub>2</sub> de cinco famílias de irmãos completos foram genotipados com estes marcadores. O número de meioses informativas totais por loco variou de 232 a 862 e o de meioses informativas de fase conhecida de 0 a 764. A ordem dos marcadores nos cromossomos coincidiu com a do mapa consenso da galinha, com exceção dos marcadores ADL0147 e MCW0213 do cromossomo 13 que tiveram sua ordem invertida. O número reduzido de meioses informativas de fase conhecida para o marcador ADL0147 (150) pode ser apontado como uma possível causa para a inversão, além da relativa proximidade entre os dois marcadores envolvidos na inversão (10,5 cM).

**Palavras-chave:** *Gallus gallus*, locos de características quantitativas (QTLs), mapas genéticos, marcadores microssatélites, melhoramento animal

## INTRODUCTION

Economically important traits are genetically controlled by many genes, and the genetic linkage map (LM) construction is essential for mapping quantitative trait loci (QTL). Microsatellite markers are commonly used in animal genetic mapping, due to easy amplification by PCR, a high polymorphic degree, and their codominant characteristic (Ferreira & Grattapaglia, 1998).

Several aspects regarding chickens makes this species extremely well suited to experiments aimed at the localization of QTLs, such as a short generation interval, the ability to generate large full-sib pedigrees, and the ease of obtaining large quantities of DNA from the nucleated blood cell (Groenen et al., 1998). Currently 2,306 loci on 53 linkage groups from a total size of 4,200 cM were identified in chickens (Schmid et al., 2005). Two of three reference populations are actively used for LM: East Lansing (EL) and Wageningen (WAU) (Schmid et al., 2005). However, the first consensus chicken LM (Groenen et al., 2000) was constructed based on these three populations. The first of them, Compton (C), was genotyped with 100 markers on 18 linkage groups, covering 585 cM (Bumstead & Palyga, 1992). The second map used the EL population and 98 markers in 19 linkage groups, generating a linkage map of 590 cM (Levin et al., 1994). And the last population, WAU, used 430 markers, comprising 28 linkage groups containing 3,062 cM (Groenen et al., 1998). The consensus linkage maps is constantly reviewed and updated, helping as a source for the scientific community.

Our group is conducting the whole chicken genome scanning to identify QTLs in an  $F_2$  population generate by crossing a broiler line and a layer line. However, before QTL mapping analyses it is necessary to identify informative markers and construct the LM. A linkage map based on genotyping information from 27 microsatellite markers positioned on chromosomes 6, 7, 8, 11 and 13 in the intercross is reported here, and will be used for mapping QTL controlling performance and carcass traits.

## MATERIAL AND METHODS

### Experimental population

To generate an  $F_2$  population, seven males from a broiler line (TT) were mated to seven females from a layer line (CC). Each male was mated to three unrelated females selected randomly to generate 21  $F_1$  families, with approximately 100 chicks per family, in 17 incubations, totalizing 2,063  $F_2$  animals. The construction of the genetic linkage map of chromosomes

6, 7, 8, 11 and 13, used the five informative families indicated in a previous selective genotyping study of chromosomes 1 to 5 in this population (Baron, 2004; Nones et al., 2005; Ruy et al., 2005). TT was selected for various generations for growth related traits, such as body weight, whereas CC was selected for egg production traits (Nones et al., 2006).

### DNA isolation

Animal's blood samples were collected in tubes containing EDTA 10% and stored in a  $-70^\circ\text{C}$  freezer. Blood samples were extracted from brachial vein in parents, and through bleeding at slaughter in  $F_1$  and  $F_2$  animals. Genomic DNA isolation was conducted using the *DNAzol*<sup>®</sup> reagent following manufacturer's protocol with minor modifications. DNA concentration in each sample was assessed by spectrophotometer and standardized to a final concentration of  $20 \text{ ng } \mu\text{L}^{-1}$ .

### Genotyping

Fluorescent primers were used in PCR to amplify DNA fragments, provided by *United States Poultry Genome Project* and *Roslin Institute, UK*. Fragment lengths were determined by the automatic sequencer *MEGABace1000* (GE HealthCare<sup>®</sup>). Each PCR sample was prepared using  $100 \text{ ng } \mu\text{L}^{-1}$  of DNA,  $4.0 \text{ mmol L}^{-1} \text{ MgCl}_2$ ,  $50 \text{ mmol L}^{-1} \text{ KCl}$ ,  $10 \text{ mmol Tris-HCl}$  (pH 8.5),  $400 \text{ } \mu\text{M}$  of each dNTP, 5 U Taq DNA Polymerase and 5 pmol of each primer, totalizing final volume of 25  $\mu\text{L}$ . The program used in the thermocycler for PCR reactions consisted in an initial denaturation at  $95^\circ\text{C}$  for 5 minutes, and 30 cycles of: 1 minute at  $95^\circ\text{C}$ , 1 minute at 50 to  $67^\circ\text{C}$  (annealing temperature) depending on primers sequence, and an extension of 1 minute at  $72^\circ\text{C}$ . After the 30 cycles, an extension at  $72^\circ\text{C}$  for 10 minutes was conducted.

Three to four markers were combined according to amplicon size and primer fluorescence. Samples were precipitated and resuspended with  $4.75 \text{ } \mu\text{L}$  loading solution and  $0.25 \text{ } \mu\text{L}$  internal standard *ET-ROX400*. Genotyping data were checked and corrected manually. A total of 28 markers (out of 51 tested) were used to genotype approximately 459  $F_2$  individuals from five full-sib families, 10 parental and 8  $F_1$  animals on chromosomes 6, 7, 8, 11 and 13 (4, 8, 7, 4 and 5, respectively) were genotyped.

### Linkage analysis

Genotyping data of the 28 informative markers were used in the linkage analysis, employing the CRIMAP software, version 2.4 (Green et al., 1990), which uses maximum likelihood procedures to estimate recombination fractions and the Kosambi function to convert them to map distances in centiMorgans (cM).

Initially, the *Twopoint* option of CRIMAP was used for two-point linkage analysis in which the recombination rate between each of the two most informative linked markers of a chromosome and each one of the other markers was estimated. The order of loci was obtained with the *Build* option. A pair of highly polymorphic linked loci was chosen to start marker's ordering. The other loci were progressively added to the map, placed in each possible position with respect to the loci already ordered. The best position was based on the highest  $\log_{10}$  likelihood and  $\text{LOD} > 3.0$ .

The order of different loci was checked using the *Flips2* option, to look for an erroneous order. Finally the *Chrompic* option was used to verify every combination event amount order markers and to identify errors and potential double-crossing overs. The maps were draw with the MapChart software version 2.1(Voorrips, 2002).

## RESULTS AND DISCUSSION

From 16 markers tested on chromosome 6, 11 could not be used because no amplification products were obtained, and one (ADL0040) was not informative (only one allele). Four markers were used in the construction of the linkage group. On chromosome 7, eight markers were informative, out of 15 tested. Six markers did not generate amplification products, and marker ADL0180 showed only one allele. On chromosome 8, 12 markers were tested, but only seven were informative. The leptin receptor (LEPR) SNP marker identified and genotyped by Ninov et al. (2006) was informative in three full-sib families, presenting two alleles. Chromosomes 11 and 13 counted with four and six markers respectively, and only for MCW0322 on chromosome 13 no amplification products were ob-

tained. Table 1 lists all markers used on these five chromosomes and their positions (cM) on the consensus map (Schmid et al., 2005). A total of 28 informative markers were employed on the construction of the linkage maps and the respective number of informative meioses and of phase-known informative meioses are shown on Table 2.

Map distances between markers given in this paper are sex-averaged distances in cM. An initial  $\text{LOD}_{\text{score}} = 0.0$  was accepted in *Twopoint* option to estimate the recombination rate between the pairs of markers. Recombination fractions between all markers from each of the five chromosomes were calculated.

The linkage group of chromosome 6 (Figure 1) covered 38 cM out of 146 cM from the consensus map (Schmid et al., 2000), and the average distance between markers was 12.7 cM, varying from 2.8 to 20 cM. Marker positions on the linkage group were concentrated in the central region, leaving gaps on chromosomes extremities. An effort to increase marker density in gap regions was made, but no informative markers were identified. A larger number of informative markers on these regions would contribute to a complete QTL analysis.

Linkage map of chromosome 7 (Figure 1) had 131.2 cM and average distance of markers was 18.7 cM (varying from 5.9 to 35.4 cM). In comparison, the consensus map (Schmid et al., 2000) had 165 cM and 19 fixed position markers were typed. The low information content of marker ADL0107 precluded the inclusion of this marker with a  $\text{LOD} > 3.0$ . A  $\text{LOD} = 2.0$  was accepted to allow linkage between markers ADL0107 (33% informative meioses phase known compared to total meioses, Table 2) and ADL0279. The markers followed the same order of the consensus map.

Table 1 - Markers selected for linkage map construction of chromosomes 6, 7, 8, 11 and 13.

Chromosome	Marker (Position cM)
6	ADL0323 (0.0); LEI0196 (26.8); MCW0176 (27); *ADL0142 (29); LEI0093 (33.6); *ADL0377 (47); ADL0138 (55); LEI0092 (67); LEI0097 (74); LEI0212 (80); ADL0040 (82); *MCW0250 (86); ROS0062 (88); *ROS0003 (91); LEI0192 (115); ABR0323 (146)
7	*LEI0064 (0.0); *ABR0326 (30); *ADL0107 (51); MCW0201 (79); MCW0183 (86); *ADL0279 (92); ROS0019 (101); *MCW0236 (109); ADL0180 (109); *ADL0109 (117); LEI0158 (120); MCW0092 (117-140); MCW0316(127); *ADL0315 (140); *ADL0169 (175)
8	*ABR0322 (0.0-14); MCW0305 (15); *MCW0095 (26); ADL0171 (26-27); MCW0100 (40-50); *ADL0154 (46); *ABR0345 (56); ADL0301 (80); *LEPR (89.6); LEI0136 (70-105); *ADL0172 (80-105); *MCW0351 (105)
11	*LEI0143 (0.0); *ADL0123 (22); *ADL0210 (54); *MCW0230 (88)
13	*MCW0213 (22); *ADL0147 (32); *LEI0251 (47); *MCW0110 (59); MCW0322 (67); *MCW0104 (74)

\*Informative markers

Table 2 - Informative microsatellite markers used in the linkage analysis, respective number of alleles, number of informative meioses and phase-known informative meioses.

Marker	GGA	Number of alleles	Number of informative meioses	Number of informative meioses (phase-known)
ROS0062	6	03	372	372
ROS0003	6	04	702	702
ADL0377	6	04	269	269
ADL0142	6	04	232	159
LEI0064	7	06	764	764
ABR0326	7	05	862	555
ADL0107	7	02	293	97
ADL0279	7	05	858	360
MCW0236	7	03	451	272
ADL0109	7	03	326	326
ADL0315	7	03	251	251
ADL0169	7	02	357	357
ABR0322	8	04	698	346
MCW0095	8	03	438	438
ADL0154	8	04	700	544
ABR0345	8	04	611	507
ADL0172	8	03	630	428
MCW0351	8	03	345	241
LEPR	8	02	396	0
LEI0143	11	03	463	325
ADL0123	11	04	620	620
ADL0210	11	04	478	478
MCW0230	11	04	608	432
ADL1047	13	03	307	150
MCW0213	13	06	606	606
LEI0251	13	05	602	430
MCW0110	13	05	612	612
MCW0104	13	05	626	446

Means of total and phase-known informative meioses of chromosome 8 were 545 and 358, respectively. Three out of seven markers are fixed in the consensus map and followed the same order of the consensus map (ADL0154, ABR0345 e MCW0351). The number of phase known informative meioses for MCW0095 (not fixed) was 100%. The remaining markers showed variation from 68% to 83% for phase known informative meioses compared to total meioses, with the exception of marker ABR0322 that showed 49.6%.

The average marker interval on chromosome 8 was 14.9 cM (Figure 1), varying between 0.5 and 40.3 cM, and the total length was 89.6 cM. The largest gap was between markers ABR0345 and ADL0172.

Linkage map of chromosome 11 (Figure 1) was in good agreement with the consensus linkage

map, showed the same marker order and a total length of 105.5 cM. The average distance between adjacent markers was 35.2 cM. The consensus map length was 88 cM. Differences in length could be explained by the use of different population crosses and number of markers (four were used in the present study, comparing to nine on the consensus map). These factors result in different estimates of recombination rates that contribute to map divergences.

The linkage map of chromosome 13 showed 57 cM, whereas the consensus map had 74 cM. An inversion of marker positions occurred between markers ADL0147 and MCW0213. In the study by Jennen et al. (2004) the linkage map for chromosome 13 had 54.8 cM and the first and last markers were MCW0104 and MCW0213, similar to this study.




Figure 1 - Linkage group of chromosome 6 and linkage maps of chromosomes 7, 8, 11 and 13, and their updated linkage consensus maps (Schmid et al., 2000).

Marker ADL0147, involved in the order inversion, presented a low number of phase known informative meioses compared to other markers (Table 2). This can make it difficult to determine the order of the marker in the linkage map. Nevertheless, the other marker involved in the order inversion (MCW0213) was highly informative. Additionally, the interval between these two markers was small (10.5 cM), which may have caused difficulties in determin-

ing the relative position of these two markers in the linkage map.

Linkage maps were generated using a high number of F<sub>2</sub> animals from the experimental population, resulting in an average number of 517 informative meioses. In WAU resource population maps, the average was 400 for 10 half-sib families with approximately 46 F<sub>2</sub> animals per family (Groenen et al., 1998). The EL and C resource population maps (52 and 56



backcrosses) showed even lower number of informative meioses, varying from 20 to 50, respectively. Therefore, the resource population used in this study provided a high number of informative meioses, generating reliable linkage maps for QTL mapping. The average numbers of informative meioses and of phase-known informative meioses of each chromosome (6, 7, 8, 11 and 13) were: 393.7 and 375.5; 520.2 and 372.8; 545.4 and 357.7; 542.2 and 463.7; 550.6 and 448.8, respectively. The addition of microsatellite markers, especially in the intervals that exceeded 20 cM, would greatly aid in the saturation and utility of this genetic map for QTL mapping.

## CONCLUSIONS

The procedures used on linkage map construction for this resource population showed valuable results when compared with data from the chicken consensus map. Average numbers of informative meioses were high, indicating that results were reliable.

## ACKNOWLEDGEMENTS

The current study was funded by Embrapa/PRODETAB and Fapesp (Fundação de Amparo à Pesquisa do Estado de São Paulo) that has also granted scholarships to Marcel Ambo, R.L.R. Campos, C. Boschiero, K. Nones and M.F. do Rosário. A scholarship from CNPq was granted to A.S.A.M.T. Moura. L.L. Coutinho is a recipient of a research productivity scholarship from CNPq.

## REFERENCES

- BARON, E.E. Identificação de QTLs nos cromossomos 2 e 4 que controlam características de desempenho e carcaça em aves (*Gallus gallus*). Piracicaba: USP/ESALQ, 2004. 96p. Tese (Doutorado).
- BUMSTEAD, N.; PALLYGA, J. A preliminary linkage map of the chicken genome. *Genomics*, v.13, p.690-697, 1992.
- FERREIRA, M.E.; GRATTAPAGLIA, D. **Introdução ao uso de marcadores em análise genética**. 3 ed. Brasília: CENARGEN, 1998. 220p.
- GREEN, P.; FALLS, K.; CROOKS, S. **CRI-MAP Program version 2.4**. St. Louis: Washington University School of Medicine, 1990.
- GROENEN, M.A.M.; CROOIJMANS, R.P.M.A.; VEENENDAAL, A.; CHENG, H.H.; SIWEK, M.; POEL, J.J. van der A comprehensive microsatellite linkage map of the chicken genome. *Genomics*, v.49, p.265-274, 1998.
- GROENEN, M.A.M.; CHENG, H.H.; BUMSTEAD, N.; BENKEL, B.F.; BRILES, W.E.; BURKE, T.; BURT, D.W.; CRITTENDEN, L.B.; DODGSON, J.; HILLEL, J.; LAMONT, S.J.; LEON, A.P.; SOLLER, M.; TAKAHASHI, H.; VIGNAL, A. A consensus linkage map of the chicken genome. *Genome Research*, v.10, p.137-147, 2000.
- JENNEN, D.G.J.; VEREIJKEN, A.L.J.; BOVENHUIS, H.; CROOIJMANS, R.P.M.A.; VEENENDAAL, A.; POEL, J.J. van der; GROENEN, M.A.M. Detection and localization of quantitative trait loci affecting fatness in broilers. *Poultry Science*, v.83, p.295-301, 2004.
- LEVIN, I.; SANTANGELO L.; CHENG, H.; CRITTENDEN, L. B.; DODGSON, J.B.; An autosomal genetic linkage map of the chicken. *Journal of Heredity*, v.85, p.79-85, 1994.
- NINOV, K.; LEDUR, M.C.; NONES, K.; CAETANO, A.R.; COLDEBELLA, A.; BERTOL, T.M.; COUTINHO, L.L. Mining of polymorphisms in the leptin receptor gene in two chicken lines and their association with performance and carcass traits. In: INTERNATIONAL CONFERENCE ON ANIMAL GENETICS, 30., Porto Seguro, 2006. **Proceedings**. Belo Horizonte: CBRA, 2006. CD-ROM.
- NONES, K.; LEDUR, M.C.; RUY, D.C.; BARON, E.E.; MOURA, A.S.A.M.T.; COUTINHO, L.L. Genetic linkage map of chicken chromosome 1 from a Brazilian resource population. *Scientia Agricola*, v.62, p.12-17, 2005.
- NONES, K.; LEDUR, M. C.; RUY, D.C.; BARON, E.E.; MELO, C.M.R.; MOURA, A. S.A.M.T.; ZANELLA, E.L.; BURT, D.W.; COUTINHO, L.L. Mapping QTLs on chicken chromosome 1 for performance and carcass traits in a broiler x layer cross. *Animal Genetics*, v.37, p.95-100, 2006.
- RUY, D.C. NONES, K.; BARON, E.E.; LEDUR, M.C.; MELO, C.M.R.; AMBO, M.; CAMPOS, R.L.R.; COUTINHO, L.L. Strategic marker selection to detect quantitative trait loci in chicken. *Scientia Agricola*, v.62, p.111-116, 2005.
- SCHMID, M.; NANDA, I.; GUTTENBACH, M.; STEINLEIN, C.; HOEHN, H.; SCHARTL, M.; HAAF, T.; WEIGEND, S.; FRIES, R.; BUERSTEDDE, J.-M.; WIMMERS, K.; BURT, D.W.; SMITH, J.; A'HARA, S.; LAW, A.; GRIFFIN, D.K.; BUMSTEAD, N.; KAUFMAN, J.; THOMPSON, P.A.; BURKE, T.A.; GROENEN, M.A.M.; CROOIJMANS, R.P.M.A.; VIGNAL, A.; FILLON, V.; MORRISON, M.; PITEL, F.; TIXIER-BOICHARD, M.; LADJALI-MOHAMMEDI, K.; HILLEL, J.; MAKI-TANILA, A.; CHENG, H.H.; DELANY, M.E.; BURNSIDE, J.; MIZUNO, S. First report on chicken genes and chromosomes 2000. *Cytogenetics and Cell Genetics*, v.90, p.169-218, 2000.
- SCHMID, M.; NANDA, I.; HOEHN, H.; SCHARTL, M.; HAAF, T.; BUERSTEDDE, J.M.; ARAKAWA, H.; CALDWELL, R.B.; WEIGEND, S.; BURT, D.W.; SMITH, J.; GRIFFIN, D.K.; MASABANDA, J.S.; GROENEN, M.A.; CROOIJMANS, R. P.; VIGNAL, A.; FILLON, V.; MORISSON, M.; PITEL, F.; VIGNOLES, M.; GARRIGUES, A.; GELLIN, J.; RODIONOV, A.V.; GALKINA, S.A.; LUKINA, N.A.; BEN-ARI, G.; BLUM, S.; HILLEL, J.; TWITO, T.; LAVI, U.; DAVID, L.; FELDMAN, M.W.; DELANY, M.E.; CONLEY, C.A.; FOWLER, V.M.; HEDGES, S.B.; GODBOUT, R.; KATYAL, S.; SMITH, C.; HUDSON, Q.; SINCLAIR, A.; MIZUNO, S. Second report on chicken genes and chromosomes 2005. *Cytogenetic and Genome Research*, v.109, p.415-479, 2005.
- VOORRIPS, R.E. MapChart: software for the graphical presentation of linkage maps and QTLs. *The Journal of Heredity*, v.93, p.77-78, 2002.

Received July 26, 2007

Accepted February 15, 2008