

*Disclosure statement:* The author has declared no conflicts of interest.

K. KNOBLOCH<sup>1</sup>

<sup>1</sup>*Plastic, Hand and Reconstructive Surgery, Hannover Medical School, Hannover, Germany*  
Accepted 4 September 2008

Correspondence to: K. Knobloch, Plastic, Hand and Reconstructive Surgery, Hannover Medical School, Carl-Neuberg-Str. 130625, Hannover, Germany. E-mail: kknobi@yahoo.com

- 1 Rees JD, Lichtwark GA, Wolman RL, Wilson AM. The mechanism for efficacy of eccentric loading in Achilles tendon injury; an *in vivo* study in humans. *Rheumatology* 2008;47:1493–7.
- 2 Mahieu NN, McNair P, Cools A, D'Haen C, Vandermeulen K, Witvrouw E. Effect of eccentric training on the plantar flexor muscle-tendon tissue properties. *Med Sci Sports Exerc* 2008;40:117–23.
- 3 Langberg H, Ellingsgaard H, Madsen T *et al.* Eccentric rehabilitation exercises increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scand J Med Sci Sports* 2007;17:298–9.
- 4 Shalabi A, Kristoffersen-Wiberg M, Aspelin P, Movin T. Immediate Achilles tendon response after strength training evaluated by MRI. *Med Sci Sports Exerc* 2004;36:1841–6.
- 5 Ohberg L, Alfredson H. Effects of neovascularisation behind the good results with eccentric training in chronic mid-portion tendinosis? *Knee Surg Sports Traumatol Arthrosc* 2004;12:465–70.
- 6 Knobloch K, Kraemer R, Lichtenberg A *et al.* Achilles tendon and paratendon microcirculation in midportion and insertional tendinopathy in athletes. *Am J Sports Med* 2006;34:92–7.
- 7 Knobloch K. Eccentric training in Achilles tendinopathy: is it harmful to tendon microcirculation? *Br J Sports Med* 2007;41:e2.
- 8 Knobloch K, Kraemer R, Jagodzinski M, Zeichen J, Meller R, Vogt PM. Eccentric training decreases paratendon capillary blood flow and preserves paratendon oxygen saturation in chronic Achilles tendinopathy. *J Orthop Sports Phys Ther* 2007;37:269–76.
- 9 Cook JL, Bass SL, Black JE. Hormone therapy is associated with smaller Achilles tendon diameter in active postmenopausal women. *Scand J Med Sci Sports* 2007;17:128–32.
- 10 Knobloch K, Schreibermueller L, Meller R, Busch KH, Spies M, Vogt PM. Superior Achilles tendon microcirculation in tendinopathy among symptomatic female versus male patients. *Am J Sports Med* 2008;36:509–14.

*Rheumatology* 2009;48:203

doi:10.1093/rheumatology/ken423

Advance Access publication 16 November 2008

### Comment on: The mechanism for efficacy of eccentric loading in Achilles tendon injury; an *in vivo* study in humans: reply

SIR, We would like to thank Dr Knobloch [1] for his kind comments regarding our paper concerning the importance of studying the underlying mechanism of efficacy of eccentric exercises [2].

Regarding the title of our paper we stand by the title used. We have directly compared the physiology of eccentric and concentric exercises as used in common protocols for the management of Achilles injury and we do not believe the title to be misleading. Indeed we believe it medically prudent to start with healthy subjects in a study such as this. We do of course agree that it would be desirable to extend the study to injured tendons, and we allude to this in the discussion where we pass comment on how the results may alter in tendinopathy of the Achilles tendon (Discussion, paragraph 4 [2]).

Knobloch makes reference to several papers which he believes address the issue of the therapeutic mechanism of eccentric loading. Whilst many of these papers do shed light on changes observed in Achilles tendons over time (such as the excellent paper published by Langberg and co-workers on the rates of collagen synthesis) [3], we believe his comment misses the central point of our paper. Our paper directly compares eccentric and concentric loading of the Achilles tendon. Only in one of the papers quoted

by Knobloch is eccentric loading actually compared with concentric loading (the paper by Shalabi *et al.* [4]). This paper compared MRI-determined volume of the Achilles tendon in both concentric and eccentric programmes; it found an increase in tendon size with both exercise regimes but no significant difference in tendon volume between the two exercise regimes and does not explain why eccentric exercises are superior. Furthermore one of the papers quoted by Knobloch does not even involve an exercise protocol [5]. We believe our paper remains the only paper to both directly compare eccentric and concentric loading and find a possible explanation for the efficacy of eccentric loading exercises.

The question of gender and tendinopathy is extremely interesting. Certainly there is evidence from the work of Cook and colleagues [6] that HRT may be protective against tendinopathy and it is interesting to compare this to rates of anterior cruciate ligament injury which are increased in women. In our study, we found no difference physiologically in the tendon loading patterns of our male and female subjects but we agree the roles of gender and tendinopathy are worthy of further research.

*Disclosure statement:* The authors have declared no conflicts of interest.

J. D. REES<sup>1,2</sup>, G. A. LICHTWARK<sup>1,3,4</sup>, R. L. WOLMAN<sup>1</sup>,  
A. M. WILSON<sup>1,3</sup>

<sup>1</sup>*Department of Sport and Exercise Medicine, Royal National Orthopaedic Hospital, Stanmore,* <sup>2</sup>*Defence Medical Rehabilitation Centre, Headley Court, Epsom,* <sup>3</sup>*Structure and Motion Laboratory, The Royal Veterinary College, Hatfield, UK and* <sup>4</sup>*School of Physiotherapy and Exercise Science, Griffith University, Brisbane, Australia*

Accepted 3 October 2008

Correspondence to: J. D. Rees, Defence Medical Rehabilitation Centre, Headley Court, Epsom, Surrey KT18 6JN, UK.  
E-mail: j.rees@doctors.org.uk

- 1 Knobloch K. Comment on: The mechanism for efficacy of eccentric loading in Achilles tendon injury; an *in vivo* study in humans. *Rheumatology* 2009;48:202–3.
- 2 Rees JD, Lichtwark GA, Wolman RL, Wilson AM. The mechanism for efficacy of eccentric loading in Achilles tendon injury; an *in vivo* study in humans. *Rheumatology* 2008;47:1793–7.
- 3 Langberg H, Ellingsgaard H, Madsen T *et al.* Eccentric rehabilitation exercises increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scand J Med Sci Sports* 2007;17:298–9.
- 4 Shalabi A, Kristoffersen-Wiberg M, Aspelin P, Movin T. Immediate Achilles tendon response after strength training evaluated by MRI. *Med Sci Sports Exerc* 2004;36:1841–6.
- 5 Knobloch K, Kraemer R, Lichtenberg A *et al.* Achilles tendon and paratendon microcirculation in midportion and insertional tendinopathy in athletes. *Am J Sports Med* 2006;34:92–7.
- 6 Cook JL, Bass SL, Black JE. Hormone therapy is associated with smaller Achilles tendon diameter in active postmenopausal women. *Scand J Med Sci Sports* 2007;17:128–32.

*Rheumatology* 2009;48:203–204

doi:10.1093/rheumatology/ken440

Advance Access publication 11 December 2008

### Comment on: Hypovitaminosis D among rheumatology outpatients in clinical practice

SIR, Mouyis *et al.* [1] highlighted the high prevalence of unrecognized hypovitaminosis D amongst rheumatology patients. The new, long-acting bisphosphonates, such as zoledronic acid, are being increasingly used in patients with osteoporosis or Paget's disease to improve acceptability and compliance. Patients with pre-existing vitamin D deficiency who receive intravenous bisphosphonates may develop severe hypocalcaemia. This has been described in patients with underlying malignancy and hypercalcaemic bone pain treated with intravenous