

2  
3 **Integrating Chemical Analysis and Bioanalysis to Evaluate the Contribution of Wastewater**  
4 **Effluent on the Micropollutant Burden in Small Streams**

5  
6 Peta A. Neale<sup>1,2,†</sup>, Nicole A. Munz<sup>3,4,†</sup>, Selim Aït-Aïssa<sup>5</sup>, Rolf Altenburger<sup>6</sup>, François Brion<sup>5</sup>, Wibke  
7 Busch<sup>6</sup>, Beate I. Escher<sup>1,2,6,7\*</sup>, Klára Hilscherová<sup>8</sup>, Cornelia Kienle<sup>9</sup>, Jiří Novák<sup>8</sup>, Thomas-Benjamin  
8 Seiler<sup>10</sup>, Ying Shao<sup>10</sup>, Christian Stamm<sup>3</sup>, Juliane Hollender<sup>3,4</sup>

9  
10 <sup>1</sup>Smart Water Research Centre, School of Environment, Griffith University, Southport QLD 4222,  
11 Australia

12 <sup>2</sup>The University of Queensland, National Research Centre for Environmental Toxicology (Entox),  
13 Brisbane QLD 4108, Australia

14 <sup>3</sup>Eawag, Swiss Federal Institute of Aquatic Science and Technology, CH-8600, Dübendorf,  
15 Switzerland

16 <sup>4</sup>Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, 8092 Zürich, Switzerland

17 <sup>5</sup>Institut National de l'Environnement Industriel et des Risques INERIS, Unité d'Ecotoxicologie,  
18 60550 Verneuil-en-Halatte, France

19 <sup>6</sup>UFZ – Helmholtz Centre for Environmental Research, 04318 Leipzig, Germany

20 <sup>7</sup>Eberhard Karls University Tübingen, Environmental Toxicology, Center for Applied Geosciences,  
21 72074 Tübingen, Germany

22 <sup>8</sup>Masaryk University, Research Centre for Toxic Compounds in the Environment (RECETOX),  
23 Kamenice 753/5, 62500 Brno, Czech Republic

24 <sup>9</sup>Swiss Centre for Applied Ecotoxicology Eawag-EPFL, Überlandstrasse 133, CH-8600, Dübendorf,  
25 Switzerland

26 <sup>10</sup>Department of Ecosystem Analysis, Institute for Environmental Research, RWTH Aachen  
27 University, 52074 Aachen, Germany

28  
29  
30 **Science of the Total Environment** <http://dx.doi.org/10.1016/j.scitotenv.2016.10.141>

31 ©2016. This manuscript version is made available under the CC-BY-NC-ND 4.0  
32 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

33  
34 \*corresponding author: [beate.escher@ufz.de](mailto:beate.escher@ufz.de); Ph: +49 341 235 1285

35 †Both Neale and Munz contributed equally to the manuscript and are co-first authors

36 **Abstract**

37 Surface waters can contain a range of micropollutants from point sources, such as wastewater  
38 effluent, and diffuse sources, such as agriculture. Characterizing the source of micropollutants is  
39 important for reducing their burden and thus mitigating adverse effects on aquatic ecosystems. In  
40 this study, chemical analysis and bioanalysis were applied to assess the micropollutant burden  
41 during low flow conditions upstream and downstream of three wastewater treatment plants  
42 (WWTPs) discharging into small streams in the Swiss Plateau. The upstream sites had no input of  
43 wastewater effluent, allowing a direct comparison of the observed effects with and without the  
44 contribution of wastewater. Four hundred and five chemicals were analyzed, while the applied  
45 bioassays included activation of the aryl hydrocarbon receptor, activation of the androgen receptor,  
46 activation of the estrogen receptor, photosystem II inhibition, acetylcholinesterase inhibition and  
47 adaptive stress responses for oxidative stress, genotoxicity and inflammation, as well as assays  
48 indicative of estrogenic activity and developmental toxicity in zebrafish embryos. Chemical  
49 analysis and bioanalysis showed higher chemical concentrations and effects for the effluent  
50 samples, with the lowest chemical concentrations and effects in most assays for the upstream sites.  
51 Mixture toxicity modeling was applied to assess the contribution of detected chemicals to the  
52 observed effect. For most bioassays, very little of the observed effects could be explained by the  
53 detected chemicals, with the exception of photosystem II inhibition, where herbicides explained the  
54 majority of the effect. This emphasizes the importance of combining bioanalysis with chemical  
55 analysis to provide a more complete picture of the micropollutant burden. While the wastewater  
56 effluents had a significant contribution to micropollutant burden downstream, both chemical  
57 analysis and bioanalysis showed a relevant contribution of diffuse sources from upstream during  
58 low flow conditions, suggesting that upgrading WWTPs will not completely reduce the  
59 micropollutant burden, but further source control measures will be required.

60

61 **Keywords:** wastewater; micropollutant; chemical analysis; bioassays; surface water; mixture  
62 modeling

63

64

65 **1. Introduction**

66 Surface waters can contain a wide range of micropollutants, including pesticides, pharmaceuticals,  
67 personal care products and industrial compounds (Loos et al., 2009; Moschet et al., 2015), which  
68 have the potential to adversely impact exposed ecosystems (Malaj et al., 2014; Stalter et al., 2013).  
69 To mitigate the effect of micropollutants on the aquatic environment, it is important to identify their  
70 sources, which can be either from point sources, such as wastewater effluent discharges, or diffuse  
71 sources, such as agriculture (Eggen et al., 2014; Maletz et al., 2013). This can help to inform  
72 solutions to reduce the concentration and bioactive fraction of micropollutants, i.e., the  
73 micropollutant burden, in surface waters, which may include upgrading wastewater treatment plants  
74 (WWTPs) or regulatory changes, such as banning certain chemicals (Schwarzenbach et al., 2006).

75 Water quality monitoring programs, such as requested under the European Union Water  
76 Framework Directive (European Commission, 2011; European Commission, 2012), typically focus  
77 on chemical analysis, which can provide useful information about the concentration and type of  
78 chemicals present in a sample. However, targeted chemical analysis alone has some limitations. It is  
79 unable to detect unidentified chemicals and transformation products or account for the mixture  
80 effects that can occur between the many compounds present in water. For a comprehensive view of  
81 the micropollutant burden, chemical analysis should be combined with bioanalysis. While bioassays  
82 cannot identify individual chemicals, they can provide information about the joint effect of all  
83 bioavailable active chemicals present in a sample, with more potent chemicals showing a greater  
84 effect in the assay (Escher and Leusch, 2012; Prasse et al., 2015; Wernersson et al., 2015). The  
85 development of bioanalytical tools for water monitoring requires adequate choice of biological  
86 endpoints and quality measures (Altenburger et al., 2015; Busch et al., 2016), with applied  
87 bioanalytical test batteries ideally including assays indicative of induction of xenobiotic  
88 metabolism, endocrine disruption, reactive modes of action, adaptive stress responses and  
89 cytotoxicity (Escher et al., 2014).

90 The complementary approach of chemical analysis and bioanalysis has been applied to  
91 monitor water quality and to evaluate WWTP and advanced water treatment plant efficiency  
92 (Creusot et al., 2014; Jállová et al., 2013; Margot et al., 2013; Tang et al., 2014). Applying bioassays  
93 and chemical analysis in parallel overcomes the limitations associated with the individual  
94 approaches and can reveal the presence of potent undetected chemicals and identify chemicals that  
95 contribute to the observed effect (Escher and Leusch, 2012). Mixture toxicity modeling can be used  
96 to determine the fraction of the observed effect that can be explained by detected chemicals using  
97 the bioanalytical equivalent concentration (BEQ) approach (Neale et al., 2015a). Bioanalytical  
98 equivalent concentrations from chemical analysis ( $BEQ_{chem}$ ) are calculated using the detected  
99 chemical concentration and their relative potency, which can be compared to bioanalytical

100 equivalent concentrations from bioassays ( $BEQ_{bio}$ ). For example, detected chemicals can often  
101 explain a high percentage of estrogenic activity (Leusch et al., 2014; Murk et al., 2002), while only  
102 a small fraction of non-specific effects or adaptive stress responses can typically be explained (Tang  
103 et al., 2013; Yeh et al., 2014).

104 In this study, both chemical analysis and bioanalysis were applied to assess the  
105 micropollutant burden in small streams upstream and downstream of three WWTPs, with the  
106 upstream sites not being affected by inputs of treated wastewater. The water samples were collected  
107 under low flow conditions to minimize the impact of diffuse sources. The analyzed chemicals were  
108 primarily pharmaceuticals and pesticides, with the other analyzed chemicals including biocides,  
109 food additives, illicit drugs, industrial chemicals and estrogens. The biological effects were  
110 evaluated using a suite of *in vitro* assays, which represent different cellular toxicity pathways,  
111 including xenobiotic metabolism, receptor-mediated effects, adaptive stress responses and  
112 cytotoxicity, as well as whole organism assays with algae and zebrafish embryos. Assays indicative  
113 of xenobiotic metabolism, such as activation of the aryl hydrocarbon receptor (AhR), and adaptive  
114 stress responses, such as the oxidative stress response, can respond to a range of compounds with  
115 different modes of action (Martin et al., 2010; US EPA, 2015). In contrast, assays indicative of  
116 receptor-mediated effects can provide information about the presence of chemicals with a common  
117 specific mode of action. For example, hormone-mediated effects including activation of the  
118 estrogen receptor (ER) and activation of the androgen receptor (AR) can detect natural and  
119 synthetic hormones, as well as other environmental endocrine disrupting compounds, which are  
120 often associated with wastewater (Vethaak et al., 2005). Further, assays indicative of inhibition of  
121 photosystem II (PSII) and of acetylcholinesterase (AChE) are more suitable to detect chemicals of  
122 an agricultural origin as they can detect PSII inhibiting herbicides (Escher et al., 2008a) and  
123 organophosphate and carbamate insecticides (Hamers et al., 2000), respectively.

124 The current study aimed to assess the impact of wastewater effluent on the micropollutant  
125 burden in small streams using a complementary chemical analysis and bioanalysis approach. Four  
126 hundred and five chemicals were analyzed and the applied test battery included assays indicative of  
127 activation of AhR, activation of the AR, activation of the ER, PSII and algal growth inhibition,  
128 AChE inhibition, mutagenicity and adaptive stress responses for oxidative stress, genotoxicity and  
129 inflammation, as well as assays indicative of estrogenic activity and developmental toxicity in  
130 zebrafish embryos. A mass balance approach was used to calculate the fraction of effluent  
131 downstream based on both chemical analysis and bioanalysis, while mixture toxicity modeling was  
132 applied to assess whether the detected chemicals were contributing to the biological effect.

133  
134

135 **2. Materials and Methods**

136 *2.1. Sampling*

137 Three WWTPs, Birmensdorf, Muri and Reinach, in Switzerland were the focus of the study. The  
138 location of the study sites is shown in Figure S1 of the Supplementary Material, with information  
139 about treatment processes, catchment size and land use provided in Table S1. All selected sites had  
140 an average dry weather flow rate greater than 10 L/s, had no input of treated wastewater upstream  
141 and greater than 20% wastewater input downstream during low flow conditions. For further details  
142 and exact sampling locations refer to Burdon et al. (2016) and Stamm et al. (2016). Within each site  
143 there was comparable land use (Table S1), morphology and river bank vegetation upstream and  
144 downstream and all WWTPs applied biological activated sludge treatment. At each site, grab water  
145 samples were collected in glass bottles upstream and downstream of the WWTP during low flow  
146 conditions, as well as from the effluent stream, on the 19<sup>th</sup> May 2014. Eighteen liters of water were  
147 collected at both the upstream and downstream sites, while 10.5 L of effluent was sampled, with all  
148 samples collected within an hour at each site. The upstream samples were collected directly before  
149 the WWTP, while the downstream samples were collected approximately 50 to 300 m from the  
150 effluent discharge once complete mixing was obtained, which was assessed by conductivity  
151 measurements. Nanopure water was used as the process control. All samples were stored in the dark  
152 at 4°C or -20°C prior to extraction, with samples stored at 4°C extracted within 96 h of sampling.

153

154 *2.2. Chemical Analysis*

155 400 individual chemicals in the water samples were analyzed using online solid phase extraction  
156 (SPE) liquid chromatograph-high resolution tandem mass spectrometry (LC-HR-MS/MS) based on  
157 the method described in Huntscha et al. (2012) with some modifications. Briefly, the water samples  
158 were thawed at room temperature and filtered with 0.7 µm pore size glass fiber filters (Whatman,  
159 United Kingdom). Chemicals sorbed to particulate matter were not considered in this study, with  
160 samples for bioanalysis also filtered prior to extraction. Effluent samples were diluted by a factor of  
161 4 with nanopure water to minimize matrix effects. After the addition of internal standards to  
162 account for any loss during sample preparation, 20 mL aliquots were automatically enriched using  
163 SPE, with the pH of the aliquots automatically adjusted to pH 7 with 80 µL of 0.5 M citrate buffer  
164 prior to enrichment. The online SPE cartridges were manually packed with 9 mg of Strata X-CW,  
165 Strata X-AW and Isolute Env+ at a ratio of 1:1:1.5 (Phenomenex, Switzerland; Biotage, Sweden) at  
166 the bottom and 9 mg of Oasis HLB (Waters, US) as the top layer in the enrichment flow direction.  
167 Elution was performed in back-flush mode with methanol containing 0.1% formic acid. Five  
168 additional estrogenic compounds were analyzed in the effluent samples using a dedicated LC-  
169 MS/MS method (Vermeirssen et al., 2005). The samples for estrogenic analysis were enriched using

170 LiChrolut EN RP-18 SPE cartridges (Merck, Germany) based on the protocol in Margot et al.  
171 (2013). The list of analyzed chemicals is shown in Table S2. Detection limits are provided in ng/L,  
172 but the detected chemicals were converted to molar units for mixture toxicity modeling.

173

### 174 *2.3. Sample Extraction for Bioanalysis*

175 To extract a wide range of contaminants from the water samples for bioanalysis, multi-layer SPE  
176 cartridges were manually filled with three layers: 1) 200 mg of EnviCarb, 2) 350 mg of Strata X-  
177 CW, Strata X-AW and Isolute Env+ at a ratio of 1:1:1.5 (Phenomenex, Switzerland; Biotage,  
178 Sweden), and 3) 200 mg of Oasis HLB (Waters, US) as the first material in the enrichment flow  
179 direction. To extract the large water volume needed for bioanalysis, several cartridges were  
180 extracted in parallel to enrich 7.5 L of effluent and 15 L of surface water upstream and downstream,  
181 respectively. The same volume of nanopure water was extracted for the process control. The  
182 extraction protocol was based on Kern et al. (2009) with some modifications. Briefly, the pH of the  
183 water samples was adjusted to 6.5-6.7 prior to filtration with 0.7  $\mu\text{m}$  pore size glass fiber filters (47  
184 mm diameter, Whatman, United Kingdom). The SPE cartridges were conditioned with 5 mL  
185 methanol followed by 10 mL nanopure water. For each cartridge 1.5 L of upstream or downstream  
186 sample or 0.5 L of effluent sample was extracted by passing the sample under vacuum. After  
187 drying, the cartridges were inverted to elute them in the opposite flow direction in three steps with 6  
188 mL of ethyl acetate/methanol (50:50) containing 0.5% ammonia, 3 mL of ethyl acetate/methanol  
189 (50:50) containing 1.7% formic acid and 2 mL of methanol. The combined neutral extracts were  
190 evaporated at 40°C under a stream of nitrogen and split into the corresponding volumes for  
191 bioanalysis, then evaporated to dryness prior to shipping. The composition of the offline SPE  
192 cartridge used for bioanalysis differed from the online SPE cartridges used for chemical analysis  
193 due to the higher volume requirements, but for comparison 14 of the analyzed chemicals were also  
194 analyzed in the offline SPE extracts after the addition of internal standard. The properties of the  
195 analyzed compounds included a range of octanol-water partition coefficients ( $\log K_{\text{OW}}$  -1.0 to 5.0)  
196 and speciation. Some variability was observed between the two methods for some compounds in the  
197 low pM concentration range, such as atrazine, but overall, the comparison indicates that the  
198 chemical concentration in the extracts used for bioanalysis was representative of the analyzed  
199 chemical concentration (Figure S2). In addition, samples for the AChE inhibition assay and the  
200 combined algae assay were enriched using LiChrolut EN RP-18 SPE cartridges (Merck, Germany)  
201 based on the protocol in Margot et al. (2013).

202

203

204

205 *2.4. Bioanalysis*

206 The water extracts were analyzed in 11 bioassays, representing 13 different endpoints. A summary  
207 of the applied bioassays can be found in Table 1, with detailed information provided in Section S1.

209 *2.5. Data Evaluation*

210 The data were evaluated using linear concentration-effect curves to determine the concentration  
211 causing 10% effect ( $EC_{10}$ ) or effect concentration causing an induction ratio of 1.5 ( $EC_{IR1.5}$ ) (Escher  
212 et al., 2014). The zebrafish embryo acute toxicity assay was tested at a single concentration and %  
213 effect was reported. For the adaptive stress response endpoints no maximum effect can be reached  
214 and the effect is reported as an induction ratio (IR). Concentration-effect curves for adaptive stress  
215 responses are typically linear up to an IR of 5 and therefore the  $EC_{IR1.5}$  is interpolated from a linear  
216 regression through IR 1, the control effect (Escher et al., 2012). For endpoints where the effects can  
217 be converted to % maximum effect, the concentration-effect curves typically have log-sigmoidal  
218 shapes and can be fitted with probit, logit or Weibull models (Scholze et al., 2001). Since log-  
219 sigmoidal concentration-effect curves may be considered linear in the lower portion of the curve,  
220 responses up to 30 to 40% effect can be plotted using linear concentration-effect curves (Escher et  
221 al., 2014). Thus, for consistency, in the current study, all assays were evaluated using linear  
222 concentration-effect curves with the equations described in detail in Escher et al. (2014). As the  
223 concentration of all chemicals in the sample was unknown, units of relative enrichment factor  
224 (REF) were used, which take into account sample enrichment by SPE and dilution in the assay  
225 (Escher and Leusch, 2012).

226 For mixture toxicity modeling,  $BEQ_{bio}$  was calculated using the EC values of the reference  
227 compound (ref) and the matching EC value of the sample (Equation 1). The error associated with  
228  $BEQ_{bio}$  was determined using error propagation.

$$BEQ_{bio} = \frac{EC_{10}(ref)}{EC_{10}(sample)} \text{ or } \frac{EC_{IR1.5}(ref)}{EC_{IR1.5}(sample)}$$

230 (1)

231  
232 To assess the contribution of the detected chemicals to the biological effect,  $BEQ_{bio}$  can be  
233 compared to  $BEQ_{chem}$ . This involves calculating the relative effect potency ( $REP_i$ ) of the detected  
234 chemicals using the EC value of the reference compound and the EC value of the detected chemical  
235 (i) (Equation 2). EC values were collected from the peer-reviewed literature or the US EPA  
236 ToxCast database (US EPA, 2015). EC values from the literature were usually provided as  $EC_{50}$   
237 values, while raw fluorescence data was available in the ToxCast MySQL database, allowing

238  $EC_{IR1.5}$  values to be calculated for the adaptive stress response assays using linear concentration-  
239 effect curves.  $EC_{10 \text{ absolute}}$  values for the activation of 10% maximum effect in the AR assay were  
240 derived using the 50% activity concentration ( $AC_{50}$ ) values provided in the ToxCast database and  
241 the hill top of the concentration-effect curve (Equation 3).

242

$$REP_i = \frac{EC_{50}(\text{ref})}{EC_{50}(i)} \text{ or } \frac{EC_{IR1.5}(\text{ref})}{EC_{IR1.5}(i)} \text{ or } \frac{EC_{10}(\text{ref})}{EC_{10}(i)}$$

243

(2)

$$\log EC_{10 \text{ absolute}} = \log AC_{50} + \log \left( \frac{10\%}{\text{Hill top-10\%}} \right)$$

244

(3)

245

246 The detected chemical concentration ( $C_i$ ) in molar units and the calculated  $REP_i$  (also derived from  
247 molar ratios) were used to determine  $BEQ_{\text{chem}}$  (Equation 4).

248

$$BEQ_{\text{chem}} = \sum_i^n REP_i \cdot C_i$$

249

(4)

250

251 To assess whether the chemical concentration or biological effects at the downstream sites deviated  
252 from what would be expected based on pure physical mixing of upstream water and WWTP  
253 effluent, the mean fraction of wastewater effluent downstream ( $f_{\text{eff}}$ ) was calculated using  $C_i$  or  
254  $BEQ_{\text{bio}}$  for the upstream, downstream and effluent samples (Equation 5). For chemical analysis only  
255 chemicals detected in all three samples (upstream, effluent and downstream) were used to calculate  
256  $f_{\text{eff}}$ , with the mean  $f_{\text{eff}}$  based on individual chemicals reported for each site. In some cases, there was  
257 no substantial difference in the detected concentration in the samples when considering analytical  
258 uncertainty, leading to a  $f_{\text{eff}}$  outside of the range of 0 to 1. These values, which were all in the low  
259 ng/L range, were excluded from the  $f_{\text{eff}}$  calculation. Likewise for  $f_{\text{eff}}$  derived from  $BEQ_{\text{bio}}$ , only  
260 those bioassays that were above the detection limit in all three samples were used. Therefore  
261 activation of AR and algal growth inhibition could not be included. The uncertainties associated  
262 with the  $f_{\text{eff}}$  estimates were quantified by error propagation of the measurement uncertainty, as  
263 described in detail in Section S2.

264

265



$$f_{\text{eff}} = \frac{C_{i, \text{down}} - C_{i, \text{up}}}{C_{i, \text{effluent}} - C_{i, \text{up}}} \text{ or } f_{\text{eff}} = \frac{\text{BEQ}_{\text{bio, down}} - \text{BEQ}_{\text{bio, up}}}{\text{BEQ}_{\text{bio, effluent}} - \text{BEQ}_{\text{bio, up}}}$$

266

(5)

267

### 268 3. Results and Discussion

#### 269 3.1. Chemical Analysis

270 Four hundred chemicals were analyzed in the upstream and downstream samples, while a subset of  
 271 78 of the 400 chemicals were also analyzed in the wastewater effluent. Further, 5 estrogenic  
 272 chemicals were only analyzed in the effluent samples, as these compounds are often present at or  
 273 below the analytical limit of quantification in surface water. In total, 191 chemicals were detected at  
 274 least once, with the detected concentrations provided in Table S3. Overall, 57 chemicals were  
 275 detected at least once in the upstream samples and 185 chemicals were detected at least once in the  
 276 downstream samples. Despite the fact that fewer chemicals were analyzed, the highest percentage  
 277 of detected chemicals and the highest total chemical concentrations were found in the effluent  
 278 sample at all three sites (Figure 1).

279 Based on the 78 chemicals analyzed in all samples, the sum concentration in the effluent  
 280 samples ranged from 74.7 nM (16.2 µg/L, 58 compounds) to 103 nM (23.1 µg/L, 62 compounds),  
 281 while the sum concentration in the downstream samples ranged from 18.5 nM (4.11 µg/L, 63  
 282 compounds) to 24.6 nM (5.47 µg/L, 65 compounds). The upstream samples had the lowest sum of  
 283 chemical concentrations and the lowest number of detected chemicals, with sum concentrations  
 284 ranging from 0.37 nM (0.10 µg/L, 16 compounds) to 2.18 nM (0.67 µg/L, 20 compounds). The  
 285 most prevalent chemical class in the effluent was corrosion inhibitors, followed by pharmaceuticals.  
 286 Pharmaceuticals and corrosion inhibitors were also the predominant chemicals detected downstream  
 287 with pesticide concentrations up to two orders of magnitude lower. The upstream chemical profile  
 288 differed from the effluent and downstream samples, with pesticides contributing to 43 to 90% of the  
 289 sum of chemical concentrations upstream (Figure S3). It should be noted that the concentration of  
 290 many pesticides was similar in both the upstream and downstream samples, but the presence of  
 291 other chemical classes meant that the contribution of pesticides to the total micropollutant  
 292 concentration downstream was low (1.3 to 13%). The difference in chemical profiles and sum of  
 293 concentrations upstream and downstream clearly shows the influence of wastewater effluent, as  
 294 well as other micropollutant sources (e.g. agriculture), on the receiving streams.

295

#### 296 3.2. Bioanalysis

297 The EC values, provided in units of REF, for all samples are shown in Table S4 and Figure S4,  
 298 along with the concentration-effect curves in Figure S5. To demonstrate the applicability of linear

299 concentration-effect curves, EC<sub>10</sub> values calculated from linear concentration-effect curves at low  
300 effect levels (<40%) were compared to EC<sub>10</sub> values determined by evaluating the same data using  
301 log-sigmoidal concentration-effect curves for the activation of ER (MELN), activation of AR and 2  
302 h PSII inhibition assays in Figure S6. Reasonable agreement between the two evaluation methods  
303 was observed, supporting the use of linear concentration-effect curves in the current study (Figure  
304 S6). Cell viability was measured in parallel to oxidative stress, p53 and NF-κB induction (Figures  
305 S7-S9) and only concentrations less than the cytotoxicity EC<sub>10</sub> were included in the linear  
306 concentration-effect assessment to determine EC<sub>IR1.5</sub>. Cytotoxicity masked induction for all samples  
307 in the p53 response assay (Figure S8), so it was not possible to derive EC<sub>IR1.5</sub> values. Previous  
308 studies have also found a small window between p53 induction and cytotoxicity for the assay  
309 (Neale et al., 2015c; Yeh et al., 2014). Cytotoxicity was also evaluated for the activation of AhR,  
310 activation of ER and activation of AR assays (data not shown) and only non-cytotoxic values were  
311 reported below.

312         Activation of ER (MELN) and NF-κB response assays were among the most responsive  
313 assays, with mutagenicity being the least responsive endpoint; effects were observed in the  
314 Birmensdorf and Reinach downstream samples only. The measurement of activation of ER using  
315 the human MELN assay identified WWTP effluents as a major source of estrogenic compounds to  
316 the aquatic environment at all studied sites. The estrogenic activities measured were similar among  
317 the three WWTP effluents and were systematically higher than at the upstream and downstream  
318 sites, the latter being clearly impacted by effluent discharges. In addition to activation of ER,  
319 activation of AR was also assessed, but was less responsive, with only the effluent samples and  
320 Muri downstream having a response in the assay. The MDA-kB2 cell line also contains  
321 glucocorticoid receptors (GR) (Wilson et al., 2002), but exposure of the samples in the presence of  
322 AR antagonist flutamide abolished the effect of the active samples, indicating that the observed  
323 effects are really androgenic and not related to activation of GR. Previous studies have also shown  
324 low or no AR activation in water samples compared to ER activation (Leusch et al., 2014; Scott et  
325 al., 2014). Further, as some environmental estrogenic compounds are also antiandrogens, this could  
326 lead to an underestimation of AR activation (Sohoni and Sumpter, 1998).

327         While this study further confirmed the usefulness of reporter gene estrogenic assays to  
328 identify and characterize sources of estrogenic compounds in aquatic systems, it is also important to  
329 determine whether fish can be affected after exposures to such levels of estrogenic contamination.  
330 For example, Sonavane et al. (2016) recently showed fish-specific estrogenic responses in some  
331 environmental samples when comparing human and zebrafish *in vitro* ER activation assays. As a  
332 consequence, the samples were also tested in the transgenic *cyp19a1b*-GFP zebrafish embryo assay.  
333 At the upstream and downstream sites, where quantifications of the ER activation using the MELN

334 assay were low, no induction of the brain aromatase was observed. Conversely, Muri effluent  
335 induced a concentration-dependent induction of the ER-regulated brain aromatase gene showing  
336 that the level of estrogenic contamination was sufficient to induce an estrogenic response in the  
337 developing brain of fish. In contrast, no estrogenic activity was recorded in zebrafish for the  
338 Birmensdorf and Reinach effluents, despite all three effluents having similar ER activation  
339 responses in the MELN assay. This was due to the occurrence of mortality caused by the  
340 Birmensdorf and Reinach effluents after 96 h exposure at REFs as low as 4.5 and 4.8, respectively.  
341 These data highlight the relevance of using whole organism assays complementary to reporter gene  
342 assays, as it allows the identification of samples that can induce both estrogenic effects and/or  
343 developmental toxicity in fish.

344 All samples had an effect in the zebrafish embryo acute toxicity assay, though there was no  
345 significant difference in lethal effect between all sites after 120 h (Figure S10). Due to volume  
346 demands, the assay was only run at a single concentration, so EC values could not be determined.  
347 The applied REFs ranged from 25 to 50, which were higher than the maximum REF applied in the  
348 transgenic *cyp19a1b*-GFP zebrafish assay, where mortality was observed in the Birmensdorf and  
349 Reinach effluents at lower REFs (4.4 and 5.8, respectively). The result was also unusual as very  
350 little effect was observed at 24 and 48 h, though some sublethal effects, including malformations  
351 and reduced blood circulation, were observed at 48 h (Figure S10).

352 The majority of assays showed a clear trend of highest response in the effluent samples, with  
353 a reduced effect in the downstream samples and the lowest effect in the upstream samples. This  
354 result was expected and fits well with the detected chemical concentrations in Figure 1. One  
355 exception was the NF- $\kappa$ B response assay at Muri, where upstream was the most potent sample,  
356 followed by downstream and then effluent. The NF- $\kappa$ B response assay has only recently been  
357 applied for water quality monitoring (Escher et al., 2014; Neale et al., 2015a) and it is still unclear  
358 which environmental compounds activate this assay. For example, of the 191 detected chemicals,  
359 135 were included in the ToxCast database and all were either inactive or cytotoxic in the NF- $\kappa$ B  
360 response assay (US EPA, 2015). It should be pointed out that the NF- $\kappa$ B response assay in the  
361 ToxCast database is based on the ME180 cervical cancer cell line, while the THP-1 monocytic cell  
362 line was used in the current study. Thus, before this assay is used further for water quality  
363 monitoring, more work is required to better understand which environmental compounds are  
364 inducing a response in this assay.

365 In any case, the Muri upstream sample appears to be more potent than the other upstream  
366 samples, despite having a comparable sum chemical concentration (0.98 nM, 29 chemicals), with a  
367 higher response in the oxidative stress response assay and low level mortality in the *cyp19a1b*-GFP  
368 zebrafish embryo assay, which was not observed at the other upstream sites. Further, the algal

369 growth assay is of particular interest at the Muri upstream sample. For all other sites, algal growth  
370 inhibition was less sensitive than PSII inhibition and this fits with previous results in the literature  
371 where it was shown that PSII inhibiting herbicides typically dominate the biological effect on  
372 photosynthesis (Escher et al., 2008a; Tang and Escher, 2014). However, the opposite is observed at  
373 Muri upstream, with increased algal growth inhibition, which suggests that the contaminants are not  
374 only PSII inhibiting herbicides, but also potentially other herbicides with different modes of action,  
375 such as amino acid biosynthesis inhibition, or chemicals that have a non-specific effect on the algae.  
376 This example demonstrates the value of applying bioanalytical tools complementary to chemical  
377 analysis for water quality monitoring as they are able to indicate the presence of potent unidentified  
378 chemicals.

379 As well as containing micropollutants, environmental waters may also contain dissolved  
380 organic carbon (DOC), which may interfere with bioassays. For example, co-extracted DOC has  
381 previously been shown to cause experimental artefacts in the enzymatic AChE inhibition assay,  
382 with concentrations as low as 2 mg of carbon per liter ( $\text{mg}_C/\text{L}$ ) suppressing the agonist response  
383 (Neale and Escher, 2013). The DOC extraction efficiency of LiChrolut EN RP-18 SPE cartridges,  
384 which were used for the AChE inhibition assay, has not been assessed previously, but the  
385 commonly used Oasis HLB SPE cartridge can co-extract between 40-70% DOC (Neale and Escher,  
386 2013). The DOC concentration in the studied water samples ranged from 2.2 to 11.8  $\text{mg}_C/\text{L}$ , with  
387 samples enriched 500 to 1000 times by SPE. If only 10% of the DOC was co-extracted by SPE, this  
388 would yield a DOC concentration from 220 to 590  $\text{mg}_C/\text{L}$  in the extracts. After dilution in the assay,  
389 the lowest applied REF of 5.2 would still have a co-extracted DOC concentration of 2.6  $\text{mg}_C/\text{L}$ .  
390 This suggests that DOC is most likely interfering with the AChE inhibition assay and therefore the  
391 assay will not be used further for mixture toxicity modeling. The effect of DOC has been previously  
392 assessed for cell based assays, with negligible effect found in agonist mode (Neale and Escher,  
393 2014), though DOC can potentially interfere with assays run in antagonist mode (Neale et al.,  
394 2015b). Additional sample pre-treatment steps, such as fractionation, may help to reduce  
395 experimental artifacts associated with DOC (Ouyang et al., 2016).

### 396 397 *3.3. Assessing the Fraction of Wastewater Effluent Downstream*

398 The fraction of wastewater effluent downstream ( $f_{\text{eff}}$ ) was determined based on both chemical  
399 analysis and bioanalysis. Assuming pure mixing without any elimination during the small spatial  
400 and temporal range the same  $f_{\text{eff}}$  would be expected based on both chemical analysis and  
401 bioanalysis. For chemical analysis, mean  $f_{\text{eff}}$  were calculated from 10 to 11 individual chemicals  
402 that were detected in the upstream, effluent and downstream samples at each site (Table 2, Figure  
403 S11). Based on the individual chemicals at each site, mean  $f_{\text{eff}}$  ranged from  $0.20 \pm 0.06$  to  $0.30 \pm 0.10$ ,

404 which indicates a substantial influence of effluent on the downstream site. The chemical classes of  
405 the individual chemicals used to calculate mean  $f_{\text{eff}}$  included pesticides, pharmaceuticals, corrosion  
406 inhibitors and food additives.

407 For the  $f_{\text{eff}}$  calculated from bioassays, the EC values were converted to  $\text{BEQ}_{\text{bio}}$  (Table S5),  
408 which provides the concentration of a reference compound in molar units that would have the same  
409 response as the sample extract. Figure 2 shows a good agreement between the mean  $f_{\text{eff}}$  calculated  
410 from individual chemicals and  $f_{\text{eff}}$  calculated from  $\text{BEQ}_{\text{bio}}$  for activation of ER, 2h PSII inhibition  
411 and oxidative stress response. For example,  $f_{\text{eff}}$  for 2 h PSII inhibition ranged from  $0.22\pm 0.13$  to  
412  $0.26\pm 0.21$ , which fits within the range of  $f_{\text{eff}}$  calculated for individual PSII herbicides (0.15 to 0.45)  
413 (Table 2). In contrast,  $f_{\text{eff}}$  for activation of AhR at Muri and Reinach were higher than mean  $f_{\text{eff}}$   
414 based on chemical analysis, with  $f_{\text{eff}}$  up to  $0.87\pm 0.47$  at Reinach. However, there was high  
415 uncertainty associated with the calculation of  $f_{\text{eff}}$  for  $\text{BEQ}_{\text{bio}}$  for the AhR assay (coefficients of  
416 variance ranging from 56 to 258%). This was due to the small differences seen between the  
417  $\text{BEQ}_{\text{bio,up}}$  and  $\text{BEQ}_{\text{bio,down}}$  as well as  $\text{BEQ}_{\text{bio,up}}$  and  $\text{BEQ}_{\text{bio,eff}}$  for AhR at all sites resulting in very  
418 small numbers for both the nominator and denominator in Equation 5, with associated large errors  
419 of the resulting  $f_{\text{eff}}$ .

420 Further, it should be noted that grab samples were used in this study to derive  $f_{\text{eff}}$ . Therefore,  
421 potential daily variations in effluent discharge may alter  $f_{\text{eff}}$ , though the fact that  $f_{\text{eff}}$  was similar for  
422 all three sites supports the application of the current approach under low flow conditions.  
423 Nevertheless, overall we conclude that there is a reasonable agreement of the mass balance over  
424 upstream, downstream and effluent locations for  $\text{BEQ}_{\text{bio}}$  and the chemical analysis. This has been  
425 previously observed for chemical analysis, with Fairbairn et al. (2016) finding that the downstream  
426 composition could be generally explained by physical mixing of upstream river water and  
427 wastewater effluent using a mass balance approach.

428

### 429 3.4. Mixture Toxicity Modeling

430 Mixture toxicity modeling was applied to determine if the detected chemicals were contributing  
431 significantly to the observed effect. Between 2 and 26 EC values were found in the literature or  
432 ToxCast database for the detected chemicals in the studied assays, respectively. Out of the 191  
433 detected chemicals, 135 and 142 chemicals were included in the ToxCast database for the oxidative  
434 stress response and activation of AR assays, respectively (Table S6). For the oxidative stress  
435 response assay, 26 of the detected chemicals were active, 109 were inactive and no information was  
436 available for 56 chemicals, while only 6 were active in the activation of AR assay, with 136 inactive  
437 and no information available for 49 chemicals. As discussed above, none of the detected chemicals  
438 in the ToxCast database had a response in the NF- $\kappa$ B response assay so it was not possible to

439 determine  $BEQ_{chem}$ . Large screening datasets with known active and inactive chemicals were not  
440 available for the other studied assays, with available EC values instead collected from the literature.  
441 The available EC values were used to calculate  $REP_i$  values (Table S7), which, along with detected  
442 chemical concentrations, were used to calculate  $BEQ_{chem}$  (Table 3).  $BEQ_{bio}$  and  $BEQ_{chem}$  were  
443 compared to assess what fraction of the effect could be explained by detected chemicals (Table 3),  
444 with the percent contribution of individual detected chemicals shown in Figure 3 for activation of  
445 AhR, activation of ER (MELN), activation of AR, 2 h PSII inhibition, algal growth inhibition and  
446 oxidative stress response. The contribution of detected chemicals to 24 h PSII inhibition is shown in  
447 Figure S12. Overall, the availability of effect data for the analyzed compounds presented a  
448 limitation for mixture toxicity modeling.

449 By comparing  $BEQ_{bio}$  and  $BEQ_{chem}$ , up to 30% of AhR activation at the Birmensdorf site  
450 could be explained by the fungicide propiconazole, with the herbicide terbuthylazine contributing to  
451 12% of effect in the Muri effluent. With the exception of the insecticide diazinon, the literature EC  
452 values were determined using an AhR assay based on a mouse model, while an AhR assay based on  
453 a rat model was used in the current study. Consequently, differences in species sensitivity and  
454 selectivity may potentially limit the utility of mixture toxicity modeling for this assay.

455 Numerous studies have shown that a significant fraction of ER activation in wastewater and  
456 surface water can be explained by chemicals including natural and synthetic hormones,  
457 alkylphenols and phytoestrogens (Murk et al., 2002; Neale et al., 2015a; Rutishauser et al., 2004).  
458 However, only a small fraction of the effect could be explained for the ER activation assay in the  
459 current study as the chemical analysis focused primarily on emerging contaminants, such as  
460 pharmaceuticals and pesticides, rather than natural and synthetic estrogenic contaminants, with only  
461 five estrogenic chemicals analyzed in the effluent samples. Three estrogenic chemicals, estrone,  
462 bisphenol A and 4-nonylphenol, were detected in the effluent and could explain only up to 0.4% of  
463 the observed effect. Potent estrogenic chemicals  $17\alpha$ -ethinyl estradiol and  $17\beta$ -estradiol were also  
464 analyzed in wastewater effluent, but were not detected above the limit of quantification. The  
465 detection limits for  $17\alpha$ -ethinyl estradiol and  $17\beta$ -estradiol were in the low ng/L range (0.3 and 0.1  
466 ng/L, respectively), though the detection limit for  $17\alpha$ -ethinyl estradiol is still over an order of  
467 magnitude higher than the proposed environmental quality standard of 0.035 ng/L (European  
468 Commission, 2012). The measured  $BEQ_{bio}$  values are similar to previously measured  $BEQ_{bio}$  values  
469 for surface water and wastewater (Jugan et al., 2009) and shows that the activation of ER assay is a  
470 sensitive tool to detect the presence of estrogenic compounds and can be used complementary to  
471 chemical analysis that is often not sensitive enough for the low effect thresholds. In comparison,  
472 few studies have assessed the contribution of the detected chemicals to the activation of AR. Bellet  
473 et al. (2012) found that detected steroidal hormones, none of which were analyzed in the current

474 study, could only explain up to 5.5% of AR activity in raw wastewater. In the current study, EC  
475 values were available for six of the detected chemicals in the ToxCast database, with only up to  
476 0.4% of AR activity explained.

477 The majority of PSII inhibition at 2 h was explained by ten of the detected chemicals, with  
478 the herbicides diuron and terbuthylazine mostly contributing to the effect. Previous studies have  
479 also shown that detected herbicides can often explain the majority of PSII inhibition (Bengtson  
480 Nash et al., 2006; Escher et al., 2011) as only PSII herbicides will have an effect in the assay after  
481 the short exposure time. At all three sites, a lower fraction of the effect could be explained in the  
482 upstream samples, which could suggest the presence of additional undetected PSII inhibitors.  
483 Thirteen further PSII inhibitors, including bromacil and hexazinone, were analyzed in the upstream  
484 and downstream samples, but were not detected above the limit of quantification; however, it is  
485 possible that the presence of low concentrations of these compounds could still contribute to the  
486 effect. Further, herbicide transformation products may also contribute to the effect; however, as can  
487 be seen in Table S7, transformation products, such as atrazine-desethyl and atrazine-desisopropyl,  
488 are often less potent than their parent compounds. In contrast to PSII inhibition, only up to 73% of  
489 algal growth inhibition was explained by 7 detected compounds, which were all PSII inhibiting  
490 herbicides. Thus, the presence of other detected herbicides, such as mecoprop and metolachlor, may  
491 have contributed to the effect on growth rate, but EC values were not available in the studied assay.  
492 EC values for up to 26 of the detected chemicals were available in the ToxCast database, though  
493 only 1.9% of the oxidative stress response could be explained. This discrepancy has been observed  
494 previously for wastewater, surface water and swimming pool water (Escher et al., 2014; Neale et  
495 al., 2015a; Yeh et al., 2014) and demonstrates that many compounds can induce the oxidative stress  
496 response. For example, 22.6% of the 7522 analyzed chemicals in the ToxCast database were active  
497 in the oxidative stress response assay (US EPA, 2015). As different chemical classes can induce  
498 oxidative stress, it was possible to see different effect profiles upstream and downstream. For  
499 example, the pharmaceutical diclofenac only contributed to the effect in the effluent and  
500 downstream samples, while the fungicide propiconazole contributed more upstream than  
501 downstream at the Birmensdorf site. The herbicide metolachlor contributed to 1.5% and 0.7% of the  
502 oxidative stress response in the Muri effluent and downstream samples, respectively, with minimal  
503 contribution in the Muri upstream sample, indicating that wastewater discharge is the likely source.

504

#### 505 **4. Conclusions**

506 The combination of chemical analysis and bioanalysis proved to be a valuable complementary  
507 approach to monitor the micropollutant burden in the aquatic environment. Bioanalysis provided

508 information about the mixture effects of additional chemicals in the samples, while the chemical  
509 analysis showed differences in the chemical pollution profiles at the different sampling locations.

510 Mixture toxicity modeling was performed to assess the contribution of detected chemicals to  
511 the observed effect, with the fraction explained varying for the different assays. The lack of effect  
512 data for the detected micropollutants in the different assays was a major limitation and future work  
513 should focus on fingerprinting the effect of common water pollutants including the question of  
514 which chemicals and which biological endpoints a comprehensive effect analysis would encompass.

515 Further, this study shows that micropollutants, including pharmaceuticals, pesticides and  
516 corrosion inhibitors, are being discharged into small streams at nanomolar concentrations, with  
517 WWTPs as one of the main sources during low flow conditions. While less contaminated, the  
518 upstream sites were far from pristine, with agriculture contributing to the detected chemicals, as  
519 indicated by elevated concentrations of some pesticides detected, which was also reflected in the  
520 bioanalytical results. To our knowledge we have for the first time expanded a mass balance model  
521 (Equation 5) used to determine the fraction of effluent to the overall volume of the creeks from  
522 chemical analysis to bioassays. Results on the mass balance between up- and downstream samples  
523 in relation to input by effluent were consistent between chemical analysis and bioassays. Both  
524 chemical analysis and bioanalysis suggest that improved wastewater treatment technology will not  
525 completely reduce the micropollutant burden, which renders additional source control measures also  
526 necessary. Further, this study illustrates the relevance of the complementary approach to identify  
527 and characterize micropollutant sources.

528

## 529 **Acknowledgements**

530 The project SOLUTIONS is supported by the European Union Seventh Framework Programme  
531 (FP7-ENV-2013-two-stage Collaborative project) under grant agreement number 603437. Field and  
532 analytical work of EcoImpact was funded by Eawag, the Ecotox Centre and the Swiss Federal  
533 Office for the Environment (FOEN). Neale was supported by the National Health and Medical  
534 Research Council (NHMRC) – European Union Collaborative Research Grant (APP1074775). The  
535 research at RECETOX was also supported by the Ministry of Education, Youth and Sports of the  
536 Czech Republic (projects LO1214 and LM2011028). Brion and Aït-Aïssa were also supported by  
537 P181 and P190-Ecotoxicologie programmes from the French Ministry of Ecology. Shao was also  
538 supported by a personal grant from the scholarship program of the Chinese Scholarship Council.  
539 The authors thank Fabian Deuber, Laura Melo, Marta Reyes, Urs Schönenberger, Heinz Singer and  
540 Barbara Spycher (all Eawag) for support in collection and analysis of the water samples, the SPE  
541 and the GIS maps, as well as the whole EcoImpact team. Barbara Ganser, Sereina Gut, Beatrice  
542 Läubli, Janine Mazenauer (all Ecotox centre) are thanked for performing the AChE and algal



543 bioassays and Nadzeya Homazava (Ecotox Centre) is thanked for performing SPE as well as  
544 estrogen analysis. Janet Krüger (UFZ) is thanked for assistant with the zebrafish embryo acute  
545 toxicity assay and Nicolas Creusot and Benjamin Piccini (both INERIS) are thanked for their help  
546 with the ER assays. Henner Hollert (RWTH) is thanked for providing his experience with the Ames  
547 fluctuation test and for help in planning the experiments.

548

## 549 **References**

550 Altenburger, R., Ait-Aissa, S., Antczak, P., Backhaus, T., Barcelo, D., Seiler, T.-B., Brion, F.,  
551 Busch, W., Chipman, K., Lopez de Alda, M., de Aragao Umbuzeiro, G., Escher, B.I.,  
552 Falciani, F., Faust, M., Focks, A., Hilscherova, K., Hollender, J., Hollert, H., Jaeger, F.,  
553 Jahnke, A., Kortenkamp, A., Krauss, M., Lemkine, G.F., Munthe, J., Neumann, S.,  
554 Schymanski, E.L., Scrimshaw, M., Segner, H., Slobodnik, J., Smedes, F., Kughathas, S.,  
555 Teodorovic, I., Tindall, A.J., Tollefsen, K.E., Walz, K.-H., Williams, T.D., Van den Brink,  
556 P.J., van Gils, J., Vrana, B., Zhang, X., Brack, W., 2015. Future water quality monitoring -  
557 Adapting tools to deal with mixtures of pollutants in water resource management. *Sci. Total*  
558 *Environ.* 512, 540-551.

559 Balaguer, P., Francois, F., Comunale, F., Fenet, H., Boussioux, A.M., Pons, M., Nicolas, J.C.,  
560 Casellas, C., 1999. Reporter cell lines to study the estrogenic effects of xenoestrogens. *Sci.*  
561 *Total Environ.* 233, 47-56.

562 Bellet, V., Hernandez-Raquet, G., Dagnino, S., Seree, L., Pardon, P., Bancon-Montiny, C., Fenet,  
563 H., Creusot, N., Ait-Aissa, S., Cavailles, V., Budzinski, H., Antignac, J.-P., Balaguer, P.,  
564 2012. Occurrence of androgens in sewage treatment plants influents is associated with  
565 antagonist activities on other steroid receptors. *Water Res.* 46, 1912-1922.

566 Bengtson Nash, S.M., Goddard, J., Muller, J.F., 2006. Phytotoxicity of surface waters of the  
567 Thames and Brisbane River estuaries: a combined chemical analysis and bioassay approach  
568 for the comparison of two systems. *Biosens. Bioelectron.* 21, 2086-93.

569 Brion, F., Le Page, Y., Piccini, B., Cardoso, O., Tong, S.-K., Chung, B.-C., Kah, O., 2012.  
570 Screening estrogenic activities of chemicals or mixtures *in vivo* using transgenic (*cyp19a1b*-  
571 GFP) zebrafish embryos. *PLoS ONE.* 7, e36069.

572 Burdon, F.J., Reyes, M., Alder, A.C., Joss, A., Ort, C., Räsänen, K., Jokela, J., Eggen, R.I.L.,  
573 Stamm, C., 2016. Environmental context and magnitude of disturbance influence trait-  
574 mediated community responses to wastewater in streams. *Ecol. Evol.* 6, 3923–3939.

- 575 Busch, W., Schmidt, S., Kühne, R., Schulze, T., Krauss, M., Altenburger, R., 2016. Micropollutants  
576 in European rivers: A mode of action survey to support the development of effect-based  
577 tools for water monitoring. *Environ. Toxicol. Chem.* 35, 1887-1899.
- 578 Creusot, N., Ait-Aissa, S., Tapie, N., Pardon, P., Brion, F., Sanchez, W., Thybaud, E., Porcher, J.-  
579 M., Budzinski, H., 2014. Identification of synthetic steroids in river water downstream from  
580 pharmaceutical manufacture discharges based on a bioanalytical approach and passive  
581 sampling. *Environ. Sci. Technol.* 48, 3649-3657.
- 582 Eggen, R.I.L., Hollender, J., Joss, A., Schaerer, M., Stamm, C., 2014. Reducing the discharge of  
583 micropollutants in the aquatic environment: The benefits of upgrading wastewater treatment  
584 plants. *Environ. Sci. Technol.* 48, 7683-7689.
- 585 Ellman, G.L., Courtney, K.D., Andres, V., Featherstone, R.M., 1961. A new and rapid colorimetric  
586 determination of acetylcholinesterase activity. *Biochem. Pharmacol.* 7, 88-90.
- 587 Escher, B.I., Allinson, M., Altenburger, R., Bain, P.A., Balaguer, P., Busch, W., Crago, J.,  
588 Denslow, N.D., Dopp, E., Hilscherova, K., Humpage, A.R., Kumar, A., Grimaldi, M.,  
589 Jayasinghe, B.S., Jarosova, B., Jia, A., Makarov, S., Maruya, K.A., Medvedev, A., Mehinto,  
590 A.C., Mendez, J.E., Poulsen, A., Prochazka, E., Richard, J., Schifferli, A., Schlenk, D.,  
591 Scholz, S., Shiraish, F., Snyder, S., Su, G.Y., Tang, J.Y.M., van der Burg, B., van der  
592 Linden, S.C., Werner, I., Westerheide, S.D., Wong, C.K.C., Yang, M., Yeung, B.H.Y.,  
593 Zhang, X.W., Leusch, F.D.L., 2014. Benchmarking organic micropollutants in wastewater,  
594 recycled water and drinking water with *in vitro* bioassays. *Environ. Sci. Technol.* 48, 1940-  
595 1956.
- 596 Escher, B.I., Bramaz, N., Mueller, J.F., Quayle, P., Rutishauser, S., Vermeirssen, E.L.M., 2008a.  
597 Toxic equivalent concentrations (TEQs) for baseline toxicity and specific modes of action as  
598 a tool to improve interpretation of ecotoxicity testing of environmental samples. *J. Environ.*  
599 *Monitor.* 10, 612-621.
- 600 Escher, B.I., Bramaz, N., Quayle, P., Rutishauser, S., Vermeirssen, E.L.M., 2008b. Monitoring of  
601 the ecotoxicological hazard potential by polar organic micropollutants in sewage treatment  
602 plants and surface waters using a mode-of-action based test battery. *J. Environ. Monitor.* 10,  
603 622-631.

- 604 Escher, B.I., Dutt, M., Maylin, E., Tang, J.Y.M., Toze, S., Wolf, C.R., Lang, M., 2012. Water  
605 quality assessment using the AREc32 reporter gene assay indicative of the oxidative stress  
606 response pathway. *J. Environ. Monitor.* 14, 2877-2885.
- 607 Escher, B.I., Lawrence, M., Macova, M., Mueller, J.F., Poussade, Y., Robillot, C., Roux, A.,  
608 Gernjak, W., 2011. Evaluation of contaminant removal of reverse osmosis and advanced  
609 oxidation in full-scale operation by combining passive sampling with chemical analysis and  
610 bioanalytical tools. *Environ. Sci. Technol.* 45, 5387-5394.
- 611 Escher, B.I., Leusch, F.D.L., 2012. *Bioanalytical tools in water quality assessment*. IWA  
612 Publishing, London.
- 613 European Commission. 2011 Common implementation strategy for the water framework directive  
614 (2000/60/EC). Guidance document No. 27. Technical guidance for deriving environmental  
615 quality standards, Brussels.
- 616 European Commission. 2012 Proposal for a Directive of the European Parliament and the Council  
617 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the  
618 field of water policy, COM/2011/0876 final - 2011/0429 (COD), Brussels.
- 619 Fairbairn, D.J., Arnold, W.A., Barber, B.L., Kaufenberg, E.F., Koskinen, W.C., Novak, P.J., Rice,  
620 P.J., Swackhamer, D.L., 2016. Contaminants of emerging concern: Mass balance and  
621 comparison of wastewater effluent and upstream sources in a mixed-use watershed. *Environ.*  
622 *Sci. Technol.* 50, 36-45.
- 623 Hamers, T., Molin, K.R.J., Koeman, J.H., Murk, A.J., 2000. A small-volume bioassay for  
624 quantification of the esterase inhibiting potency of mixtures of organophosphate and  
625 carbamate insecticides in rainwater: Development and optimization. *Toxicol. Sci.* 58, 60-67.
- 626 Huntscha, S., Singer, H.P., McArdell, C.S., Frank, C.E., Hollender, J., 2012. Multiresidue analysis  
627 of 88 polar organic micropollutants in ground, surface and wastewater using online mixed-  
628 bed multilayer solid-phase extraction coupled to high performance liquid chromatography-  
629 tandem mass spectrometry. *J. Chromatogr. A.* 1268, 74-83.
- 630 Jállová, V., Jarošová, B., Bláha, L., Giesy, J.P., Ocelka, T., Grabic, R., Jurčíková, J., Vrana, B.,  
631 Hilscherová, K., 2013. Estrogen-, androgen- and aryl hydrocarbon receptor mediated  
632 activities in passive and composite samples from municipal waste and surface waters.  
633 *Environ. Int.* 59, 372-383.

- 634 Jin, L., Gaus, C., Escher, B.I., 2015. Adaptive stress response pathways induced by environmental  
635 mixtures of bioaccumulative chemicals in dugongs. *Environ. Sci. Technol.* 49, 6963-6973.
- 636 Jugan, M.L., Oziol, L., Bimbot, M., Huteau, V., Tamisier-Karolak, S., Blondeau, J.P., Levi, Y.,  
637 2009. *In vitro* assessment of thyroid and estrogenic endocrine disruptors in wastewater  
638 treatment plants, rivers and drinking water supplies in the greater Paris area (France). *Sci.*  
639 *Total Environ.* 407, 3579-3587.
- 640 Kern, S., Fenner, K., Singer, H.P., Schwarzenbach, R.P., Hollender, J., 2009. Identification of  
641 transformation products of organic contaminants in natural waters by computer-aided  
642 prediction and high-resolution mass spectrometry. *Environ. Sci. Technol.* 43, 7039-7046.
- 643 Kinani, S., Bouchonnet, S., Creusot, N., Bourcier, S., Balaguer, P., Porcher, J.-M., Ait-Aissa, S.,  
644 2010. Bioanalytical characterisation of multiple endocrine- and dioxin-like activities in  
645 sediments from reference and impacted small rivers. *Environ. Pollut.* 158, 74-83.
- 646 Leusch, F.D.L., Khan, S.J., Gagnon, M.M., Quayle, P., Trinh, T., Coleman, H., Rawson, C.,  
647 Chapman, H.F., Blair, P., Nice, H., Reitsema, T., 2014. Assessment of wastewater and  
648 recycled water quality: A comparison of lines of evidence from *in vitro*, *in vivo* and  
649 chemical analyses. *Water Res.* 50, 420-431.
- 650 Loos, R., Gawlik, B.M., Locoro, G., Rimaviciute, E., Contini, S., Bidoglio, G., 2009. EU-wide  
651 survey of polar organic persistent pollutants in European river waters. *Environ. Pollut.* 157,  
652 561-568.
- 653 Malaj, E., von der Ohe, P.C., Grote, M., Kühne, R., Mondy, C.P., Usseglio-Polatera, P., Brack, W.,  
654 Schäfer, R.B., 2014. Organic chemicals jeopardize the health of freshwater ecosystems on  
655 the continental scale. *PNAS.* 111, 9549-9554.
- 656 Maletz, S., Floehr, T., Beier, S., Kluemper, C., Brouwer, A., Behnisch, P., Higley, E., Giesy, J.P.,  
657 Hecker, M., Gebhardt, W., Linnemann, V., Pinnekamp, J., Hollert, H., 2013. *In vitro*  
658 characterization of the effectiveness of enhanced sewage treatment processes to eliminate  
659 endocrine activity of hospital effluents. *Water Res.* 47, 1545-1557.
- 660 Margot, J., Kienle, C., Magnet, A., Weil, M., Rossi, L., de Alencastro, L.F., Abegglen, C.,  
661 Thonney, D., Chevre, N., Scharer, M., Barry, D.A., 2013. Treatment of micropollutants in  
662 municipal wastewater: Ozone or powdered activated carbon? *Sci. Total Environ.* 461, 480-  
663 498.

- 664 Martin, M.T., Dix, D.J., Judson, R.S., Kavlock, R.J., Reif, D.M., Richard, A.M., Rotroff, D.M.,  
665 Romanov, S., Medvedev, A., Poltoratskaya, N., Gambarian, M., Moeser, M., Makarov, S.S.,  
666 Houck, K.A., 2010. Impact of environmental chemicals on key transcription regulators and  
667 correlation to toxicity end points within EPA's ToxCast program. *Chem. Res. Toxicol.* 23,  
668 578-590.
- 669 Moschet, C., Vermeirssen, E.L.M., Singer, H., Stamm, C., Hollender, J., 2015. Evaluation of in-situ  
670 calibration of Chemcatcher passive samplers for 322 micropollutants in agricultural and  
671 urban affected rivers. *Water Res.* 71, 306-317.
- 672 Murk, A.J., Legler, J., van Lipzig, M.M.H., Meerman, J.H.N., Belfroid, A.C., Spenkelink, A., van  
673 der Burg, B., Rijs, G.B.J., Vethaak, D., 2002. Detection of estrogenic potency in wastewater  
674 and surface water with three *in vitro* bioassays. *Environ. Toxicol. Chem.* 21, 16-23.
- 675 Nagy, S.R., Sanborn, J.R., Hammock, B.D., Denison, M.S., 2002. Development of a green  
676 fluorescent protein-based cell bioassay for the rapid and inexpensive detection and  
677 characterization of Ah receptor agonists. *Toxicol. Sci.* 65, 200-210.
- 678 Neale, P.A., Ait-Aissa, S., Brack, W., Creusot, N., Denison, M.S., Deutschmann, B., Hilscherova,  
679 K., Hollert, H., Krauss, M., Novak, J., Schulze, T., Seiler, T.-B., Serra, H., Shao, Y., Escher,  
680 B.I., 2015a. Linking *in vitro* effects and detected organic micropollutants in surface water  
681 using mixture-toxicity modeling. *Environ. Sci. Technol.* 49, 14614-14624.
- 682 Neale, P.A., Escher, B.I., 2013. Coextracted dissolved organic carbon has a suppressive effect on  
683 the acetylcholinesterase inhibition assay. *Environ. Toxicol. Chem.* 32, 1526-1534.
- 684 Neale, P.A., Escher, B.I., 2014. Does co-extracted dissolved organic carbon cause artefacts in cell-  
685 based bioassays? *Chemosphere.* 108, 281-288.
- 686 Neale, P.A., Escher, B.I., Leusch, F.D.L., 2015b. Understanding the implications of dissolved  
687 organic carbon when assessing antagonism *in vitro*: An example with an estrogen receptor  
688 assay. *Chemosphere.* 135, 341-346.
- 689 Neale, P.A., Stalter, D., Tang, J.Y.M., Escher, B.I., 2015c. Bioanalytical evidence that chemicals in  
690 tattoo ink can induce adaptive stress responses. *J. Hazard. Mater.* 296, 192-200.
- 691 OECD. 2013. Test No. 236: Fish Embryo Acute Toxicity (FET) Test, OECD Guidelines for the  
692 Testing of Chemicals, Section 2. OECD Publishing, Paris.

- 693 Ouyang, X., Leonards, P.E.G., Tousova, Z., Slobodnik, J., de Boer, J., Lamoreet, M.H., 2016.  
694 Rapid screening of acetylcholinesterase inhibitors by effect-directed analysis using LC x LC  
695 fractionation, a high throughput *in vitro* assay, and parallel identification by time of flight  
696 mass spectrometry. *Anal. Chem.* 88, 2353-2360.
- 697 Pavlíková, N., Blahova, L., Klan, P., Bathula, S.R., Sklenar, V., Giesy, J.P., Blaha, L., 2012.  
698 Enantioselective effects of alpha-hexachlorocyclohexane (HCH) isomers on androgen  
699 receptor activity *in vitro*. *Chemosphere.* 86, 65-69.
- 700 Prasse, C., Stalter, D., Schulte-Oehlmann, U., Oehlmann, J., Ternes, T.A., 2015. Spoilt for choice:  
701 A critical review on the chemical and biological assessment of current wastewater treatment  
702 technologies. *Water Res.* 87, 237-270.
- 703 Reifferscheid, G., Maes, H.M., Allner, B., Badurova, J., Belkin, S., Bluhm, K., Brauer, F.,  
704 Bressling, J., Domeneghetti, S., Elad, T., Fluckiger-Isler, S., Grummt, H.J., Guertler, R.,  
705 Hecht, A., Heringa, M.B., Hollert, H., Huber, S., Kramer, M., Magdeburg, A., Ratte, H.T.,  
706 Sauerborn-Klobucar, R., Sokolowski, A., Soldan, P., Smital, T., Stalter, D., Venier, P.,  
707 Ziemann, C., Zipperle, J., Buchinger, S., 2012. International round-robin study on the Ames  
708 fluctuation test. *Environ. Mol. Mutagen.* 53, 185-197.
- 709 Rutishauser, B.V., Pesonen, M., Escher, B.I., Ackermann, G.E., Aerni, H.R., Suter, M.J.F., Eggen,  
710 R.I.L., 2004. Comparative analysis of estrogenic activity in sewage treatment plant effluents  
711 involving three *in vitro* assays and chemical analysis of steroids. *Environ. Toxicol. Chem.*  
712 23, 857-864.
- 713 Scholze, M., Boedeker, W., Faust, M., Backhaus, T., Altenburger, R., Grimme, L.H., 2001. A  
714 general best-fit method for concentration-response curves and the estimation of low-effect  
715 concentrations. *Environ. Toxicol. Chem.* 20, 448-457.
- 716 Schwarzenbach, R.P., Escher, B.I., Fenner, K., Hofstetter, T.B., Johnson, C.A., von Gunten, U.,  
717 Wehrli, B., 2006. The challenge of micropollutants in aquatic systems. *Science.* 313, 1072-  
718 1077.
- 719 Scott, P.D., Bartkow, M., Blockwell, S.J., Coleman, H.M., Khan, S.J., Lim, R., McDonald, J.A.,  
720 Nice, H., Nuggeoda, D., Pettigrove, V., Tremblay, L.A., Warne, M.S.J., Leusch, F.D.L.,  
721 2014. An assessment of endocrine activity in Australian rivers using chemical and *in vitro*  
722 analyses. *Environ. Sci. Pollut. Res.* 21, 12951-12967.

- 723 Sohoni, P., Sumpter, J.P., 1998. Several environmental oestrogens are also anti-androgens. *J.*  
724 *Endocrinol.* 158, 327-339.
- 725 Sonavane, M., Creusot, N., Maillot-Maréchal, E., Péry, A., Brion, F., Aït-Aïssa, S., 2016.  
726 Zebrafish-based reporter gene assays reveal different estrogenic activities in river waters  
727 compared to a conventional human-derived assay. *Sci. Total Environ.* 550, 934-939.
- 728 Stalter, D., Magdeburg, A., Quednow, K., Botzat, A., Oehlmann, J., 2013. Do contaminants  
729 originating from state-of-the-art treated wastewater impact the ecological quality of surface  
730 waters? *PLoS ONE.* 8, e60616.
- 731 Stamm, C., Räsänen, K., Burdon, F.J., Altermatt, F., Jokela, J., Joss, A., Ackermann, M., Eggen,  
732 R.I.L., 2016. Unraveling the impacts of micropollutants in aquatic ecosystems: cross-  
733 disciplinary studies at the interface of large-scale ecology. *Adv. Ecol. Res.* In Press,  
734 <http://dx.doi.org/10.1016/bs.aecr.2016.07.002>
- 735 Tang, J.Y.M., Buseti, F., Charrois, J.W.A., Escher, B.I., 2014. Which chemicals drive biological  
736 effects in wastewater and recycled water? *Water Res.* 60, 289-299.
- 737 Tang, J.Y.M., Escher, B.I., 2014. Realistic environmental mixtures of micropollutants in surface,  
738 drinking, and recycled water: Herbicides dominate the mixture toxicity towards algae.  
739 *Environ. Toxicol. Chem.* 33, 1427-1436.
- 740 Tang, J.Y.M., McCarty, S., Glenn, E., Neale, P.A., Warne, M.S.J., Escher, B.I., 2013. Mixture  
741 effects of organic micropollutants present in water: Towards the development of effect-  
742 based water quality trigger values for baseline toxicity. *Water Res.* 47, 3300-3314.
- 743 US EPA. 2015 Interactive Chemical Safety for Sustainability (iCSS) Dashboard v2,  
744 <http://actor.epa.gov/dashboard/>, Accessed 5th Jan 2016.
- 745 Vermeirssen, E.L.M., Korner, O., Schonenberger, R., Suter, M.J.F., Burkhardt-Holm, P., 2005.  
746 Characterization of environmental estrogens in river water using a three pronged approach:  
747 Active and passive water sampling and the analysis of accumulated estrogens in the bile of  
748 caged fish. *Environ. Sci. Technol.* 39, 8191-8198.
- 749 Vethaak, A.D., Lahr, J., Schrap, S.M., Belfroid, A.C., Rijs, G.B.J., Gerritsen, A., de Boer, J.,  
750 Bulder, A.S., Grinwis, G.C.M., Kuiper, R.V., Legler, J., Murk, T.A.J., Peijnenburg, W.,  
751 Verhaar, H.J.M., de Voogt, P., 2005. An integrated assessment of estrogenic contamination

752 and biological effects in the aquatic environment of The Netherlands. *Chemosphere*. 59,  
753 511-524.

754 Wernersson, A.-S., Carere, M., Maggi, C., Tusil, P., Soldan, P., James, A., Sanchez, W., Dulio, V.,  
755 Broeg, K., Reifferscheid, G., Buchinger, S., Maas, H., Van Der Grinten, E., O'Toole, S.,  
756 Ausili, A., Manfra, L., Marziali, L., Polesello, S., Lacchetti, I., Mancini, L., Lilja, K.,  
757 Linderoth, M., Lundeberg, T., Fjällborg, B., Porsbring, T., Larsson, D.J., Bengtsson-Palme,  
758 J., Förlin, L., Kienle, C., Kunz, P., Vermeirssen, E., Werner, I., Robinson, C.D., Lyons, B.,  
759 Katsiadaki, I., Whalley, C., den Haan, K., Messiaen, M., Clayton, H., Lettieri, T., Carvalho,  
760 R.N., Gawlik, B.M., Hollert, H., Di Paolo, C., Brack, W., Kammann, U., Kase, R., 2015.  
761 The European technical report on aquatic effect-based monitoring tools under the water  
762 framework directive. *Env. Sci. Eur.* 27, 1-11.

763 Wilson, V.S., Bobseine, K., Lambright, C.R., Gray, L.E., 2002. A novel cell line, MDA-kb2, that  
764 stably expresses an androgen- and glucocorticoid-responsive reporter for the detection of  
765 hormone receptor agonists and antagonists. *Toxicol. Sci.* 66, 69-81.

766 Yeh, R.Y.L., Farre, M.J., Stalter, D., Tang, J.Y.M., Molendijk, J., Escher, B.I., 2014. Bioanalytical  
767 and chemical evaluation of disinfection by-products in swimming pool water. *Water Res.*  
768 59, 172-184.

769



770 Table 1: Summary of studied bioassays (REF: relative enrichment factor).

Endpoint	Assay	Method reference	Positive reference compound	EC value	Positive reference	Maximum REF
					compound EC <sub>10</sub> or EC <sub>IR1.5</sub> (M)	
Activation of AhR	CAFLUX	Nagy et al. (2002)	2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin (TCDD)	EC <sub>10</sub>	$(1.24 \pm 0.22) \times 10^{-12}$	100
Activation of ER (MELN)	MELN	Balaguer et al. (1999); Kinani et al. (2010)	17 $\beta$ -Estradiol	EC <sub>10</sub>	$(1.74 \pm 0.44) \times 10^{-12}$	84.7
Activation of ER (zebrafish)	cyp19a1b-GFP	Brion et al. (2012)	17 $\beta$ -Estradiol	EC <sub>IR1.5</sub>	$(1.29 \pm 0.07) \times 10^{-10}$	17
Activation of AR	MDA-kB2	Wilson et al. (2002); Pavlíková et al. (2012)	Dihydrotestosterone (DHT)	EC <sub>10</sub>	$(3.12 \pm 0.50) \times 10^{-11}$	50
AChE inhibition	AChE inhibition	Ellman et al. (1961); Escher et al. (2008b)	Parathion	EC <sub>10</sub>	$(1.80 \pm 0.16) \times 10^{-8}$	333
2 and 24 h PSII inhibition, algal growth inhibition	Combined algae assay with <i>Pseudokirchneriella subcapitata</i>	Escher et al. (2008a)	Diuron	EC <sub>10</sub>	2 h: $(2.58 \pm 0.81) \times 10^{-9}$ 24 h: $(2.68 \pm 0.45) \times 10^{-9}$ Growth: $(1.62 \pm 0.50) \times 10^{-8}$	267
Mutagenicity (TA98 + S9)	Ames fluctuation test	Reifferscheid et al. (2012)	2-Aminoanthracene	EC <sub>10</sub>	$(7.52 \pm 2.57) \times 10^{-8}$	400
Oxidative stress response	ARE- <i>bla</i>	Neale et al. (2015a)	tert-Butylhydroquinone (tBHQ)	EC <sub>IR1.5</sub>	$(2.44 \pm 0.24) \times 10^{-6}$	125
p53 response	p53RE- <i>bla</i>	Neale et al. (2015c)	Mitomycin	EC <sub>IR1.5</sub>	$(4.53 \pm 0.15) \times 10^{-8}$	125
NF- $\kappa$ B response	NF- $\kappa$ B- <i>bla</i>	Jin et al. (2015)	Tumor necrosis factor alpha	EC <sub>IR1.5</sub>	$(2.00 \pm 0.40) \times 10^{-2\dagger}$	125

Mortality	Zebrafish embryo toxicity test	OECD (2013)	(TNF $\alpha$ ) 3,4-Dichloroaniline	-	-	50
-----------	--------------------------------	-------------	--	---	---	----

771 NB: No EC value could be derived for the zebrafish embryo toxicity test as it was tested at a single concentration only.

772 †in units of  $\mu\text{g/L}$

773 Table 2: The fraction of wastewater effluent downstream ( $f_{\text{eff}}$ ) for individual chemicals detected at  
 774 the three sites.

<b>Chemical</b>	<b>Birmensdorf</b>	<b>Muri</b>	<b>Reinach</b>
1H-Benzotriazole	0.22		0.31
5-Methyl-1H-benzotriazole	0.23	0.23	0.32
Atrazine		0.21	
Atrazine-desethyl		0.45	
Azoxystrobin			0.24
Carbendazim	0.23		0.36
Caffeine		0.21	
Cyprodinil	0.20		0.03
Diazinon	0.22		0.30
Diuron			0.35
Epoxyconazole		0.13	
Hydrochlorothiazide			0.30
Mecoprop	0.07	0.20	0.44
Metolachlor		0.23	
Propiconazole	0.29		
Simazine	0.15	0.22	
Sucralose			0.30
Sulfamethoxazole		0.23	
Tebuconazole		0.24	
Telmisartan	0.15		
Terbutylazine	0.25	0.23	0.35
<i>Mean ± standard deviation</i>	<i>0.20 ± 0.06</i>	<i>0.23 ± 0.08</i>	<i>0.30 ± 0.10</i>
<i>Number of individual chemicals (n)</i>	<i>10</i>	<i>11</i>	<i>11</i>

775

776 Table 3: BEQ<sub>bio</sub> and BEQ<sub>chem</sub> values in molar units (M) for all samples in the different assays with percent effect that can be explained by the detected  
 777 chemicals (% effect).

		Birmensdorf			Muri			Reinach		
		Upstream	Effluent	Down stream	Upstream	Effluent	Down stream	Upstream	Effluent	Down stream
Activation of AhR	BEQ <sub>bio</sub>	1.04×10 <sup>-13</sup>	1.96×10 <sup>-13</sup>	1.16×10 <sup>-13</sup>	9.43×10 <sup>-14</sup>	3.10×10 <sup>-13</sup>	2.00×10 <sup>-13</sup>	8.17×10 <sup>-14</sup>	2.38×10 <sup>-13</sup>	2.12×10 <sup>-13</sup>
	BEQ <sub>chem</sub>	3.10×10 <sup>-14</sup>	1.57×10 <sup>-14</sup>	2.61×10 <sup>-14</sup>	3.96×10 <sup>-16</sup>	3.88×10 <sup>-14</sup>	8.97×10 <sup>-15</sup>	1.83×10 <sup>-16</sup>	3.01×10 <sup>-15</sup>	5.92×10 <sup>-16</sup>
	% effect	29.7	8.0	22.5	0.4	12.5	4.5	0.2	1.3	0.3
Activation of ER (MELN)	BEQ <sub>bio</sub>	9.48×10 <sup>-13</sup>	7.28×10 <sup>-12</sup>	1.88×10 <sup>-12</sup>	1.65×10 <sup>-12</sup>	1.55×10 <sup>-11</sup>	3.49×10 <sup>-12</sup>	5.59×10 <sup>-13</sup>	1.53×10 <sup>-11</sup>	3.13×10 <sup>-12</sup>
	BEQ <sub>chem</sub>	N/A	2.27×10 <sup>-14</sup>	N/A	N/A	2.08×10 <sup>-14</sup>	N/A	N/A	5.56×10 <sup>-14</sup>	N/A
	% effect	-	0.3	-	-	0.1	-	-	0.4	-
Activation of ER (zebrafish)	BEQ <sub>bio</sub>	<9.84×10 <sup>-12</sup>	Mortality	<9.74×10 <sup>-12</sup>	<9.74×10 <sup>-12</sup>	7.89×10 <sup>-11</sup>	<9.77×10 <sup>-12</sup>	<7.60×10 <sup>-12</sup>	Mortality	<9.82×10 <sup>-12</sup>
	BEQ <sub>chem</sub>	N/A	3.90×10 <sup>-13</sup>	N/A	N/A	3.72×10 <sup>-13</sup>	N/A	N/A	1.06×10 <sup>-12</sup>	N/A
	% effect	-	-	-	-	0.5	-	-	-	-
Activation of AR	BEQ <sub>bio</sub>	<1.43×10 <sup>-12</sup>	8.89×10 <sup>-12</sup>	<1.42×10 <sup>-12</sup>	<1.42×10 <sup>-12</sup>	1.07×10 <sup>-11</sup>	2.97×10 <sup>-12</sup>	<1.25×10 <sup>-12</sup>	4.22×10 <sup>-11</sup>	<1.43×10 <sup>-12</sup>
	BEQ <sub>chem</sub>	1.60×10 <sup>-16</sup>	1.57×10 <sup>-14</sup>	3.32×10 <sup>-15</sup>	1.10×10 <sup>-16</sup>	3.80×10 <sup>-14</sup>	7.26×10 <sup>-15</sup>	2.09×10 <sup>-16</sup>	1.71×10 <sup>-15</sup>	5.01×10 <sup>-15</sup>
	% effect	-	0.2	-	-	0.4	0.2	-	0.04	-
2 h PSII Inhibition	BEQ <sub>bio</sub>	4.38×10 <sup>-11</sup>	5.53×10 <sup>-10</sup>	1.58×10 <sup>-10</sup>	2.61×10 <sup>-11</sup>	1.11×10 <sup>-9</sup>	2.87×10 <sup>-10</sup>	3.35×10 <sup>-11</sup>	2.84×10 <sup>-10</sup>	9.83×10 <sup>-11</sup>
	BEQ <sub>chem</sub>	1.96×10 <sup>-11</sup>	5.21×10 <sup>-10</sup>	1.30×10 <sup>-10</sup>	1.18×10 <sup>-11</sup>	1.20×10 <sup>-9</sup>	2.89×10 <sup>-10</sup>	2.01×10 <sup>-11</sup>	1.92×10 <sup>-10</sup>	8.91×10 <sup>-11</sup>
	% effect	44.9	94.3	82.3	45.1	108	101	59.9	67.3	90.7
24 h PSII Inhibition	BEQ <sub>bio</sub>	4.92×10 <sup>-11</sup>	7.97×10 <sup>-10</sup>	2.13×10 <sup>-10</sup>	3.06×10 <sup>-11</sup>	2.40×10 <sup>-9</sup>	7.97×10 <sup>-10</sup>	5.84×10 <sup>-11</sup>	4.72×10 <sup>-10</sup>	1.79×10 <sup>-10</sup>
	BEQ <sub>chem</sub>	1.68×10 <sup>-11</sup>	6.96×10 <sup>-10</sup>	1.60×10 <sup>-10</sup>	1.30×10 <sup>-11</sup>	1.39×10 <sup>-9</sup>	3.32×10 <sup>-10</sup>	1.78×10 <sup>-11</sup>	2.72×10 <sup>-10</sup>	1.28×10 <sup>-10</sup>
	% effect	34.2	87.3	75.1	42.4	57.6	41.7	30.6	57.7	71.4
Algal growth	BEQ <sub>bio</sub>	<1.80×10 <sup>-10</sup>	1.33×10 <sup>-9</sup>	4.54×10 <sup>-10</sup>	3.19×10 <sup>-10</sup>	6.73×10 <sup>-9</sup>	1.53×10 <sup>-9</sup>	<1.80×10 <sup>-10</sup>	1.24×10 <sup>-9</sup>	3.46×10 <sup>-10</sup>

inhibition	BEQ <sub>chem</sub>	3.04×10 <sup>-11</sup>	9.71×10 <sup>-10</sup>	2.20×10 <sup>-10</sup>	1.68×10 <sup>-11</sup>	1.83×10 <sup>-9</sup>	4.39×10 <sup>-10</sup>	2.65×10 <sup>-11</sup>	3.98×10 <sup>-10</sup>	1.98×10 <sup>-10</sup>
	% effect	-	73.0	48.6	5.30	27.2	28.7	-	32.2	57.2
Oxidative	BEQ <sub>bio</sub>	4.06×10 <sup>-8</sup>	1.43×10 <sup>-7</sup>	7.43×10 <sup>-8</sup>	5.89×10 <sup>-8</sup>	2.08×10 <sup>-7</sup>	1.09×10 <sup>-7</sup>	4.23×10 <sup>-8</sup>	2.73×10 <sup>-7</sup>	1.18×10 <sup>-7</sup>
Stress	BEQ <sub>chem</sub>	2.02×10 <sup>-10</sup>	6.51×10 <sup>-10</sup>	2.82×10 <sup>-10</sup>	3.53×10 <sup>-11</sup>	3.92×10 <sup>-9</sup>	9.06×10 <sup>-10</sup>	4.56×10 <sup>-11</sup>	5.77×10 <sup>-10</sup>	1.88×10 <sup>-10</sup>
Response	% effect	0.5	0.5	0.4	0.1	1.9	0.8	0.1	0.2	0.2

---

778

## List of Figures

**Figure 1:** Sum chemical concentration detected at each site (nM) with the percentage of analyzed chemicals detected at each site (open diamonds). Four hundred compounds were analyzed in the upstream and downstream sites and 83 compounds were analyzed in the effluents.

\*steroidal hormone estrone and two of the industrial chemicals, 4-nonylphenol and bisphenol A, were only analyzed in the effluents.

**Figure 2:** Fraction of wastewater effluent downstream ( $f_{\text{eff}}$ ) calculated using  $\text{BEQ}_{\text{bio}}$  values (error bars indicate standard deviation calculated using error propagation) compared to the mean  $f_{\text{eff}}$  based on individual chemicals (solid black lines, with dashed lines indicating standard deviation).

**Figure 3:** Percent effect explained by individual detected chemicals for A) activation of AhR, B) activation of ER (MELN), C) activation of AR, D) 2 h PSII inhibition, E) algal growth inhibition and F) oxidative stress response. Note the different scales for the y-axes.

\*estrogenic compounds were only measured in the effluent samples.

**Figure 1**




Figure 2






Figure 3

