

# Contaminant Exchange rates in estuaries – new formulae accounting for advection and dispersion

Fernando P. Andutta<sup>1,2</sup>, Peter V. Ridd<sup>1</sup>, Eric Deleersnijder<sup>3,4</sup>, David Prandle<sup>5</sup>

<sup>1</sup> School of Engineering and Physical sciences, James Cook University, Townsville QLD 4811, Australia.

<sup>2</sup> School of Physical, Environmental and Mathematical Sciences, University of New South Wales at Australian Defence Force Academy UNSW-ADFA, ACT 2600, Australia.

<sup>3</sup> Université catholique de Louvain (UCL), Institute of Mechanics, Materials and Civil Engineering (IMMC), 4 Avenue Georges Lemaître, Bte L4.05.02, B-1348 Louvain-la-Neuve, Belgium

<sup>4</sup> Université catholique de Louvain (UCL), Earth and Life Institute (ELI), Georges Lemaître Centre for Earth and Climate Research (TECLIM), 3 Place Louis Pasteur, Bte L4.03.08, B-1348 Louvain-la-Neuve, Belgium

<sup>5</sup> National Oceanography Centre, Joseph Proudman Building, 6 Brownlow Street, Liverpool L3 5DA, UK.

\* Corresponding author. Email: [fernando.andutta@my.jcu.edu.au](mailto:fernando.andutta@my.jcu.edu.au); [f.andutta@adfa.edu.au](mailto:f.andutta@adfa.edu.au); [andutta@usp.br](mailto:andutta@usp.br)

**Abstract** – The transport timescales of water in an estuary are important measurements for monitoring pollution threats to the estuarine ecosystem. In this study we re-evaluated the application of simple analytical solutions to estimate these timescales and found that the Land Ocean Interaction Coastal Zone model (LOICZ) uses similar equation as from the Fresh Water Fraction model, and thus often resulting in shortened transport timescales. Therefore, the LOICZ model is neither based upon the well-known Knudsen relation nor Fischer formulation. Three transport timescales, namely water renewal, residence time and exposure time were calculated using analytical solutions for a range of estuaries worldwide. The analytical results were compared with available estimates of residence times from numerical models. The theoretical formulation from the LOICZ, the fresh water fraction model, and a newly proposed modified LOICZ model were used to calculate water renewal. Residence times and exposure times were calculated using the Constituent-oriented Age and Residence time Theory (CART). The modified LOICZ model was found to be the most comparable to residence times from numerical models, with  $r^2 \sim 0.7$ . In addition to the proposed modified LOICZ model (which uses Fischer formulation), we have developed an

advection-dispersion timescale diagram. This graphic conceptual model provides a visual representation of the relative contribution of advective and dispersive processes to water renewal for different estuaries. Estuaries can be categorized as either dominated by dispersion, dominated by advection, or having dispersion and advection of similar magnitude.

Keywords: LOICZ model; CART model; Fresh water fraction model; water renewal, residence time; exposure time; return coefficient.

## 1. Introduction

The transport timescales of water in an estuary are important measurements to analyse and estimate pollution threats to the estuarine ecosystem (Lucas et al., 2009; McLusky and Elliott, 2004; Wolanski, 2007). There are a number of defined transport timescales, namely (1) the flushing time, (2) the age, (3) the residence time, (4) the exposure time, and (5) the renewal time. The flushing time is defined as the time taken for a concentration to decrease to  $1/e$  ( $\sim 0.37$ ) of its initial value, i.e. the time necessary for  $\sim 63\%$  of a concentration (e.g. passive tracer particles) to cross the open boundary of an aquatic system (Ketchum, 1951; Dyer, 1973; Monsen et al., 2002; Deleersnijder et al., 2006; Valle-Levinson, 2010). The age is the time necessary for a water parcel to travel from a defined inlet boundary (e.g. the salinity intrusion limit or the tidal intrusion limit) to another specific location (e.g. the estuary mouth; Monsen et al., 2002). The residence time is generally defined as the time necessary for a water particle to exit the domain for the first time (Bolin and Rodhe 1973; Zimmerman 1976; Takeoka 1984; Buffoni et al., 1997; Falco et al., 2000; Poulain and Hariri, 2013). The residence time therefore varies spatially, depending on the starting location and time. For these timescale calculations, once water particles leave the estuary they are disregarded. In practice, however, some particles may return to the estuary with reversing tidal currents, after first exiting (Monsen et al., 2002). To incorporate this process in a timescale, exposure time is used. This can be defined as the total time that a parcel of water spends inside an aquatic environment (Delhez, 2006; Andutta et al., 2012; de Brye et al. 2012; Andutta et al., 2013). The difference between the exposure time and residence time depends on the circulation in coastal waters; for instance, swift longshore currents decrease the difference between the exposure and residence times (Wolanski, 2007). Previously, residence time had been defined as the time required for the volume of an estuary to be replaced with new water from both the ocean and the river (e.g. Dyer, 1973); we propose to call this the renewal time, as the definition of residence time has now changed. Transit time is also important (Bolin and Rodhe 1973; Zimmerman 1976), and may be defined as the time taken for a particle to cross from the inflow to the outflow open boundary of the domain. Thus the

transit time for some conditions would be similar to both the water renewal time and the residence time in the upstream location of an estuary.

These transport timescales (Fig. 1) are determined by the hydrodynamics of the system (Monsen et al., 2002; Delhez, 2006; Sheldon and Alber, 2006; Delhez and Deleersnijder, 2006; Lowe et al., 2009; Andutta et al., 2012; Andutta et al., 2013). Their precise values remain elusive, however, because the hydrodynamics are comprised not only of the water circulation, which is generally well known and can usually be modeled reliably, but also all of the unresolved processes, which are relatively unknown and generally assumed to amount to turbulent mixing.

Figure 1– Preferred position.

Estimating transport timescales, in even the simplest 1-D estuarine models, has proven to be a challenge to modelers. For an estuary located between  $x = L_I$  (the upstream limit) and  $x = L_0$  (the estuary mouth), where the estuarine segment length  $L = L_I - L_0$ , the residence time can be estimated from models using virtual passive particles. In the simplest case of cross-sectional homogeneity, and in the absence of baroclinic currents, a 1-D model may be sufficient; in which case the tracers have a concentration,  $C$ , that follows the equation

$$A \frac{\partial C}{\partial t} + Q \frac{\partial C}{\partial x} = \frac{\partial}{\partial x} \left( AK \frac{\partial C}{\partial x} \right), \quad (1)$$

where  $A(x)$  is the cross-sectional area of the estuary,  $t$  is the time,  $x$  is the distance along-channel,  $Q$  is the flow rate, and  $K$  is the along-channel mean dispersion. This mean dispersion is due to numerous processes in addition to turbulence that affects the timescales of interest (i.e. multiple tidal cycles). To calculate the residence time, the modeler seeds the model with tracers so that  $C = 1$  at  $t = 0$  everywhere at  $L_0 < x < L_I$ . Eq. (1) must be solved to calculate  $C$  as a function of  $x$  and  $t$ ; from this solution the various timescales can be calculated. None of these steps, however, are straightforward. Firstly, a hydrodynamic model is required to provide the data for  $Q$  as a function of  $x$  and  $t$ . Secondly, the modeler must also provide a closed-form formula or a model to estimate  $K$  as a function of  $x$  and  $t$ . Thirdly, the modeler must specify the open boundary conditions for  $C$  at  $x = L_I$  and  $x = L_0$ . To calculate the residence time, the open boundary conditions are in their most simple form, namely  $C=0$  at  $x = L_I$  and  $x = L_0$  (e.g. Delhez and Deleersnijder 2006). To compute the exposure time, the open boundary conditions for  $C$  at  $x = 0$  and  $x = L$  are unknown a priori; the usual practice is for the modeler to extend the model domain offshore ( $x > L$ ) in 2-D or 3-D, at a distance far enough that the estuary does not influence the adjacent circulation. The model should

105 also be extended upstream into the river (see e.g. de Brye et al. 2012). This requires knowledge of  
 106 the coastal circulation, which subsequently requires knowledge of the oceanic circulation offshore.  
 107 In the simplest case, three independent parameters control this timescale: the residual velocity,  $u =$   
 108  $Q_R / A$ , where  $Q_R$  is often assumed to be the river discharge and  $A$  is the cross-sectional area; the  
 109 estuarine segment length,  $L$ ; and the along-channel eddy dispersion coefficient,  $K$ . Two timescales  
 110 result therefrom (Fischer et al., 1979), namely an advective timescale,  $T_1 = L/u = V/Q_R$ , and a  
 111 dispersive timescale,  $T_2 = L^2/K$ . The relative importance of these two timescales is determined by  
 112 their ratio, which is the Peclet number  $P_E = T_2/T_1 = u L/K$ . While  $P_E$  is used in estuarine  
 113 classification schemes (Prandle, 2009), as yet no formula has been proposed to estimate the  
 114 residence time of an estuary as a function of  $P_E$ . The dispersive timescale,  $T_2$ , may be expressed as  
 115  $T_2 = P_E V/Q_R$ , taking into consideration the estuarine volume.

116 In view of these complexities, modelers have moved away from analytical solutions and  
 117 have increasingly used numerical models in 2-D and 3-D. Such models are elegant, but they still  
 118 carry uncertainties in terms of the hydrodynamic conditions and fate of tracers at the open  
 119 boundaries. Additionally, for quantifying dispersion, adjusting the horizontal dispersion coefficient  
 120 at the sub-grid scale is still a challenge (Fischer, 1969, 1974 and 1976). Although there are an  
 121 increasing number of physical oceanographers modelling estuaries, residence time has been  
 122 estimated with calibrated numerical models for only a few estuaries, while simple box models have  
 123 been applied to over 200 estuaries.

124 In view of this, we propose to re-evaluate the usefulness of 1-D and box estuarine models.  
 125 Historically, simple zero-dimension box models were proposed to estimate the replacement time of  
 126 the estuarine volume,  $V$ , which was then called the residence time (Swaney et al., 2011) and  
 127 turnover time (Sheldon and Alber 2006), and is now termed the renewal time. Because, under  
 128 many conditions, water renewal time would be similar to residence time at the inflow open  
 129 boundary (i.e. upstream estuarine location), the residence time results from calibrated numerical  
 130 models would be useful to verify the efficiency of simple models. Probably the two most well-  
 131 known models of this type are the tidal flushing box model, and the gravitational circulation model  
 132 (Ketchum, 1951; Dyer, 1973; Officer, 1976). The tidal flushing box model assumes that the estuary  
 133 is flushed at each tidal cycle by the tidal prism,  $V_p$ , i.e. the amount of water that exits the estuary at  
 134 ebb tide (Ketchum, 1951; Harleman, 1966; Officer, 1976; Dyer, 1973; Zimmerman, 1988; Luketina,  
 135 1998; Sanford, 1992; Solis and Powell, 1999; Sheldon and Alber, 2006; Figure 2b). Thus, a fraction,  
 136  $r$ , of the estuary water is renewed by ocean water at each tidal cycle,

$$137 \quad r = V_p / V. \quad (2)$$

138  
 139  
 140  
 141  
 142  
 143  
 144  
 145

140 The renewal time,  $T$ , was calculated as

$$141 \quad T = \tau/r, \quad (3)$$

143 where  $\tau$  is the tidal period (e.g. 0.5 d for solar semi-diurnal tide and 1 d for solar diurnal tide).

145 In an attempt to provide information on the along-channel variation of the renewal time, this  
 146 model was later improved by segmenting the estuary, with the assumption that the volume of water  
 147 that exits a segment at ebb tide completely replaces the water in the downstream segment (Wood,  
 148 1979). Such models do not account for the baroclinic circulation that is typical of partially mixed  
 149 estuaries (Figure 2c). The gravitational circulation model calculates the renewal time as the ratio  
 150 between the volume of fresh water and the inflow rate. The inflow rate is the sum of the oceanic  
 151 inflow of water plus the riverine inflow (Figure 2c; Dyer, 1973); tidal mixing is neglected, and the  
 152 oceanic inflow is calculated from the salt conservation equation by Knudsen (1900), which is

$$153 \quad Q_{OUT} = Q_{IN} + Q_R. \text{ Where } Q_{OUT} = Q_R \frac{S_{IN}}{S_0 - S_E} \text{ and } Q_{IN} = Q_R \frac{S_{OUT}}{S_0 - S_E}.$$

154 coastal waters, and the average salinity in the estuary, respectively. In contrast, the salinity balance  
 155 equation of Fischer (1979) states that the downstream residual transport of salt balances the  
 156 upstream dispersion (diffusion) of salt, i.e.  $Q\bar{S} = -KA \frac{\Delta S}{L}$ , where  $\Delta S = (S_0 - S_{UP})$ ,  $S_{UP}$  is the  
 157 salinity at the upstream estuarine segment (i.e. salinity at the upstream inflow open boundary), and  
 158  $\bar{S}$  is the mean salinity of the estuarine segment considered. If one assumes the length of the  
 159 estuarine segment to be the length of the entire estuary, therefore  $\bar{S} = S_E$ .

160 The fresh water renewal time calculated using the gravitational circulation model, which is  
 161 also called fresh water fraction model,  $T_F$  (Hansen and Rattray, 1965; Dyer, 1973; Wolanski, 2007)  
 162 is,

$$163 \quad T_F = \frac{V}{Q_R} F_R = \frac{V}{Q_R \left( \frac{S_0 - \bar{S}}{S_0 - \bar{S}} \right)} = \frac{V}{Q_R + Q_R \frac{\bar{S}}{S_0 - \bar{S}}}, \quad (4)$$

164 where  $T_F$  is the time taken to renew the portion of fresh water in the estuary,  $F_R = (S_0 - \bar{S})/S_0$  is  
 165 the fresh water fraction (Dyer, 1973), and  $Q_R$  is the river discharge (or total residual outflow). It is  
 166 important to note that the fresh water fraction model estimates the time to renew only the fresh  
 167 water portion, which is smaller than the volume of the whole domain. Therefore, the fresh water  
 168 fraction model would result in a shorter timescale than the residence time of particles released near  
 169

171 the estuary head. Previous simple box models neglected either baroclinic circulation or tidal mixing,  
 172 and the fresh water fraction model has always been criticized for lacking seawater inflow and  
 173 miscalculating transport timescales. Sheldon and Alber (2006) demonstrated that the fresh water  
 174 fraction model incorporates the effect of seawater inflow, which is due to the gravitational  
 175 circulation effect (Figure 2c). In general, many researchers dislike formulations that cast advective  
 176 fluxes of the gravitational circulation in terms of along-channel diffusivity or similar simplified  
 177 assumptions. These physical processes, i.e. dispersion, gravitational circulation etc., are governed  
 178 by different physics.

179 Similar to the fresh water fraction model, the well-known LOICZ model (Smith et al., 2005  
 180 and 2010; Crossland et al., 2005; Swaney et al., 2011) includes both advective and exchange flow  
 181 transport (Figure 2d). Sheldon and Alber (2006) demonstrated that the LOICZ model does not  
 182 consider the salt conservation equation proposed by Knudsen (1900), and this manuscript shows  
 183 that the LOICZ solution does not properly apply the salinity balance proposed by Fischer (1979).  
 184 Therefore, there are two proposed methods that might be considered by the LOICZ scientific  
 185 community.

186 For the LOICZ model, rainfall, groundwater and evaporation effects can also be added if  
 187 required, and are incorporated into the residual flow  $Q_R$ . For a vertically well-mixed estuary,  
 188 oceanic water enters the estuary at a rate,  $Q_D$  (the exchange flow), with a salinity that exceeds the  
 189 initial estuarine salinity,  $S_R$ , by the amount  $\Delta S$ .  $Q_D$  applied in the LOICZ model is different from the  
 190 well-known salinity balance equations by Knudsen (1900) and Fischer et al. (1979).

192 Figure 2– Preferred position.

194 The water renewal of an estuary depends upon the residual flow, e.g. river discharge  $Q_R$ , and  
 195 the exchange flow  $Q_D$ . The general formulation to estimate water renewal is:

$$197 \quad T = V / (Q_R + Q_D), \quad (5)$$

199 where  $Q_D$  is defined as the exchange flux between the estuary and the coastal areas. We note that  
 200 this term, i.e.  $Q_D$ , is called  $V_X$  in the LOICZ model (Gordon et al., 1996); however,  $Q_D$  seems more  
 201 appropriate to represent a term denoting a large dispersive contribution to water renewal. Sheldon  
 202 and Alber (2006) presented a good argument for applying a different method to estimate the  
 203 exchange flow,  $Q_D$ . They used the Knudsen relation.

204 In summary, there are two well-known competing definitions for the salt balance, and they  
 205 rely upon different physics. Knudsen relation assumes that steady salinity is balanced by residual  
 206 circulation and gravitational circulation (i.e. used in the fresh water fraction model), while in the

207 second definition (i.e. from Fischer formulation for the salinity balance), steady salinity is balanced  
 208 by residual circulation and dispersive processes. This manuscript presents results of water renewal  
 209 calculated using both methods, and compares them with results from numerical models.

210 Advection and dispersion of salt are balanced in a steady-state along an estuarine system,  
 211 giving the salt-balance equation by Fischer et al. (1979),

$$212 \quad Q_D = Q_R S_E / (S_O - S_{UP}). \quad (6)$$

213 In the Knudsen relation, the inputs of freshwater  $Q_R$  and seawater  $Q_D$  have salinities of 0  
 214 and  $S_O$ , respectively. In order to balance the water volume, an outflow of magnitude  $(Q_R + Q_D)$  must  
 215 exit the system, therefore the salt balance is:

$$216 \quad Q_R 0 + Q_D S_O = (Q_R + Q_D) S_E, \quad (7)$$

217 which results in the exchange flow,

$$218 \quad Q_D = Q_R S_E / (S_O - S_E). \quad (8)$$

219 The difference between equation (6) and (8) is due to the different assumptions regarding the  
 220 salinity balance, i.e. Knudsen relation and Fischer formulation. If one applies the salinity  
 221 conservation equation proposed by Knudsen relation, Eq. 5 reduces to Eq. 4, which is the fresh  
 222 water fraction model. Sheldon and Alber (2006) demonstrated that the LOICZ model does not apply  
 223 Knudsen relation for the salinity balance properly, because the salinity of the outflow would be  $S_E$ .  
 224 The LOICZ model considers the salinity balance in the outer half of the estuary, which would result  
 225 in a mass balance problem in the Knudsen relation.

226 The LOICZ method assumes neither Knudsen relation nor Fischer formulation for the  
 227 salinity balance. Here it is demonstrated that the LOICZ model applies a salinity balance slightly  
 228 different from the balance proposed by Fischer et al. (1979). For the LOICZ model the exchange  
 229 flow is,

$$230 \quad Q_D \Delta S = Q_R S_R, \quad (9)$$

231 the estuarine outflow salinity is,

$$232 \quad S_R = 0.5 (S_O + S_E) \quad (10)$$

and mean salinity in the outer estuary (see, Gordon, et al., 1996; Dupra et al., 2001; Swaney et al., 2011) is,

$$\Delta S = (S_O - S_E) \quad (11)$$

The total water inflow rate for the estuary is thus equal to  $Q_R + Q_D$  (see Eq. 5). The same water flux leaves the estuary; thus the renewal time  $T_{LOICZ}$  is,

$$T_{LOICZ} = V / (Q_R + Q_D) = V / (Q_R + Q_R S_R / (S_O - S_E)). \quad (12)$$

The LOICZ box model is simple and easy to use, and has been applied to about 200 estuaries worldwide (Swaney et al., 2011). However, box models only capture part of the physics, and they give no information on the spatial distribution of the renewal time in an estuary. To obtain this spatial information, it is necessary to use numerical models or to apply box models using a reasonable number of boxes along the estuary. However, a drawback of this approach is that it commonly requires the modeller to assume values of the turbulent dispersive coefficient, which is often unknown *a priori*.

In this paper, water renewal, residence time and exposure time were calculated using simple formulations, and then compared with the residence times calculated using numerical models for a few estuaries where data were available. Water renewal times calculated using a modified LOICZ model were compared with results from the original LOICZ model and the fresh water fraction model (Gordon et al., 1996; Yanagi, 2000; Smith et al., 2005; Newton and Icely, 2007; Breitburg et al., 2009a,b; Smith et al., 2010; Swaney et al., 2011). The LOICZ model and the modified LOICZ model share very similar assumptions, which include representing the estuarine concentration with a single value. Mean residence and exposure times were calculated using CART theory (Constituent-oriented Age and Residence time Theory). In addition to comparing timescale estimates of these methods, we also applied a new advection-dispersion diagram that quantifies the relative contribution of the advective and dispersive timescales to water renewal.

## 2. The LOICZ model

The LOICZ model includes both advective and an exchange flow transport. Dispersive residual transport is calculated as an exchange flow between the estuary and coastal waters. This exchange flow is calculated by assuming the salinity gradient in the seaward half of the estuary, i.e.



from the estuarine mouth ( $x = L_0$ ) to  $x = L/2$ , and considering geometric features of the estuary, including sewage, groundwater, rainfall, evaporation, freshwater runoff, as well as the salinity in the estuary and in coastal waters. The inflow water would be a result of processes such as fresh water discharge, rainfall, groundwater etc. However, to simplify the model we neglect all terms aside from river discharge. The LOICZ box model calculates the average residence time  $T_{LOICZ}$  using Eq.12. Using Eq. 10, Eq. 12 becomes,

$$T_{LOICZ} = \frac{V}{Q_R + Q_R \frac{0.5(S_E + S_O)}{S_O - S_E}} \quad (13)$$

exchange flow

The LOICZ model depends on accurate measurements of the salinity at the boundaries of the domain in order to properly estimate the exchange flow ( $Q_D$ ). The basic assumptions associating  $Q_D$  with the salinity balance may be invalid if the salinity gradient is too small (Swaney et al., 2011).

Eq. 13 may be written as,

$$T_{LOICZ} = \frac{VL}{Q_R L + \left( \frac{0.5(S_E + S_O)}{S_O - S_E} \right) Q_R L} \quad (14)$$

The salinity balance equation of Fischer et al. (1979) can be used in this case to estimate the amount of dispersion applied into the LOICZ model, and assuming the salinity at the mouth  $S(L)$  to be  $S(L) \approx S_O$ ,

$$Q_R L = AK^{LOICZ} \frac{(S_O - S_E)}{0.5(S_O + S_E)} \quad (15)$$

where  $K^{LOICZ}$  is the dispersive coefficient applied in the salinity balance in the LOICZ model, and  $A$  is the mean cross-sectional area. It will be further shown that the LOICZ model often results in a large dispersion coefficient because it accounts for a salinity balance near the estuarine mouth. From Eqs. 14 and 15,

$$T_{LOICZ} = \frac{VL}{Q_R L + AK^{LOICZ}} \quad (16)$$

In Eq. 14 the influence of the dispersive timescale on the water renewal is parameterized by  $\gamma_{LOICZ}$ , which is defined as,

$$\gamma_{LOICZ} = \left[ \frac{0.5(S_E + S_O)}{S_O - S_E} \right]. \quad (17)$$

Using the advective and dispersive timescales (Fischer et al. 1979), as previously defined, i.e.

$$T_1 = \frac{V}{Q_R}, \quad \text{and} \quad T_2^{LOICZ} = \frac{L^2}{K^{LOICZ}}. \quad (18)$$

Eqs. 16 and 18 become,

$$\frac{1}{T_{LOICZ}} = \frac{1}{T_1} + \frac{1}{T_2^{LOICZ}}. \quad (19)$$

To our knowledge, Eq. 19 has not previously been derived for the LOICZ model. This equation shows the importance of the advective and dispersive timescales for water renewal.

### 3. The modified LOICZ model

For the modified LOICZ model, we propose a modified exchange flow that takes into consideration the average dispersion, which is calculated using Fischer formulation (Fischer et al., 1979). Field data indicate that water renewal,  $T$ , decreases with increased dispersion (Uncles et al., 2002). The along-channel dispersive coefficient,  $K$ , can be estimated from salinity data along the whole estuarine segment,  $L$  (Fischer et al., 1979).  $K$  can be estimated from the salinity balance equation because all the other terms in that equation are measureable. Other methods exist that can be used to estimate  $K$  from salinity measurements, e.g. that of Hansen and Rattray (1965), which involves choosing values of  $K$  to fit the observed distribution of salinity with the analytical solution. Additional methods also exist to estimate the salinity balance and dispersion (Knudsen, 1900; MacCready, 1998; MacCready, 2004; MacCready, 2011; MacCready and Banas, 2011), which could be used to further quantify the exchange flux applied in our method.

The proposed method is also based upon Fick's first law, which relates the diffusive (dispersive) flux to the concentration under the assumption of steady state. The magnitude of the flux is proportional to the concentration gradient, from high to low. Consider a large number of particles to represent a concentration,  $C$ , which are initially deployed at the upstream location of the estuarine segment,  $L$ . Horizontal dispersion would result in a random walk in two dimensions, i.e.

341 along and across the estuary. However, the movement of particles across the estuary is limited by  
 342 solid boundaries, and thus dispersion of particles would be predominately along the estuary, and  
 343 the dispersive flux of particles would have two orientations (upstream and downstream). The mean  
 344 square displacement  $\sigma^2$  of particles (in one dimension) would be  $\sigma^2 = 2K_d\Delta t$ . Note that this  
 345 random displacement of particles (towards the upstream or downstream boundaries), depends upon  
 346 the small time step,  $\Delta t$ , and the dispersive coefficient,  $K_d$ . The coefficient  $K_d$  determines the  
 347 random walk within the time step  $\Delta t$ , this coefficient is not the characteristic dispersion  $K$  of the  
 348 whole estuarine segment  $L$ . In fact, it is a much smaller dispersion ( $K_d \ll K$ ). To avoid an  
 349 overestimate of the downstream movement due to dispersion, the mean square downstream  
 350 displacement is assumed to be  $\Delta d^2 = \alpha\sigma^2$ , where the reducing factor is considered to be  $\alpha = 1/2$ .  
 351 By assuming the displacement to be a constant under steady state conditions, the finite additivity  
 352 property of displacements satisfies,  $L = \sum_{j=0}^n \Delta d(\Delta t_j) = \Delta d_1 + \Delta d_2 + \dots + \Delta d_n = n\Delta d = n\sqrt{K_d\Delta t}$ . The  
 353 dispersive timescale in this case would be  $T_2 = n\Delta t$ , but also assumes the form  $T_2 = L^2/K$ , and thus  
 354 suggests that the dispersion coefficient  $K$  is the finite additivity of the dispersion in each of the  $n$   
 355 sub-segments of length  $\Delta d$ , i.e.  $K = nK_d$ . The mean dispersive flux across the area,  $A$ , is  
 356 calculated using the estuarine segment length and the dispersive timescale, i.e.  $Q_D = LA/T_2$ .  
 357 Additionally, it can be expressed in terms of  $K$  or  $K_d$ , i.e.  $Q_D = KA/L = nK_dA/L$ . From Fischer  
 358 formulation, the dispersion,  $K$ , can be estimated and applied into the dispersive flux relation,

$$359 \quad Q_D \sim KA/L \quad (20)$$

360 and as shown before,

$$361 \quad Q_D = Q_R S_E / (S_O - S_{UP}) \quad (21)$$

362 where  $S_{UP}$  is in the range  $0 \leq S_{UP} < S_E$ . Here  $S_{UP}$  is defined at the upstream end of the domain, and  
 363 depends upon the estuarine segment length,  $L$ . The length,  $L$ , can be shorter than, but cannot exceed  
 364 the length of the maximum salinity intrusion; otherwise it would lead to a misapplication of the  
 365 model. It can be seen when comparing Eqts. (9-10) and Eq (21) that the LOICZ method uses a  
 366 larger exchange flow,  $Q_D$ , than the modified LOICZ model. Thus we propose to use Eq. (21) as the  
 367 basis of the modified LOICZ model. This gives water renewal time,  $T_P$ , as,  
 368  
 369  
 370

$$T_P = \frac{VL}{(Q_R L + AK)} = \frac{VL}{Q_R L + \left(\frac{S_E}{S_O - S'}\right) Q_R L} = \frac{V P_E}{Q_R (1 + P_E)}, \quad (22)$$

where  $P_E = uL/K$  is the Peclet number, and

$$\gamma_P = S_E / (S_O - S'), \quad (23)$$

$\gamma_P$  determines the contribution to water renewal from the dispersive timescale, and is calculated differently from  $\gamma_{LOICZ}$  in the LOICZ model (Eq. 19). Using the dispersive and advective timescales, Equation 22 becomes,

$$\frac{1}{T_P} = \frac{1}{T_a} + \frac{1}{T_b}, \quad (24)$$

where  $T_a = V/Q_R$  and  $T_b = VP_E/Q_R$  are the advective and dispersive timescales, respectively, as calculated using the river flow  $Q$  ( $\text{m}^3 \text{s}^{-1}$ ). The contribution of advection to the total water renewal,  $\theta$  ( $0 \leq \theta \leq 1$ ), is given by

$$\theta = T_P/T_a = Q_R/(Q_R + Q_D) \quad (25)$$

#### 4. The difference between the water renewal time calculated from the LOICZ model and the modified LOICZ model

There are some differences between the original formula used in the LOICZ model and the newly modified LOICZ model. These are parameterized by  $\gamma = \gamma_{LOICZ} / \gamma_P$ , which is the ratio between the exchange flow in the LOICZ model (Officer, 1980; Swaney et al., 2011) and the suggested new method. Assuming a linear salinity gradient along an estuary, the salinity at the mouth is related to salinity in the estuary by  $S_O = 2S_E$ , salinity at the maximum salinity intrusion is  $S_{UP} = 0$ , and thus the ratio  $\gamma / \gamma_P = 3$ . Therefore, for a nearly linear salinity distribution along an estuary, the dispersive timescales for the LOICZ model are expected to be three times larger than for the modified LOICZ model. The salinity balance used to calculate the so called exchange flow in the LOICZ model is described in (Fig. 3).

402 Figure 3– Preferred position.

403  
1

404  
3

405  
5

406  
7

407  
9

408  
11

409  
13

410  
15

411  
17

412  
19

413  
21

414  
23

415  
25

416  
27

417  
29

418  
31

419  
33

420  
35

421  
37

422  
39

423  
41

424  
43

425  
45

426  
47

427  
49

428  
51

429  
53

430  
55

431  
57

432  
59

433  
61

434  
63

435  
65

436  
67

437  
69

438  
71

439  
73

440  
75

441  
77

442  
79

In the LOICZ model, the exchange flux is calculated slightly differently from that proposed by Fischer et al. (1979), see Figure 3. The exchange flow from the LOICZ model is calculated using the salinity balance from  $x = L/2$  to  $x = L$ . Our approach was to estimate exchange flux from measurements of salinity along the estuary, using Fischer formulation,  $Q_D \sim AK/L$  ( $\text{m}^3 \text{s}^{-1}$ ). This exchange flux,  $Q_D$ , therefore accounts for the dispersive process at steady state along the whole estuarine segment of length  $L$ .

To compare the contributions of the river discharge and the exchange flow to water renewal, we assume a linear variation of the along-channel salinity gradient,  $S_E = 0.5S_O$  and  $S_{UP} = 0$ . Thus equations 14 and 22 become respectively,

$$T_{LOICZ} = \frac{V}{Q_R + \underbrace{1.5Q_R}_{\text{exchange flow}}}, \quad (26)$$

and

$$T_P = \frac{V}{Q_R + \underbrace{0.5Q_R}_{\text{exchange flow}}}, \quad (27)$$

The exchange flow in the LOICZ formula (Eq. 26) is much larger than the exchange flow in the modified model (Eq. 27). Additionally, in the LOICZ model, the exchange flow contributes 50% more to water renewal than river discharge does. Conversely, Eq. 27 gives an exchange flow contribution that is half of the advective term.

## 5. The CART analytical model

The Constituent-oriented Age and Residence time Theory (CART, [www.climate.be/cart](http://www.climate.be/cart)) consists of general partial differential problems from which a number of timescales may be derived at any time and location so as to assess water and contaminant exchange rates. For instance, the age, residence time and exposure time have been estimated numerically in realistic semi-enclosed domains such the New York Bight (Zhang et al 2010) and the Scheldt Estuary (de Brauwere et al. 2011, de Brye et al. 2012). Analytical solutions obtained in an idealised setup may also be of use. Accordingly, a one-dimensional flow with constant hydrodynamical and geometrical features shall be considered, allowing the exact solutions to be constructed for the residence and exposure times of passive tracer particles, which may also be regarded as tagged water parcels. These idealised CART timescales will be compared with the water renewal timescales ensuing from the original and

434 modified LOICZ models, as well as those obtained from numerical models, for estuaries where such  
 435 estimates are available.

436 Consider an infinite pipe ( $-\infty < x < \infty$ ) with a constant section  $A$ , in which there is a steady-  
 437 state, one-dimensional current whose volumetric flow rate is denoted  $Q_R$ . The upstream and  
 438 downstream boundaries of the domain of interest, i.e. the idealised estuary, are located at  $x = L_1$  and  
 439  $x = L_0$ , respectively. The length of the estuary, its volume and the water flow are given by,

$$440 \quad L = L_0 - L_1, \quad V = AL \quad \text{and} \quad U = Q_R / A = LQ_R / V.$$

441 The general differential problem from which the residence time may be estimated was  
 442 established by Delhez et al. (2004). For the one-dimensional steady-state problem to be dealt with,  
 443 the residence time  $\varphi(x)$  satisfies the equation (Delhez and Deleersnijder 2006, Blaise et al. 2010)

$$445 \quad \frac{d}{dx} \left( AK \frac{d\varphi}{dx} + Q_R \varphi \right) = -A \quad (28)$$

446 under the boundary conditions,

$$449 \quad \varphi(L_1) = 0 = \varphi(L_0), \quad (29)$$

450 where the positive constant,  $K$ , denotes the along-flow diffusivity. It must be stressed that (28) is  
 451 not an advection-dispersion-reaction equation; if it were an equation of this type, the advective term  
 452 would be  $-Q_R \varphi$  rather than  $+Q_R \varphi$ . In fact, the equation for the residence time is the adjoint of the  
 453 relevant transport equation (Delhez et al. 2004, Delhez 2006). The physical meaning of the  
 454 residence time mathematically defined above is as follows: the average time taken for water or  
 455 passive tracer particles initially located in the interval  $[x, x + \delta x]$  to reach one of the open  
 456 boundaries of the domain for the first time (i.e.  $x = L_1$  or  $x = L_0$ ) tends to  $\varphi(x)$  as  $\delta x \rightarrow 0$ . As all of  
 457 the flow parameters are assumed to be constant, the solution to (28)-(29) is easily derived:

$$460 \quad \varphi(x) = \frac{V}{Q_R} \left( 1 - \frac{\xi}{L} \right) + \frac{V}{Q_R} \left( \frac{e^{-Pe} - e^{-Pe\xi/L}}{1 - e^{-Pe}} \right) \quad (30)$$

461 where  $\xi = x - L_1$  and the dimensionless parameter

$$464 \quad Pe = \frac{UL}{K} \quad (31)$$

may be regarded as the Peclet number of the flow under consideration. The latter is the ratio of the dispersive timescale  $T_2 = L^2 / K$  to the advective timescale  $T_1 = L / U$ .

After crossing an open boundary, a particle may re-enter the domain at a later stage (e.g. Spivakovskaya, et al. 2007, Delhez and Deleersnijder, 2010). This is not accounted for by the residence time, which is related to the time taken to hit an open boundary for the first time. To take into account the possibility of particles returning into the domain, the concept of exposure time was introduced (Monsen, et al. 2002, Delhez, et al. 2004, Delhez, 2006). This timescale, which is denoted hereinafter  $\Theta(x)$ , is defined in the domain of interest and its surrounding environment, and is the solution of the equation (Delhez et al. 2004, Delhez and Deleersnijder 2006)

$$\frac{d}{dx} \left( AK \frac{d\varphi}{dx} + Q_R \varphi \right) = \begin{cases} -A\varphi, & L_1 < x < L_0 \\ 0, & -\infty < x \text{ or } L_0 < x < \infty \end{cases} \quad (32)$$

The solution thereof is

$$-\infty < x \leq L_1 \quad : \quad \Theta(x) = \frac{V}{Q_R} \quad (33a)$$

$$L_1 \leq x \leq L_0 \quad : \quad \Theta(x) = \frac{V}{Q_R} \left( 1 - \frac{\xi}{L} \right) + \frac{V}{Q_R} \left( \frac{1 - e^{-Pe\xi/L}}{Pe} \right) \quad (33b)$$

$$L_0 \leq x < \infty \quad : \quad \Theta(x) = \frac{V}{Q_R} \left( \frac{e^{Pe} - 1}{Pe} e^{-Pe\xi/L} \right) \quad (33c)$$

The physical meaning of the exposure time is as follows: the average time that water or passive tracer particles initially located in the interval  $[x, x + \delta x]$  will spend in the domain of interest ( $L_1 < x < L_0$ ) tends to  $\Theta(x)$  as  $\delta x \rightarrow 0$ .

Figure 4– Preferred position.

Figure 4 displays the profile of the residence and exposure time for various values of the Peclet number. For high values of the Peclet number, the boundary layer for the concentration,  $C$ , is

developed in the vicinity of the upstream boundary of the domain (Delhez and Deleersnijder 2006, Blaise et al. 2010). The cause thereof may be summarized as follows: the greater the relative importance of advection, the less likely it is that dispersion will cause a water particle to hit the upstream boundary of the domain ( $x = L_0$ ). In accordance with elementary physical intuition, the exposure time is larger than the residence time ( $L_1 \leq \xi/L \leq L_0$ ).

It is appropriate to assess the propensity of water particles to return into the domain after hitting one of its open boundaries for the first time. For this, we used the approach of Arega et al. (2008) and de Brauwere et al. (2011), which leads to the dimensionless number termed “return coefficient”. This is defined as

$$r(x) = \frac{\Theta(x) - \phi(x)}{\Theta(x)}. \quad (34)$$

The values of this coefficient (Eq. 34) fall within the interval [0,1]: the larger this value, the more likely it is that water particles will re-enter the domain after hitting one of its open boundaries for the first time. Accordingly, particles that never return into the domain are characterized by a zero return coefficient, while particles returning most often are associated with a value of  $r$  that is close to unity (Fig 5).

Figure 5– Preferred position.

From Equations 30 and 33, the domain-averaged residence time and exposure times obey the relations,

$$\bar{\phi} = \frac{V}{Q_R} \left( \frac{1}{2} \right) + \frac{V}{Q_R} \left( \frac{1}{e^{Pe} - 1} - \frac{1}{Pe} \right) \quad (35)$$

and

$$\bar{\Theta} = \frac{V}{Q_R} \left( \frac{1}{2} \right) + \frac{V}{Q_R} \left( \frac{1}{Pe} - \frac{1 - e^{-Pe}}{Pe^2} \right). \quad (36)$$

These global timescales can also be regarded as the residence and exposure times of an arbitrarily large number of water particles that initially are uniformly distributed over the domain. The first terms in Equations 35 and 36 are the mean advective timescales, while the second terms cause the decrease and increase to the final residence time and exposure time.

## 6. Results and discussions



528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

The CART model, the LOICZ model, the fresh water fraction model, and a newly modified LOICZ model were applied to the estuaries (Appendix 1 to 7), namely the Curimataú, Caravelas and Peruípe estuaries in Brazil (Miranda et al., 2005 and 2006; Andutta et al., 2006), the York River and Hudson estuaries in the USA (Sandifer, 1973; Haas, 1977; Shen and Haas, 2004; Warner et al., 2005 and 2010), the Conwy and Mersey estuaries in the UK (Bowden and Gilligan, 1971; Turrell et al., 1996; Wu et al., 2005), and the Scheldt in France-Belgium-Netherlands (de Brauwere et al., 2011; de Brye et al., 2012). Water renewal times were estimated using the LOICZ model, the modified LOICZ model and the fresh water fraction model, while the CART theory was applied to estimate the mean residence and exposure times.

From Table 1 it can be seen that water renewal times calculated by the LOICZ method and the fresh water fraction model are always shorter than those calculated from the modified LOICZ model,  $T_P$ . Results from the fresh water fraction model,  $T_{FRAC}$ , were often similar to results from the original LOICZ model,  $T_{LOICZ}$ . Renewal times from the modified LOICZ model usually compare well with the exposure times calculated by the CART analytical model (see Table 2). Timescales from the LOICZ model are shorter than those from the modified LOICZ model, due to the larger dispersive coefficient applied in the original model (explained in section 4). Additionally, timescales from the fresh water fraction model are shorter than those from the proposed LOICZ model, because of the larger gravitational circulation contribution to water renewal.

For the Hudson River the water renewal at neap tides ranges between 6.3 and 7.8 days for  $T_P$ , and ranges between 4.3 and 5.3 days using the LOICZ model, while water renewal from the fresh water fraction model ranges between 5.9 and 7.4 days for neap tides, and between 6.7 and 8.5 days for spring tides.

The difference in  $T$  between the original and the proposed LOICZ models comes from the dispersive timescale component, which in this case ranges from 31.4 – 43.0 days using the modified LOICZ model, and from 9.4 – 12.1 days using the original LOICZ model. This difference in the dispersive timescales depends upon on the coefficient,  $K$ , which was estimated to range between 1880 to 2404  $\text{m s}^{-2}$  for the original LOICZ model, and 545 to 828  $\text{m s}^{-2}$  for the modified LOICZ. Water renewal results from the modified LOICZ model compare favourably with exposure time estimates from the CART formula, however they are slightly underestimated compared with the numerical results for the Hudson Estuary (Warner et al., 2010), which give a mean residence time for neap tides of ~ 8-10 days and for spring tides ~ 8-9 days.

The water renewal time in our proposed method is a combination of advective and dispersive timescales, and the difference between  $T_P$  and  $T_{LOICZ}$  depends on the factor  $\gamma$ . This is especially the case when the advective timescale is relatively large compared to the dispersive

563 timescale, and thus the dispersive timescales dominate the renewal time. For the Mersey estuary,  
 564 the advective timescale is very large compared with the dispersive timescale, and the water renewal  
 565 calculated by the original LOICZ model is considerably less than that of the modified LOICZ  
 566 model. Conversely, for the Hudson estuary during neap tides, the fresh water discharge is important,  
 567 and thus the advective timescale is smaller than the dispersive timescale,  $T_{LOICZ}$  approaches  $T_P$ .

568 The residence times calculated by the CART formula were mostly smaller than the results  
 569 for water renewal calculated using the LOICZ model, the fresh water fraction model, and the  
 570 modified LOICZ model. In contrast, CART exposure times were always larger than the residence  
 571 times from numerical models; this is because of how these timescales are defined. CART exposure  
 572 times are, however, more comparable to the water renewal times in most cases. Larger exposure  
 573 times result from the small values of the Peclet number (Table 2). Therefore, high values of the  
 574 Peclet number indicate that although particles may reach the open boundary quickly, they may  
 575 return to the system for long periods (e.g. Curimataú, Caravelas, Peruípe and Mersey estuaries).

576 The results of these different approaches (i.e.  $T_P$ ,  $T_{LOICZ}$ ,  $T_{FRAC}$ , and CART formula) were  
 577 compared with available numerical results (Table 2) for the Hudson Estuary (Warner et al., 2010),  
 578 the Caravelas and Peruípe estuaries (Andutta, 2011), the Mersey Estuary (Yuan et al., 2007), the  
 579 Scheldt Estuary (de Brauwere et al., 2011; de Brye et al., 2012), and the York River Estuary (Shen  
 580 and Haas, 2004). The models had previously been calibrated and validated with salinity  
 581 measurements, and hence, may be assumed to accurately reproduce advective and dispersive  
 582 transport phenomena for the domains in which they were used.

583 The numerical results from (Warner et al., 2010) showed that for the Hudson Estuary  
 584 (within 45 km of the mouth) the mean residence time for neap tides was ~8-10 days and for spring  
 585 tides was ~8-9 days. The results calculated for  $T_P$  (see Table 1) are slightly lower than the numerical  
 586 model results for neap tides, and relatively close for spring tides (i.e 6.3-7.8 days for neap tides and  
 587 7.9-9.8 for spring tides). As mentioned previously, for the Hudson Estuary the LOICZ model and  
 588 the fresh water fraction model yield a water renewal (~5 days) *ca.* 40% lower than the residence  
 589 time from numerical models. For the Caravelas and Peruípe estuaries in Brazil, the residence time  
 590 calculated from numerical models varied in the ranges of 4.2-10.3 and 1.5-2.5 days, respectively.  
 591 For the Mersey and Scheldt estuaries, the residence time varied between 0.7-4 and 5-70 days,  
 592 respectively. For the York River Estuary, the residence time varied between 11.3-33.3 and 18.1-  
 593 59.3 days for high and mean flow, respectively. Except for the Mersey Estuary, the results from  
 594 numerical models agreed well with the water renewal estimates from the modified LOICZ model  
 595 (i.e. the model using Fischer formulation for the salinity balance), and were slightly higher than the  
 596 water renewal estimates from the original LOICZ model, the fresh water fraction model, and the  
 597 residence time estimates calculated by the CART formula.

598

599

600 Table 1– Preferred position.

601 Table 2– Preferred position.

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

The modified LOICZ model provided the best fit with results of average maximum residence time from numerical models (Fig. 6a), with  $r^2 \sim 0.7$ . The slope was calculated to be 0.52, indicating that the modified LOICZ model yields water renewal slightly larger than residence times from numerical models at upstream locations. Although the compared timescales have slightly different definitions, this method offers a simple way to easily estimate the residence time for which minimal oceanographic data are available (i.e. geometry, river discharge and salinity measurements).

Linear correlation of water renewal times from both the original LOICZ model and the fresh water fraction model with the residence times from numerical models resulted in  $r^2 < 0.30$ . Linear correlation between residence time from numerical models and the time of exposure from CART formula resulted in  $r^2$  smaller than 0.20.

The dispersive contribution to water renewal (i.e.  $1 - \theta$ ) showed a strong exponential correlation against results of the return coefficient (Fig. 6b), with  $r^2 = 0.85$ , indicating that estuaries dominated by dispersion may have long periods of return for water particles.

Figure 6– Preferred position.

The relative contribution to water renewal for a particular estuary, from the advective and dispersive processes, can be visualized in the Advection-Dispersion Diagram (Figure 7), which is generated using equations 24 and 25. The parameter  $\theta$ , represented by straight lines, indicates the relative advective contribution to water renewal varying in the range  $0 \leq \theta \leq 1$ . The diagonal line  $\theta = 0.5$ , separates the areas where transport is dominated by dispersion (diagram lower zone,  $\theta < 0.5$ ) and advection (diagram upper zone,  $\theta > 0.5$ ). Although the results in figure 7 were obtained from table 1, this diagram synthesises the results into a single image. This allows an easy graphical comparison to be made between different estuaries, and between different conditions for a particular estuary. It can be seen that for the Curimataú Estuary, dispersion dominated over advection at spring tides, while advection dominated over dispersion at neap tides. The absolute, but not the relative, estimates of these timescales varied somewhat depending on which of the two methods for estimating  $K$  was used (e.g. Hansen-Rattray or Fischer formulations). For the Hudson Estuary, during both neap and spring tides, the water renewal was predominantly caused by advection, 0.72

633  $< \theta < 0.82$ . In the Conwy Estuary there was a slight dominance of advection during neap tides  $\theta \sim$   
 634 0.63, and dispersion during spring tides  $\theta \sim 0.45$ . For the Mersey, York, Caravelas and Peruípe  
 635 estuaries, dispersion dominated over advection (*ca*  $\theta < 0.50$ ), while for the Scheldt Estuary,  
 636 advection and dispersion contributed nearly equally to water renewal  $\theta \sim 0.50$ .

637 In the Conwy Estuary, where the river inflow was nearly constant, the relative contribution  
 638 of dispersion increased from neap to spring tides. This occurred because dispersion increases with  
 639 increasing values of the ratio of the tidal oscillation ( $R$ ) over the mean depth ( $h$ ), i.e.  $R/h$  (Uncles et  
 640 al., 2002). This effect might be more apparent if the advective timescale, due to river discharge, is  
 641 large, i.e. during low river discharge rates. Estuaries with high ratio values  $R/h$  may have a greater  
 642 dispersive contribution, e.g. the Curimataú Estuary ( $R \sim 2.5$  m and  $h \sim 6$  m), the Caravelas ( $h \sim 6.5$   
 643 m) and Peruípe ( $h \sim 7.5$  m) estuaries with the maximum tidal range ( $R$ ) of  $\sim 2.5$  meters, and the  
 644 Mersey estuary ( $h \sim 16$  m) forced by a tidal range ( $R$ ) of up to  $\sim 10.5$  meters. Therefore, a larger  
 645 dispersive contribution may result in a larger time of exposure (see Table 2), and thus a return  
 646 coefficient close to unity. Field data suggest that water renewal,  $T$ , decreases with increasing tidal  
 647 range, and increases with increasing estuary length (Uncles et al., 2002).

648

649 Figure 7– Preferred position.

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

## 7. Summary and Conclusions

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

653 We have developed a new method to quantify the relative contribution of advection and  
 654 dispersion to water renewal in estuaries, using simple measurements of the river flow  $Q_r$  ( $\text{m}^3 \text{s}^{-1}$ ),  
 655 the estuary volume  $V$  ( $\text{m}^3$ ) and length  $L$  (m), estimates of the mean salinity  $S_E$ , and the salinity at the  
 656 inflow  $S'$  (at,  $x = L_I$ ) and outflow  $S_O$  (at,  $x = L_O$ ) boundaries. Such data are available for most  
 657 estuaries worldwide. From this method, estuaries may be categorized using the Advection-  
 658 Dispersion diagram (Figure 7); they can be divided into those which are dominated by dispersion  
 659 (e.g. Mersey Estuary), those dominated by advection (e.g. Hudson Estuary), and those for which  
 660 dispersion and advection are of similar magnitude (e.g. Conwy Estuary). The model is applicable to  
 661 estuaries that have simple geometries and that lack significant baroclinic circulation.

662 Sheldon and Alber (2006) noted that the application of  $S_R$  was inconsistent with the  
 663 compartmentalization for the LOICZ model, of which the salinity balance was inconsistent with the  
 664 application of both Knudsen relation and Fischer formulation. Using  $S_R$  resulted in a higher salinity  
 665 exportation by  $Q_D$  (also called  $V_X$  in LOICZ model), which resulted in a short water renewal time.  
 666 We showed that that the Land Ocean Interaction Costal Zone model uses similar equation as from  
 667 the Fresh Water Fraction model, and thus often resulting in shortened transport timescales.

668           The proposed modified LOICZ method can be used to study the impact of spring-neap tidal  
669 fluctuations and seasonal variations of river flow and tides on water renewal. For example, in the  
670 Curimataú, high river flows occur mostly in summer during the wet season, whereas low river flows  
671 occur mainly in winter. For such systems, the river discharge may remain nearly constant for  
672 several weeks, and the relative contribution of dispersion and advection to the estuarine residence  
673 time would change along the spring-neap tidal cycle. In contrast, results of the Conwy estuary  
674 showed a nearly equal contribution of advection and dispersion at neap and spring tides. In the  
675 Hudson estuary, the residence time was dominated by advection, while dispersion dominated in the  
676 Caravelas, Peruípe, Mersey and Scheldt estuaries. .

677           Timescales calculated using the LOICZ model were always smaller than those from the  
678 modified LOICZ model and from hydrodynamic models. The reason being the method used to  
679 calculate the exchange flow in the LOICZ model; based upon Fischer formulation the exchange  
680 flow is calculated based upon the salt balance in the first half of the estuary, near the mouth, in  
681 which most dispersion is expected to occur. We know, however, that this large dispersion  
682 considered in the exchange flow would result in an underestimation of the water renewal time. The  
683 difference in the exchange flow between the LOICZ model and our proposed method depends upon  
684 the factor  $\gamma$ , which determines the dispersion contribution to the water renewal time. By using the  
685 equation of salt balance, proposed by Fischer et al. (1979), we have developed a new exchange flow  
686 that may be used in the LOICZ model, and which reduces the dispersion coefficient applied to the  
687 estuary. In addition, it has been demonstrated that the LOICZ model uses neither Knudsen relation  
688 nor Fischer formulation for the salinity balance.

689           Results from CART theory have shown that the exposure times were comparable to the  
690 residence times estimated using numerical models, while the mean residence times calculated using  
691 the CART formula were usually lower than numerical model estimates. In most cases, the water  
692 renewal times estimated using the newly proposed modified LOICZ model provided the best fit  
693 with the mean maximum residence time from hydrodynamic models, with  $r^2 = 0.70$ . The return  
694 coefficient calculated from CART theory showed good correlation with the dispersion contribution,  
695 with  $r^2 = 0.66$ . Our study adds value to the LOICZ formulation as we have made improvements to  
696 estimate the water renewal timescales.

## 697 698 **Acknowledgements**

699  
700           This study was supported by the IPRS fellowship to FPA. The authors greatly acknowledge  
701 kind assistance of Dr. L. de Miranda, Dr. E. Wolanski and Dr. L. Lucas. Eric Deleersnijder is an  
702 honorary research associate with the Belgian Fund for Scientific Research (FNRS) and his

703 contribution to the present study was achieved in the framework of the Interuniversity Attraction  
 704 Pole TIMOTHY, which was funded by BELSPO (www.belspo.be) under contract IAP6.13.

705  
 706

## 706 **References**

707  
 708

708 Andutta, F.P., Miranda, L.B., Castro, B.M., Fontes, R.F.C., 2006. Numerical Simulation of the  
 709 Hydrodynamic in the Curimataú Estuary, RN Brazil. *Oceanography and Global Changes*,  
 710 SP-Brazil, 545–558.

711 Andutta, F.P., 2011. The estuarine system of the Caravelas and Peruípe rivers (Bahia):  
 712 Observations, simulations, residence time, and advective and diffusive processes, PhD  
 713 Thesis. IO-USP, Oceanography Institute of the University of São Paulo, 121p.

714 Andutta, F.P., Kingsford, M.J., Wolanski, E., 2012. ‘Sticky water’ enables the retention of larvae in  
 715 a reef mosaic. *Estuarine, Coastal and Shelf Science* 101, 54–63.

716 Andutta, F.P., Ridd, P.V., Wolanski, E., 2013. Age and the flushing time in the Great Barrier Reef  
 717 coastal waters. *Continental Shelf Research* 53, 11–19.

718 Arega, F., Armstrong, S., Badr, A.W., 2008. Modeling of residence time in the East Scott Creek  
 719 Estuary, South Carolina, USA. *Journal of Hydro-environment Research* 2, 99–108.

720 Blaise, S., de Brye, B., de Brauwere, A., Deleersnijder, E., Delhez, E.J.M., Comblen, R., 2010.  
 721 Capturing the residence time boundary layer - Application to the Scheldt Estuary, *Ocean  
 722 Dynamics* 60, 535–554.

723 Bolin, B., H. Rodhe, 1973. A note on the concepts of age distribution and transit time in natural  
 724 reservoirs, *Tellus* 25, 58–62.

725 Bowden, K.F. and Gilligan, R.M., 1971. Characteristic features of estuarine circulation as  
 726 represented in the Mersey Estuary. *Limnology and Oceanography* 16(3), 490–502.

727 Breitburg, D.L., Hondorp, D.W., Davias, L.A., Diaz, R.J., 2009a. Hypoxia, nitrogen, and fisheries:  
 728 integrating effects across local and global landscapes. *Annual Review of Marine Science* 1,  
 729 329–349.

730 Breitburg, D.L., Davias, L.A., Limburg, K.E., Swaney, D.P., 2009b. Linking Nutrients, Hypoxia,  
 731 Fisheries, and Fishes: Interim Report from a Workshop supported by LOICZ. LOICZ Inprint  
 732 2009/2,13–14.

733 Buffoni, G., Falco, P., Griffa A., Zambianchi, E., 1997. Dispersion processes and residence times in  
 734 a semi-enclosed basin with recirculating gyres: An application to the Tyrrhenian Sea,  
 735 *Journal of Geophysical Research* 102(C8), 18699–18713, doi:10.1029/96JC03862.

736 Crossland, C.J., Kremer, H.H., Lindeboom, H.J., Marshall Crossland, J.I., Le Tissier, M.D.A.,  
 737 2005. *Coastal Fluxes in the Anthropocene*. Springer-Verlag, Berlin, 231 pp.

738  
 739  
 740  
 741  
 742  
 743  
 744  
 745

- 738 Deleersnijder, R., Beckers, J.M., Delhez, E.J.M., 2006. On the behaviour of the residence time at  
739 the bottom of the mixed layer. *Environmental Fluid Mechanics* 6, 541–547.
- 740 Delhez, E.J.M., Heemink, A.W., Deleersnijder, E., 2004. Residence time in a semi-enclosed domain  
741 from the solution of an adjoint problem, *Estuarine, Coastal and Shelf Science* 61, 691–702.
- 742 Delhez, E.J.M., 2006. Transient residence and exposure times. *Ocean Science* 2, 1–9.
- 743 Delhez, E.J.M. and Deleersnijder, E., 2006. The boundary layer of the residence time field, *Ocean  
744 Dynamics* 56, 139–150.
- 745 Delhez, E.J.M., Deleersnijder, E., 2010. Residence time and exposure time of sinking  
746 phytoplankton in the euphotic layer, *Journal of Theoretical Biology* 262, 505–516.
- 747 de Brauwere, A., de Brye, B., Blaise, S., Deleersnijder, E., 2011. Residence time, exposure time and  
748 connectivity in the Scheldt Estuary. *Journal of Marine Systems* 84, 85–95.
- 749 de Brye, B., de Brauwere, A., Gourgue, O., Delhez, E.J.M., Deleersnijder, E., 2012. Water renewal  
750 timescales in the Scheldt Estuary. *Journal of Marine Systems* 94, 74–86.
- 751 Dyer, K.R., 1973. *Estuaries: a Physical Introduction*. Wiley, London.
- 752 Dupra, V., Smith, S.V., Marshall Crossland, J.I. and Crossland, C.J., 2001. Estuarine systems of  
753 sub-Saharan Africa: carbon, nitrogen and phosphorus fluxes. LOICZ Reports & Studies No.  
754 18, i + 83 pages, LOICZ, Texel, The Netherlands.
- 755 Falco, P., Griffa, A., Poulain, P.-M., Zambianchi, E., 2000. Transport properties in the Adriatic Sea  
756 as deduced from drifter data, *Journal of Physical Oceanography* 30, 2055–2071.
- 757 Fischer, H.B., 1969. Cross-sectional time scales and dispersion in estuaries. *Proc. Congr. Int. Assoc.  
758 Hydraul. Res.*, 13<sup>th</sup> 3, 173–80.
- 759 Fischer, H.B., 1974. Numerical modelling of dispersion in estuaries. *Int. Symp. On Discharge of  
760 Sewage from Sea Outfalls*, London. Pap. 37, 1–8.
- 761 Fischer, H.B., 1976. Mixing and dispersion in estuaries. *Annual Rev. Fluid Mechanics* 8, 107–133.
- 762 Fischer, H.B., List, E.Y., Koh, R.C.Y., Imberger, J., Brooks, N.H., 1979. *Mixing in Inland and  
763 Coastal Waters*. Academic Press, New York, 483 pp.
- 764 Gordon, Jr.D.C., Boudreau, P.R., Mann, K.H., Ong, J.-E., Silvert, W.L., Smith, S.V. Wattayakorn,  
765 G., Wulff, F., Yanagi, T., 1996.. LOICZ biogeochemical modelling guidelines. *Land-Ocean  
766 Interactions in the Coastal Zone*, LOICZ Reports and Studies 5. Texel, The Netherlands.
- 767 Haas, L.W., 1977. The effect of the spring-neap tidal cycle on the vertical salinity structure of the  
768 James, York and Rappahannock Rivers, Virginia, U.S.A. *Estuarine and Coastal Marine  
769 Science* 5, 485–496.
- 770 Hansen, D.V. and Rattray, Jr.M., 1965. Gravitational circulation in straits and estuaries. *Journal of  
771 Marine Research* 23(1), 102–122.
- 772 Harleman, D.R.F., 1966. Pollution in estuaries. In *Estuary and Coastline Hydrodynamics*,

- 773 *Engineering Society Monographs*, Chap. 14. McGraw-Hill, New York.
- 774 Ketchum, B.H., 1951. The exchanges of fresh and salt water in tidal estuaries. *Journal of Marine*  
 775 *Research* 10, 18–38.
- 776 Knudsen, M. 1900. Ein Hydrographische Lehrsatz, *Annalen der Hydrographie und Marinen*  
 777 *Meteorologie* 28, 316–320.
- 778 Lowe, R. J., Falter, J. L., Monismith, S. G., Atkinson, M. J., 2009. A numerical study of circulation  
 779 in a coastal reef lagoon system, *Journal of Geophysical Research* 114(C06022), 1–18.
- 780 Lucas, L.V., Thompson, J.K., Brown, L.R., 2009. Why are diverse relationships observed between  
 781 phytoplankton biomass and transport time? *Limnology and Oceanography* 54, 381–390.
- 782 Luketina, D., 1998. Simple tidal prism models revisited. *Estuarine Coastal and Shelf Science* 46,  
 783 77–84.
- 784 MacCready, P., Banas, N.S., 2011. Residual Circulation, Mixing, and Dispersion. In: Wolanski E  
 785 and McLusky DS (eds.) *Treatise on Estuarine and Coastal Science*, Vol 2, pp. 75–89.  
 786 Waltham: Academic Press.
- 787 MacCready, P., 2011. Calculating Estuarine Exchange Flow Using Isohaline Coordinates, *Journal*  
 788 *of Physical Oceanography* 41, 1116–1124.
- 789 MacCready, P., 2004. Toward a Unified Theory of Tidally-Averaged Estuarine Salinity Structure,  
 790 *Estuaries* 27(4), 561–570.
- 791 MacCready, P., 1998. Estuarine Adjustment to Changes in River Flow and Tidal Mixing 29, 708–  
 792 726.
- 793 McLusky, D.S., Elliott, M., 2004. *The Estuarine Ecosystem: Ecology, Threats and Management*.  
 794 OUP, Oxford, 214 pp.
- 795 Miranda, L.B., Bérghamo, A.L., Castro, B.M., 2005. Interactions of river discharge and tidal  
 796 modulation in a tropical estuary, NE, Brazil. *Ocean Dynamics* 55, 430–440.
- 797 Miranda, L.B., Bérghamo, A.L., Silva, C.A.R., 2006. Dynamics of a Tropical Estuary: Curimataú  
 798 River, NE Brazil. *Journal of Coastal Research* SI 39, 697–701.
- 799 Monsen, N.E., Cloern, J.E., Lucas, L.V., Monismith, S.G., 2002. A Comment on the Use of  
 800 Flushing Time, Residence Time, and Age as Transport Timescales. *Limnology and*  
 801 *Oceanography* 47, 154–1553.
- 802 Newton, A., Icely, J., 2007. Land Ocean Interactions in the Coastal Zone, LOICZ: Lessons from  
 803 Banda Aceh, Atlantis, and Canute. *Estuarine, Coastal and Shelf Science* 77, 181–184.
- 804 Officer, C.B., 1976. *Physical Oceanography of Estuaries (and Associated Coastal Waters)*. Wiley,  
 805 London.
- 806 Officer, C.B., 1980. Box models revisited, pp. 65–114. In P. Hamilton and K. B. Macdonald (eds.),  
 807 *Estuarine and Wetland Processes with Emphasis on Modeling*. Plenum Press, New York.



- 808 Poulain, P.-M. and Hariri, S., 2013. Transit and residence times in the surface Adriatic Sea as  
809 derived from drifter data and Lagrangian numerical simulations, *Ocean Science Discussions*  
810 10, 197–217, doi:10.5194/osd-10-197-2013.
- 811 Prandle, D., 2009. *Estuaries: Dynamics, Mixing, Sedimentation and Morphology*. Cambridge  
812 University Press, Cambridge, 236 pp.
- 813 Sandifer, P.A., 1973. Distribution and abundance of Decapod Crustacean larvae in the York.  
814 Chesapeake Science 14(4), 235–257.
- 815 Sanford L.P., Boicourt, W.C., Rives, S.R., 1992. Model for estimating tidal flushing of small  
816 embayments. *Journal of Waterway, Port, Coastal, and Ocean Engineering* 118, 635–654.
- 817 Schettini, C.A.F., Miranda, L.B., 2010. Circulation and suspended particulate matter transported in  
818 a tidally dominated estuary: Caravelas Estuary, Bahia, Brazil. *Brazilian Journal of*  
819 *Oceanography* 58(10), 1–11.
- 820 Sheldon, J.E., Alber M., 2006. The Calculation of Estuarine Turnover Times Using Freshwater  
821 Fraction and Tidal Prism Models: A Critical Evaluation. *Estuaries and Coasts* 29, 133–146.
- 822 Shen, J., Haas, L., 2004. Calculating and the residence time in the York River using three-  
823 dimensional model experiments. *Estuarine, Coastal and Shelf Sciences* 61, 449–461.
- 824 Smith, S.V., Buddemeier, R.W., Wulff, F., Swaney, D.P., 2005. C, N, P Fluxes in the Coastal Zone.  
825 Chapter 3 in: Crossland, C.J.; Kremer, H.H.; Lindeboom, H.J.; Marshall Crossland, J.I.; Le  
826 Tissier, M.D.A. (eds). *Coastal Fluxes in the Anthropocene. The Land-Ocean Interactions in*  
827 *the Coastal Zone Project of the International Geosphere-Biosphere Programme*. Springer-  
828 Verlag, Berlin Heidelberg.
- 829 Smith, S.V., D.P. Swaney, D.P., Talaue-McManus, L. 2010. Carbon–Nitrogen–Phosphorus Fluxes in  
830 the Coastal Zone: The LOICZ Approach to Global Assessment. Chapter 14 (pp 575-586) in:  
831 *Carbon and Nutrient Fluxes in Continental Margins A Global Synthesis Series: Global Change -*  
832 *The IGBP Series*. Liu, K.-K.; Atkinson, L.; Quiñones, R.; Talaue-McManus, L. (Eds.) 2010,  
833 XXVIII, 744 pp.
- 834 Solis, R.S. Powell, G.L., 1999. Hydrography, mixing characteristics, and residence times of Gulf of  
835 Mexico estuaries, p. 29–61. In T. S. Bianchi, J. R. Pennock, and R. R. Twilley (eds.),  
836 *Biogeochemistry of Gulf of Mexico Estuaries*. John Wiley and Sons, New York.
- 837 Spivakovskaya, D., Heemink, A.W., Deleersnijder, E., 2007. The backward Ito method for the  
838 Lagrangian simulation of transport processes with large space variations of the diffusivity,  
839 *Ocean Science* 3, 525–535.
- 840 Swaney, D.P., Smith S.V., Wulff, F., 2011. The LOICZ Biogeochemical Modeling Protocol and its  
841 Application to Estuarine Ecosystems. In: Wolanski E and McLusky DS (eds.) *Treatise on*  
842 *Estuarine and Coastal Science* 9, 135–159. Waltham: Academic Press.

- 843 Takeoka, H., 1984. Fundamental concepts of exchange and transport timescales in a coastal sea,  
844 Continental Shelf Research 3, 311-326.
- 845 Turrell, W.R., Brown, J., Simpson, J.H., 1996. Salt Intrusion and Secondary Flow in a Shallow,  
846 Well-mixed Estuary. Estuarine, Coastal and Shelf Science 42, 153–169.
- 847 Uncles, R.J., Stephens, J.A., Smith, R.E., 2002. The dependence of estuarine turbidity on tidal  
848 intrusion length, tidal range and residence time. Continental Shelf Research 22, 1835–1856.
- 849 Valle-Levinson, A., 2010. Contemporary Issues in Estuarine Physics. Cambridge University Press,  
850 United Kingdom, 315 pp.
- 851 Yanagi, T., 2000. Simple method for estimating  $V_x$  from mixing equations in a 1-dimensional, steady-  
852 state system for LOICZ biogeochemical modelling. Pages 108-110 in Dupra, V., Smith, S.V.,  
853 Marshall Crossland, J.I. and Crossland, C.J. 2000. Estuarine systems of the East Asia region:  
854 carbon, nitrogen and phosphorus fluxes. LOICZ Reports and Studies 16, LOICZ, Texel, The  
855 Netherlands, 127 pp.
- 856 Yuan, D., Lin, B., Falconer, R.A., 2007. A modelling study of residence time in a macro-tidal  
857 estuary. Estuarine Coastal and Shelf Sciences 71, 401–411.
- 858 Warner, J.C., Geyer, W.R., Lerczak, J.A., 2005. Numerical modeling of an estuary: A  
859 comprehensive skill assessment. Journal of Geophysical Research 110(C05001), 1–13.
- 860 Warner, J.C., Geyer, R.W. and Arango, H.G., 2010. Using a composite grid approach in a complex  
861 coastal domain to estimate estuarine residence time. Computers and Geosciences Computers  
862 & Geosciences 36, 921–935.
- 863 Wolanski, E., 2007. Estuarine Ecohydrology. Elsevier, Amsterdam, 157 pp.
- 864 Wood, T.J., 1979. A modification of existing simple segmented tidal prism models of mixing in  
865 estuaries. Estuarine, Coastal and Shelf Science 8, 339–347.
- 866 Wu, Y., Falconer, R.A, Lin, B., 2005. Modelling tracer metal concentration distribution in estuarine  
867 waters. Estuarine, Coastal and Shelf Science 64, 699–709.
- 868 Zhang W.G., Wilkin J.L., Schofield, O.M.E., 2010. Simulation of water age and residence time in  
869 New York Bight. Journal of Physical Oceanography 40, 965–982.
- 870 Zimmerman, J.T.F., 1976. Mixing and flushing of tidal embayments in the western Dutch Wadden  
871 Sea. Part I: Distribution of salinity and calculation of mixing timescales, Netherlands  
872 Journal of Sea Research 10, 149-191.
- 873 Zimmerman, J.T.F., 1988. Estuarine residence times, p. 75–84. In B. Kjerfve (ed.), Hydrodynamics  
874 of Estuaries, Volume 1: Estuarine Physics. CRC Press, Boca Raton, Florida.

Table 1 – Water renewal timescale (days) of the Curimataú, Caravelas, Peruípe, Hudson, Conwy, Scheldt and York estuaries. For the Curimataú Estuary, A denotes the dispersive timescale using  $K$  calculated from Fisher's formula while B uses Hansen-Rattray's formula. The advection contribution to water renewal is  $\theta = T_P / T_a$  and  $\theta_{LOICZ} = T_{LOICZ} / T_2^{LOICZ}$ . The coefficient  $k^{LOICZ}$  was estimated using  $K^{LOICZ} = L Q_R 0.5(S_O + S_E) / A(S_O - S_E)$ , and  $k$  was estimated using  $K = L Q_R S_E / A(S_O - S')$ .

Curimataú Estuary										
Conditions	$T_2$	$T_2^{LOICZ}$	$T_1$	$T_P$	$T_{LOICZ}$	$T_{FRAC}$	$\Theta$	$\Theta_{LOICZ}$	$k(m^2 s^{-1})$	$k^{LOICZ}(m^2 s^{-1})$
(A) neap tides	2.4-4.7	0.5-1.0	1.4-2.4	0.9-1.6	0.5-0.9	0.6-1.2	0.64-0.67	0.36-0.38	245-480	735-1600
(B) neap tides	2.3-4.6	0.5-1.0	1.4-2.4	0.9-1.6	0.5-0.9	0.6-1.2	0.50-0.67	0.36-0.38	250-500	-
(A) spring tides	1.8-3.3	0.5-1.4	4.3-7.1	1.2-2.2	0.5-1.4	0.5-1.6	0.30-0.31	0.11-0.20	352-656	653-2267
(B) spring tides	1.7-2.7	0.5-1.4	4.3-7.1	1.2-2.0	0.5-1.4	0.5-1.6	0.27-0.28	0.11-0.20	435-700	-
Caravelas Estuary										
neap tides	9.5-70.5	3.2-45.8	55.6-388.9	8.1-59.7	3.0-40.9	3.1-43.2	0.14-0.15	0.05-0.11	24-179	37-552
spring tides	7.2-51.9	3.0-32.7	55.6-388.9	6.4-45.8	2.9-30.2	2.9-31.4	0.11-0.12	0.05-0.08	31-227	49-525
Peruípe Estuary										
neap tides	1.8-4.4	0.3-2.7	2.3-13.9	0.5-3.3	0.3-2.3	0.3-2.5	0.22-0.24	0.13-0.17	66-427	107-1035
spring tides	0.3-1.7	0.1-1.4	2.3-13.9	0.2-1.5	0.1-1.2	0.1-1.3	0.09-0.11	0.04-0.09	173-1071	212-1938
Hudson Estuary										
neap tides	31.4-43.0	9.4-12.1	7.9-9.5	6.3-7.8	4.3-5.3	5.9-7.4	0.80-0.82	0.53-0.54	545-745	1935-2484
spring tides	36.8-28.3	9.7-12.5	11.0-13.3	7.9-9.8	4.2-5.2	6.7-8.5	0.72-0.74	0.38-0.39	637-828	1880-2419
Conwy Estuary										
neap tides	157.5-175	49.5-54.9	89.9-98.7	56.8-63.1	31.8-35.3	38.7-43.0	0.63-0.64	0.35-0.36	9.5-10.4	30.25-32.95
spring tides	100-111.2	36.7-40.7	120.7-134	54.8-60.8	28.1-31.2	31.9-35.3	0.45-0.46	0.23-0.24	14.9-16.3	40.63-44.27
Mersey Estuary										
average conditions	4.9-13.4	1.9-7.6	27.4-72.4	4.2-11.3	1.8-6.9	1.8-7.2	0.16-0.17	0.07-0.10	304-1070	534-2771
Scheldt Estuary										
average conditions	146-207	9.7-26.8	161-214.4	76.4-105.4	9.2-23.8	9.5-25.2	0.47-0.49	0.06-0.11	581-826	2996-9268
York River Estuary										
high flow	45.0-81.0	17.7-36.0	109.3-161.9	31.9-54.0	15.2-29.9	16.4-32.4	0.28-0.33	0.14-0.18	316-568	974-1053
mean flow	112.2-201.9	44.2-89.8	272.6-403.9	79.5-134.6	38.0-73.4	40.9-80.8	0.29-0.33	0.14-0.18	127-228	390-422

Table 2 – The mean residence time ( $\bar{\phi}_{CART}$  in days), mean exposure time ( $\bar{\Theta}_{CART}$  in days), and the return coefficient of the Curimataú, Caravelas, Peruípe, Hudson, Conwy, Mersey, Scheldt and York estuaries. The Peclet number  $P_E = uL/K$  was calculated using the coefficient  $K$  from Fisher’s formula (see Table 1), which was used in our method.  $\bar{\phi}_a$  and  $\bar{\Theta}_a$  correspond to the mean advective timescale in the residence time and exposure time, respectively.

Curimataú Estuary								
Conditions	$P_E$	$\bar{\phi}_a$ and $\bar{\Theta}_a$	$\bar{\phi}_b$	$\bar{\Theta}_b$	$\bar{\phi}_{CART}$	$\bar{\Theta}_{CART}$	Return coefficient	Residence time from Numerical Model
neap tides	1.02-3.27	0.7-1.2	-0.6	0.5	0.1-0.6	1.2-1.7	0.65-0.92	-
spring tides	0.25-0.76	2.2-3.6	-2.1 to -3.1	2.0-2.8	0.1-0.5	4.1-6.4	0.92-0.98	-
Caravelas Estuary								
neap tides	0.02-1.25	27.8-194.5	-158.0 to -27.7	27.6-137.2	0.1-36.3	55.4-331.7	0.89-0.99	4.2-10.3
spring tides	0.02-0.98	27.8-194.5	-163.2 to -27.7	27.6-143.9	0.1-21.3	55.4-338.3	0.89-0.99	
Peruípe Estuary								
neap tides	0.04-1.47	1.7-7.0	-5.3 to -1.1	1.1-4.5	0.1-1.6	2.3-11.5	0.86-0.96	1.5-2.5
spring tides	0.02-0.56	1.7-7.0	-6.3 to -1.2	1.1-5.8	0.1-0.7	2.3-12.8	0.86-0.95	
Hudson Estuary								
neap tides	3.32-5.47	4.0-4.8	-2.1 to -1.6	1.4-1.7	1.9-3.1	5.4-6.2	0.50-0.65	8-10
spring tides	2.13-3.34	5.5-6.7	-3.5 to -3.7	2.9-3.0	1.8-3.2	8.5-9.6	0.67-0.79	8-9
Conwy Estuary								
neap tides	1.62-1.93	45.0-49.4	-34.4 to -33.3	28.0-28.5	11.6-15.0	73.0-77.9	0.81-0.84	-
spring tides	0.76-0.91	60.4-67.0	-57.0 to -52.8	47.6-50.6	7.6-10.0	107.9-117.6	0.92-0.93	-
Mersey Estuary								
Average conditions	0.16-0.21	13.7-36.2	-20.0 to -43.7	19.5-63.9	0.6-1.6	40.0-109.2	0.98-0.99	0.7-4
Scheldt Estuary								
Average conditions	0.91-0.97	80.5-107.2	-90.1 to -68.5	60.8-79.5	12.0-17.1	141.3-186.7	0.91-0.92	5-70
York River Estuary								
high flow	0.28-0.74	54.7-81.0	-52.1 to -71.1	49.9-64.2	2.5-9.9	104.5-145.1	0.93-0.98	11.3-33.3
mean flow	0.28-0.74	136.4-201.9	-130 to -177.3	124.4-160.1	6.3-24.7	260.7-362.1	0.93-0.98	18.1-59.3



## Appendix 1

Features of the Curimataú Estuary, Brazil (Miranda et al., 2005; Miranda et al., 2006; Andutta et al., 2006).

Feature	Curumataú Estuary
mean estuary depth $h$ (m)	6-7
mean estuary width $l$ (m)	250-350
mean estuary cross-sectional area $A$ (m <sup>2</sup> )	1500-2450
longitudinal distance $L$ (m)	10000
estuary volume $V$ (m <sup>3</sup> ) x10 <sup>6</sup>	15-24.5
river discharge $Q$ (m <sup>3</sup> s <sup>-1</sup> )	120 (measured in neap tides)
	40 (measured in spring tides)
gradient of salinity $dS/L$ (psu m <sup>-1</sup> )	neap tides $3.0 \times 10^{-3}$
	spring tides $1.3 \times 10^{-3}$
estuarine salinity $S_E$	$S_{E(neap)}$ 15-18
	$S_{E(spring)}$ 28-32
salinity at mouth $S_O$	$S_{O(neap)}$ 30
	$S_{O(spring)}$ 36
salinity at 10 km away from the mouth; $S'$	$S_{UP(neap)}$ 0
	$S_{UP(spring)}$ 23
dispersion $K$ (m <sup>2</sup> s <sup>-1</sup> ) Hansen-Rattray's formula	$K_{(neap)}$ 250-800
	$K_{(spring)}$ 435-1400
dispersion $K$ (m <sup>2</sup> s <sup>-1</sup> ) Fisher's formula	$K_{(neap)}$ 490-960
	$K_{(spring)}$ 703-1313

Estimative of horizontal dispersion $K$ to the Curimataú Estuary using Hansen-Rattray formula and vertical steady state salinity profiles of Miranda et al., 2005				
Parameters	Neap tide		Spring tide	
residual velocity- $u$ (m s <sup>-1</sup> )	0.057		0.016	
salinity at ( $x = 10$ km) - $S_{UP}$	0		23	
salinity at mouth - $S_O$	30		36	
longitudinal distance - $L$ (m)	10000		10000	
position of the vertical profile $x$ (m)	5000		5000	
non-dimensional number ( $v$ )	0.65		1	
vertical viscosity - $N_z$ (m <sup>2</sup> s <sup>-1</sup> )	$1 \times 10^{-4}$		$5 \times 10^{-4}$	
vertical difusivity - $K_z$ (m <sup>2</sup> s <sup>-1</sup> )	$1 \times 10^{-3}$		$1 \times 10^{-3}$	
wind strees - (Nm <sup>-2</sup> )	0		0	
average depth - $h$ (m)	6	7	6	7
estuarine salinity - $S_E$	17.0-19	20.3-23.5	29.0-30.0	30.3-32.5

## Appendix 2

Features of the Caravelas Estuary, Brazil (Schettini and Miranda 2010; Andutta, 2011).

Feature	Caravelas Estuary
mean estuary depth $h$ (m)	6-7
mean estuary width $l$ (m)	600-800
mean estuary cross-sectional area $A$ (m <sup>2</sup> )	$3.6-5.6 \times 10^3$
longitudinal distance $L$ (m)	~12000
estuary volume $V$ (m <sup>3</sup> ) $\times 10^6$	43.2-67.2
river discharge $Q$ (m <sup>3</sup> s <sup>-1</sup> )	$Q \sim 2-9$
gradient of salinity $dS/L$ (psu m <sup>-1</sup> )	neap tides $4.75 \times 10^{-4}$ spring tides $3.75 \times 10^{-4}$
estuarine salinity $S_E$	$S_{E (neap)}$ 32-34 $S_{E (spring)}$ 33-34
salinity at mouth $S_O$	$S_{O (neap)}$ 35.9 $S_{O (spring)}$ 36.0
salinity at 10 km away from the mouth; $S_{UP}$	$S_{UP (neap)}$ 30.2 $S_{UP (spring)}$ 31.5

### Appendix 3

Features of the Peruípe Estuary, Brazil (Schettini and Miranda 2010; Andutta, 2011).

Feature	Peruípe Estuary
mean estuary depth $h$ (m)	7-8
mean estuary width $l$ (m)	400-600
mean estuary cross-sectional area $A$ (m <sup>2</sup> )	2.8-4.8 x10 <sup>3</sup>
longitudinal distance $L$ (m)	~5000
estuary volume $V$ (m <sup>3</sup> ) x10 <sup>6</sup>	14-24
river discharge $Q$ (m <sup>3</sup> s <sup>-1</sup> )	Q ~ 20-70
gradient of salinity $dS/L$ (psu m <sup>-1</sup> )	neap tides 1.6x10 <sup>-3</sup>
	spring tides 3.6 x10 <sup>-4</sup>
estuarine salinity $S_E$	$S_{E(neap)}$ 26-28
	$S_{E(spring)}$ 29-30
salinity at mouth $S_O$	$S_{O(neap)}$ 31.6
	$S_{O(spring)}$ 32.0
salinity at 10 km away from the mouth; $S_{UP}$	$S_{UP(neap)}$ 23.4
	$S_{UP(spring)}$ 28.5



## Appendix 4

Features of the Hudson Estuary, USA (Warner et al., 2005).

Feature	Hudson Estuary
mean estuary depth $h$ (m)	7.5-8.5
mean estuary width $l$ (m) $\times 10^3$	1.55-1.65
mean estuary cross-sectional area $A$ (m <sup>2</sup> ) $\times 10^3$	11.625-14.025
longitudinal distance $L$ (m)	45000
estuary volume $V$ (m <sup>3</sup> ) $\times 10^6$	523-631
river discharge $Q$ (m <sup>3</sup> s <sup>-1</sup> )	770 (measured in neap tides)
	550 (measured in spring tides)
gradient of salinity $dS/L$ (psu m <sup>-1</sup> )	neap tides $4.7 \times 10^{-4}$
	spring tides $4.9 \times 10^{-4}$
estuarine salinity $S_E$	$S_{E(neap)}$ 7.5-8.5
	$S_{E(spring)}$ 13-14
salinity at mouth $S_O$	$S_{O(neap)}$ 34
	$S_{O(spring)}$ 36
salinity at 45 km away from the mouth; $S_{UP}$	$S_{UP(neap)}$ 0
	$S_{UP(spring)}$ 0

## Appendix 5

Features of the Conwy Estuary, UK (Turrell et al., 1996).

Geometry features by segments						
segment	distance from Deganwy Narrows (km)	length $L$ (km)	width $l$ (m)	depth $h$ (m)	area $\times 10^3 A$ (m <sup>2</sup> )	volume $\times 10^6 V$ (m <sup>3</sup> )
a)	2.44-3.93	1.5	530	5.0-5.5	2.7-2.9	$V_a = 4.0-4.3$
b)	3.93-7.86	3.9	430	4.5-5.0	1.9-2.2	$V_b = 7.6-8.4$
c)	7.86-9.51	1.7	225	4.0-4.5	900-1013	$V_c = 1.5-1.7$
d)	9.51-11.14	1.6	120	3.5-4.0	420-480	$V_d = 0.7-0.8$
e)	11.14-14.32	3.2	90	3.0-3.5	270-315	$V_e = 0.9-1.0$
total	2.44-14.32	11.9	246-274	5.0	1228-1368	$V_{total} = 14.6-16.2$
Features of salinity by segments						
segment	Volume normalized	$\Delta S$ (neap)	$S_o$ (neap)	$\Delta S$ (spring)	$S_o$ (spring)	
a)	$V_e/V_{total} \sim 0.27$	26.5-28.0	$S_a \sim 27.3$	30.0-32.0	$S_a \sim 31.0$	
b)	$V_b/V_{total} \sim 0.52$	15.0-26.5	$S_b \sim 20.8$	24.2-30.0	$S_b \sim 27.1$	
c)	$V_c/V_{total} \sim 0.10$	10.5-15.0	$S_c \sim 12.8$	20.5-24.2	$S_c \sim 22.4$	
d)	$V_d/V_{total} \sim 0.05$	5.0-10.5	$S_d \sim 7.8$	19.0-20.5	$S_d \sim 19.8$	
e)	$V_e/V_{total} \sim 0.06$	0.0-5.0	$S_e \sim 2.5$	13.9-19.0	$S_e \sim 16.5$	
total	$V_{total}/V_{total} = 1$	0.0-28.0	$S_{total} \sim 20.0$	13.9-32.0	$S_{total} \sim 26.7$	

**Notes:** To estimate each segment mean width it was used the ruler tool from Google Earth. The mean salinity of the entire system was calculated from the volume normalizer expression:

$$S_{total} = S_a \cdot \frac{V_a}{V_{total}} + S_b \cdot \frac{V_b}{V_{total}} + \dots + S_f \cdot \frac{V_f}{V_{total}} .$$

Feature	Conwy Estuary
mean estuary depth $h$ (m)	5.0
mean estuary width $l$ (m)	246-274
mean estuary cross-sectional area $A$ (m <sup>2</sup> ) $\times 10^3$	1.23-1.34
longitudinal distance $L$ (m)	11880
estuary volume $V$ (m <sup>3</sup> ) $\times 10^6$	14.6-16.2
river discharge $Q$ (m <sup>3</sup> s <sup>-1</sup> )	1.9 (measured in neap tides) 1.4 (measured in spring tides)
gradient of salinity $dS/L$ (psu m <sup>-1</sup> )	neap tides $2.4 \times 10^{-3}$ spring tides $1.5 \times 10^{-3}$
estuarine salinity $S_E$	$S_{(neap)}$ 19.19 $S_{(spring)}$ 26.46
salinity at mouth $S_O$	$S_{O(neap)}$ 34 $S_{O(spring)}$ 36
salinity at 11.88 km away from the mouth; $S_{UP}$	$S_{UP(neap)}$ 0 $S_{UP(spring)}$ 14

## Appendix 6

Features of the Mersey Estuary, UK (Bowden and Gilligan, 1971; Wu et al., 2005; Yuan et al., 2007).

Geometry features by segments						
Segment	Distance from Mersey entrance (km)	Length $L$ (km)	Width $l$ (m)	Depth $h$ (m)	Area $\times 10^3 A$ ( $m^2$ )	Volume $V \times 10^6$ ( $m^3$ )
a)	0-4.4	4.4	1200	15-17	18.7-21.2	$V_a$ 233-264
b)	4.41-7.86	3.7	1200	15-17	17.9-20.3	$V_b$ 214-243
c)	7.86-12.83	5	2400	15-17	35.7-40.5	$V_c$ 850-963
d)	12.83-20	7.2	3500	15-17	52.5-59.5	$V_d$ 184-208
total	0-20	20	1400	15-17	15.7-17.7	$V_{total}$ 313-355
Features of salinity by segments						
Segment	Volume normalized		$\Delta S$	$S_o$		
a)	$V_a/V_{total} \sim 0.07$		29.4-30.1	$S_a \sim 29.75$		
b)	$V_b/V_{total} \sim 0.07$		28.7-29.4	$S_b \sim 29.05$		
c)	$V_c/V_{total} \sim 0.27$		27.8-28.7	$S_c \sim 28.25$		
d)	$V_d/V_{total} \sim 0.59$		24.9-27.8	$S_d \sim 26.35$		
total	$V_{total}/V_{total} = 1$		24.9-30.1	$S_{total} \sim 27.29$		

**Notes:** To estimate each segment mean width it was used the ruler tool from Google Earth. The mean salinity of the entire system was calculated from the volume normalizer expression:

$$S_{total} = S_a \cdot \frac{V_a}{V_{total}} + S_b \cdot \frac{V_b}{V_{total}} + \dots + S_f \cdot \frac{V_f}{V_{total}}.$$

Feature	Mersey Estuary
mean estuary depth $h$ (m)	15-17
mean estuary width $l$ (m)	1445
mean estuary cross-sectional area $A$ ( $m^2$ ) $\times 10^3$	15.7-17.8
longitudinal distance $L$ (m)	20000
estuary volume $V$ ( $m^3$ ) $\times 10^6$	313-355
river discharge $Q$ ( $m^3 s^{-1}$ )	50-150 (range)
gradient of salinity $dS/dx$ (psu $m^{-1}$ )	$2.6 \times 10^{-4}$
estuarine salinity $S_E$	27-28
salinity at mouth $S_O$	30
salinity at 20 km upstream from the mouth; $S_{UP}$	25

## Appendix 7

Features of the Scheldt Estuary, (de Brauwere et al., 2011; de Brye et al., 2012).

<b>Geometry features by segments</b>					
Segment	Length $L$ (km)	Width $l$ (m)	Depth $h$ (m)	Cross-sectional Area $A$ $10^3$ (m <sup>2</sup> )	Volume $V$ $10^3$ (m <sup>3</sup> )
1	7950	372.5	9.0	3.35	$V_1 = 26649$
2	8300	370.2	10.4	3.85	$V_2 = 31957$
3	9600	666.0	9.3	6.19	$V_3 = 59460$
4	5100	559.3	10.1	5.65	$V_4 = 28807$
5	9700	779.9	9.0	7.02	$V_5 = 69830$
6	5950	2754.8	8.0	22.04	$V_6 = 131128$
7	5700	2533.2	6.0	15.20	$V_7 = 86636$
8	5300	2702.9	7.4	20.00	$V_8 = 106008$
9	5900	2257.2	11.9	26.86	$V_9 = 158476$
10	6900	5033.6	9.0	45.30	$V_{10} = 312588$
11	6200	4889.5	9.9	48.41	$V_{11} = 300120$
12	12100	4088.7	12.0	49.06	$V_{12} = 593684$
13	13300	4791.6	13.7	65.65	$V_{13} = 873080$
Total	102000	2808.2	9.7	27.24	2778423

<b>Feature</b>	<b>Scheldt Estuary</b>
mean estuary depth $h$ (m)	9.7
mean estuary width $l$ (m)	2808.2
mean estuary cross-sectional area $A$ (m <sup>2</sup> ) $\times 10^3$	27.24
longitudinal distance $L$ (m)	102000
estuary volume $V$ (m <sup>3</sup> ) $\times 10^6$	2778
river discharge $Q$ (m <sup>3</sup> s <sup>-1</sup> )	150-200 (average range)
gradient of salinity $dS/L$ (psu m <sup>-1</sup> )	$1.0 \times 10^{-4}$
estuarine salinity $S_E$	30-32
salinity at mouth $S_O$	34
salinity at 102 km away from the mouth; $S_{UP}$	5

## Appendix 8

Features of the York River Estuary, (Sandifer, 1973; Hass, 1977; Shen and Haas, 2004).

Feature	York River Estuary
mean estuary depth $h$ (m)	8-10
mean estuary width $l$ (m)	2700-3200
mean estuary cross-sectional area $A$ ( $\text{m}^2$ ) $\times 10^3$	21.6-32.0
longitudinal distance $L$ (m)	47000
estuary volume $V$ ( $\text{m}^3$ ) $\times 10^6$	1015-1504
river discharge (Mattaponi and Pamunkey rivers) $Q$ ( $\text{m}^3\text{s}^{-1}$ )	43.1 (average)
river discharge (Mattaponi and Pamunkey rivers) $Q$ ( $\text{m}^3\text{s}^{-1}$ )	107.5 (high)
estuarine salinity $S_E$	16-17
salinity at mouth $S_O$	20
salinity at 47 km away from the mouth; $S_{UP}$	12-13

FIGURE 1




Figure 1– Illustration of the concept of residence and exposure times for a single water particle in the domain, between the upstream,  $x = L_1$ , and downstream,  $x = L_0$ , open boundaries. Residence time and exposure time are  $\bar{\phi} = t_1$  and  $\bar{\Theta} = t_1 + (t_3 - t_2)$ , respectively. Timescales regarding a single particle are introduced for pedagogical purposes only, and are actually physically meaningless.

**FIGURE 2**




Figure 2 – (a) A 1-D estuary model; (b) the tidal prism box model (side view); (c) the gravitational circulation box model (side view); (d) the LOICZ box model for a vertically well-mixed estuary (plan view).

**FIGURE 3**




Figure 3 – Conceptual description of a generic salinity distribution along an estuary from ( $x = L_1$ ) to the mouth ( $x = L_0$ ). The salinity  $S_{UP}$  at  $x = L_1$ , and ( $0 \leq S_{UP} < S_E$ ).



**FIGURE 4**




Figure 4 – Representation of the residence time  $\phi$  (solid curve) and the exposure time  $\Theta$  (dashed curve) as a function of the distance  $x$  to the upstream boundary of the domain. The timescales are normalised by means of the advective timescale  $T_1$ . It is noteworthy that the exposure time, as opposed to the residence time, is defined including areas outside the domain of interest ( $L_1 \leq x/L \leq L_0$ ).

**FIGURE 5**




Figure 5 – Representation for various values of the Peclet number of the return coefficient as defined by formula (34).

**FIGURE 6**




Figure 6 – (a) Scatter plot of the mean water renewal timescale,  $T_P$ , against the residence time at the upstream location from numerical models. (b) Scatter plot of the mean dispersive contribution to water renewal  $(1 - \theta)$  against the mean return coefficient calculated using CART's formula.




Figure 7- The position of estuaries on the advection-diffusion diagram to indicate the relative contribution to the water renewal  $T_P$  by the advective ( $T_1$ ) and dispersive ( $T_2$ ) timescales using a logarithmic scale. Subscript (n) and (s) indicate neap and spring tide conditions. For the Curimataú Estuary (C),  $K$  was estimated from Hansen-Rattray's formula. Hudson, Caravelas, Peruípe, and Conwy estuaries are denoted (H), (CA), (P), and (C), respectively. For the Mersey (M), and Scheldt (S) estuaries, results are from an average range. For the York River Estuary the conditions are for high river flow ( $Y_H$ ), and mean river flow ( $Y_M$ ).