

I. Darren Grice,^a Ian D. Jenkins,^b
W. Ken Busfield,^c Karl A. Byriel^d
and Colin H. L. Kennard^{e*}^aInstitute for Glycomics, Griffith University,
Gold Coast, Queensland, 4215, Australia,^bNatural Product Discovery, Griffith University,
Nathan, Brisbane, Queensland, 4111, Australia,^cSchool of Science, Griffith University, Nathan,
Brisbane, Queensland, 4111, Australia, ^dCentre
for Drug Design and Development, The
University of Queensland, Brisbane,
Queensland, 4072, Australia, and ^eChemistry
Department, School of Molecular and Microbial
Sciences, The University of Queensland,
Brisbane, Queensland, 4072, Australia

Correspondence e-mail: C.Kennard@uq.edu.au

Key indicators

Single-crystal X-ray study

T = 293 K

Mean $\sigma(\text{C}-\text{C}) = 0.008 \text{ \AA}$

R factor = 0.039

wR factor = 0.142



Data-to-parameter ratio = 15.2

For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.**tert-Butyl diphenylphosphinate**Molecules of the title compound, $\text{C}_{16}\text{H}_{19}\text{O}_2\text{P}$, pack simply in the orthorhombic space group $Pbca$, the dominant molecular feature being a tetrahedral P atom attached to two phenyl rings, a phosphate O atom and a tertiary butyloxy group.

Received 19 October 2004

Accepted 12 November 2004

Online 20 November 2004

CommentThe title compound, (I), was part of an investigation using the radical scavenger 1,1,3,3-tetramethyl-2,3-dihydro-1*H*-isoindol-2-yloxy to study the reaction of phosphorus-centred radicals with alkenes and alkynes (Busfield *et al.*, 1995, Bottle *et al.*, 1994).There is an intramolecular short distance between H26 and O4 of 2.46 Å, with an angle of 106° for C26–H26...O4. Reaction of the title compound with polyether ligands has been reported by Albanese *et al.* (2001).**Experimental**The title compound was obtained in 16% yield from the reaction of diphenylphosphine (100 mg), di-*tert*-butyl peroxyoxalate (DTBP) (58 mg) and benzene (2.5 ml) (solvent) in the presence of the radi-**Figure 1**

The molecular structure of the title compound, (3), with displacement ellipsoids drawn at the 50% probability level.

cal scavenger 1,1,3,3-tetramethyl-2,3-dihydro-1*H*-isoindol-2-yloxy (103 mg), along with five other phosphorus-containing compounds. Reaction mixtures were degassed using repeated freezing/evacuating/thawing cycles on a high vacuum line, then sealed under vacuum in glass and heated for 10 half lives of the initiator, DTBP (68 minutes at 333 K). The reaction mixture was then separated by HPLC (Whatman Partisil 10-ODS-3 500 x 10 mm C18) using an isocratic 80:20 methanol/water, 4.0 ml/min method.

The isolated phosphorus-containing reaction products were eluted from the reversed-phase HPLC column in the following order.

(1) Diphenylphosphinic acid (0.4%), identical to an authentic sample by HPLC, m.p. 466–468 K. (Edmundson, 1986).

(2) Diphenylphosphine oxide (4.8%), identical to an authentic sample by HPLC and NMR.

(3) *tert*-Butyl diphenylphosphinate (16%). ^1H NMR δ (250 MHz, CDCl_3) 1.51, *s*, 9H, (CH_3)₃; 7.42, *m*, 6H, 7.79, *m*, 4H, aromatic H. ^{13}C NMR δ (62.8 MHz, CDCl_3) 30.8, $^3J_{\text{PC}}$ 18.5, (CH_3)₃; 83.4, C(CH_3)₃; 128.2, $^3J_{\text{PC}}$ 13.2, *meta* C; 129.1, *para* C; 131.3, $^2J_{\text{PC}}$ 10.7, *ortho* C; 134.8, J_{PC} 138.5, *ipso* C. ^{31}P NMR (121 MHz, CDCl_3) 9.94. Structural confirmation by X-ray structural analysis.

(4) 2-(Diphenylphosphinyl)-1,1,3,3-tetramethyl-2,3-dihydro-1*H*-isoindole (6.3%), white solid, m.p. 458–461 K (Found: C, 77.3; H, 6.5; N, 3.6. $\text{C}_{24}\text{H}_{26}\text{NOP}$ requires C, 76.8; H, 6.9; N, 3.7%). ^1H NMR δ (250 MHz, CDCl_3) 1.56, *br s*, 12H, 4CH₃; 7.04, *m*, 2H, H4, H7; 7.23, *m*, 2H, H5, H6; 7.44, *m*, 6H, 7.83, *m*, 4H, Phenyl; ^{13}C NMR δ (62.8 MHz, CDCl_3) 32.2, *br s*, 4 \times ring CH₃; 68.6, C1, C3; 120.9, C4, C7; 127.5, C5, C6; 127.8, *d*, $^2J_{\text{PC}}$ 15.7, *d*, *ortho* C; 131.5, *s*, *para* C; 133.4, $^3J_{\text{PC}}$ 10.7, *d*, *meta* C; 133.7, $^1J_{\text{PC}}$ 138.1, *d*, *ipso* C; 145.9, C3a, C7a. ^{31}P NMR δ (121 MHz, CDCl_3) 9.70.

(5) 2-(Diphenylphosphinyloxy)-1,1,3,3-tetramethyl-2,3-dihydro-1*H*-isoindole (34.3%), identical to an authentic sample (Busfield *et al.*, 1995).

(6) The material, 1,1,3,3-tetramethyl-2,3-dihydro-1*H*-isoindol-2-yloxy diphenylphosphinite (12%), was unstable, decomposing almost immediately in warmed aqueous THF to give by HPLC the free radical scavenger and diphenylphosphine oxide. High resolution MS failed to give a parent ion and the material was too unstable for elemental analysis. However, the following spectroscopic data were obtained: ^1H NMR δ (250 MHz, CDCl_3) 1.36, *br s*, 12H, 4CH₃; 7.18–7.30, *m*, 8H, H4–H7, 4*H*-phenyl; 7.47, *m*, 6H, Phenyl; ^{13}C NMR δ (62.8 MHz, CDCl_3) 29.9, 30.0, *d*, 4 \times ring CH₃; 120.0, C4, C7; 128.1, *d*, $^3J_{\text{PC}}$ 6.97, *meta* C; 128.6, *s*, *para* C; 130.0, *d*, $^2J_{\text{PC}}$ 22.1, *ortho* C; 143.7, 144.0, C3a, C7a. ^{31}P NMR δ (121 MHz, CDCl_3) 86.90. C1, C3, C5 and C6 from the isoindole moiety and the two *ipso* C atoms of the phenyl rings were not detected in the ^{13}C NMR spectrum. Percentage yields are based on moles of diphenylphosphine.

Sosnovsky & Zaret (1969) prepared the title compound by reacting diphenylchlorophosphine with *tert*-butyl hydroperoxide in pyridine. Another synthesis was reported by Sosnovsky *et al.* (1970).

Crystal data

$\text{C}_{16}\text{H}_{19}\text{O}_2\text{P}$
 $M_r = 274.28$
 Orthorhombic, *Pbca*
 $a = 11.530$ (2) Å
 $b = 17.118$ (2) Å
 $c = 15.285$ (2) Å
 $V = 3016.8$ (7) Å³
 $Z = 8$
 $D_x = 1.208$ Mg m⁻³

Mo $K\alpha$ radiation
 Cell parameters from 25 reflections
 $\theta = 10.0$ – 12.0°
 $\mu = 0.18$ mm⁻¹
 $T = 293$ (2) K
 Needle, colourless
 $0.40 \times 0.13 \times 0.12$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer
 ω – 2θ scans
 Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\text{min}} = 0.954$, $T_{\text{max}} = 0.979$
 2631 measured reflections
 2631 independent reflections

802 reflections with $I > 2\sigma(I)$
 $\theta_{\text{max}} = 25.0^\circ$
 $h = 0 \rightarrow 13$
 $k = 0 \rightarrow 20$
 $l = 0 \rightarrow 18$
 25 standard reflections
 frequency: 120 min
 intensity decay: 1%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.142$
 $S = 0.90$
 2631 reflections
 173 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0552P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.20$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³
 Extinction correction: *SHELXL97*
 Extinction coefficient: 0.0030 (4)

Table 1

Selected geometric parameters (Å, °).

P1–O1	1.476 (3)	P1–O4	1.569 (3)
O1–P1–O4	118.00 (18)	O1–P1–C11	112.3 (2)
O1–P1–C21	113.3 (2)	O4–P1–C11	106.40 (19)
O4–P1–C21	99.7 (2)	C21–P1–C11	105.7 (2)

The crystal selected was a poor diffractor and only 30% of the intensities were significantly greater than background. H atoms were positioned geometrically (C–H = 0.93–0.96 Å) and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2$ or 1.5 times $U_{\text{eq}}(\text{C})$.

Data collection: *SDP* (Frenz, 1985); cell refinement: *SDP*; data reduction: *WinGX* (Farrugia, 1999); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON98* (Spek, 1988); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the Australian Research Council, the University of Queensland and Griffith University for financial support for the purchase of the CAD-4 diffractometer.

References

- Albanese, D., Landini, D. & Maia, A. (2001). *J. Org. Chem.* **66**, 3249–3252.
 Bottle, S., Busfield, W. K., Grice, I. D., Heiland, K., Jenkins, I. D., Meutermaans, W. & Monteiro, M. (1994). *Prog. Pac. Polym. Sci.* **48**, 85–97.
 Busfield, W. K., Grice, I. D., & Jenkins, I. D. (1995). *Aust. J. Chem.* **48**, 625–634.
 Edmundson, R. S., (Ed) (1986). In *Dictionary of Organophosphorus Compounds*. London: Chapman and Hall.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Frenz, B. A. (1985). *Enraf–Nonius SDP-Plus Structure Determination Package*. Version 3.0. Enraf–Nonius, Delft, The Netherlands.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Sheldrick, G. M. (1997). *SHELXS97 and SHELXL97*. University of Göttingen, Germany.
 Spek, A. L. (1988). *J. Appl. Cryst.* **21**, 578–579.
 Sosnovsky, G. & Zaret, E. H. (1969). *J. Org. Chem.* **34**, 968–970.
 Sosnovsky, G., Zaret, E. H. & Schmitt, K. D. (1970). *J. Org. Chem.* **35**, 336–340.