

A comparison of the hand hygiene knowledge, beliefs and practices of Greek nursing and medical students

TF van de Mortel,
MHlthSc. FCN, FRCNA
Faculty of Health & Applied Sciences,
Southern Cross University*,
Lismore, 2480 NSW
Australia
Email: tvandemo@scu.edu.au
Ph: 61-2-66203305
Fax: 61-2-66203022

E. Apostolopoulou
PhD
A. Prof.,
Faculty of Nursing,
University of Athens#,
Greece

G. Petrikkos,
PhD
A. Prof.,
Faculty of Medicine#

Abstract

Profession influences healthcare workers' hand hygiene (HH) adherence. Greek nursing and medical students were surveyed to determine if there were cross-disciplinary differences in HH education, assessment, knowledge, beliefs and practices. Nursing students' HH knowledge was significantly higher than that of medical students. Nursing students reported significantly more positive HH practices and beliefs, and received more HH education and assessment than medical students. Improving undergraduate HH education may improve graduates' HH knowledge, beliefs and practices.

Introduction

Studies indicate that healthcare workers' adherence to hand hygiene (HH) guidelines is poor (~ 40%) and that physician status is a risk factor for non-adherence.¹ Disciplinary differences in HH education and assessment during undergraduate training may impact on graduates' behaviour upon entering the workforce.

Several studies have examined aspects of healthcare students' HH beliefs or practices. Duration of clinical experience,^{2,3} gender,² the example of mentors,² and perceptions of handwashing benefits, barriers and severity of infectious diseases⁴ significantly influenced self-reported^{3,4} or observed HH compliance.² Gaps in students' HH knowledge were also identified by Sangkard⁴ and Mann and Wood.⁵ Sangkard's survey of nursing students' infection control knowledge in relation to Human Immunodeficiency Virus infection included a short handwashing quiz with simple true/false questions. Students' scores on this quiz ranged from 68-71%. Mann and Wood reported that medical students' average score on a hand hygiene quiz was 52%. However, these studies can't be used to make cross-disciplinary comparisons as the questions on the respective surveys were very different.

Thus the objectives of this study were to:

- determine if the HH knowledge, beliefs, practices, education and assessment of undergraduate Greek nursing and medical students differed by discipline.
- use this information to inform HH education and assessment in the undergraduate curriculum.

Methods

A survey was conducted at the University of Athens, Greece, utilising a translated version of the HH Questionnaire,⁶ which elicited information on demographics, HH teaching and assessment, and HH knowledge (13 questions based on the CDC HH guidelines).¹ The questionnaire also measured HH beliefs (20 items), practices (14 items), and perceptions of the importance of HH in the curriculum (3 items) on 5-point Likert scales. The theoretical framework, scale items, questionnaire development and validation are described in van de Mortel.⁶

The questionnaire was translated into Greek by a bilingual infection control specialist, back-translated by an accredited company to ensure accuracy, and distributed to all final year nursing (n=90) and medical (n=60) students. The courses were four and six years in duration, respectively. Participation was voluntary, and responses were confidential. Ethics approval was obtained from the relevant Ethics Committees.

SPSS 16.0 for Mac was used to conduct analyses. The Cronbach's alpha values of the beliefs, practices and importance scales were 0.79, 0.74 and 0.71. A Chi² test was used to assess discipline differences in gender proportions. Descriptive statistics were calculated. Preliminary assumption testing was carried out prior to multivariate analysis. A one-way between-groups MANOVA was performed to investigate discipline differences in three domains (table 1). A Mann-Whitney test was used to examine the negatively skewed variable 'rate importance of HH from 1–10'.

Results

The response rates of nursing and medical students were 85.6% and 36%. Sixty-two percent of medical students and 82% of nursing students were female ($\chi^2 = 2.70(1)$; $p = 0.100$). Medical students were significantly older than nursing students (\bar{x} 25.20 years \pm 0.33 vs. 23.62 \pm 0.53)($t=-2.53(85,2)$; $p=0.013$) and had spent longer on practicum (\bar{x} 58.88 weeks \pm 9.68 vs. 36.39 \pm 1.11)($p=0.035$).

There were significant differences between disciplines on the combined dependent variables in each group; nursing students scored higher on all variables (table 1). Knowledge score, beliefs score, practices score, the frequency of HH assessment, the number of strategies used to teach HH, students' perceptions of teaching effectiveness, and perceptions of the importance of HH in the curriculum, were significantly different when considered alone.

Table 1. Differences in HH outcomes, education and perceptions by discipline.

Variable	Discipline	Range (mean \pm s.e.m)	F value (df)	p value	Partial eta ²
Outcomes			14.40 (3, 94)	0.000*	0.32 (L)
Knowledge	Nursing	3-11 (8.84 \pm 0.19)	35.16 (1, 96)	0.000*	0.27 (L)
	Medicine	3-11 (6.14 \pm 0.52)			
Beliefs (HBS)	Nursing	3.00-4.68 (3.92 \pm 0.05)	12.94 (1, 96)	0.000*	0.12 (M)
	Medicine	3.05-4.40 (3.52 \pm 0.08)			
Practices (HHPI)	Nursing	4.00-5.00 (4.75 \pm 0.03)	4.63 (1, 96)	0.034*	0.05 (S)
	Medicine	4.00-5.00 (4.59 \pm 0.08)			
Importance HH 1-10	Nursing	7-10 (9.60 \pm 0.08)		0.24	
	Medicine	5-10 (9.29 \pm 0.27)			
Education			13.85(2, 94)	0.000*	0.23 (L)
HH assessment frequency	Nursing	1-15 (7.13 \pm 0.42)	23.58 (1, 95)	0.000*	0.20 (L)
	Medicine	0-11 (2.86 \pm 0.74)			
No. teaching strategies	Nursing	3-13 (10.78 \pm 0.25)	12.56 (1, 95)	0.001*	0.12 (M)
	Medicine	0-13 (8.19 \pm 1.04)			
Perceptions			8.13 (2, 94)	0.000*	0.15 (L)
Effectiveness of teaching strategies	Nursing	0.46-3.38 (2.92 \pm 0.08)	12.30 (1, 95)	0.001*	0.12 (M)
	Medicine	0 -2.85 (2.43 \pm 0.20)			
Importance HH in curriculum (HIS)	Nursing	2-5 (3.89 \pm 0.07)	6.88 (1, 95)	0.004*	0.09 (M)
	Medicine	1-5 (3.23 \pm 0.37)			

L = large; M = moderate; S = small effect size; *significant at alpha < 0.05

Nursing students found lectures, tutorials, textbooks and lecture notes significantly more effective than medical students (table 2). Teaching in the clinical setting occurred most frequently and was highly rated by both groups. Nursing and medical students

were assessed on HH most frequently in the clinical setting (2.12 ± 0.18 vs 1.57 ± 0.37). Nursing students were assessed significantly more often using written ($\bar{x} 1.86 \pm 0.15$ vs 0.19 ± 0.15 ; $t=7.97$ (62.6); $p = 0.000$) and verbal assessment ($\bar{x} 1.42 \pm 0.16$ vs 0.43 ± 0.16 ; $t=4.36$ (61.5); $p = 0.000$), and in the simulated clinical setting ($\bar{x} 1.71 \pm 0.15$ vs 0.67 ± 0.23 ; $t=3.45$ (94); $p = 0.001$).

Table 2. Percentage of students taught HH using a particular method and perceived effectiveness (mean \pm s.e.m). 1 = ineffective, 4 = highly effective.

Teaching method	Nursing	Medicine	Significance
Lectures*	90.9% (2.84 \pm 0.09)	57.1% (1.67 \pm 0.26)	$t=4.77(80)$; $p= 0.000$
Tutorials*	96.1% (3.46 \pm 0.10)	71.4% (2.27 \pm 0.27)	$t=4.94(87)$; $p= 0.000$
Clinical setting	97.4% (3.43 \pm 0.09)	81.0% (3.18 \pm 0.25)	
Demonstration	96.1% (3.24 \pm 0.10)	76.2% (2.63 \pm 0.27)	
Practical laboratories	93.5% (3.11 \pm 0.11)	57.1% (3.17 \pm 0.17)	
Videos	81.8% (2.75 \pm 0.10)	47.6% (2.20 \pm 0.29)	
Textbooks*	96.1% (2.76 \pm 0.10)	81.0% (2.00 \pm 0.17)	$t=3.38(89)$; $p= 0.001$
Lecture notes*	89.6% (2.58 \pm 0.10)	61.9% (1.69 \pm 0.24)	$t=3.40(80)$; $p= 0.001$
Computer simulations	35.1% (2.41 \pm 0.19)	42.9% (2.22 \pm 0.28)	
Internet	53.2% (2.32 \pm 0.13)	47.6% (1.90 \pm 0.23)	
Research articles	68.8% (2.74 \pm 0.12)	57.1% (2.08 \pm 0.23)	
Published guidelines	94.8% (2.96 \pm 0.13)	71.4% (2.53 \pm 0.24)	
Posters	84.4% (3.02 \pm 0.15)	61.9% (2.85 \pm 0.32)	
Mean no. teaching strategies	9.95 (\pm 0.23)	7.52 (\pm 0.97)	

*significant at $\alpha < 0.004$

Discussion

Nursing students had greater HH knowledge, more positive beliefs and practices, and considered HH more important in their curriculum than medical students. Nursing

students also received more HH education, rated their education as more effective and received more frequent HH assessment than medical students despite the longer duration of the medical degree. Instruction on HH was an elective topic for medical students; if they didn't enrol in the electives there was limited emphasis on HH in their course. As assessment often drives learning⁷ this may have impacted on medical students' knowledge and beliefs. Additionally, Calabro et al.⁸ demonstrated that a single educational infection control intervention for medical students did not result in long-term information retention, indicating that repeated exposure may be necessary for retention of key information.

The results suggest that an increased emphasis on HH education and assessment in the undergraduate curriculum may improve students' HH knowledge, beliefs and practices and make the practice culture more positive towards HH. Contextualising HH education and assessment in the clinical setting may also improve learning outcomes as both student groups rated learning in the clinical environment as highly effective; the benefit of contextualisation in improving learning outcomes is supported by the literature.⁹

The study was limited by the small medical student sample (increasing the likelihood that the null hypothesis will be wrongly accepted), and the fact that only one higher education institution was sampled. Additionally, self-reported HH adherence can be substantially higher than observed adherence,² although this is not always the case.¹⁰ The longer duration of medical students' training may have impaired recall of their HH education and assessment, although it also offered more opportunities for both. Finally, without manipulating variables of interest and observing the effects upon outcomes, it is not possible to state with certainty that one variable is influencing another. Other unmeasured factors may have influenced outcomes.

Recommendations

Increasing the emphasis upon HH in the undergraduate curriculum through frequent HH education and assessment, particularly in the clinical setting, may improve students' HH knowledge, beliefs and practices and facilitate a positive practice culture towards HH. Studies with a larger and more diverse sample are needed to confirm these results.

1. Centers for Disease Control. Guideline for hand hygiene in healthcare settings. *MMWR* 2002;51(RR-16): 1-48.
2. Snow M, White GL, Alder SC, Stanford, JB. Mentor's hand hygiene practices influence student's hand hygiene rates. *Am J Infect Control* 2006;34:18-24.
3. Sangkard, K. Assessment of nursing students' knowledge about infection control: implications for nursing education. Doctoral thesis: University of Iowa; 1991.
4. Karraffa J. Handwashing practices of university students: development of an instrument to test the Health Belief Model. Doctoral thesis: Southern Illinois University; 1989.
5. Mann CM, Wood A. How much do medical students know about infection control? *J Hosp Infect* 2006;64(4): 366-70.
6. van de Mortel, T.F. Development of an instrument to assess healthcare students' hand hygiene knowledge, beliefs and practices. *Aust J Adv Nurs* 2009;26(3): 9-16.
7. Gibbs G, Simpson C. Does your assessment support your students' learning? *Learning and Teaching in Higher Education* 2004-2005;Issue 1: 3-31.
8. Calabro K, Bright K, Kouzekanani K. Long-term effectiveness of infection control training among fourth-year medical students. *Med Ed Online* 2000; Retrieved March 10, 2009 from <http://www.med-ed-online.org/res00009.htm>

9. Cordova DI, Lepper MR. Intrinsic motivation and the process of learning: Beneficial effects of contextualization, personalization, and choice. *J Educ Psychol* 1996; 88(4): 715-730.
10. Larson EL, Aiello AE, Carcillo JP. Assessing nurses hand hygiene practices by direct observation or self-report. *J Nurs Meas* 2004; 12(1): 77-85.