

**Impact of Centre-of-Mass Acceleration on Perceived Exertion, the Metabolic Equivalent and Heart Rate Reserve in Triathlete Spin Cycling: A Pilot Study**

Author

Evans, SA, James, DA, Rowlands, D, Lee, JB

Published

2022

Journal Title

Journal of Human Kinetics

Version

Version of Record (VoR)

DOI

[10.2478/hukin-2022-0004](https://doi.org/10.2478/hukin-2022-0004)

Rights statement

© 2022 Stuart A Evans, Daniel A James, David Rowlands, James B Lee, published by Sciendo. This work is licensed under the Creative Commons Attribution 4.0 International License.

Downloaded from

<http://hdl.handle.net/10072/420987>

Griffith Research Online

<https://research-repository.griffith.edu.au>



# Impact of Centre-of-Mass Acceleration on Perceived Exertion, the Metabolic Equivalent and Heart Rate Reserve in Triathlete Spin Cycling: A Pilot Study

by

Stuart A Evans<sup>1</sup>, Daniel A James<sup>1</sup>, David Rowlands<sup>2</sup>, James B Lee<sup>1</sup>

*Indoor spin cycling has gained popularity as a training modality for triathletes. Part of its appeal is that it can form a component of a structured periodised training program and provide an alternative to outdoor cycling. Indices of physiological components (i.e., the metabolic equivalent, caloric cost, perceived exertion) and changes in the body position can be inferred by wearable technology such as an accelerometer. This pilot study aimed to investigate the relationship between the rating of perceived exertion, heart rate reserve, and the metabolic equivalent between the whole body centre of mass acceleration using a sacrum mounted triaxial accelerometer during 20 minutes of 6 varied power conditions of indoor spin cycling. Compared with other conditions, cycling at a steady state (>152-205 W) resulted in extremely large effects (> 0.9) in mediolateral acceleration and the rating of perceived exertion ( $p < 0.0001$ ). The relationship between the body position (aerodynamic to drops) induced significant changes in anteroposterior acceleration magnitude ( $p < 0.0001$ ), although moving from drops to the aerodynamic position was not significant despite a large increase in heart rate reserve and extremely large effects of perceived exertion. The rating of perceived exertion scale and the metabolic equivalent comparative to the whole body centre of mass acceleration magnitude and power displayed a strong correlation ( $r = 0.865$ ). An individually determined whole body centre of mass accelerations combined with perceived exertion, the metabolic equivalent and heart rate reserve could potentially contribute to improved indoor triathlete spin cycling performance.*

**Key words:** spin cycling, trunk centre of mass, accelerometer, fatigue, MET, triathlon.

## Introduction

Specialised indoor bicycles manufactured using a weighted flywheel at the front are used in indoor cycling exercises called spinning. Cycling indoors using an immobile spin bike has gained popularity as a training modality for triathletes. Part of its appeal is that indoor spin cycling can form part of a structured and periodised training program and provide an alternative to outdoor cycling when faced with environmental extremes of cold, hot and/or hypoxic environments. In this regard, indoor spin cycling allows triathletes to preserve cardiovascular fitness levels and training loads.

In cycling, the lower extremities are responsible for producing energy imparted to the bicycle. To mediate this effect, triathletes continuously adjust the force and timing produced relative to the pedal position to obtain a specific self-selected pace (de Melo Dos Santos et al., 2017). A change in the body position may alter cycling variability due to both fatigue and mechanical factors (Padulo et al., 2012).

A characteristic of efficient movement control in cycling is that of trunk stability. Cycling requires trunk stabilisation in order to balance the bicycle (McDaniel et al., 2005). A forward shift of the whole body centre of mass (CoM) means that

<sup>1</sup> - SABEL Labs, College of Health and Human Science, Charles Darwin University, Darwin, 0810 Australia.

<sup>2</sup> - School of Engineering, Griffith University, Nathan, Queensland, Australia.

the cyclist's body is less supported by the saddle and would require additional stabilisation from the trunk. Similarly, a change in the cycling position when moving from the drop handlebars to an aerodynamic position would alter the whole body CoM. A low level of core trunk muscle strength or stability can cause additional upper body movement. In this instance, the capacity of the lumbar-pelvic-hip complex to control trunk movement and preserve trunk stability can be compromised. Therefore, the core musculature may influence the kinematics and load-bearing capacity of the knee by determining what loads are transmitted from the trunk. Taken together, poor trunk strength or stability combined with increases in power output could cause excessive CoM acceleration leading to inefficient movement. The idea being that higher intensity levels are related with larger mediolateral force swaying (Rannama et al., 2017) with strenuous cycling decreasing stability in the anteroposterior direction (Wiest et al., 2011). Correspondingly, an increase in the workload necessitates additional upper body stabilisation (McDaniel et al., 2005).

During incremental, non-steady state exercise, a point is reached at which a participant's ventilation shows a non-linear increase, which is termed the ventilatory threshold (Ghosh, 2004). Relative to cycling, this threshold implies an effort that exceeds the steady state. In steady state cycling (i.e., changes to intensity vary only negligibly over a specified time), the forces involved in pedalling largely fluctuate throughout the crank cycle (Gregor, 2000). While these designs have been utilised to inform decision-making as to the effects of indoor cycling, studies involving steady state indoor cycling are limited in comparison.

A multi-method approach, including intensity measured as relative (i.e., as a percentage of maximal oxygen uptake,  $VO_{2max}$ ) or absolute (as metabolic equivalent of tasks, METs), where one MET is defined as the resting metabolic rate (RMR), can be employed. Yet, this presents limitations as without direct measurements, the RMR is usually replaced by a 1-MET reference value of  $3.5 \text{ ml O}_2\text{kg}^{-1}\text{min}^{-1}$  (Ainsworth et al., 2011). Absolute and relative energy expenditure – EE (kcal, kJ, MET) can be estimated by caloric cost. Although notwithstanding its own limitations, the MET is a method to indicate and compare the

absolute intensity and energy expenditure of different physical activities (American College of Sports Medicine, 2008); thus, the MET is a measurement of exertion intensity (Pollock et al., 1998) and has been used alongside the rating of perceived exertion (RPE) during graded cycle ergometer tests (Kasimary Çakir et al., 2012). The MET and RPE may be attractive due to their non-invasive nature, low cost, squad-wide concurrent measurement capability and time efficiency.

The relationship between the RPE and physiological markers such as the heart rate (HR) has been strongly correlated (Scherr et al., 2013) and often demonstrated during passive estimation and perceptually regulated procedures (Garcin et al., 1998). Measurement of the RPE remains of interest because the scale also predicts oxygen uptake ( $VO_2$ ) (Wong et al., 2011). In cycling, the RPE is usually applied either as a means of estimating pre-selected power output or producing appropriate power output for a pre-selected RPE (Myles and Maclean, 1987). This method assumes that participants can adjust power output to match numerically anchored verbal expressions of effort.

The use of the RPE in the prescription of alternated power protocols during indoor spin cycling assumes a comparable relationship between indices of physiological mechanisms inclusive of the heart rate and heart rate reserve (HRR) usually measured during continuous laboratory tests. These relationships appear to follow identical trends for continuous and intermittent exercises with the same power output (Edwards et al., 1972). Yet, the influence of whole body CoM acceleration on these variables remains largely unexplored in spin cycling, particularly in the triathlete population.

An objective in cycling is to eliminate all unwanted postural movement and reduce unused force. This unwanted force measurement is possible using accelerometers. Advances in wearable technology have made the accelerometer the device of choice for measuring frequency, duration and intensity of activity (Sallis and Saelens, 2000) as they provide an objective assessment of energy expenditure (Crouter et al., 2018). The accelerometer presents as a tool to provide evidence for coaches and triathletes to monitor the trunk position via accelerations to the whole body CoM whilst performing varied spin

cycling protocols during training and competition. Consequently, quantification of CoM acceleration can be detected. Therefore, the aims of this pilot study were to (1) identify any differences between two commonly used triathlete bicycle racing positions (aerodynamic and drops), which may cause variations to the whole body CoM; and (2) investigate the association between the whole body CoM, RPE, MET and HRR during steady state indoor spin cycling. Quantifying these variables could provide a better understanding of the functional significance of the whole body CoM in order for new insights into postural control mechanisms.

## Methods

This observational pilot study recorded accelerations to the whole body CoM in recreational triathletes during a characteristic spin cycling training session which was conducted in the triathlete's natural training environment.

### Participants

Four triathletes (3 males, 1 female) (mean  $\pm$  standard deviation [SD];  $31 \pm 8.8$  years,  $77 \pm 8.5$  kg,  $178 \pm 0.59$  cm,  $8.7 \pm 3.71$  weekly training hours) were recruited through contact with the local triathlon club (Greensborough Triathlon Club). All participants were asymptomatic of illness and free from any acute or chronic injury, as established by the American College of Sports Medicine (2008) participant activity readiness questionnaire (PAR-Q). To be included participants had to be cycling at least 50 km a week and competing in triathlon for a minimum of 12 months. Participants were excluded if they were sensitive to double sided tape or had an injury that might affect their riding kinematics. Written informed consent was provided, which was approved by Charles Darwin University Ethics Committee (HREC19028).

### Procedures

The magnitude of whole body CoM acceleration was continuously measured by a single triaxial accelerometer (52 x 30 x 12 mm, mass 23 g; resolution 16-bit, full-scale range 16 g, sampling at 100 Hz: SABEL Labs, Darwin, Australia) (James et al., 2011) which was attached between the L5 and S1 spinous process (Kavanagh et al., 2006) using double sided elastic adhesive tape (Medtronic Australasia Pty Ltd, Macquarie, NSW). This position was selected as it

is the closest external point to the whole body CoM and the lowest point where a single measuring device can monitor left and right lower limb kinematic data (Winter et al., 2016). Each participant performed one indoor 20 min cycle which consisted of six different power output conditions (Watts) and two body positions (DP = drops, AO = aerodynamic) (Table 1).

The DP position is described as hands placed onto the bicycle drop bars with slight elbow flexion and trunk inclination. The AO position is described as using tri (aerodynamic) – bars with elbows on triathlon bars with a greater horizontal torso than the DP position (Figure 1).

Participants completed the cycling conditions at their customary training location. The room temperature was standardised and controlled at 19°C using a climate control system. Cycling was conducted in the evening with participants cooled with two ventilated fans and permitted 250 ml of water. For the bicycle set up, participants were asked to configure the handlebar and saddle position to their preferred settings on a Schwinn Carbon Blue bicycle (Schwinn Bicycles, Dorel Industries, Inc, Washington, USA) which consisted of a belt drive (122 x 109 x 51 cm). Strap-in pedals were used to replicate a typical training setting in order to limit potential confounding effects. Power output was displayed via the digital display unit on the bicycle which was maintained using visual feedback. When a change in power was required (i.e. at three minute epochs), participants were verbally instructed to adjust gear ratios that would sustain the level of intensity required. To ensure familiarity, power output was based on a typical training session. Participants were required to refrain from any strenuous exercise for 24 hours before testing. Prior to commencement, participants performed a self-selected 2 min warm-up at their freely chosen power output. Participants were asked to verbally signify exertion using the Borg 6-20 RPE scale (Borg, 1998) at the end of each epoch. Heart rate reserve (HRR) was estimated by extrapolation of the RPE by 10 to obtain individual beats per minutes (Borg, 1974). All participants had previous experience of RPE perceptual scaling. The accelerometer was controlled wirelessly by the principal author via a standard Hewlett Packard PC using a comprehensive MATLAB

Toolkit. This allowed for control of multiple accelerometers providing no restrictions during data capture. Data were subsequently downloaded from the accelerometers using a SABEL Sense software program (SABEL Sense 1.2\_x64, SABEL Labs) via a CSV file. Prior to cycling, the accelerometer was calibrated (Lee et al., 2018). The accelerometer was synchronised at the start and the end of each three minute epoch where the longitudinal (vertical), mediolateral and anteroposterior axis aligned with X, Y and Z, respectively. Data were recorded continuously.

### Data analysis

Cycling conditions and corresponding power changes were identified in the raw accelerometer data. Longitudinal acceleration (CoMLN =  $\sqrt{X^2}$ ) was used to identify a change in posture and was identified when the acceleration magnitude began increasing towards its largest peak. Mediolateral acceleration (CoMML =  $\sqrt{Y^2}$ ) was used to identify pedal strokes. For each epoch, whole body CoM acceleration magnitude (CoMtotal =  $\sqrt{X^2 + Y^2 + Z^2}$ ) was then hand scored to obtain a 60 s average in order to attain a true reflection of steady state cycling. To obtain the MET compendium codes to be included in our study, researchers from this paper reviewed the 2011 Compendium of Physical activities (Ainsworth et al., 2011) to select the description which best matched the activities. The MET values were subsequently classified as: (1) conditioning exercise bicycling, stationary, 8.8 MET (101–160 W, vigorous effort); (2) conditioning exercise bicycling, stationary, 11 MET (161–200 W, vigorous effort); (3) conditioning exercise bicycling, stationary, 14.0 MET (201–270 W, very vigorous effort). To convert mph to kilometres per hour (km/h) a conversion factor of 1.609344 was applied to ensure consistency with the digital display screen on the Schwinn bicycle. To obtain a 60 s average, the caloric cost was calculated according to Equation 1:

$$MET = 3.5 \times w (kg) / 200$$

#### Equation 1

where MET is multiplied by oxygen uptake ( $VO_2$ ) of approximately  $3.5 \text{ O}_2 \text{ kg}^{-1} \text{ min}^{-1} \text{ ml/kg/min}^{-1}$  multiplied by weight in kilograms.

### Statistical Analysis

Prior to any inferential statistical analyses,

descriptive statistics were checked for normality using quantile–quantile plots. Statistical analysis was completed using Analyse-it (Leeds, United Kingdom, version 4.92). Results correspond to the mean  $\pm$  SD. The analysed variables were: (1) mean value of whole body CoM acceleration magnitude; (2) mean value of the MET and caloric cost; (3) mean value of the RPE; and (4) the contribution of both drops and aerodynamic position to all analysed variables. Values were tested using 95% confidence intervals (CI) and threshold values classified as 0.1–0.3 (small), >0.3–0.5 (moderate), >0.5–0.7 (large), >0.7–0.9 (very large) and >0.9 (extremely large) (Hopkins et al., 2009). A paired-sample *t* test assessed differences in variables with an alpha ( $\alpha$ ) level set at 5% ( $p < 0.05$ ). Two SD represents the expected change in whole body CoM acceleration given a  $2 \pm$  SD change in power, or otherwise the change from a typically low ( $-1$  SD). The Pearson's product-moment correlation coefficient (*r*) measured the strength of the relationship between the dependent variable of the MET and both the whole body CoM acceleration and distinct magnitudes (i.e., longitudinal, CoMLN) along with the RPE and HRR.

### Results

All participants completed the six cycling conditions. Effect sizes revealed mostly small differences in both longitudinal and anteroposterior acceleration ( $< 0.2$ ) throughout all conditions. The exception was found from epoch 8–11 min to 11–14 min performed at 152–205 W with extremely large effects ( $> 0.9$ ) for all acceleration magnitudes as well as the RPE and HRR, the latter with greater variability given the disparity in standard deviation (Table 2).

There was a large effect in longitudinal acceleration as power increased to 206–246 W (epoch 14–17 min) with a slight decrease in mediolateral and anteroposterior acceleration compared to 11–14 min. There were no significant differences between whole body CoM acceleration and MET ( $t = -1.50, p \leq 0.05$ ) and RPE ( $t = -1.50, p \leq 0.05$ ). In relation to the RPE, when participants reported an RPE 13 they were cycling at higher power output ( $>206$ –246 W) which resulted in extremely large effects. Switching from the drops to the aero hand position (2–5 min to 5–8 min) caused a decrease in triaxial CoM accelerations

with an inverse relationship seen in the RPE and HRR (difference between means 5 and 27.5, respectively). When participants changed from the aero to the drops hand position (minutes 14 through to minutes 20) an extremely large effect occurred in the anteroposterior direction ( $> 0.9$ ) (Figure 2).

There were strong correlations between whole body CoM acceleration, MET and RPE ( $r = 0.865$ ,  $p \leq 0.0001$ ) that yielded the equation:

$$\text{MET} = 238.7 + 0.605 \text{ RPE} - 4.191$$

whole body CoM acceleration  $\text{m/s}^2$

Replacing MET for caloric cost resulted in an equally strong correlation ( $r = 0.853$ ,  $p \leq 0.0001$ ) (Figure 3).

## Discussion

This study investigated the relationship between whole body CoM acceleration, energy expenditure determined by the metabolic equivalent (MET), the rating of perceived exertion (RPE) and heart rate reserve (HRR) during a typical triathlete indoor spin cycling training session. Power and the body position were varied to identify differences in acceleration patterns. The major experimental finding was that of extremely large effects pertaining to mediolateral acceleration during six minutes of steady-state cycling at 152-205 W as well as concurrent increases in longitudinal and anteroposterior CoM acceleration along with the RPE and the MET (Table 2). Additionally, switching from an aero to drops hand position induced extremely large effects in anteroposterior acceleration. Together, these combined variables

may affect steady state triathlete spin cycling performance of longer duration.

Variations in mediolateral acceleration were most salient at epochs 8–11 min to 11–14 min which corresponded to constant power output of  $>152$ -205 W. Although this period of steady state cycling implies a sustainable, moderate effort which is below the lactate threshold (Ghosh, 2004), the effect of individual training status should be considered. In this instance, individual training volume and hours dedicated to cycling could have influenced results. It is also possible that gender-related differences in the pelvic tilt could arise, in part, from the bicycle set-up that was used. In particular, participants were asked to configure the handlebar and saddle position to their preferred settings, meaning that the change in the body position (i.e., from drops to aero) may not have been correctly configured, which may have required the female participant to rotate further forward when moving to the drops position. Along this line, gender-related differences in pelvic orientation when changing the body position have been observed with females exhibiting a greater anterior tilt in the drops position compared with males (Sauer et al., 2008). This could explain the high dispersion of standard deviation observed in the current study, noticeable in both longitudinal and anteroposterior magnitudes relative to increases in power. As the absolute magnitude of CoM acceleration would likely change between genders, further research is warranted to assess changes in performance settings.

**Table 1**

*Protocol to assess body CoM acceleration, RPE, MET, HRR at 6 spinning conditions.*


*FCPO = freely chosen power output; DP = drops position; AO = aerodynamic position*

Epoch (minutes)	2-5 min	5-8 min	8-11 min	11-14 min	14-17 min	17-20 min
Power (Watts)	FCPO	<150	>152-205	>152-205	>206-246	FCPO
Position	DP	AO	AO	AO	AO	DP

**Table 2**  
*Mean ± SD and effect size for CoM acceleration magnitude (m/s<sup>2</sup>), RPE, MET and HRR, stratified by epoch and condition*

	<b>**Epoch 2-5 min (FCPO)</b>		<b>*Epoch 5-8 min &lt;150 w</b>		<b>Threshold Effect Size (ES)</b>
<b>Direction</b>	<b>Mean</b>	<b>± SD</b>	<b>Mean</b>	<b>± SD</b>	<b>Descriptor</b>
X	25.19	3.28	24.50	3.31	-0.2 (small)
Y	1.35	1.36	0.74	0.62	-0.5 (moderate)
Z	12.83	6.39	12.06	5.51	-0.1 (small)
RPE	6.75	0.50	11.75	1.89	>0.9 (extremely large)
HRR (bpm)	67.5	5.00	95.00	17.32	
MET	6.00		6.00		
	<b>**Epoch 5-8 mins &lt;150 w</b>		<b>*Epoch 8-11 mins 152-205 w</b>		
X	24.50	3.31	24.74	4.10	<0.1 (small)
Y	0.74	0.62	0.77	0.59	<0.1 (small)
Z	12.06	5.51	14.34	8.46	<0.1 (small)
RPE	11.75	1.89	11.75	1.89	<0.1 (small)
HRR (bpm)	95.00	17.32	117.5	18.93	
MET	6.00		8.00		
	<b>*Epoch 8-11 mins 152-205 w</b>		<b>*Epoch 11-14 mins 152-205 w</b>		
X	24.74	4.10	24.90	4.07	>0.9 (extremely large)
Y	0.77	0.59	1.24	1.05	>0.9 (extremely large)
Z	14.34	8.46	14.39	8.17	>0.9 (extremely large)
RPE	11.75	1.89	12.25	1.50	>0.9 (extremely large)
HRR (bpm)	117.5	18.93	122.5	15.00	
MET	8.00		8.00		
	<b>*Epoch 11-14 mins 152-205 w</b>		<b>*Epoch 14-17 mins &lt;206-246 w</b>		
X	24.90	4.07	25.42	4.29	>0.9 (extremely large)
Y	1.24	1.05	0.53	0.40	>0.9 (extremely large)
Z	14.39	8.17	12.86	8.43	>0.9 (extremely large)
RPE	12.25	1.50	12.50	1.00	>0.9 (extremely large)
HRR (bpm)	122.5	15.00	125.00	10.00	
MET	8.00		10.00		
	<b>*Epoch 14-17 mins &lt;206-246 w</b>		<b>**Epoch 17-20 FCPC</b>		
X	25.42	4.29	25.14	3.68	<0.1 (small)
Y	0.53	0.40	0.50	0.48	<0.1 (small)
Z	12.86	8.43	14.20	7.15	>0.9 (extremely large)
RPE	12.50	1.00	12.50	1.73	<0.1 (small)
HRR (bpm)	125.00	10.00	125.00	17.32	
MET	10.00		8.00		

*\*Epochs 5-8min to 14-17min performed in an aerodynamic position;  
 \*\* performed in a drops position. bpm = beats per minute*




**Figure 1**

(a)

(b)

(a) Participant cycling in the aerodynamic position; (b) Schwinn Carbon bicycle used in the study




**Figure 2**

Comparison of the drops against the aerodynamic position at four cycling conditions for body CoM acceleration ( $m/s^2$ ), MET, caloric cost (CC) and RPE

† Significant at  $<0.005$





**Figure 3**

Total group average per epoch and power showing CoM acceleration ( $m/s^2$ ), MET, CC = caloric cost; and RPE

In the current study, participants exhibited small effects in mediolateral acceleration when moving from the aero to drops hand position towards the end of the protocol. Regarding this change, as power went from high values of 206-246 W in aero cycling to freely chosen power in the drops, it is feasible that participants deliberately selected a lower power range as the protocol neared conclusion to anchor the perceptual exertion range (Figures 2 and 3). If, instead, the differences were attributable to anatomical factors, these did not seem to be captured by simple measures of acceleration magnitude. However, anteroposterior trunk acceleration could explain intraindividual differences in CoM motion because of the high magnitudes observed during the final minutes of cycling. This potentially represents the gross motion of the participant's CoM around the anteroposterior axis. Equally, the possibility of poor trunk strength cannot be ruled out. A low level of core muscle strength can cause greater upper movement (Rannama et al., 2017). This would suggest a destabilising effect of the trunk and inefficient movement patterns leading to increases in whole body CoM acceleration.

The data in the current study suggest that triathletes increased whole body CoM acceleration magnitude during constant power as evidenced through minutes 8-11 to 14-17, respectively. The absolute increase in whole body CoM acceleration magnitudes combined with the RPE, the MET and HRR suggest an increase in participants' work. Thus, the power transfer from the trunk to the lower body would increase with power due to an increase in hip reaction force, a result consistent with that found by others (Costes et al., 2015; Mestdagh, 1998). This warrants the need to quantify postural deviations to determine the influence of stability as it is currently unknown whether consistent trends would emerge for extended duration of spin cycling. Along this line, a structured strength and conditioning program with the focus on core trunk and hip strength may also reduce these effects.

In order to address whether whole body CoM acceleration was affected by changes in the body position and power, values were obtained at freely chosen power output at the commencement and conclusion of the protocol. When the stabilising role of the trunk is considered, reduced

trunk lean can occur early as muscle fatigue can precede changes in trunk lean and trunk range of motion (Dingwell et al., 2008). Similarly, a significant compensatory effect on movement kinematics without alterations in pedalling would occur (Abt et al., 2007). Thus, it is possible that a similar attributable effect occurs through whole body acceleration of the CoM which may contribute to greater acceleration magnitudes and the RPE with a corresponding reduction in power.

The concomitant increases in the MET, RPE and HRR put increased emphasis on the blended physiological and temporal kinematics relationship. This is comparable to that reported by Zinoubi et al. (2018) and supports the premise that the RPE is associated with physiological variables (Kasimay Çakir et al., 2012). Although this increase was consistent until <206–246 W at which point the RPE levelled to a freely chosen cadence at minutes 17–20, an increase in the workload would theoretically lead to higher activation of the majority of muscles and therefore CoM trunk acceleration. As the RPE increases with fatigue (Rietjens et al., 2005), taken together with an increase in whole body CoM acceleration and power, it is plausible that excessive accelerations of the CoM could provide a technique to investigate postural changes during frequent spin cycle training sessions. Future research should determine variables which reflect muscular, whole body CoM acceleration and RPE status. Nevertheless, it is important to note that the RPE is affected not only by physical stress, but also psychological stress (Kumar et al., 2013), and therefore, the changes seen in HRR in the present study may also provide information not only on whether the athlete is physically able to train, but their mental readiness (i.e., holistic psychophysiological athlete preparedness). Furthermore, at submaximal levels exertion could be due to neurological alterations owing to reduced force despite increases in perception of effort (Noakes and St Clair Gibson, 2004). However, care must be taken when interpreting these results as data from the current study show the effect of a single cycling session with no direct measure of neuromuscular fatigue markers obtained.

This study provides information relevant for plausible associations between triathlete steady state spin cycling, whole body CoM

acceleration, HRR, RPE and the MET. Although observable trunk movement can be indicative of an incorrect saddle position, it would be anticipated that increases in the RPE and the MET would ensue due to incremental increases in power. Notably, if participants increased power a concurrent increase in muscular force would occur which would conceivably increase the amount of mechanical work. Although insufficient to completely explain the increase in the RPE or to indicate its effects on whole body CoM acceleration, spin cycling performance and postural control, the use of musculoskeletal modelling to simulate pedalling and body acceleration during higher intensities may provide greater insights.

Whereas Galy et al. (2003) required cycling to fatigue, compared to the moderately short and non-fatiguing protocol used in this pilot study, it is challenging to draw conclusions. The use of steady state spin cycling negated participants performing a fatiguing bout of cycling, as might be experienced in a triathlon race. Whilst diligence must be applied to assess how kinematics might change with greater levels of intensity, this information, along with systematic evaluation of triaxial CoM acceleration distributions, could prove useful for training interventions.

### **Practical implications**

The findings of this study have practical implications for coaches and triathletes. The quantification of energy expenditure based on accelerometric data in triathletes during a representative training session could be a useful and cost-effective metric to use in a real-time setting. This will ensure coaches and athletes can monitor and adjust whole body trunk CoM motion in accordance with the RPE, HRR and the MET efficiently throughout training.

### **Conclusions**

An accelerometer, in combination with the RPE, the MET and HRR, may provide a practical, non-invasive, accessible and economically valid tool to monitor whole body CoM acceleration in triathletes. The use of a simple and highly accessible accelerometer tool means coaches and athletes can adopt the methods used within the study at minimal expense and without extensive training. This pilot study sets the foundation for further evaluation of

a sensor-based approach to trunk CoM acceleration in triathlon cycling in order to take

the measures reported here towards greater reductions in trunk accelerations.

## Acknowledgements

The authors would like to thank the triathletes from Greensborough Triathlon Club who volunteered for this study.

## References

- Abt, J., Smoliga, J., Brick, M., Jolly, J.T., Lephart, S.M., & Fu, F.H. (2007). Relationship between cycling mechanics and core stability. *Journal of Strength and Conditioning*, 21, 1300-1304.
- Ainsworth, B.E., Haskell, W.L., Whitt, M.C., Irwin, M.L., Swartz, A.M., Strath, S.J., O'Brien, W.L., Bassett, D.R. Jr., Schmitz, K.H., Emplaincourt, P.O., Jacobs, D.R. Jr., & Leon, A.S. (2011). Compendium of Physical Activities: a second update of codes and MET values. *Medicine in Science & Sports Exercise*, 43, 1575-81. doi: 10.1097/00005768-200009001-00009
- American College of Sports Medicine (2008). *ACSM's guidelines for exercise testing and prescription*. Lippincott Williams & Wilkins.
- Borg, G.A. (1998). *Borg's perceived exertion and pain scales*. Human Kinetics.
- Borg, G.A., & Noble, B.J. (1974). Perceived exertion. *Exercise and Sports Science Reviews*, 2, 131-53.
- Costes, A., Turpin, N., & Villeger, D.A. (2015). Reduction of the saddle vertical force triggers the sit-stand transition in cycling. *Journal of Biomechanics*, 12, 2998-3003.
- Crouter, S.E., Flynn Oody, J., & Bassett, D.R. Jr. (2018). Estimating physical activity in youth using an ankle accelerometer. *Journal of Sports Science*, 36(19), 2265-2271. doi: 10.1080/02640414.2018.1449091.
- de Melo Dos Santos, R., Costa, F., Saraiva, T.S, & Callegari, B. (2017). Muscle fatigue in participants of indoor cycling. *Muscles, Ligaments and Tendons Journal*, 7(1), 173-179. doi: 10.11138/mltj/2017.7.1.173
- Dingwell, J.B., Joubert, J.E., Diefenthaler, F., & Trinity, J.D. (2008). Changes in muscle activity and kinematics of highly trained cyclists during fatigue. *IEEE Trans Biomed Eng*, 55, 2666-2674.
- Edwards, R.H.T., Melcher, A., Hesser, C.M., Wigertz, O., Ekelund, L.G. (1972). Physiological correlates of perceived exertion in continuous and intermittent exercise with the same average power output. *European Journal of Clinical Investigation*, 2(2), 108-14. doi: 10.1111/j.1365-2362.1972.tb00578.x
- Galy, O., Manetta, J., & Coste, O. (2003). Maximal oxygen uptake and power of lower limbs during a competitive season in triathletes. *Scandinavian Journal of Medicine and Science in Sports*, 13, 185-193.
- Garcin M., Vautier, J.F., Vandewalle, H., Wolff, M., & Monod, H. (1998). Ratings of perceived exertion (RPE) during cycling exercises at constant power output. *Ergonomics*, 1998; 41: 1500-1509.
- Ghosh, A.K. (2004). Anaerobic threshold: its concept and role in endurance sport. *Malaysian Journal of Medicine and Science*, 11, 24-36.
- Gregor, R.J. (2000). *Biomechanics of cycling*. Philadelphia: Williams & Wilkins.
- Hopkins, W., Marshall, S., Batterham, A., & Hanin, J. (2009). Progressive statistics for studies in sports medicine and exercise science. *Medicine and Science in Sports and Exercise*, 41(1), 3-13. doi: 10.1249/MSS.0b013e31818cb278.
- James, D.A., Leadbetter, R.I., Neeli M.R, Burkett, B.J., Thiel, D.V., & Lee, J.B. (2011). An integrated swimming monitoring system for the biomechanical analysis of swimming strokes. *Sports Technology*, 4, 141-150.
- Kasimary Çakir Ö., Cakir, B., Erzençin, O.U., Kurtel, H. (2012). Rating of perceived exertion in cycle ergometer: effect of maximal capacity difference. *Turkish Journal of Medical Sciences*, 42(5), 906-912.
- Kavanagh, J., Morrison, S., James, D.A., & Barrett, R. (2006). Reliability of segmental accelerations measured using a new wireless gait analysis system. *Journal of Biomechanics*, 39(15), 13. doi: 10.1016/j.jbiomech.2005.09.012.
- Kumar, Y., Agarwal, V., & Gautam, S. (2013). Heart rate variability during examination stress in medical students. *International Journal of Physiology*, 1(1), 83-86.
- Lee, J.B., Wheeler, K., & James, D.A. (2018). *Wearable sensors in sport: a practical guide to usage and implementation*. Springer.

- McDaniel, J., Subudhi, A., & Martin, J.C. (2005). Torso stabilization reduces the metabolic cost of producing cycling power. *Canadian Journal of Applied Physiology*, 30(4), 433–441. doi:10.1139/h05-132
- Macdonald, J.H., Farina, D., & Marcora, S.M. (2008). Response of electromyographic variables during incremental and fatiguing cycling. *Medicine and Science in Sport and Exercise*, 40(2), 335–344. doi: 10.1249/mss.0b013e31815b491e
- Mestdagh, Vey de K. (1998). Personal perspective: in search of an optimum cycling posture. *Applied Ergonomics*, 29(5), 325–334.
- Myles, W.S., & Maclean, D. (1987). A comparison of response and production protocols for assessing perceived exertion. *European Journal of Applied Physiology*, 55(6), 585–7. doi: 10.1007/BF00423201
- Noakes, T. D., & St Clair Gibson, A. (2004). Logical limitations to the “catastrophe” models of fatigue during exercise in humans. *British Journal of Sports Medicine*, 38(5), 648–9. doi: 10.1136/bjism
- Padulo, J., Di Capua, R., & Viggiano, D. (2012). Pedaling time variability is increased in dropped riding position. *European Journal of Applied Physiology*, 112, 3161–5.
- Pollock, M.L., Gaesser, G.A., Butcher, J.D., Després, J.P., Dishman, R.K., Franklin, B.A., & Garber, C.E. (1998). The recommended quantity and quality of exercise for developing and maintaining fitness in healthy adults. *Medicine and Science in Sports and Exercise*, 30(6), 975–91.
- Rannama, I., Pedak, K., Bazanov, B., & Port, K. (2017). Cycling specific postural stability during incremental exercise: The relationship with cyclists functional movement screen score. *Journal of Human Sport and Exercise*, 12, 83–95.
- Rietjens, G.J., Kuipers, H., Adam, J.J., Saris, W.H., van Breda, E., van Hamont, D., & Keizer, H.A. (2005). Physiological, biochemical and psychological markers of strenuous training-induced fatigue. *International Journal of Sports Medicine*, 26(1), 16–26. doi: 10.1055/s-2004-817914
- Sallis, J.F., & Saelens, B.E. (2000). Assessment of physical activity by self-report: status, limitations, and future directions. *Research Quarterly for Exercise and Sport*, 71 Suppl S1–S14. doi: 10.1080/02701367.2000.11082780
- Sauer, J.L., Potter, J.J., Weissbarr, C.L., Ploeg, H.L., & Thelen, D.G (2008). Influence of gender, power, and hand position on pelvic motion during seated cycling. *Medicine and Science in Sports and Exercise*, 39(12), 2204–11. doi:10.1249/mss.0b013e3181568b66
- Scherr, J., Wolfarth, B., Christle, J.W., Pressler, A., Wagenpfeil, S., & Halle, M. (2013). Associations between Borg’s rating of perceived exertion and physiological measures of exercise intensity. *European Journal of Applied Physiology*, 113(1), 147–55. doi: 10.1007/s00421-012-2421-x
- Wiest, M.J., Diefenthaler, F., Mota, C.B., & Carpes, F.P. (2011). Changes in postural stability following strenuous running and cycling. *Journal of Physical Education in Sport*, 11(4), 406–413.
- Winter, S.C., Gordon, S.J., Lee, J.B., & Leadbetter, R.I. (2016). Validation of a single inertial sensor for measuring running kinematics overground during a prolonged run. *Journal of Fitness Research*, 5, 14–23
- Wong, D.P., Carling, C., Chaouachi, A., Dellal, A., Castagna, C., Chamari, K., & Behm, D.G. (2011). Estimation of oxygen uptake from heart rate and ratings of perceived exertion in young soccer players. *Journal of Strength and Conditioning Research*, 25(7), 1983–8. doi: 10.1519/JSC.0b013e3181e4ffe6
- Zinoubi, B., Zbidi, S., Vandewalle, H., Chamari, K., & Driss, T. (2018). Relationships between rating of perceived exertion, heart rate and blood lactate during continuous and alternated-intensity cycling exercises. *Biology of Sport*, 35(1), 29–37. <https://doi.org/10.5114/biolsport.2018.70749>

**Corresponding author:****Stuart Evans**

SABEL Labs, College of Health and Human Science,  
Charles Darwin University, Darwin, 0810 Australia.  
E-mail: [stuart@qsportstechnology.com](mailto:stuart@qsportstechnology.com)