

Correction to: Characterization of biofilm-forming capacity and resistance to sanitizers of a range of E.coli O26 pathotypes from clinical cases and cattle in Australia

Author

Lajhar, Salma A, Brownlie, Jeremy, Barlow, Robert

Published

2018

Journal Title

BMC Microbiology

Version

Version of Record (VoR)

DOI

[10.1186/s12866-018-1202-z](https://doi.org/10.1186/s12866-018-1202-z)

Rights statement

© The Author(s). 2018 Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Downloaded from

<http://hdl.handle.net/10072/388282>

Griffith Research Online

<https://research-repository.griffith.edu.au>

CORRECTION

Open Access



Correction to: Characterization of biofilm-forming capacity and resistance to sanitizers of a range of E.coli O26 pathotypes from clinical cases and cattle in Australia

Salma A. Lajhar^{1,2*}, Jeremy Brownlie¹ and Robert Barlow²

Correction

1. On page 4 of the original publication [1], the correct sentence should read:
Quantitative measurement was performed by dissolving CV stained pellicle in 4 ml of 85% ethanol and the OD₅₇₀ was measured using microplate reader at 200 µl per well.
2. Table 2 of this original publication contained some errors. The updated Table 2 is published in this correction article.

Received: 22 May 2018 Accepted: 12 June 2018

Published online: 19 June 2018

Reference

1. Lajhar SA, et al. Characterization of biofilm-forming capacity and resistance to sanitizers of a range of E.coli O26 pathotypes from clinical cases and cattle in Australia. *BMC Microbiol.* 2018;18:41. <https://doi.org/10.1186/s12866-018-1182-z>.

* Correspondence: salmaibsais.lajhar@griffithuni.edu.au

¹School of Environment and Science, Griffith University, Brisbane, QLD, Australia

²Present address: CSIRO Agriculture and Food, 39 Kessels Rd, Coopers Plains, Brisbane, QLD 4108, Australia



Table 2 Biofilm formation on polystyrene microtiter plates, stainless steel coupons, glass slides and pellicle formation at the air-liquid interface

Pathotypes	Isolates No.	Biofilm mass on polystyrene at incubation time of			Biofilm mass on ^b		Pellicle formation ^c
		24 h	48 h	72 h	SS	GS	
EHEC	EC1A	0.024 ± 0.004 ^a	0.076 ± 0.015	0.049 ± 0.007	1	1	0.061 ± 0.011
	EC1113B	0.029 ± 0.005	0.119 ± 0.022	0.191 ± 0.039	3	3	1.884 ± 0.259
	EC1643B	0.047 ± 0.006	0.251 ± 0.033	1.126 ± 0.153	3	3	3.595 ± 0.191
	EC1857	0.036 ± 0.006	0.063 ± 0.011	0.107 ± 0.016	3	3	0.856 ± 0.078
	EC217	0.054 ± 0.009	0.100 ± 0.015	0.043 ± 0.008	2	1	0.025 ± 0.009
	EC3455	0.066 ± 0.008	0.025 ± 0.005	0.060 ± 0.007	3	3	1.228 ± 0.145
	EC3522	0.060 ± 0.008	0.058 ± 0.010	0.086 ± 0.011	1	1	0.039 ± 0.034
	EC3547A	0.043 ± 0.003	0.236 ± 0.014	0.503 ± 0.030	2	2	0.839 ± 0.113
	EC3652B	0.056 ± 0.010	0.102 ± 0.014	0.104 ± 0.016	3	3	1.875 ± 0.103
	EC3659B	0.026 ± 0.009	0.058 ± 0.009	0.093 ± 0.014	3	3	1.652 ± 0.152
	EC3671A	0.016 ± 0.005	0.091 ± 0.010	0.125 ± 0.017	3	3	1.554 ± 0.167
	EC3738B	-0.010 ± 0.003	0.032 ± 0.013	0.093 ± 0.027	1	1	0.145 ± 0.030
	EC3743A	0.038 ± 0.006	0.087 ± 0.011	0.075 ± 0.010	3	3	1.218 ± 0.087
	EC4	0.023 ± 0.004	0.045 ± 0.009	0.024 ± 0.005	2	1	0.097 ± 0.023
	EC478B	0.086 ± 0.011	0.127 ± 0.011	0.133 ± 0.011	2	2	0.480 ± 0.061
	EC674	0.011 ± 0.005	0.064 ± 0.013	0.130 ± 0.023	3	3	1.660 ± 0.151
	EC7B	0.082 ± 0.014	0.109 ± 0.013	0.230 ± 0.052	3	3	1.397 ± 0.091
	EC4158QH1	0.047 ± 0.005	0.082 ± 0.015	0.069 ± 0.011	1	1	0.092 ± 0.042
	EC4159QH2	0.049 ± 0.007	0.089 ± 0.013	0.055 ± 0.007	2	1	0.035 ± 0.025
	EC4160QH3	0.098 ± 0.015	0.034 ± 0.008	0.044 ± 0.008	2	3	2.257 ± 0.099
	EC3213QH34	0.036 ± 0.006	0.072 ± 0.007	0.171 ± 0.017	3	3	1.079 ± 0.046
	EC4161QH4	0.075 ± 0.012	0.060 ± 0.010	0.061 ± 0.009	2	1	0.050 ± 0.021
	EC4162QH5	0.001 ± 0.009	0.088 ± 0.020	1.021 ± 0.154	3	3	2.882 ± 0.255
	EC4163QH6	0.017 ± 0.003	0.135 ± 0.030	1.469 ± 0.304	2	3	1.848 ± 0.127
EC4164QH7	0.053 ± 0.006	0.223 ± 0.025	0.245 ± 0.025	2	1	0.092 ± 0.034	
EC4165QH8	0.070 ± 0.008	0.127 ± 0.019	0.156 ± 0.026	1	1	0.065 ± 0.037	
EC4166QH9	0.061 ± 0.007	0.051 ± 0.010	0.070 ± 0.010	3	3	2.746 ± 0.163	
pEHEC	EC801	0.040 ± 0.005	0.119 ± 0.017	0.121 ± 0.020	2	1	0.150 ± 0.042
	EC3983A	0.033 ± 0.006	0.057 ± 0.011	0.074 ± 0.010	2	1	0.081 ± 0.024
	EC3989A	0.063 ± 0.008	0.638 ± 0.036	1.402 ± 0.203	3	3	3.859 ± 0.138
aEPEC	EC3435A	0.044 ± 0.009	0.041 ± 0.007	0.060 ± 0.007	0	1	0.104 ± 0.064
	EC3457	0.110 ± 0.020	0.134 ± 0.023	0.135 ± 0.014	1	1	-0.001 ± 0.030
	EC3610A	0.019 ± 0.003	0.017 ± 0.004	0.033 ± 0.005	1	0	0.018 ± 0.023
	EC3727A	0.037 ± 0.005	0.034 ± 0.007	0.119 ± 0.021	0	1	0.268 ± 0.051
	EC3735A	0.046 ± 0.011	0.017 ± 0.011	0.108 ± 0.019	0	1	0.135 ± 0.027
	EC3768A	0.035 ± 0.012	0.078 ± 0.017	0.206 ± 0.028	0	1	0.128 ± 0.020
	EC4013A	0.048 ± 0.018	0.027 ± 0.011	0.101 ± 0.014	0	1	0.177 ± 0.032
	EC4039A	0.006 ± 0.003	0.060 ± 0.009	0.048 ± 0.007	1	0	-0.005 ± 0.013
NTEC	EC3536B	0.052 ± 0.005	0.224 ± 0.021	0.475 ± 0.034	3	3	0.966 ± 0.158
	EC3946A	0.052 ± 0.006	0.313 ± 0.056	0.404 ± 0.034	3	3	0.587 ± 0.127
Negative control	Lb broth	0.080 ± 0.0009	0.078 ± 0.001	0.071 ± 0.0004	0	0	

^aValues are shown as mean of biofilm production ± standard error on polystyrene surfaces, SS: stainless steel, GS: glass slide at 25 °C. According to the biofilm mass quantified with crystal violet staining assay at 570 nm isolates were labelled as the following: low, medium and thick biofilm formers

^bVisible biofilms on stainless steel and glass slides and was scored as 0: no visible biofilm, scored on a scale from 1 to 3 to a thick biofilm at the air-liquid

^cThe presence and absence of visible pellicles biofilms was scored visually before staining with CV. Strains with visible pellicles at the air-liquid interface had $OD_{570} \geq 0.48$