

Permanent, long-acting and short-acting reversible contraceptive methods use among women in Bangladesh: an analysis of Bangladesh Demographic and Health Survey 2017–2018 data

Author

Sharif, Azaz Bin, Hasan, Md Tamzid, Naziat, Md Habib, Zerine, Tahmina, Kundu, Satyajit

Published

2023

Journal Title

BMJ Open

Version

Version of Record (VoR)

DOI

[10.1136/bmjopen-2023-073469](https://doi.org/10.1136/bmjopen-2023-073469)

Rights statement

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ. <http://creativecommons.org/licenses/by-nc/4.0/>





Downloaded from

<http://hdl.handle.net/10072/430111>

Griffith Research Online

<https://research-repository.griffith.edu.au>

BMJ Open Permanent, long-acting and short-acting reversible contraceptive methods use among women in Bangladesh: an analysis of Bangladesh Demographic and Health Survey 2017–2018 data

Azaz Bin Sharif ^{1,2}, Md Tamzid Hasan ², Md. Habib Naziat ^{2,3}, Tahmina Zerine ^{2,4}, Satyajit Kundu ^{1,5}

To cite: Sharif AB, Hasan MT, Naziat MH, *et al.* Permanent, long-acting and short-acting reversible contraceptive methods use among women in Bangladesh: an analysis of Bangladesh Demographic and Health Survey 2017–2018 data. *BMJ Open* 2023;**13**:e073469. doi:10.1136/bmjopen-2023-073469

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2023-073469>).

Received 06 March 2023
Accepted 26 June 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Satyajit Kundu;
satyajitnfs@gmail.com

ABSTRACT

Objectives This study aims to explore the factors associated with the permanent and long-acting reversible contraceptive (LARC) method use compared with short-acting reversible contraceptive (SARC) methods among Bangladeshi ever-married women aged 15–49 years.

Design Cross-sectional study.

Setting We used data from Bangladesh Demographic Health Survey (BDHS) 2017–2018.

Participants A total of 9669 Bangladeshi reproductive-aged women who gave information on contraception use were the study participants. A multilevel multinomial logistic regression model was employed where the SARC method was considered as the base category and the cluster was considered as level-2 factor.

Results Among the contraceptive users in Bangladeshi women, about 83.48% used the SARC method, while 11.34%, and 5.18% used permanent and LARC methods, respectively. Compared with SARC, women with no formal education and only primary education who were non-Muslims, and had parity of ≥ 3 had a higher likelihood of using both permanent and LARC methods. Women from the age group of 25–34 years (adjusted relative risk ratio (aRRR): 7.03, 95% CI: 4.17 to 11.85) and 35–49 years (aRRR: 12.53, 95% CI: 7.27 to 21.58) who were employed (aRRR: 1.19, 95% CI: 1.00 to 1.40), had media access (aRRR: 1.24, 95% CI: 1.03 to 1.49), gave birth in last 5 years (aRRR: 1.40, 95% CI: 1.11 to 1.76), whose contraception decision solely made by their husband (aRRR: 7.03, 95% CI: 5.15 to 9.60) and having high decision-making power (aRRR: 2.12, 95% CI: 1.62 to 2.77) were more likely to use permanent contraceptive methods. We observed that women from households with richer (aRRR: 0.65, 95% CI: 0.45 to 0.93) and richest (aRRR: 0.38, 95% CI: 0.23 to 0.63) wealth quintiles were less likely to use LARC methods.

Conclusions This study identified that women with no/less education, non-Muslims, and having parity of ≥ 3 were more likely to use both permanent and LARC methods than SARC methods. Targeted interventions could be developed and implemented to promote personalised contraceptive use.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This study used the most recent nationally representative data set; therefore, the study results could be generalisable across the country and other similar settings.
- ⇒ The cluster effect on the factors associated with the contraception method usage in Bangladesh was accounted for using a multilevel modelling approach, which is the proper analytical strategy for this type of hierarchical data.
- ⇒ We identified the factors associated with different types of modern contraceptive methods among Bangladeshi women.
- ⇒ Cross-sectional study design was one of the limitations which prevented us from drawing a causal association between the outcome and explanatory variables.
- ⇒ Another limitation was that a few important factors related to the respondents' health were not included in this study due to unavailability and incomplete information of those variables in the data set.

INTRODUCTION

Unintended pregnancy is a major cause of increasing population burden¹ and other maternal and neonatal difficulties in many developing countries like Bangladesh² and sub-Saharan Africa (SSA).³ Significant efforts have been made to lower fertility rates and unplanned pregnancies.⁴ Family planning (FP) is one of the most effective methods for preventing unintended pregnancies and reducing maternal and newborn mortality rates.⁵ One of the primary predictors of fertility transition and FP in emerging nations, contraceptive use is one of the proximal determinants of fertility.⁶ Despite the high efficiency of long-acting reversible contraceptives (LARCs), the rise in the use of contraceptives for FP has been largely driven by

short-acting reversible contraceptive (SARC) techniques.⁷ Compared with SARC methods, (LARCs) offer uninterrupted protection for 3–12 years against undesired and unexpected pregnancy.⁸ LARCs (including intrauterine devices (IUD) and hormonal implants) are more cost-efficient, oestrogen-free, extremely safe and highly effective regardless of user compliance after insertion than SARCs.^{9–11} Reports from Sweden¹² and the USA¹³ have shown that LARCs provide significant economic advantages over short-term solutions.

Globally, there were an estimated 358 000 maternal deaths in 2008, and developing nations continued to account for 99% (355 000) of these deaths; on the other hand, 87% (313 000) of all maternal deaths worldwide occurred in SSA and South Asia.¹⁴ To reduce the maternal mortality ratio and support families and nations in achieving their health-related goals, it is critical to provide affordable, acceptable and accessible contraceptive options.¹⁵ According to several indicators, the maternal mortality ratio may have decreased by more than 25% because of effective FP efforts.⁵ Bangladesh's contraceptive prevalence rate (CPR), which increased from 8% in 1975 to 62% in 2014, impressively increased sevenfold in less than 40 years.¹⁶

The CPR is crucial in determining how FP policies affect demographics.¹⁷ Therefore, it is essential to understand that fertility depends not only on the prevalence of contraceptive usage but also on the efficacy of that use and adherence of the users.¹⁷ Most maternal and newborn deaths can be avoided with established strategies to ensure that every pregnancy is desired with contemporary contraception and every delivery is safe.¹⁸ Compared with SARCs, LARCs provide better contraceptive effectiveness, lowering the need for abortion treatment.¹⁹ However, everyone may not prefer LARC, and factors other than effectiveness in preventing pregnancy are also important in making contraceptive decisions. While expanding access to LARC for those who choose to use such methods is important, FP providers should disclose women's true contraceptive preferences. In addition, educational materials should include information on all contraceptive method options, and they should be effectively used.

Using contraceptive methods has increased globally. About 49% of reproductive aged women (15–49 years) were using any type of contraception, an increase from 42% in 1990.²⁰ Bangladesh's current contraceptive usage prevalence (62%)²¹ is higher than the global prevalence, and than that of other South Asian nations like India (58%),²² Nepal (50%),²³ Pakistan (35%)²⁴ and Afghanistan (23%).²⁵ When looking at the LARCs, not more than 15% of women use LARCs globally.²⁶ The prevalence of LARC usage varies widely throughout Europe, for instance; although it is 2.9% in Poland, it is 16% in France, which is greater than the global percentage of women who use LARCs.²⁷ The USA has historically lagged well behind Europe,²⁸ while Asia and Latin America showed a high prevalence of using LARC procedures.²⁹

Higher education of both parents, having more than three children, women having used LARC in the past and women's strong knowledge and favourable attitudes towards LARC were revealed to be major drivers of usage in a study in Nigeria.³⁰ According to a previous study, women who select permanent contraception may differ considerably from those who choose SARC based on age or socioeconomic status. Superior socioeconomic position may be linked to more frequent use of permanent contraception and LARC.⁷ Women's autonomy is crucial to improve health-related outcomes like lower total fertility,³¹ and studies conducted in SSA have shown that women's autonomy impacts modern contraception use.³²

Bangladesh is now the fifth most populated nation in Asia and the eighth most populous country overall, with an estimated 165.16 million people.³³ The government of Bangladesh has decentralised healthcare facilities to ensure better access to maternity and FP services among the rural community and to promote LARC and permanent contraceptive methods more strongly towards the population in addition to all modern contraceptive methods, in an effort to lessen the burden of the country's population.³⁴ Given the significance of permanent methods to lower fertility rates and LARCs to ensure high efficacy rates, it is essential to comprehend the factors that influence their use among sexually active women to improve reproductive health outcomes and increase the uptake of permanent and LARC methods.^{4 35} Although there is a need for a better understanding of the factors associated with using permanent and LARC methods, it has not been given much attention in the Bangladeshi context. Therefore, we aimed to identify the factors associated with permanent and LARC methods used among women in Bangladesh using the most recent Bangladesh Demographic Health Survey (BDHS) 2017–2018 data. By expanding access to, and distribution of permanent and LARC methods especially among those who prefer those methods, we anticipate that the results of this study might help Bangladesh to considerably reduce unwanted pregnancies, maternal mortality and morbidity.

METHODS

Data source and study design

The current study used nationally representative cross-sectional data from the BDHS 2017–2018. The survey included urban and rural households from all administrative divisions of Bangladesh. A two-stage cluster sampling was used to select samples. At the first stage, enumeration areas (EAs) were selected with probability proportional to sizes (675 in 2017–2018 BDHS). In the second stage, on average, 30 households were selected from each EA (cluster) using systematic sample selection. Information on the survey methodology, sampling design and sampling technique is detailed in the BDHS 2017–2018 survey report.²¹ According to the birth record (IR) file, 20 127 women aged between 15 and 49 were interviewed to gather information on




Figure 1 Flow chart of the participants selection from Bangladesh Demographic and Health Survey 2017–2018 data (IR file). EAs, enumeration areas; HHs, households; IR, individual record.

contraception use. Among them, 8282 participants who did not use contraception were excluded. In addition, 1958 cases who were using contraceptives other than modern methods were also excluded. After deleting 172 missing cases, the final analyses included a total of 9715 (unweighted) cases (weighted, N=9669) (figure 1).

Outcome measure

Types of contraceptive methods used by the reproductive-aged women in Bangladesh were the primary outcome variable of interest. Only the women using modern contraceptive methods were included in this study. We categorised the modern contraceptive methods into three categories: permanent contraceptive method (male and female sterilisation), LARC method (IUD and implants/norplants) and SARC method (condoms, injections, pills and emergency contraception).⁶ The outcome variable was coded as ‘1’, ‘2’ and ‘3’ for the use of permanent, LARC and SARC methods, respectively.

Explanatory variables

According to the guidance of reviewed literature^{4 36–43} and the availability of the variables in BDHS 2017–2018, several explanatory variables were included in this current analysis. They are the age of women, education of women, employment status of women, husband/partner’s education, religion, body mass index (BMI), media exposure, current breastfeeding status, parity, birth history in last 5 years, history of a terminated pregnancy, hearing information about FP from media, the decision for using contraception, knowledge of ovulatory cycle, women decision-making power, household wealth status, place of residence and administrative division. The mothers’ BMI was divided into four categories based on the WHO guidelines: underweight ($<18.5\text{ kg/m}^2$), normal ($18.5\text{--}24.9\text{ kg/m}^2$), overweight ($25.0\text{--}24.9\text{ kg/m}^2$) and obese ($\geq 30.0\text{ kg/m}^2$).⁴⁴ Mothers were asked how frequently they watch television, listen to the radio and read the newspaper to estimate their media exposure. Media exposure was reclassified as ‘less than once a week’

**Table 1** Explanatory variables along with their categories

Sl. no.	Variables	Categories
1	Age of women (years)	15–24, 25–34, 35–49
2	Education of women	No education, primary, secondary, higher
3	Women employment status	Working, not working
4	Husband's education	No education, primary, secondary, higher
5	Religion	Muslim, non-Muslim
6	Body mass index	Underweight, normal, overweight, obese
7	Media exposure	Never/less than a week, at least once a week
8	Currently breast feeding	Yes, no
9	Parity	None, 1–2, 3 or more
10	Birth in the last 5 years	No birth, gave birth
11	Ever had a terminated pregnancy	Yes, no
12	Heard about family planning from media	Yes, no
13	Decision for using contraception	Mainly respondents, mainly husband/partner, joint decision
14	Knowledge of ovulatory cycle	Yes, no
15	Women decision-making power	High, moderate, low power
16	Household wealth status	Poorest, poorer, middle, richer, richest
17	Place of residence	Rural, urban
18	Administrative division	Barisal, Chittagong, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur, Sylhet

or 'at least once a week'. Knowledge of the ovulatory cycle was assessed by asking whether women properly knew about the fertile period during their ovulatory cycle.²¹ The household wealth status was calculated based on the ownership of different household assets using principal component analysis (PCA).⁴⁵ Women's involvement in decisions about their healthcare, large household purchases and visits to family/relatives were used to estimate their decision-making power. These components were further condensed using PCA into a more generalised set of weights that rate 'decision-making power' on a scale of 0–100. Individual scores were then allocated to low, medium and high categories based on the standardised z-scores.³⁸ The explanatory variables and their categories have been listed in [table 1](#).

Statistical analysis

Both unweighted and weighted frequencies and percentages were calculated to show the background characteristics of the study participants. Considering the complex survey of BDHS, we used the 'svy' command in Stata V.16.0 (StataCorp, College Station, Texas, USA) for assigning the sample weight to adjust for clustering effect and sample stratification. Since the BDHS 2017–2018 used a two-stage stratified cluster sampling involving a hierarchical composition, a multilevel regression model would be an appropriate technique to consider that accounts for complex survey design-related variation.⁴⁶ Thus, we used the multilevel multinomial regression model to identify the factors associated with the contraception method, where all the explanatory variables were included in one model to adjust for the confounding effects and clusters (EAs) were considered as level-2 factor. Generalised structural equation modelling (GSEM) using the 'gsem' command in Stata was used to execute the analysis. The fixed effects of various explanatory variables and random effects at the cluster level were estimated. Multicollinearity among explanatory variables was checked using the variance inflation factor. The adjusted relative risk ratio (aRRR) and 95% CI were interpreted, and statistical significance was considered when a p value was less than 0.05.

Patient and public involvement

No patient involved.

RESULTS

Background characteristics of study participants

A total of 9669 (weighted) married women were included in our study. Among the contraceptive users in Bangladeshi women, about 83% (95% CI: 82.7% to 84.2%) of participants used the SARC method, while 11.3% (95% CI: 10.72% to 11.99%) and 5.2% (95% CI: 4.75% to 5.64%) of women used permanent and LARC methods, respectively. In this study, 40.9% of the women belong to the 25–34 age group and 32.5% of the women from 35 to 49 age group while only 26.61% were from the 15–24 age group. Of 9669 women, 14.66% had no formal education, and only 11.93% had higher education. About more than half (52%) of women were unemployed, and the majority of the participants were Muslim (89.2%). Majority 58% of women and 38% had one to two and three or more living children, respectively, and 47% gave birth in the last 5 years of the survey, where 4.2% have no children. Among the sample, 19.4% of women had a history of terminated pregnancy. About 83.5% of the study participants did not hear about FP from any media, and 66% of women were unaware of the ovulatory cycle. In decision-making for contraception use, 77.6% was taken jointly, while 6.5% was taken solely by the partner. About 31% of the participants had low women decision-making power, while only 8% of women had high power regarding decision-making. A large number of participants (70.4%) were from rural areas, while the highest number of

participants were from the Dhaka division (25.6%), and the lowest number was from the Sylhet division (5%). All the explanatory variables were significantly associated with the contraceptive method usage in the bivariate χ^2 analysis (all $p < 0.001$) (table 2).

Factors associated with contraceptive method use

Compared with women who had higher education, those having no education (aRRR: 2.93, 95% CI: 1.88 to 4.58) or only primary education (aRRR: 2.19, 95% CI: 1.44 to 3.33) were more likely to use permanent contraceptive methods compared with SARC method. Similarly, the use of the LARC method was also higher than the SARC method among women with no education (aRRR: 2.35, 95% CI: 1.32 to 4.18) or only primary education (aRRR: 1.77, 95% CI: 1.04 to 2.99). Compared with women aged 15–24 years, the older women aged 25–34 years (aRRR: 7.03, 95% CI: 4.17 to 11.85) and 35–49 years (aRRR: 12.53, 95% CI: 7.27 to 21.58) were more willing to use permanent method compared with SARC method. Unemployed women (aRRR: 1.19, 95% CI: 1.00 to 1.40) had a 19% more likelihood of using the permanent method than the SARC method. Compared with the Muslim women, non-Muslims were more likely to use both permanent (aRRR: 1.35, 95% CI: 1.04 to 1.76) and LARC (aRRR: 1.73, 95% CI: 1.27 to 2.35) methods than the SARC methods. Women having no/less exposure to media compared with those who were exposed at least once a week (aRRR: 1.24, 95% CI: 1.03 to 1.49) were 24% more willing to use the permanent method than the SARC methods (table 3).

When looking at the parity of the women, those having three or more children had a higher likelihood of using both permanent (aRRR: 3.10, 95% CI: 1.30 to 7.41) and LARC methods (aRRR: 3.65, 95% CI: 1.87 to 7.92) compared with SARC methods in comparison with those having no children. Women who gave no birth in the last 5 years were 40% (aRRR: 1.40, 95% CI: 1.11 to 1.76) more likely to use the permanent contraceptive methods than the SARC methods. When looking at the decision-making for contraceptive use, we found when the decision was only by husband/partner (aRRR: 7.03, 95% CI: 5.15 to 9.60) and joint decision (aRRR: 1.84, 95% CI: 1.47 to 2.30), the likelihood of using the permanent methods was 7.03 times and 1.84 times higher, respectively, compared with the decision taken by the women own. Women with high decision-making power were 2.12 times (aRRR: 2.12, 95% CI: 1.62 to 2.77) more likely to use permanent methods compared with those with low decision-making power. For women from richest households, the likelihood of using LARC methods was 62% lower (aRRR: 0.38, 95% CI: 0.23 to 0.63) than those from the poorest households (table 3).

DISCUSSION

This study found that the factors associated with permanent method use versus SARC include the age of women,

education of women, women's employment status, religion, BMI, media exposure, parity, birth in the last 5 years, the decision to use contraception and women's decision-making power while factors associated with the use of LARC versus SARC methods were the education of women, religion, parity, the decision to use contraception and household wealth status.

Among the contraceptive users, 83.48% of the participants used the SARC method and 11.34% and 5.18% used the permanent and LARC methods, respectively. SARC methods were among the most popular in Bangladesh, except for older women, among the modern methods.²¹ The key rationales for SARC use on a large scale were strategic behaviour change communication programmes involving the integration of three major communication channels—mass media, interpersonal and community channels—as well as the introduction of new brands of contraceptives, an effective commodity supply chain and continuing and strengthening domiciliary services.⁴⁷ Another study also identified that women who chose to use SARC over LARC significantly believed that LARC is not effective in preventing pregnancy.⁴⁸ Not everyone prefers LARC, and factors other than effectiveness in preventing pregnancy are also important in making contraceptive decisions. While expanding access to LARC for those who choose to use such methods is important, FP providers should disclose women's true contraceptive preferences. In addition, educational materials should include information on all contraceptive method options.

In line with other studies in North West Ethiopia⁴⁹ and Central Ethiopia,⁵⁰ the study indicated a significant relationship between age and the usage of permanent contraceptive methods among the increasingly aged women compared with the younger ones. The older women could have more children and were more motivated to restrict their pregnancies, which might be the most probable reason for this finding.⁴⁹ Also, the fact that couples generally do not begin using permanent FP strategies until they have the desired number of children, which generally coincides with older ages and larger female parities.⁵¹

Women with no formal education or less education were more likely to use both permanent and LARC methods than those with higher education, consistent with findings from Kenya,⁵² and Ethiopia.⁵³ Previous studies reported that the highest fertility rates are found in women with less education, and also they begin having children sooner. Therefore, they are more willing to attain their desired children earlier than the higher educated women.^{54 55} Hence, as they age, they may be more likely to use the permanent and LARC methods to prevent pregnancies, which might explain this finding. Besides, educated women might have better knowledge of the menstrual cycle and management of SARC methods, and hence they are more willing to use SARC methods than the permanent and LARC methods.

Our study showed a significant association between women's employment status and their use of the permanent contraceptive method. Unemployed women were

Table 2 Distribution of using contraception methods among women in Bangladesh by explanatory variables (N=9669)

Variables	Weighted n (%)	Contraceptive method (weighted n (%), row)		
		Permanent method	LARC method	SARC method
Overall % (95% CI)	9669	11.34(10.72 to 11.99)	5.18(4.75 to 5.64)	83.48(82.73 to 84.21)
Age of women				
15–24 years	2573 (26.61)	19 (0.74)	121 (4.69)	2433 (94.57)
25–34 years	3953 (40.88)	351 (8.89)	216 (5.47)	3386 (85.64)
35–49 years	3143 (32.51)	726 (23.09)	164 (5.22)	2254 (71.70)
Education of women				
No education	1417 (14.66)	353 (24.87)	100 (7.05)	965 (68.08)
Primary	3133 (32.41)	446 (14.23)	187 (5.96)	2500 (79.81)
Secondary	3965 (41.00)	243 (6.12)	190 (4.79)	3532 (89.09)
Higher	1153 (11.93)	55 (4.79)	24 (2.11)	1074 (93.10)
Women employment status				
Working	5014 (48.15)	629 (12.55)	295 (5.88)	4090 (81.57)
Not working	4656 (51.85)	467 (10.03)	206 (4.42)	3982 (85.54)
Husbands' education				
No education	2190 (22.65)	421 (19.24)	158 (7.20)	1611 (73.56)
Primary	3258 (33.70)	362 (11.11)	198 (6.07)	2698 (82.82)
Secondary	2727 (28.21)	220 (8.07)	107 (3.92)	2400 (88.00)
Higher	1493 (15.45)	93 (6.21)	38 (2.57)	1362 (91.22)
Religion				
Muslim	8629 (89.24)	953 (11.04)	423 (4.91)	7253 (84.05)
Non-Muslim	1040 (10.76)	144 (13.81)	77 (7.44)	819 (78.75)
Body mass index				
Underweight	1048 (10.84)	123 (11.70)	64 (6.06)	862 (82.25)
Normal	5592 (57.84)	586 (10.47)	304 (5.43)	4703 (84.10)
Overweight	2437 (25.21)	292 (12.00)	107 (4.39)	2038 (83.61)
Obese	591 (6.11)	96 (16.17)	27 (4.51)	469 (79.32)
Media exposure				
Never/less than a week	4178 (43.21)	492 (11.77)	273 (6.54)	3413 (81.69)
At least once a week	5491 (56.79)	605 (11.01)	227 (4.14)	4659 (84.85)
Currently breast feeding				
Yes	2494 (25.80)	108 (4.32)	150 (6.00)	2237 (89.68)
No	7175 (74.20)	988 (13.78)	351 (4.89)	5835 (81.33)
Parity				
None	405 (4.19)	7 (1.73)	8 (1.98)	390 (96.29)
1–2	5585 (57.76)	291 (5.20)	283 (5.06)	5011 (89.73)
3 or more	3679 (38.05)	799 (21.71)	218 (5.92)	2663 (72.37)
Birth in the last 5 years				
No birth	5104 (52.79)	852 (16.69)	243 (4.75)	4009 (78.56)
Gave birth	4565 (47.21)	244 (5.35)	258 (5.66)	4063 (88.99)
Ever had a terminated pregnancy				
Yes	1872 (19.36)	262 (13.97)	99 (5.27)	1512 (80.76)
No	7797 (80.64)	835 (10.71)	402 (5.16)	6560 (84.14)

Continued

Table 2 Continued

Variables	Weighted n (%)	Contraceptive method (weighted n (%), row)		
		Permanent method	LARC method	SARC method
Heard about family planning from media				
Yes	1594 (16.48)	139 (8.72)	73 (4.61)	1381 (86.67)
No	8076 (83.52)	957 (11.85)	427 (5.29)	6691 (82.85)
Decision for using contraception				
Mainly respondents	1539 (15.92)	132 (8.58)	91 (5.93)	1316 (85.50)
Mainly husband/partner	631 (6.52)	163 (25.81)	9 (1.46)	459 (72.72)
Joint decision	7499 (77.55)	801 (10.69)	400 (5.34)	6297 (83.97)
Knowledge of ovulatory cycle				
Yes	3290 (34.03)	334 (10.16)	165 (5.02)	2791 (84.83)
No	6378 (65.97)	762 (11.95)	336 (5.26)	5280 (82.79)
Women decision-making power				
High	784 (8.11)	152 (19.39)	41 (5.26)	591 (75.35)
Moderate	5893 (60.94)	677 (11.49)	310 (5.26)	4906 (83.25)
Low	2992 (30.95)	267 (8.93)	150 (5.00)	2575 (86.06)
Household wealth status				
Poorest	1966 (20.33)	253 (12.85)	157 (7.98)	1557 (79.17)
Poorer	1939 (20.06)	239 (12.33)	120 (6.20)	1581 (81.47)
Middle	1915 (19.81)	223 (11.65)	110 (5.72)	1583 (82.63)
Richer	2011 (20.80)	212 (10.55)	75 (3.71)	1725 (85.74)
Richest	1836 (18.99)	169 (9.21)	39 (2.15)	1627 (88.64)
Place of residence				
Rural	6811 (70.44)	817 (11.99)	387 (5.68)	5608 (82.33)
Urban	2858 (29.56)	280 (9.79)	114 (3.98)	2464 (86.23)
Administrative division				
Barisal	531 (5.49)	30 (5.73)	28 (5.31)	472 (88.96)
Chittagong	1493 (15.45)	143 (9.60)	63 (4.25)	1287 (86.15)
Dhaka	2473 (25.58)	283 (11.46)	115 (4.63)	2075 (83.91)
Khulna	1136 (11.75)	132 (11.62)	59 (5.16)	945 (83.22)
Mymensingh	803 (8.30)	69 (8.62)	39 (4.89)	694 (86.49)
Rajshahi	1439 (14.88)	184 (12.78)	83 (5.77)	1172 (81.45)
Rangpur	1314 (13.59)	191 (14.53)	95 (7.20)	1028 (78.27)
Sylhet	480 (4.97)	63 (13.14)	19 (3.97)	398 (82.88)

LARC, long-acting reversible contraceptive ; SARC, short-acting reversible contraceptive .

more likely to use the permanent method, aligned to a study from Northern Ethiopia.⁵⁶ The plausible explanation might be that unemployed women in Bangladesh might experience financial hardship and lack the resources to maintain additional children. In addition, permanent contraceptive techniques provide a long-term solution to preventing future pregnancies, which might be a more cost-effective strategy. We found that, non-Muslim (Hindus, Christians, Buddhists and others) women were more likely to use both permanent and LARC methods compared with Muslim women. This conclusion was in

alignment with studies in North East Ethiopia,⁵⁷ Nepal⁵⁸ and Malawi.⁵⁹ Possible explanations for our findings could be the apparent disagreement between religious teachings and the avoidance of contraception.⁵⁹ Besides, a further qualitative investigation is warranted to know the in-depth scenario of why Muslim women are less likely to use permanent and LARC methods in Bangladesh.

Furthermore, women were more likely to use the permanent and LARC methods if exposed to the media at least once a week. A significant association with media exposure indicates a positive influence of media on

Table 3 Multilevel multinomial regression showing the factors associated with contraceptive method use among Bangladeshi women

Variables	Permanent method versus SARC method		LARC method versus SARC method	
	aRRR	95% CI	aRRR	95% CI
Fixed-effect results				
Age of women				
15–24 years (RC)	1		1	
25–34 years	7.03***	4.17 to 11.85	0.97	0.73 to 1.28
35–49 years	12.53***	7.27 to 21.58	0.87	0.60 to 1.29
Education of women				
No education	2.93***	1.88 to 4.58	2.35**	1.32 to 4.18
Primary	2.19***	1.44 to 3.33	1.77*	1.04 to 2.99
Secondary	1.24	0.84 to 1.83	1.55	0.95 to 2.53
Higher (RC)	1		1	
Women employment status				
Working (RC)	1		1	
Not working	1.19*	1.00 to 1.40	1.05	0.85 to 1.30
Husbands' education				
No education	1.14	0.79 to 1.64	1.53	0.95 to 2.47
Primary	0.89	0.63 to 1.25	1.33	0.85 to 2.08
Secondary	0.88	0.63 to 1.22	1.10	0.71 to 1.69
Higher (RC)	1		1	
Religion				
Muslim (RC)	1		1	
Non-Muslim	1.35*	1.04 to 1.76	1.73***	1.27 to 2.35
Body mass index				
Underweight	1.56**	1.21 to 2.02	1.16	0.86 to 1.55
Normal (RC)	1		1	
Overweight	0.90	0.75 to 1.08	0.95	0.75 to 1.22
Obese	1.33*	1.00 to 1.76	1.19	0.76 to 1.85
Media exposure				
Never/less than a week (RC)	1		1	
At least once a week	1.24*	1.03 to 1.49	0.87	0.69 to 1.10
Currently breast feeding				
Yes	0.75	0.56 to 1.01	1.09	0.83 to 1.43
No (RC)	1		1	
Parity				
None (RC)	1		1	
1–2	1.05	0.45 to 2.48	1.07	0.49 to 2.57
3 or more	3.10*	1.30 to 7.41	3.65**	1.87 to 7.92
Birth in the last 5 years				
No birth	1.40**	1.11 to 1.76	0.89	0.68 to 1.17
Gave birth (RC)	1		1	
Ever had a terminated pregnancy				
Yes	1.08	0.91 to 1.28	0.94	0.73 to 1.20
No (RC)	1		1	
Heard about family planning from media				

Continued

Table 3 Continued

Variables	Permanent method versus SARC method		LARC method versus SARC method	
	aRRR	95% CI	aRRR	95% CI
Yes (RC)	1		1	
No	1.04	0.83 to 1.32	0.77	0.57 to 1.02
Decision for using contraception				
Mainly respondents (RC)	1		1	
Mainly husband/partner	7.03***	5.15 to 9.60	0.34**	0.16 to 0.68
Joint decision	1.84***	1.47 to 2.30	0.98	0.76 to 1.26
Knowledge of ovulatory cycle				
Yes (RC)	1		1	
No	1.02	0.87 to 1.20	0.97	0.79 to 1.19
Women decision-making power				
High	2.12***	1.62 to 2.77	1.20	0.82 to 1.74
Moderate	1.14	0.95 to 1.37	0.95	0.76 to 1.18
Low (RC)	1		1	
Household wealth status				
Poorest (RC)	1		1	
Poorer	0.88	0.69 to 1.12	0.88	0.66 to 1.16
Middle	0.82	0.63 to 1.07	0.81	0.59 to 1.12
Richer	0.81	0.61 to 1.09	0.65*	0.45 to 0.93
Richest	0.84	0.59 to 1.20	0.38***	0.23 to 0.63
Place of residence				
Rural	0.97	0.78 to 1.21	0.88	0.66 to 1.15
Urban (RC)	1		1	
Administrative division				
Barisal	0.48**	0.31 to 0.74	0.95	0.58 to 1.55
Chittagong	0.74	0.51 to 1.07	0.80	0.50 to 1.28
Dhaka (RC)	1		1	
Khulna	1.14	0.79 to 1.63	1.01	0.63 to 1.61
Mymensingh	0.62*	0.41 to 0.92	0.89	0.55 to 1.42
Rajshahi	1.21	0.84 to 1.72	1.16	0.74 to 1.82
Rangpur	1.11	0.77 to 1.61	1.08	0.68 to 1.72
Sylhet	1.04	0.71 to 1.52	0.64	0.38 to 1.07
Random-effect results				
Cluster random effects	0.577	0.420 to 0.792	0.670	0.456 to 0.985

*p<0.05, **p<0.01, ***p<0.001.

aRRR, adjusted relative risk ratio; LARC, long-acting reversible contraceptive ; RC, reference category; SARC, short-acting reversible contraceptive .

contraception choices. Our findings coincide with those from studies conducted in SSA,⁶⁰ and Western Ethiopia.⁶¹ Women exposed to media at least once a week may obtain information regarding FP since the media can describe various methods, their advantages and where they are accessible to women, improving women's usage of contraceptive methods.⁴³

Regarding parity, women with three or more children were more likely to use both permanent and LARC

methods. According to previous studies,^{462 63} using permanent and LARCs increases as children grow.⁶⁴ One explanation could be that multiparous women frequently get FP information and counselling on contraceptive usage throughout their pregnancy counselling or that women who have previously achieved their desired fertility may take permanent contraceptives or LARCs until menopause.^{4 64} Also, there was a higher tendency for women who had not given birth in the previous 5 years to use

permanent contraceptive methods. The fact behind this finding could be that women who had not given birth in the last 5 years might have already adopted the permanent contraceptive methods.

In this study, we identified that when decision-making for contraception use was taken solely by the husband or jointly (both respondents and their husband), the use of the permanent contraception method was higher, that is, the husband's approval had a more significant influence. The earlier investigations found positive associations between contraceptive use and the husband's acceptance and shared decision.^{61 65–68} The most likely explanation could be that Bangladesh has a traditional society where women are expected to follow their husbands' advice in many aspects of life⁶⁶ and frequently require their husbands' consent before using contraception.³⁶ On the other hand, a few husbands might be unaware of the advantages of LARCs, and some might even have misapprehended that they are incompatible with the use, which might be the possible reasons for the low use of LARCs when the decision was taken by the husbands only.⁶⁵

Notably, the likelihood of using permanent contraception was higher among women with high decision-making power than those with low decision-making power. We observed a consistent result in our study that is in agreement with other studies from SSA^{69 70} and the Democratic Republic of the Congo.⁷¹ Though a previous study argued that women are more likely to prefer a readily accessible type of contraception regardless of their empowerment status,⁷⁰ an alternative statement might explain our finding that increasing women's autonomy gives them greater authority to choose whether or not to use contraception as well as which contraceptive method to use.⁴⁰

Women from affluent households tended to adopt LARC methods less than those from households with lower wealth quintiles. This result is in line with the findings from previous studies conducted in Ethiopia,⁸ France,⁷² Bangladesh and India.⁷³ Bangladesh's government has prioritised contraceptive services, and the country has a substantial budget that includes money for client repayment and provider payments, which might explain this situation.⁷⁴ Additionally, wealthy families might believe that their children could help them maintain their wealth, and they might need more family members to inherit their assets when they pass away.⁸ In contrast, poorer families might use a variety of contraceptive methods for an effective and long-lasting method of pregnancy prevention because childbirth and parenting are more expensive.⁸

Implications for public health research and policy

There are several benefits from this study regarding public health and policy. This study identified a positive association between household wealth status and LARC practices among women, where affluent women were less likely to use the LARC method than the poorest women. This association was significant and it indicates a disparity in accessing LARC and permanent methods in

Bangladesh. Further, prospective studies and qualitative exploration could be adopted to investigate why women from wealthier households are less willing to use the LARC method in Bangladesh. Along with the government, this inequity could be addressed by other relevant stakeholders. We also identified that media exposure among women significantly influenced their choice of permanent method. Given that media is an inexpensive way to convey information and spread awareness among women, it might be used objectively to promote FP by sensitising and creating awareness regarding the availability and useability of the services. To increase the permanent and LARC methods uptake, who choose to use them, we also recommend that public health campaigns and policies focusing on young and employed women and those having lower decision-making power might be created and disseminated. Finally, the findings of this study will be helpful to our policymakers in reducing maternal mortality rates and controlling fertility rates in Bangladesh and other similar settings. Besides, further trend analysis using data from previous survey waves of BDHS is recommended to look at the demographic shifts over time in contraceptive use among Bangladeshi women.

Strengths and limitations

The major strength of this study is the use of the most recent nationally representative data set; therefore, the study results could be generalisable across the country and other similar settings. The cluster effect on the factors associated with the contraception method usage in Bangladesh was taken into account in this study using a multilevel model where clusters were considered as level-2 factor, which is the proper analytical strategy for this type of hierarchical data. This study identified the factors associated with both permanent and LARC methods among Bangladeshi women. However, this study has some limitations, and one of those is that the cross-sectional study design prevented us from concluding the causal association between outcome and explanatory variables. Furthermore, some important factors related to the respondent's health were not included in this study due to the unavailability and incomplete information of those variables in the data set.

CONCLUSION

Among modern contraceptive users, about 83.48% of participants used the SARC methods, while only 11.34%, and 5.18% of women used permanent and LARC methods, respectively. This study found that women with no/less education, non-Muslims and having parity of ≥ 3 were more likely to use both permanent and LARC methods compared with SARC methods. Besides, women from higher age groups, those with media exposure, and high decision-making power were more likely to use the permanent methods, while women within the richest wealth index were less likely to use the LARC methods compared with the SARC methods. Targeted

interventions could be developed and implemented to promote personalised contraceptive use.

Author affiliations

- ¹Global Health Institute, North South University, Dhaka, Bangladesh
²Department of Public Health, North South University, Dhaka, Bangladesh
³Department of Statistics, University of Rajshahi, Rajshahi, Bangladesh
⁴Department of Oral & Maxillofacial Surgery, Marks Medical College Hospital & Dental Unit, Dhaka, Bangladesh
⁵Faculty of Nutrition and Food Science, Patuakhali Science and Technology University, Patuakhali, Bangladesh

Acknowledgements The authors of the present study greatly acknowledge the Demographic and Health Survey (DHS) for providing access to freely use their database.

Contributors SK accepts full responsibility for the work and/or the conduct of the study, had access to the data and controlled the decision to publish. ABS and SK also take responsibility for the integrity and accuracy of the data analysis. ABS and SK performed the statistical analysis. ABS, SK, TH, MHN and TZM produced the first draft of the manuscript. ABS and SK reviewed and undertook the scientific editing of the manuscript both for statistical correctness and language appropriateness. All authors read and approved the final version for publication.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open <https://dhsprogram.com/data/available-datasets.cfm> access repository. The study used data from the 2017–2018 Bangladesh Demographic and Health Survey. The data set is available at: <https://dhsprogram.com/data/available-datasets.cfm>.⁷⁵

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Azaz Bin Sharif <http://orcid.org/0000-0002-9929-1236>
 Md Tamzid Hasan <http://orcid.org/0000-0002-5812-5717>
 Md. Habib Naziat <http://orcid.org/0000-0002-6706-3052>
 Satyajit Kundu <http://orcid.org/0000-0001-9610-1479>

REFERENCES

- Gerland P, Raftery AE, Sevčiková H, *et al*. World population stabilization unlikely this century. *Science* 2014;346:234–7.
- Khan MN, Harris M, Loxton D. *Effects of unintended pregnancy on maternal Healthcare services use in Bangladesh*. 2021.
- Ameyaw EK, Budu E, Sambah F, *et al*. Prevalence and determinants of unintended pregnancy in sub-Saharan Africa: A multi-country analysis of demographic and health surveys. *PLoS One* 2019;14.
- Bolarinwa OA, Nwagbara UI, Okyere J, *et al*. Prevalence and predictors of long-acting reversible contraceptive use among sexually active women in 26 sub-Saharan African countries. *Int Health* 2022;14:492–500.
- Ahmed S, Li Q, Liu L, *et al*. Maternal deaths averted by contraceptive use: an analysis of 172 countries. *The Lancet* 2012;380:111–25.
- Das M, Anand A, Hossain B, *et al*. Inequalities in short-acting reversible, long-acting reversible and permanent contraception use among currently married women in India. *BMC Public Health* 2022;22:1264.
- Ponce de Leon RG, Ewerling F, Serruya SJ, *et al*. Contraceptive use in Latin America and the Caribbean with a focus on long-acting reversible contraceptives: prevalence and inequalities in 23 countries. *Lancet Glob Health* 2019;7:e227–35.
- Aychew EW, Bekele YA, Ayele AD, *et al*. Utilization of long-acting contraceptive methods and associated factors among married women in Farta Woreda, Northwest Ethiopia: a community-based mixed method study. *BMC Womens Health* 2022;22:533.
- Stoddard A, McNicholas C, Peipert JF. Efficacy and safety of long-acting reversible contraception. *Drugs* 2011;71:969–80.
- Shoupe D. LARC methods: entering a new age of contraception and reproductive health. *Contracept Reprod Med* 2016;1:4.
- Polis CB, Bradley SEK, Bankole A, *et al*. Typical-use contraceptive failure rates in 43 countries with demographic and health survey data: summary of a detailed report. *Contraception* 2016;94:11–7.
- Engstrand S, Kopp Kallner H. Cost of unintended pregnancy in Sweden - a possibility to lower costs by increasing LARC usage. *Contraception* 2018;97:445–50.
- Association of State and territorial health officials. Increasing access to contraception learning community. 2018. Available: <https://www.astho.org/globalassets/pdf/iac-y3-final-report.pdf> [Accessed 2 Jun 2023].
- World health organization, world bank, United Nations population Fund. *Trends in maternal mortality: 1990 to 2008 Tendances de la Mortalité Maternelle sur La Période 1990-2008: estimations Établies par L'OMS, L'UNICEF, L'UNFPA et La Banque Mondiale*. 1990. Available: <https://apps.who.int/iris/handle/10665/44423>
- Bradley SE, Croft T, Rutstein SO. The impact of contraceptive failure on unintended births and induced abortions: estimates and strategies for reduction. ICF macro Calverton Maryland Maryland 2011. 2011. Available: <https://dhsprogram.com/pubs/pdf/as22/as22.pdf> [Accessed 22 Jan 2023].
- National Institute of Population Research and Training - NIPORT/ Bangladesh, Mitra and Associates, and ICF International. Bangladesh demographic and health survey. 2014. Available: <http://dhsprogram.com/pubs/pdf/FR311/FR311.pdf>
- Bairagi R, Islam MM, Barua MK. Contraceptive failure: levels, trends and determinants in Matlab, Bangladesh. *J Biosoc Sci* 2000;32:107–23.
- Jacobstein R, Bakamjian L, Pile JM, *et al*. Fragile, threatened, and still urgently needed: family planning programs in sub-Saharan Africa. *Stud Fam Plann* 2009;40:147–54.
- Saloranta TH, Gyllenberg FK, But A, *et al*. Use of reproductive health services among women using long- or short-acting contraceptive methods - a register-based cohort study from Finland. *BMC Public Health* 2022;22.
- United Nations Department of Economic and Social Affairs. Population division. World fertility and family planning 2020: highlights (ST/ESA/SER.A/440). 2020. Available: https://www.un.org/en/development/desa/population/publications/pdf/family/World_Fertility_and_Family_Planning_2020_Highlights.pdf
- National Institute of population research and training (NIPORT), ICF. Bangladesh demographic and health survey 2017-18. 2020. Available: <https://DhsprogramCom/Publications/Publication-FR344-DHS-Final-ReportsCfm>
- Chaurasia AR. Contraceptive use in India: A data mining approach. *International Journal of Population Research* 2014;2014:1–11.
- Ministry of Health - MOH/Nepal, new ERA/Nepal, ICF. Nepal demographic and health survey 2016. Kathmandu, Nepal: MOH/Nepal, new ERA, and ICF 2017. 2017. Available: <http://dhsprogram.com/pubs/pdf/FR336/FR336.pdf>
- National Institute of population studies - NIPS/Pakistan, ICF. Pakistan demographic and health survey 2017-18. Islamabad, Pakistan: NIPS/Pakistan and ICF. 2019. Available: <http://dhsprogram.com/pubs/pdf/FR354/FR354.pdf>
- Central Statistics Organization (CSO), ICF. Afghanistan demographic and health survey 2015, Kabul, Afghanistan. Central Statistics Organization; 2017. Available: https://dhsprogram.com/Countries/Country-Main.cfm?ctry_id=71&c=Afghanistan&r=4
- Secura G. Long-acting reversible contraception: a practical solution to reduce unintended pregnancy. *Minerva Ginecol* 2013;65:271–7.
- United Nations. Contraceptive use by method 2019. New York: United Nations Department of Economic and Social Affairs, 10 December 2019.
- Sonfield A. Popularity disparity: attitudes about the IUD in Europe and the United States. *Guttmacher Institute* 2007.
- Buhling KJ, Zite NB, Lotke P, *et al*. Worldwide use of Intrauterine contraception: a review. *Contraception* 2014;89:162–73.
- Aduloju OP, Akintayo AA, Adefisan AS, *et al*. Utilization of long-acting reversible contraceptive (LARC) methods in a tertiary hospital in southwestern Nigeria: A mixed methods study. *J Obstet Gynaecol India* 2021;71:173–80.

- 31 Ahmed S, Creanga AA, Gillespie DG, *et al.* Economic status, education and empowerment: implications for maternal health service utilization in developing countries. *PLOS ONE* 2010;5.
- 32 Hogan DP, Berhanu B, Hailemariam A. Household organization, women's autonomy, and contraceptive behavior in Southern Ethiopia. *Stud Fam Plann* 1999;30:302–14.
- 33 Bangladesh Bureau of Statistics. Population and housing census 2022. 2022. Available: [https://sid.portal.gov.bd/sites/default/files/files/sid.portal.gov.bd/publications/01ad1ffe_cfef_4811_af97_594b6c64d7c3/PHC_Preliminary_Report_\(English\)_August_2022.pdf](https://sid.portal.gov.bd/sites/default/files/files/sid.portal.gov.bd/publications/01ad1ffe_cfef_4811_af97_594b6c64d7c3/PHC_Preliminary_Report_(English)_August_2022.pdf) [Accessed 22 May 2023].
- 34 El Arifeen S, Christou A, Reichenbach L, *et al.* Community-based approaches and partnerships: innovations in health-service delivery in Bangladesh. *Lancet* 2013;382:2012–26.
- 35 Hubacher D, Mavranzouli I, McGinn E. Unintended pregnancy in sub-Saharan Africa: magnitude of the problem and potential role of contraceptive implants to alleviate it. *Contraception* 2008;78:73–8.
- 36 Mutombo N, Bakibinga P. The effect of joint contraceptive decisions on the use of Injectables, long-acting and permanent methods (Ilapms) among married female (15–49) contraceptive users in Zambia: a cross-sectional study. *Reprod Health* 2014;11:1–8.
- 37 Das M, Anand A, Hossain B, *et al.* Inequalities in short-acting reversible, long-acting reversible and permanent contraception use among currently married women in India. *BMC Public Health* 2022;22:1–13.
- 38 Adedini SA, Omisakin OA, Somefun OD. Trends, patterns and determinants of long-acting reversible methods of contraception among women in sub-Saharan Africa. *PLoS One* 2019;14.
- 39 Boah M, Issah A-N, Demuyakor I, *et al.* Long-acting reversible contraceptives utilization and its determinants among married Yemeni women of childbearing age who no longer want children. *Medicine (Baltimore)* 2022;101.
- 40 Wado YD, Gurmu E, Tilahun T, *et al.* Contextual influences on the choice of long-acting reversible and permanent contraception in Ethiopia: A Multilevel analysis. *PLoS One* 2019;14.
- 41 Bhandari R, Pokhrel KN, Gabrielle N, *et al.* Long acting reversible contraception use and associated factors among married women of reproductive age in Nepal. *PLoS One* 2019;14.
- 42 Fekadu GA, Omigbodun AO, Roberts OA, *et al.* Determinants of change in long-acting or permanent contraceptives use in Ethiopia; A multivariate decomposition analysis of data from the Ethiopian demographic and health survey. *PLoS One* 2020;15.
- 43 Negash WD, Belachew TB, Asmamaw DB. Long acting reversible contraceptive utilization and its associated factors among modern contraceptive users in high fertility sub-Saharan Africa countries: a multi-level analysis of recent demographic and health surveys. *Arch Public Health* 2022;80:224.
- 44 World Health Organization. *WHO BMI Classification*. World Health Organization, 2020.
- 45 NIPORT, Mitra and Associates and Ii. *Bangladesh Demographic and Health Survey 2011*. Dhaka, Bangladesh: NIPORT, Mitra and Associates, and ICF International, 2013.
- 46 Khan HR, Shaw E. Multilevel logistic regression analysis applied to binary contraceptive prevalence data. *J Data Sci* 2011;9:93–110.
- 47 Directorate General of Family Planning. Annual report 2014, Ministry of Health & Family Welfare . 2015. Available: https://dgifp.gov.bd/sites/default/files/files/dgifp.portal.gov.bd/annual_reports/beaf0f39d_1ab9_4e08_b7ec_73f86b42dbe5/annual_rep2014.pdf
- 48 Chacko MR, Wiemann CM, Buzi RS, *et al.* Choice of postpartum contraception: factors predisposing pregnant adolescents to choose less effective methods over long-acting reversible contraception. *J Adolesc Health* 2016;58:628–35.
- 49 Bulto GA, Zewdie TA, Beyen TK. Demand for long acting and permanent contraceptive methods and associated factors among married women of reproductive age group in Debre Markos town, North West Ethiopia. *BMC Womens Health* 2014;14:46.
- 50 Haile A, Fantahun M. Demand for long acting and permanent contraceptive methods and associated factors among family planning service users. *Ethiopian Medical Journal* 2012;31–42.
- 51 Sharma V, Mohan U, Das V, *et al.* Socio demographic determinants and knowledge, attitude, practice: survey of family planning. *J Family Med Prim Care* 2012;1:43.
- 52 Amo-Adjei J, Mutua M, Mukiira C, *et al.* Fertility intentions and the adoption of long-acting and permanent contraception (LAPM) among women: evidence from Western Kenya. *BMC Womens Health* 2019;19.
- 53 Alemayehu M, Kalayu A, Desta A, *et al.* Rural women are more likely to use long acting contraceptive in Tigray region, northern Ethiopia: a comparative community-based cross sectional study. *BMC Womens Health* 2015;15.
- 54 Kasarda JD, Billy JOG, West K. *Status Enhancement and fertility: reproductive responses to social mobility and educational opportunity*. Orlando, Fla: Academic Press, 1986.
- 55 Rindfuss RR, Morgan SP, Offutt K. Education and the changing age pattern of American fertility: 1963–1989. *Demography* 1996;33:277–90.
- 56 Gebremariam A, Addissie A. Intention to use long acting and permanent contraceptive methods and factors affecting it among married women in Adigrat town, Tigray, northern Ethiopia. *Reprod Health* 2014;11:1–9.
- 57 Mare KU, Abrha E, Mohammed Yesuf E, *et al.* Factors affecting utilization of long-acting reversible contraceptives among sexually active reproductive-age women in the pastoral community of Northeast Ethiopia: a community-based cross-sectional study. *Womens Health (Lond)* 2022;18:17455057221116514.
- 58 Mehata S, Paudel YR, Dotel BR, *et al.* Inequalities in the use of family planning in rural Nepal. *BioMed Research International* 2014;2014:1–7.
- 59 Palamuleni ME, Adebowale AS. Women empowerment and the current use of long acting and permanent contraceptive: evidence from 2010 Malawi demographic and health survey. *Malawi Med J* 2014;26:63–70.
- 60 Bolarinwa OA, Afaya A, Ajayi KV, *et al.* Prevalence and factors associated with the use of long-acting reversible and permanent contraceptive methods among women who desire no more children in high fertility countries in sub-Saharan Africa. *BMC Public Health* 2022;22:2141.
- 61 Melka AS, Tekelab T, Wirtu D. Determinants of long acting and permanent contraceptive methods utilization among married women of reproductive age groups in Western Ethiopia: a cross-sectional study. *Pan Afr Med J* 2015;21:246.
- 62 Orwa J, Gatimu SM, Ngugi A, *et al.* Factors associated with use of long-acting reversible and permanent contraceptives among married women in rural Kenya: A community-based cross-sectional study in Kisii and Kilifi counties. *PLoS ONE* 2022;17.
- 63 Sambah F, Aboagye RG, Seidu A-A, *et al.* Long-acting reversible contraceptives use among adolescent girls and young women in high fertility countries in sub-Saharan Africa. *Reprod Health* 2022;19:209.
- 64 Anguzu R, Sempeera H, Sekandi JN. High parity predicts use of long-acting reversible contraceptives in the extended postpartum period among women in rural Uganda. *Contracept Reprod Med* 2018;3:6.
- 65 Gashaye KT, Tsegaye AT, Abebe SM, *et al.* Determinants of long acting reversible contraception utilization in Northwest Ethiopia: an institution-based case control study. *PLoS ONE* 2020;15.
- 66 Kamal N. The influence of husbands on contraceptive use by Bangladeshi women. *Health Policy Plan* 2000;15:43–51.
- 67 D'Exelle B, Ringdal C. Women's use of family planning services: an experiment on the husband's involvement. *Journal of Development Economics* 2022;158.
- 68 Shimeka Teferra A. Determinants of long acting contraceptive use among reproductive age women in Ethiopia: evidence from EDHS 2011. *SJPH* 2015;3:143.
- 69 Yaya S, Uthman OA, Ekholuenetale M, *et al.* Women empowerment as an enabling factor of contraceptive use in sub-Saharan Africa: a Multilevel analysis of cross-sectional surveys of 32 countries. *Reprod Health* 2018;15:214.
- 70 Adde KS, Ameyaw EK, Dickson KS, *et al.* Women's empowerment indicators and Short- and long-acting contraceptive method use: evidence from DHS from 11 countries. *Reprod Health* 2022;19:222.
- 71 Sano Y, Antabe R, Atuoye KN, *et al.* Married women's autonomy and post-delivery modern contraceptive use in the Democratic Republic of Congo. *BMC Womens Health* 2018;18.
- 72 Moreau C, Bohet A, Hassoun D, *et al.* Trends and determinants of use of long-acting reversible contraception use among young women in France: results from three national surveys conducted between 2000 and 2010. *Fertil Steril* 2013;100:451–8.
- 73 Ugaz JI, Chatterji M, Gribble JN, *et al.* Is household wealth associated with use of long-acting reversible and permanent methods of contraception? A multi-country analysis. *Glob Health Sci Pract* 2016;4:43–54.
- 74 MEASURE Evaluation. The future of long-acting and permanent methods of contraception in Bangladesh. Chapel Hill, NC: MEASURE Evaluation; 2014. Available: <https://www.measureevaluation.org/publications/fs-14-131.html> [Accessed 29 May 2023].
- 75 National Institute of Population Research and Training. Data extract from Bdir7Rfl.SAV.Bangladesh demographic and health Surveys(bdhs), version 7, BDHS and ICF. 2020. Available: <https://dhsprogram.com/data/available-datasets.cfm> [Accessed 2 Jun 2023].