

Invasive records of *Eriocheir hepuensis* Dai, 1991 (Crustacea: Brachyura: Grapsoidea: Varunidae): Implications and taxonomic considerations

Author

Naser, MD, Page, TJ, Ng, NK, Apel, M, Yasser, AG, Bishop, JM, Ng, PKL, Clark, PF

Published

2012

Journal Title

BioInvasions Records

DOI

[10.3391/bir.2012.1.1.15](https://doi.org/10.3391/bir.2012.1.1.15)

Rights statement

© The Author(s) 2012. This is an Open Access article distributed under the terms of the Creative Commons Attribution 2.0 Generic (CC BY 2.0) license (<http://creativecommons.org/licenses/by/2.0/>) which permits unrestricted distribution and reproduction in any medium, providing that the work is properly cited.

Downloaded from

<http://hdl.handle.net/10072/46953>

Griffith Research Online

<https://research-repository.griffith.edu.au>

Research Article

Invasive records of *Eriocheir hepuensis* Dai, 1991 (Crustacea: Brachyura: Grapsoidea: Varunidae): Implications and taxonomic considerations

Murtada D. Naser¹, Timothy J. Page², N.K. Ng³, Michael Apel⁴, Amaal G. Yasser¹, James M. Bishop⁵, Peter K.L. Ng³ and Paul F. Clark^{6*}

¹ Marine Biology Department, Marine Science Centre, University of Basrah, Basrah, Iraq

² Australian Rivers Institute, Griffith University, 170 Kessels Road, Nathan, Queensland 4111, Australia

³ Department of Biological Sciences, National University of Singapore, 14 Science Drive 4, Singapore 117543, Singapore

⁴ Museum Mensch und Natur, Schloss Nymphenburg, 80638 München, Germany

⁵ Kuwait Institute of Scientific Research, P.O. Box 1638, Salmiya 22017, Kuwait

⁶ Department of Zoology, The Natural History Museum, Cromwell Road, London SW7 5BD, England

E-mail: bio_mur_n@yahoo.com (MDN), penguintim@hotmail.com (TJP), nganke@nus.edu.sg (NKN), m.apel@musmn.de (MA), amaalyasser@yahoo.com (AGY), JBishop@kisir.edu.kw (JMB), peterng@nus.edu.sg (PKLN), p.clark@nhm.ac.uk (PFC)

*Corresponding author

Received: 22 November 2011 / Accepted: 30 January 2012 / Published online: 2 March 2012

Abstract

A non-ovigerous female mitten crab captured in Iraq on the 20th June 2005 was initially identified as *Eriocheir sinensis* H. Milne Edwards, 1853. More material has now been made available from Iraq and Kuwait and these specimens were compared with the extant type series of *E. sinensis* and *E. hepuensis* Dai, 1991. From this morphological study the Persian Gulf material was identified as the Hepu mitten crab, *E. hepuensis*. This is the first report of this mitten crab species outside its native range of southern China. However, because mitten crab taxonomy and systematics requires further clarification especially with respect to species of *Eriocheir* De Haan, 1835, the DNA of the Persian Gulf material was compared with a suite of GenBank COI sequences from various mitten crab taxa. The results of this extensive examination indicate that the Iraqi and Kuwaiti specimens are *E. hepuensis* and the DNA analysis indicates that mitten crabs can be assigned to three genera and six species; although the present study does discuss reports of hybridisation between three *Eriocheir* species associated with aquaculture. Also reported, and of concern, is that *E. hepuensis* may have been first collected about thirty years ago in Iraq and has now become well established within that country. As a consequence the Hepu mitten crab could, given its catadromy, disperse widely throughout the extensive riverine systems of the region.

Key words: *Eriocheir hepuensis*; invasive species; Persian Gulf; molecular and morphological analyses; mitten crab taxonomy

Introduction

Four species of “hairy” (colloquially Southeast and East Asia) or “mitten” (vernacular Europe) crabs are currently assigned to *Eriocheir* De Haan, 1835 (Brachyura: Grapsoidea: Varunidae), initially, *E. japonica* (De Haan, 1835), and *E. sinensis* H. Milne Edwards, 1853, and recently, *E. hepuensis* Dai, 1991, and *E. ogasawaraensis* Komai, Yamasaki, Kobayashi, Yamamoto and Watanabe, 2006 (Ng et al. 2008). *Eriocheir* species are indigenous to China, the Korean Peninsula and Japan (Figure 1), and are all catadromous, spending most of their lives in fresh water, but returning to higher salinity water for mating, spawning and larval development. The seaward migration is annual, occurring during the autumnal months, and after mating the adults die (semelparous). The life cycle is

completed by the upstream migration of early crab stages into freshwater.

Eriocheir sinensis has become arguably the most notorious brachyuran species on the planet. It is one of only two crabs on the world’s list of 100 most invasive aquatic invertebrates (Lowe et al. 2004). Incidentally, the other brachyuran species listed is the portunid *Carcinus maenas* (Linnaeus, 1758). The Chinese mitten crab is also cited in the handbook of alien species in Europe (DAISIE 2009: 312). This crab has successfully invaded NE Europe and has been reported from the east (Ruiz et al. 2006; Dittel and Epifanio 2009; Schmidt et al. 2009) and west coast (Cohen and Carlton 1997) of the United States and eastern Canada (Nepszy and Leach 1973; de Lafontaine 2005) with records from southern Iraq (Clark et al. 2006; Hashim 2010), and Tokyo Bay, Japan in 2004 (S. Kobayashi,




Figure 1. Known distribution of native *Eriocheir* De Haan, 1835 species.



Figure 2. Advertisement in a local newspaper promoting mitten crab on the menu of the Hôtel Plaza Athénée, Bangkok for the whole of November, 2001.

(pers. comm.; Doi et al. 2011). Furthermore, *E. sinensis* is one of the most commercially valuable crabs in East and Southeast Asia (e.g., see Peng 1986; Zhao 1988; Ng 1998; Lai et al. 1992), where the gonads, which develop during the annual downstream migration, are regarded as a delicacy (Figure 2).

Eriocheir japonica is also commercially exploited, but not to the same extent as *E. sinensis*, being consumed predominantly in Japan. The Japanese mitten crab too has been reported from outside its home range when a solitary male crab was captured in the Columbia River approximately 3km west of the Astoria-Megler Bridge, Astoria, Oregon, USA (see Jansen and Armstrong 2004). No further exotic reports have been recorded.

Recently, new mitten crab specimens from a number of localities in Iraq and Kuwait have been reported (Figure 3). Some of this material, which included several large males, has become available for detailed examination. The specimens were identified as *Eriocheir hepuensis*, the Hepu mitten crab. This is the first report of the crab outside its native range of southern China. However, because mitten crab taxonomy and systematics requires further clarification especially with respect to species of *Eriocheir* De Haan, 1835, a more extensive investigation was undertaken. This included a thorough examination of available *E. sinensis* and *E. hepuensis* material, a morphometric analysis, and the DNA barcode (*sensu* Costa et al. 2007; the mitochondrial COI gene) of an Iraqi mitten crab was compared against a suite of GenBank sequences for *Eriocheir*.

The present study reports upon the results of the morphological and genetic investigations of the mitten crabs from the Persian Gulf, and the implications for mitten crab taxonomy and systematics. Also discussed is the capacity of *E. hepuensis* for further dispersal throughout the extensive watersheds of the region.


Methods and material

Abbreviations used: ASIZ = National Zoological Museum of China, The Chinese Academy of Sciences, Beijing, People's Republic of China (previously known as Institute of Zoology, The Chinese Academy of Sciences, Beijing, People's Republic of China); coll. = collected; det. = determined by; MNHN = Muséum national d'Histoire naturelle; NHM = Natural History Museum, London; ZRC = Zoological Reference Collection, Raffles Museum of Biodiversity Research, Singapore; ZSM = Zoologische Staatssammlung München; ovig. = ovigerous; reg. = registration number.


Measurements: Carapace measurements = width × length in millimetres.

Figure 3. *Eriocheir hepuensis* specimens collected from Iraq and Kuwait.


- ★ Collected in Jan - Feb 2004 from Kuwaiti waters.
- ★ Collected in Jun 2006 from Shatt Al-Basrah Canal, Iraq (Clark et al. 2006).
- ★ Collected in Mar 2009 - Jan 2010 from Iraq (Hashim 2010).
- ★ Collected in Nov 2010 - Aug 2011 from Iraq (M.D. Naser, pers. comm.).
- ? Reports in Aug 2011 by fishermen from Iraq (M.D. Naser, pers. comm.).



E. sinensis



E. hepuensis



E. japonica

Figure 4. Dorsal carapace morphology of three mitten crab species (from Guo et al. 1997).

Literature: Only a selected synonymy for *E. sinensis* and *E. hepuensis* is provided here, but see Guo et al. (1997) for additional references.

Identification and Taxonomy

***Eriocheir sinensis* H. Milne Edwards, 1853**

(Figures 4, 5a, 6a, 7a, 8)

Eriochirus sinensis H. Milne Edwards 1853: 177.

Eriocheir sinensis – H. Milne Edwards, 1854: 146-148, Pl. 9; Figures 1, 1a-c; Peters and

Panning 1933: 1–180; Shen and Dai 1964: 127; Kim 1973: 465, text-fig. 202, Pl. 40, Figure. 154; Hwang and Mizue 1985: 12; Gu 1986: 268; Dai et al. 1986: 523; Yu and Ho 1986: 116; Adema 1991: 201, Fig. 79; Dai and Yang 1991: 523; Lai and Lu 1992: 23; Hong et al. 1993: 10, Pls. A and F; Chan et al. 1995: 301, Figure 3B; N.K. Ng et al. 1998: 493, 1999: 154; Sun et al. 2003: 592; Chu et al. 2003: 738; Tang et al. 2003: 309, 2004: 255; Chan et al. 2005: 457; Robbins et al. 2006: 33, Figures 1-2; Ng et al. 2008: 228.



Figure 5. Frontal margin of a. *Eriocheir sinensis*, 1♂ (74.0×68.5mm), NHM reg. 1993.1, River Cray, Hall Place, near Crayford, Kent, England, coll. 20 Aug. 1992; b. *E. hepuensis*, 1♂ (70.6×66.4 mm), Shatt Al-Basrah canal near the dam at 30°24'33.75"N 47°46'32.32"E, Iraq, coll. M. Naser, 30 Nov. 2010, NHM reg. 2011. 8035. Photograph taken by Phil Hurst, NHM Photo Unit.




Figure 6. Anterolateral carapace margin of a. *Eriocheir sinensis*, 1♂ (74.0×68.5mm), NHM reg. 1993.1, River Cray, Hall Place, near Crayford, Kent, England, coll. 20 Aug. 1992; b. *E. hepuensis* 1♂ (70.6×66.4 mm), Shatt Al-Basrah canal near the dam at 30°24'33.75"N 47°46'32.32"E, Iraq, coll. M. Naser, 30 Nov. 2010, NHM reg. 2011. 8035. Photograph taken by Phil Hurst, NHM Photo Unit.

Eriocheir sinensis form *rotundifrons* Panning, 1938: 109, Figure 5; Adema 1991: 201, Figure 79.

Eriocheir sinensis form *acutifrons* Panning, 1938: 109, Figure 6; Adema 1991: 201, Figure 79.





Figure 7. Fifth pereiopod, fourth ambulatory leg of a. *Eriocheir sinensis*, 1♂ (74.0×68.5mm), NHM reg. 1993.1, River Cray, Hall Place, near Crayford, Kent, England, coll. 20 Aug. 1992; b. *E. hepuensis*, 1♂ (70.6×66.4 mm), Shatt Al-Basrah canal near the dam at 30°24'33.75"N 47°46'32.32"E, Iraq, coll. M. Naser, 30 Nov. 2010, NHM reg. 2011. 8035. Photograph taken by Phil Hurst, NHM Photo Unit.

Eriocheir sinensis form *trilobata* Panning, 1938: 110, Figure 7; Adema 1991: 201, Figure 79.

Eriocheir sinensis form *rostrata* Panning, 1933: 53, Figure 22B; 1938: 110, Figure 8; Adema 1991: 201, Figure 79.

Material examined. Type material: lectotype. China. 1♀ (66.0×57.0mm), MNHN reg. B3383S, Macao, coll. M. Callery, no date. Non-type material: England. 1♂ (74.0×68.5mm), NHM reg. 1993.1, River Cray, Hall Place, near Crayford, Kent, coll. 20 Aug. 1992; 1♂ (69.0×62.0mm), NHM reg. 2003.315. China. Hong Kong Market, tied with rice grass, coll. P. Clark, Oct. 2002; 2♂ (70.9-72.8×62.5-63.3mm), 1♀ (61.6×57.0mm), NHM reg. 2011.8006-8008, Hong Kong, tied with rice grass, coll. B. Morton, Nov. 2010. In addition, see specimens listed in Guo et al. (1997).

Diagnosis. Carapace squarish, overall dorsal surface convex, regions well-defined (Figure 4). Frontal margin with four strong, sharp teeth with a narrow v-shaped median cleft (Figures 4, 5a). Anterolateral margins with four teeth including exorbital tooth (Figures 4, 6a). Third maxilliped broad; ischium, merus broad, exopod narrow. Ambulatory legs slender, long, long thick setae on anterior, posterior surfaces of carpi, propodi (Figure 7a). G1 long, slender, distal margin narrowly rounded, sloping shoulder shaped



E. sinensis *E. hepuensis* *E. japonica*

Figure 8. Morphology of the first male gonopod of three mitten crab species (from Guo et al. 1997).



Figure 9. *Eriocheir hepuensis*: distal tip of first gonopod, specimen from Kuwait (ZSM A20110200). Photograph taken by Michael Apel.



Figure 10. *Eriocheir hepuensis*: 1♂ (68.7×63.6mm) Khor Al-Gayed, East of Ras Al-Gayed, Bubiyan Isl., 29°52'N 48°21'E, Kuwait, coll. 26 January 2004, otter trawl, 4-6 m. Photograph taken by Michael Apel.

when viewed laterally, short, chitinous prominence, slightly curved dorsally outwards with subdistal lobe (Figure 8), gonopore close to distal end. Female gonopore operculate, semicircular in shape, prominent, concave dorsally.

Eriocheir hepuensis Dai, 1991

(Figures 4, 5b, 6b, 7b, 8, 9, 10)

Eriocheir japonica hepuensis Dai, 1991: 61, Figures 1-11; Dai 1993: 17.

Eriocheir hepuensis – Guo et al. 1997: 460; N.K. Ng et al. 1998: 493; N.K. Ng et al. 1999: 154; Sun et al. 2003: 592; Chu et al. 2003: 738; Tang et al. 2003: 309; Tang et al. 2004: 255; Chan et al. 2005: 457; Apel and Bishop 2006; Ng et al. 2008: 228.

Eriocheir sinensis – Chan et al. 1995: 301 (part), Figure 3D; Clark et al. 2006: 51, Figs. 2-3; Hashim 2010: 32-33, Figure 2.

Material examined. Type material: holotype. China. 1♂ (70.2×63.0mm), ASIZ reg. GX899024A), Hepu, Guangxi Province, southern China; Paratype, 1♀ (68.1×62.7mm), ASIZ reg. GX899024B, Hepu, Guangxi Province, southern China, coll. 18 Nov. 1989. Non-type material: Kuwait. 1♂ (68.7×63.6mm), ZSM reg.

ZSMA20110200, Khor Al-Gayed, East of Ras Al-Gayed, Bubiyan Isl., 29°52'N 48°21'E, coll. 26 Jan. 2004, otter trawl, 4-6 m, det. M. Apel; 1♂ (52.7×49.3mm), ZSM reg. ZSMA20110201, Khor Abdullah off Ras Al-Gayed (between Bubiyan Island and Fao Peninsula, Iraq), coll. 24 Feb. 2004, otter trawl, 4-10 m, det. M. Apel. Iraq. 1♀ (47.8×45.9mm), NHM reg. 2006.98, Shatt-Al, Basrah canal, 30°15'41.25"N, 47°48'




Figure 11. Cladogram of the four species of *Eriocheir* viz. *E. sinensis*, *E. hepuensis*, *E. japonica* and *E. ogasawaraensis*; *Neoeriocheir leptoganthus*; *Platyriocheir formosa* and *Gercarcinus weileri* (outgroup), based on 46 adult morphological characters (Ng, N.K. unpublished data).

56.91"E, coll. Ibtam Abdul-Sahib, 20 Jun. 2005, det. P. Clark as *E. sinensis*; 2♂ (71.6×68.8-73.0×67.7mm), 1♀ ovi. (75.2×72.0 mm), ZRC reg. ZRC 2011.0732, Shatt Al-Basrah canal near the dam at 30°24'33.75"N 47°46'32.32"E, coll. M. Naser, 30 Nov. 2010; 2♂ (66.6×62.6-70.6×66.4 mm), 1♀ ovi. (68.6× 65.9mm), NHM reg. 2011.8035-8037, Shatt Al-Basrah canal near the dam at 30°24'33.75"N 47°46'32.32"E, coll. M. Naser, 30 Nov. 2010. In addition, see specimens listed in Guo et al. (1997).

Diagnosis. Carapace squarish, overall dorsal surface convex, regions well-defined (Figures 4, 10). Frontal margin with four sharp, broad teeth with a broad v-shaped median cleft (Figures 4, 6b). Anterolateral margins with four teeth including exorbital tooth (Figures 4, 6b, 10). Third maxilliped broad; ischium, merus broad, exopod narrow. Ambulatory legs broad, stout, long thick setae on anterior, posterior surfaces of carpi, propodi (Figure 7b). G1 long, slender, distal margin narrowly rounded, sloping shoulder shaped when viewed laterally, short, chitinous prominence, slightly curved outwards with subdistal lobe, gonopore at ½ length of distal end (Figures 8, 9). Female gonopore operculate,

bluntly triangular in shape, prominent, slightly concave dorsally.

Molecular

Genomic DNA was extracted from the leg of one *E. hepuensis* specimen collected at Shatt Al-Basrah, Iraq, using a modified version of a CTAB-phenol/chloroform extraction (Doyle and Doyle 1987). The 5' portion of the mitochondrial cytochrome oxidase subunit I (COI) was amplified using universal COI primers LCO-1490 (5'-TGA TTT TTT GGT CAC CCT GAA GTT CA-3') and HCO-2198 (5'-GGT CAA CAA ATC ATA AAG ATA TTG G-3') (Folmer *et al.*, 1994).

PCR amplifications were 12.5µl reactions on a Geneamp PCR System 9700 (Applied Biosystems, Foster City, CA, USA) of 0.5µl template DNA, 0.4µM primers, 0.1µM dNTPs, 2µM MgCl₂, 2.5µl 10X PCR Buffer, 0.5 units of Taq polymerase (Bioline Pty Ltd, Alexandria, NSW, Australia) and the rest ddH₂O. The following cycling conditions were used: 15 cycles of 30 s at 94°C, 30 s at 40°C, 60 s at 72°C; 25 cycles of 30 s at 94°C, 30 s at 55°C, 60 s at 72°C.

BigDye v.3.1 Terminator mix (Applied Biosystems) was used for the sequencing reaction in both directions and the sequences produced on an Applied Biosystems 3130xl Genetic Analyser at the DNA Sequencing Facility at Griffith University. Sequences were edited using Sequencher 4.1.2 (Gene Codes Corporation).

The COI sequence was compared against all GenBank sequences (17 Apr. 2011) using the BLASTN search at blast.ncbi.nlm.nih.gov, and also compared against the Barcoding of Life (BOLD) online database at <http://www.barcodinglife.org>. All GenBank *Eriocheir* COI sequences were downloaded and aligned with our sequence and a minimum evolution tree created using PAUP* version 4.0 b10 (Swofford 2002) with a Kimura 2-Parameter model (K2P), and bootstrapped 1000 times. A haplotype network was constructed with TCS version 1.21 (Clement *et al.* 2000) using sequences closely related to the Iraqi specimen (>99%) to explore within-species phylogeographic patterns. Molecular distances (Kimura 2-Parameter) between and within groups were calculated using Mega version 2.1 (Kumar *et al.* 2001).




Table 1. List of characters that can be used to distinguish between *Eriocheir sinensis* and *E. hepuensis*.

| Characters/Species | <i>E. sinensis</i> | <i>E. hepuensis</i> |
|---------------------------------|--|--|
| Teeth on frontal margin | Acutely triangular, sharp and high (Figure 5a) | Broadly triangular and sometimes not as sharp, and not as high (Figure 5b) |
| Median cleft on frontal margin | Narrow v-shaped (Figure 5a) | Broad v-shaped (Figure 5b) |
| Carapace physiognomy | High and convex | High but not convex |
| Anterolateral teeth of carapace | Usually sharp and strong (Figure 6a) | Weaker and not as sharp (Figure 6b) |
| 4th leg merus | Long and slender (Figure 7a) | Short and broad (Figure 7b) |
| 4th leg propodus | Long and slender (Figure 7a) | Shorter and broader (Figure 7b) |
| Male G1 | Gonopore close to distal end | Gonopore about ½ length from distal end |
| Female gonopore | Semicircular in shape, concave dorsally | Triangular in shape, slightly concave dorsally |

Mitten Crab taxonomy and systematics

Morphology

According to Ng et al. (2008), mitten crabs comprise three genera and six species namely *Eriocheir japonica* (De Haan, 1835), *E. sinensis*, *E. hepuensis*, and *E. ogasawaraensis* Komai, Yamasaki, Kobayashi, Yamamoto and Watanabe, 2006, *Neeriocheir leptognathus* (Rathbun, 1913) and *Platyeriocheir formosa* (Chan, Hung and Yu, 1995). However, this classification and the present identification of the Hepu mitten crab specimens from the Persian Gulf may be considered as controversial to some researchers (Tang et al. 2003, 2004; Sun et al. 2003, 2005) who consider *E. hepuensis* as an invalid species.

They consider the Hepu mitten crab to be a junior synonym of *E. sinensis*. Furthermore, Chu et al. (2003) have argued for maintaining five species (*E. ogasawaraensis* was described later) within *Eriocheir*, while Tang et al. (2003) suggested that *E. japonica*, *E. sinensis* and *E. hepuensis* were conspecific and should be considered as subspecies of *E. japonica*. In contrast, studies on morphology (Guo et al. 1997; Li and Zheng 2000a), reproductive biology (Li and Zheng 2000b), geography and habitats (Li and Zheng 2001), as well as a cladistic analysis based on 46 adult morphological characters (Ng, N.K. unpublished data; Figure 11) show that the Hepu mitten crab should be recognized as a separate taxon from *E. sinensis*.

Table 2. *Eriocheir* COI sequences downloaded from GenBank that are 99% similar to Iraqi specimen.

| Area | Site | Species in original publication | GenBank accession number | Reference |
|----------------|----------------|---------------------------------|--------------------------|-----------------------------|
| Northern China | Changjiang R. | <i>E. sinensis</i> | DQ438944 | Hu and Wang unpub. |
| Central China | Feiyunjiang R. | <i>E. japonica</i> | AY640088 | Xu et al. 2009 |
| | Feiyunjiang R. | <i>E. japonica</i> | AY640089 | Xu et al. 2009 |
| | Minjiang | <i>E. sinensis</i> | AF317336 | Tang et al. 2003 |
| | Oujiang R. | <i>E. japonica</i> | AY640088 | Xu et al. 2009 |
| | Oujiang R. | <i>E. japonica</i> | AY640089 | Xu et al. 2009 |
| | Oujiang R. | <i>E. japonica</i> | FJ750320 | Xu et al. 2009 |
| | Oujiang R. | <i>E. japonica</i> | FJ750321 | Xu et al. 2009 |
| | Tongan | <i>E. japonica</i> | AY640089 | Xu et al. 2009 |
| | Tongan | <i>E. japonica</i> | AY640090 | Xu et al. 2009 |
| | Tongan | <i>E. japonica</i> | FJ750322 | Xu et al. 2009 |
| Southern China | Aotou | <i>E. japonica</i> | AY640088 | Xu et al. 2009 |
| | Aotou | <i>E. japonica</i> | AY640089 | Xu et al. 2009 |
| | Aotou | <i>E. japonica</i> | FJ750323 | Xu et al. 2009 |
| | Aotou | <i>E. japonica</i> | FJ750324 | Xu et al. 2009 |
| | Hepu | <i>E. hepuensis</i> | AF317327 | Tang et al. 2003 |
| | Hepu | <i>E. hepuensis</i> | AF317328 | Tang et al. 2003 |
| | Hepu | <i>E. hepuensis</i> | AF516699 | Chu et al. 2003 |
| | Hepu | <i>E. japonica</i> | AY640088 | Xu et al. 2009 |
| | Hepu | <i>E. japonica</i> | AY640089 | Xu et al. 2009 |
| | Hepu | <i>E. japonica</i> | FJ750326 | Xu et al. 2009 |
| | Hepu | <i>E. japonica</i> | FJ750327 | Xu et al. 2009 |
| | Hepu | <i>E. japonica</i> | FJ750328 | Xu et al. 2009 |
| | Zhujiang | <i>E. japonica</i> | AY640091 | Xu et al. 2009 |
| | Zhujiang | <i>E. japonica</i> | FJ750324 | Xu et al. 2009 |
| | Zhujiang | <i>E. japonica</i> | FJ750325 | Xu et al. 2009 |
| | Zhujiang | <i>E. japonica</i> | FJ750326 | Xu et al. 2009 |
| Taiwan | Tamshui | <i>E. japonica</i> | AY640088 | Xu et al. 2009 |
| | Tamshui | <i>E. japonica</i> | FJ750321 | Xu et al. 2009 |
| Unsure | Unsure | <i>E. hepuensis</i> | FJ455506 | Wang, Huang and Li unpub. |
| | Unsure | <i>E. japonica</i> | AF105245 | Cheng, Chen and Chen unpub. |
| | Unsure | <i>E. japonica</i> | AF105246 | Cheng, Chen and Chen unpub. |

Interestingly, Figure 11 does not support a single taxon as suggested by Tang et al. (2003, 2004). Instead this cladogram confirms the recognition of three genera as proposed by Chan et al. (2005), Komai et al. (2006) and P.K.L. Ng et al. (2008), i.e., *Eriocheir*, *Neeriocheir* Sakai, 1983, and *Platyeriocheir* N.K. Ng, Guo and P.K.L. Ng, 1999. Figure 11 also confirms that *Eriocheir* s. str. comprises four species, *E. sinensis*, *E. japonica*, *E. hepuensis* and *E. ogasawaraensis*. Moreover, *E. hepuensis* can be distinguished from *E. sinensis* based on the characters listed in Table 1, which include the shape of the frontal (Figures 4, 5) and anterolateral (Figures 4, 6)

teeth, the shape and position of the carapace ridges (Figure 4, 5, 6), the meral and propodal proportions of the fourth ambulatory leg (Figure 7) and the position of the genital pore on the male gonopod (Figures 8, 9).

Molecular

As the validity of *E. hepuensis* appears to be divisive, a DNA analysis was undertaken to further clarify mitten crab systematics. For the present study, a 659 base pair COI sequence was produced (GenBank accession number JF810991) which corresponds to positions 5343-6001

Table 3. Molecular COI average distances (Kimura 2-parameter) between *Eriocheir* groups from Figure 15a (numbers in parentheses are within group averages).

| | <i>E. japonica</i> | <i>E. formosa/recta</i> | <i>E. sinensis</i> | <i>E. hepuensis</i> | <i>E. leptognathus</i> | <i>E. japonica (Okinawa)</i> | <i>E. ogasawaraensis</i> |
|------------------------------|--------------------|-------------------------|--------------------|---------------------|------------------------|------------------------------|--------------------------|
| <i>E. japonica</i> | (0.5%) | | | | | | |
| <i>E. formosa/recta</i> | 13.6% | (0.1%) | | | | | |
| <i>E. sinensis</i> | 4.3% | 14.4% | (0.4%) | | | | |
| <i>E. hepuensis</i> | 4.2% | 14.2% | 4.7% | (0.3%) | | | |
| <i>E. leptognathus</i> | 14.0% | 16.8% | 15.3% | 14.5% | (N/A) | | |
| <i>E. japonica (Okinawa)</i> | 3.1% | 14.8% | 4.2% | 3.0% | 13.6% | (0.4%) | |
| <i>E. ogasawaraensis</i> | 4.8% | 16.1% | 6.8% | 6.3% | 14.9% | 4.9% | (0.4%) |

of the *E. hepuensis* mitochondrial genome (GenBank genome reference NC011598; Wang, Huang and Li, unpublished data). BLAST searches of GenBank and BOLD identified 21 sequences that were 99% similar (Table 2 for sequence information; Table 3 for molecular distances; Appendix 1 for other *Eriocheir* sequences), all belonging to *Eriocheir*. Most of these were derived from Xu et al. (2009), but also included sequences from Tang et al. (2003) and Chu et al. (2003) and some unpublished sequences. The final alignment of all *Eriocheir* sequences was 554 base pairs. The Iraqi specimen formed part of a strong clade (bootstrap 100%) in a minimum evolution tree (Figure 12a) that includes all 21 closely related sequences (average 0.3% K2P molecular distance within the clade). This clade contains specimens listed on GenBank as being *E. hepuensis*, *E. japonica* and *E. sinensis*. Since different authors consider these *Eriocheir* taxa in many different ways (as separate species, separate subspecies or all conspecific), these sequences are labelled on the tree as the “*E. hepuensis*” clade on the basis of their geographic location rather than as a formal taxon.

Essentially, there were seven groups in the tree (Figure 12a). *Neoeriocheir leptognathus* and *Platyeriocheir formosa* were distinct from each other and all other taxa. The remaining five groups form a strong clade, but the relationship between them is uncertain. These groups consist of 1) *E. japonica* from mainland East Asia which corresponds to the species in the strict sense as this includes the type locality of mainland Japan; 2) *E. ogasawaraensis* from the Ogasawara Islands; 3) *E. sinensis* which is the species in the strict sense as it includes the type locality of mainland northern China; 4) “*E. japonica*” from Okinawa, which may represent a cryptic species

(or a geographic lineage of one of the other species); and 5) a mix of “*E. sinensis*”, “*E. japonica*” and “*E. hepuensis*”, including the specimen from Iraq, here referred to as the “*E. hepuensis*” clade.

The mixed nature of the last group can be explained by a number of factors (which are not mutually exclusive). Firstly, the GenBank specimens may have simply been identified incorrectly. This is more likely if only juveniles were used, purchased mixed from markets and collectors, and/or the worker was inexperienced with *Eriocheir* taxonomy. Some of the taxonomic characters used to separate the different species may also be less reliable due to possible variation. Consequently, new characters may need to be considered. Table 1 presents morphological characters currently used to distinguish between *E. sinensis* from *E. hepuensis*. In any case, juveniles and immature specimens can be difficult to separate (see Guo et al. 1997). In addition, the two species are genetically close enough to hybridise and this would make definitive species level identification difficult. If sorted by DNA barcoding then the groupings would be determined by the species of the mother (because COI is a mitochondrial gene so only inherited maternally). A recent paper by Sui et al. (2009) covering six river systems using six microsatellite loci nevertheless support the species distinction of *E. japonica*, *E. sinensis* and *E. hepuensis* as recognised by Dai (1991), but does hint at possible hybridisation.

The records of *E. hepuensis* and *E. japonica* from coastal southern and central China (~21°40'N to ~32°N) and Taiwan (see cited localities in Xu et al. 2009) need to be re-examined using freshly collected wild specimens. Interestingly, *Eriocheir hepuensis* haplotypes were the only species identified from



Figure 13. Chinese art depicting cooked *Eriocheir sinensis* waiting to be eaten. By Meishuaihaozhe Zhiyou, Courtesy Jiangsu Fine Arts Publishing House.



Figure 14. Chinese mitten crabs for sale at the Turf City Seafood Restaurant, Bukit Timah Road, Singapore. Note usual price for 250gram (or above) male is \$65SD = ca. \$33US. Photograph taken by Peter Ng.

coastal southern China. In coastal central China, both *E. hepuensis* and *E. sinensis* were present. *Eriocheir sinensis* dominates northern China, with only a single *E. hepuensis* mitochondrial haplotype found in the Chiagjiang River (Hu and Wang, unpublished data). Sui et al. (2009) found nuclear DNA evidence of a split between northern and southern coastal China (Figure 1).

The sister clade to that including the Iraqi specimen contains only *E. japonica* from Okinawa (3.0% molecular distance from *E. hepuensis*). Other separate strong clades include: *E. sinensis* (coastal central and northern China from ~24°50'N to 41°10'N, and Korea, Siberia; 4.7% distance), *E. japonica* (Japan, Korea, Siberia; 4.2% distance), *E. ogasawaraensis* (Ogasawara Islands; 6.3% distance), and the more distant clades of *Platyeriocheir formosa* (14.2% distance) and *Neoeriocheir leptognathus* (14.5% distance).

A haplotype network was created using sequences from the *E. hepuensis* clade (Figure 12b). Five geographic areas were defined: Coastal China South (Hepu to Aotou), Coastal China Central (Tongan, Minjiang [see Tang et al. 2003], Feiyunjiang River, Oujiang River), Coastal China North (Chiagjiang River, Hu and Wang unpublished) and Taiwan (Tamshui). There is little obvious geographic structuring between the areas, with haplotypes shared between many areas. The Iraqi haplotype was unique and separated from the common central haplotype by two base pairs. It is not possible to narrow down the native geographic origin of the Iraqi mitochondrial haplotype because there is little geographic structuring among the areas.

Commercial exploitation and hybridisation

Eating the developing mitten crab gonads has been part of Chinese culture for many centuries dating back to the Ming Dynasty (1368-1644), when skilled craftsmen made a set of gold utensils for eating a crab that included a mallet, scissors, a shell cracker, a round salver, scoop, spoon, a long fork and combined scraper and pricker (<http://www.cultural-china.com/chinaWH/html/en/Kaleidoscope2789bye8009.html>).

Mitten crabs have also been a popular subject for beautiful brush-stroke paintings depicting the consumption of this delicacy (Figure 13). Today, mitten crabs command a high price in SE Asian restaurants (e.g., ca. \$40 for a single crab in the right condition, Figure 14) during the autumnal months when they are harvested during their



Figure 15. In season, Chinese mitten crabs tied-up in rice grass can be purchased from street markets in Hong Kong. Photograph taken by Phil Crabb, NHM Photo Unit.




Figure 16. Map indicating rapid spread of *Eriocheir sinensis* throughout NE Europe, early 20th century.

annual migration from freshwater rivers to brackish estuarine water and marketed. Typically while in season, Chinese mitten crabs can be purchased from street markets as in Hong Kong (Figure 15). However, wild Chinese populations have dramatically declined due to over-exploitation, increased demand, river pollution and irrigation schemes that have disrupted the natural migration patterns of this species (Hymanson et al. 1999). But, local and international demands for *E. sinensis* have been met by an intensive aquaculture programme and

this species has been farmed throughout China for the last 40 years (Sui et al. 2009) especially along the Yangtze valley (Jin et al. 2001; Wang et al. 2006). This industry is estimated to be worth ca. US\$ 1.25 billion annually.

Studies on growth rates for the Hepu mitten crab, revealed maximum growth at water temperatures between 23 and 28°C (N.K. Ng, unpublished data). At this range of temperatures, *E. hepuensis* growth is better than that known for *E. sinensis*, and therefore in order to enhance the stock, farmers have cross-bred different genetic populations from various parts of China, including hybridising *E. sinensis* with *E. hepuensis* (see Zhao et al. 1988; Y. Cai, pers. comm., 1999). Although offspring have been produced in these *E. sinensis* and *E. hepuensis* crosses, there are no reports of their viability (Y. Cai, pers. comm. 1999; S. Yang, pers. comm. 2000). Furthermore, some farmers in northern China have also imported southern populations (including *E. hepuensis*) for breeding programmes (A.-Y. Dai, pers. comm. 1998). However, because farmers did not recognise *E. hepuensis* and *E. sinensis* as separate species but only as geographic populations, they did not keep records as to the sources of the breeding pairs. Moreover, the breeders did not check if the F1 generation were fertile. In addition, cross breeding between *E. sinensis* and *E. japonica* has been attempted in South Korea recently (S. H. Ko, pers. comm.), but again there are no reports on the outcome of this genetic experiment. One problem resulting from this practice is that according to S. H. Ko (pers. comm. 2002) there appears to be no 'pure' South Korean populations of *E. sinensis* left in the wild. In Japan, however, no interbreeding experiments of *E. sinensis* and *E. japonica* are known to date (T. Gao 2002; S. Kobayashi, pers. comm. 2010).

As a consequence of cross breeding, the taxonomy of mitten crabs native to East Asia has become even more problematic and controversial. This is why some taxonomists do not recognise *E. hepuensis* as a valid species. Even the validity of species like *E. japonica* and *E. sinensis* has been questioned because the genetic and morphological characters do not always agree. The presumed natural distributions (see Figure 1) of the four known *Eriocheir* species have been discussed at length by Guo et al. (1997) and Komai et al. (2006). Interestingly, *E. hepuensis* is currently considered locally to be an endangered species in China (Cheng et al. 2009).

Dispersal beyond Iraq

The “invasion” and subsequent dispersal of the Chinese mitten crab in Europe has been well documented. *Eriocheir sinensis* was accidentally introduced into Germany during 1912 when a large male was captured from the River Aller, a tributary of the River Weser (Peters and Panning 1933; Panning 1939). During the 1920s and 1930s, the Chinese mitten crab spread rapidly throughout northeast Europe (Figure 16) and beyond (e.g., Harold 1935; Petit 1960; Hahtela 1963; Petit and Mizoule 1974; Ingle and Andrews 1976; Christiansen 1977; Wall and Limbert 1983; Clark 1984; Adema 1991; Zibrowius 1991; Vigneux et al. 1993; Gomoiu and Skolka 1998; Cabral and Costa 1999; Murina and Antonovsky 2001; Zaitsev and Öztürk 2001; Slynko et al. 2002; Herborg et al. 2003; Paunovic et al. 2004; Herborg et al. 2005; Mizzan, 2005; Robbins et al. 2006; Shakirova et al. 2007; Bentley 2011; Berezina et al. 2011; Clark 2011; Jazdżewski and Grabowski 2011). In addition, Chinese mitten crabs have been recorded in various river catchments at considerable distances from the coast: ~1000 km, Yangtze River, China (Schellenberg 1928); ~700-750 km, Elbe River as far as Prague, Czech Republic (Peters 1938); ~460 km, Oder River, Breslau, Poland (Herborg et al. 2003); ~512 km River Rhine Netherlands (Herborg et al. 2003) and ~2000 km, Kuibyshev Reservoir, Volga River, Russia (Shakirova et al. 2007). These data and observations are of concern because if its behaviour is comparable to *E. sinensis*, the Hepu mitten crab could now spread rapidly throughout the extensive watersheds and catchments flowing through Iraq and nearby countries.

The first actual record of a “mitten crab” in Iraq may date back to 1980 when S.D. Salman (pers. comm. in letter to JM Bishop dated 17.04.06) reported possible collection of mitten crabs from the Shatt Al-Arab and later from the “Marshes”, south of Al Fao, Shatt Al-Arab and Shatt Al-Basrah. This suggests that there has already been a long period (over 30 years) of establishment which includes ovigerous material. A rapid expansion of range could be expected as three rivers feed the Shatt Al-Arab waterway in Iraq - the Euphrates, Tigris, and Karun (see Clark et al. 2006). The origins of the Euphrates and the Tigris are in eastern Turkey and both flow to the Persian Gulf via Syria and Iraq. The Euphrates River is approximately 2800 km in length compared with the Tigris River, ca. 1800

km. Rising in the Zagros Mountains, West Iran, the Karun River flows south for about 720 km to meet the Shatt Al-Arab waterway on the Iraqi border (see Figure 3). Due to the catadromy of mitten crab species, all three watersheds would be vulnerable and these distances are within the range of those already reported for *E. sinensis*.

Origins and vectors

The *Eriocheir hepuensis* specimens reported from Kuwait (Figure 3) may just be Iraqi mitten crabs that have moved westward along the coastline or down the Shatt Al-Basrah Canal. As Kuwait has no riverine system, this species is not expected to become established in the sheikhdom. However, with regard to the origin of *E. hepuensis* from Khor Abdullah (Figure 3), it is possible that the species was introduced to the Gulf region by ship traffic. Either adult or subadult specimens may have been transported on ship hulls or larvae with ballast water. The latter mechanism has been identified as the main pathway for dispersal of *Eriocheir* species as well as for other marine invasive species (Cohen and Carlton 1997; Minchin and Gollasch 2002).

Commercial vessel traffic in the Persian Gulf is heavy and increasing. During the 365-day period from May 2006 through April 2007, 44,845 commercial vessels entered the Gulf for a daily average of 123. This represents a 40% increase over the previous year (Capt. A.M. Al-Janahi, Marine Emergency Mutual Aid Centre, Bahrain, pers. comm., 9 Oct. 2007). Many of these vessels were tankers, which are empty and discharge ballast water as they load crude oil. Although records of mitten crabs from the Gulf have only recently been published (Apel and Bishop 2006; Clark et al. 2006), their presence in the upper Gulf’s Shatt Al-Arab area has been known since 1980 (S. Salman, Basrah University, pers. comm., 17 Apr. 2006), and a “very large sample” of mitten crabs was captured by trawl some miles south of Iraq’s Fao Peninsula. Salman also reports capturing mitten crabs in the Iraqi marshes and the Shatt Al-Arab (pers. comm., 17 Apr. 2006). As the world’s demand for crude oil increases, so will tanker traffic, and re-introductions of this species or introductions of other exotic species is likely.

Conclusion

The identification of *Eriocheir hepuensis* from Kuwait and Iraq based on morphological

characters was confirmed by DNA analysis. Furthermore, these molecular data supports the view that mitten crabs comprise three genera and six species i.e., *Eriocheir japonica*, *E. sinensis*, *E. hepuensis* and *E. ogasawaraensis*, *Neoeriocheir leptognathus* and *Platyeriocheir formosa*. However, hybridisation between *E. sinensis*/*E. hepuensis* and *E. japonica*/*E. sinensis* are reported from aquaculture experiments. Furthermore, evidence presented here suggests that *E. hepuensis* may have been present for over thirty years and has now become well established in Iraq. Considering the natural distribution of *E. hepuensis* and its congeners, it appears that the Hepu mitten crab inhabits subtropical and tropical regions, while *E. sinensis* is restricted to more temperate climates (Figure 1). Therefore *E. sinensis* has successfully invaded mainly temperate regions in central and northern Europe and North America and *E. hepuensis* might be much better adapted for the subtropical climate in the Gulf region, thus implying that an even larger area of the world may be at risk of *Eriocheir* invasions.

Acknowledgements

The authors would like to thank two anonymous reviewers for their comments with regard to the original submission of our manuscript.

References

- Adema JPHM (1991) De krabben van Nederland en België (Crustacea, Decapoda, Brachyura). Nationaal Natuurhistorisch Museum, Leiden, The Netherlands, 244 pp
- Apel M, Bishop JM (2006) First record of the non-indigenous Chinese mitten crab *Eriocheir hepuensis* Dai, 1991 from the Arabian Gulf with remarks on potential impacts. – Presentation at the 1st International AEHMS Conference, on the state of the Gulf-Ecosystem: Future and Threats, Al-Ain, UAE, March 5-7 2006
- Bentley MG (2011) The global spread of the Chinese mitten crab *Eriocheir sinensis*. In: Galil BS, Clark PF, Carlton JD (eds), In the wrong place - Alien marine crustaceans: distribution, biology and impacts, invading nature - Springer Series in Invasion Ecology 6, Dordrecht, Springer, pp 107-128, http://dx.doi.org/10.1007/978-94-007-0591-3_3
- Berezina NA, Petryashev VV, Razinkovas A, Lesutienė L (2011) Alien malacostracan crustaceans in the eastern Baltic Sea: pathways and consequences. In: Galil BS, Clark PF, Carlton JD (eds), In the wrong place - Alien marine crustaceans: distribution, biology and impacts, invading nature. Springer Series in Invasion Ecology 6, Dordrecht, Springer, pp 301-322, http://dx.doi.org/10.1007/978-94-007-0591-3_10
- Cabral HN, Costa MJ (1999) On the occurrence of the Chinese mitten crab *Eriocheir sinensis* in Portugal (Decapoda, Brachyura). *Crustaceana* 72: 55-58
- Chan TY, Hung MS, Yu HP (1995) Identity of *Eriocheir recta* (Stimpson, 1858) (Decapoda: Brachyura), with description of a new crab from Taiwan. *Journal of Crustacean Biology* 15: 301-308, <http://dx.doi.org/10.2307/1548957>
- Chan TY, Ng PKL, Ng NK (2005) The nomenclature and taxonomy of *Eriocheir formosa* Chan, Hung and Yu, 1995 (Brachyura, Varunidae) from Taiwan: A rebuttal of Tang et al. (2003, 2004). *Crustaceana* 78: 457-464
- Cheng Q, Wang C, Xu J, Wang J, Yang Q (2009) Novel microsatellite markers for endangered Hepu mitten crab, *Eriocheir hepuensis*. *Conservation Genetics Resources* 1: 357-360, <http://dx.doi.org/10.1007/s12686-009-9082-z>
- Christiansen ME (1977) Kinesisk ullhåndkrabbe funnet for første gang i Norge. *Fauna* 30: 134-138
- Chu KH, Ho HY, Li CP, Chan TY (2003) Molecular phylogenetics of the mitten crab species in *Eriocheir, sensu lato* (Brachyura: Grapsidae). *Journal of Crustacean Biology* 23: 738-746, <http://dx.doi.org/10.1651/C-2347>
- Clark PF (1984) Recent records of alien crabs in Britain. *Naturalist* 109: 111-112
- Clark PF (2011) The commercial exploitation of the Chinese mitten crab, *Eriocheir sinensis* in the River Thames, London: damned if we don't and damned if we do. In: Galil BS, Clark PF, Carlton JD (eds), In the wrong place - Alien marine crustaceans: distribution, biology and impacts, invading nature - Springer Series in Invasion Ecology 6, Dordrecht, Springer, pp 537-580, http://dx.doi.org/10.1007/978-94-007-0591-3_19
- Clark PF, Abdul-Sahib IM, Al-Asadi MS (2006) The first record of *Eriocheir sinensis* H. Milne Edwards, 1853 (Crustacea: Brachyura: Varunidae) from the Basrah Area of Southern Iraq. *Aquatic Invasions* 1: 51-54
- Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. *Molecular Ecology* 9: 1657-1660, <http://dx.doi.org/10.1046/j.1365-294x.2000.01020.x>
- Cohen AN, Carlton JT (1997) Transoceanic transport mechanisms: the introduction of the Chinese mitten crab *Eriocheir sinensis* to California. *Pacific Science* 51: 1-11
- Costa FO, Dewaard JR, Boutillier J, Ratnasingham S, Dooh RT, Hajibabaei M, Hebert PDN (2007) Biological identifications through DNA barcodes: the case of Crustacea. *Canadian Journal of Fisheries and Aquatic Sciences* 64: 272-295, <http://dx.doi.org/10.1139/f07-008>
- Dai A (1991) Studies on the subspecies differentiation of the genus *Eriocheir* (Decapoda: Brachyura). *Scientific Treatise on Systematics and Evolutionary Zoology* 1: 61-71
- Dai A (1993) Systematic and biological studies on *Eriocheir japonica hepuensis* Dai, 1991, International Senckenberg Symposium Crustacea Decapoda. Frankfurt, 18-22 October 1993
- Dai A, Yang S, Song Y, Chen G (1986) Crabs of the China Seas. China Ocean Press, Beijing, People's Republic of China, pp 1-642
- Dai AY, Yang SL (1991) Crabs of the China Seas. 2nd ed. 74 pls. China Ocean Press, Beijing, People's Republic of China, 682 pp
- DAISIE (2009) Handbook of alien species in Europe. In: Invading nature – Springer series in invasion ecology, vol. 3, XXVIII, Dordrecht, Springer, 400 pp
- Dittel AI, Epifanio CE (2009) Invasion biology of the Chinese mitten crab *Eriocheir sinensis*: A brief review. *Journal of Experimental Marine Biology and Ecology* 374: 79-92, <http://dx.doi.org/10.1016/j.jembe.2009.04.012>
- Doi W, Watanabe S, Carlton JT (2011) Alien marine crustaceans of Japan: a preliminary assessment. In: Galil BS, Clark PF, Carlton JD (eds), In the wrong place - Alien marine crustaceans: distribution, biology and impacts, Invading Nature. Springer Series in Invasion Ecology 6, Dordrecht, Springer, pp 419-450, http://dx.doi.org/10.1007/978-94-007-0591-3_15
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of leaf tissue. *Phytochemistry Bulletin* 19: 11-15
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294-299

- Gomoiu M-T, Skolka M (1998) Cresterea biodiversitatii prin imigrare - noi specii in fauna Romaniei / Increase of biodiversity by immigration - new species for the Romanian fauna. "Ovidius" University Annals of Natural Science, Biology-Ecology series 2: 181-202
- GSMFC (2003) Species Summary for *Eriocheir sinensis*. - Gulf States Marine Fisheries Commission, Ocean Springs, Mississippi. http://nis.gsmfc.org/nis_factsheet.php?toc_id=132 (Accessed 29 June 2006)
- Gu J (1986) An ecological survey of *Eriocheir sinensis*. *Transactions of the Chinese Crustacean Society* 1: 268-269
- Guo JY, Ng NK, Dai AY, Ng PKL (1997) The taxonomy of three commercially important species of mitten crabs of the genus *Eriocheir* de Haan, 1835 (Crustacea: Decapoda: Brachyura: Grapsidae). *Raffles Bulletin of Zoology* 45: 445-476
- Haahela I (1963) Some new observations and remarks on the occurrence of the Mitten Crab, *Eriocheir sinensis* Milne Edwards (Crustacea, Decapoda), in Finland. *Aquilo Societas Amicorum Naturae Oulensis* 1: 9-16
- Haan W, de (1835) Crustacea. In: von Siebold PF (ed), Fauna Japonica, sive Descriptio animalium, quae in itinere per Japoniam, jussu et auspiciis superiorum, qui summum in India Batava imperium tenent, suscepto, annis 1823-1830 collegit, notis, observationibus et adumbrationibus illustravit P.F. de Siebold. Conjunctis studiis C.J. Temminck et H. Schlegel pro Vertebratis atque W. de Haan pro Invertebratis elaborata Regis aupicus edita. P. F. v. Siebold. Leiden, Lugundi-Batavorum. Decas II: 25-64, pls 9-15, 17, C, D. (For dates see Sherborn and Jentink, 1895; Holthuis, 1953 and Holthuis and T. Sakai, 1970)
- Harold CHH (1935) Thirtieth Annual Report on the Results of the Chemical and Bacteriological Examination of the London Waters for the 12 months ending 31 December 1935. Metropolitan Water Board London, 101 pp
- Hashim AA (2010) Occurrence of the Chinese mitten crab *Eriocheir sinensis* (H. Milne Edwards) in South Iraq. *Mesopotamian Journal of Marine Science* 25: 31-36
- Herborg LM, Rushton SP, Clare AS, Bentley MG (2003) Spread of the Chinese mitten crab (*Eriocheir sinensis* H. Milne Edwards) in continental Europe: analysis of a historical data set. *Hydrobiologia* 503: 21-28, <http://dx.doi.org/10.1023/B:HYDR.0000008483.63314.3c>
- Herborg L-M, Rushton SP, Clare AS, Bentley MG (2003) Spread of the Chinese mitten crab (*Eriocheir sinensis* H. Milne Edwards) in Continental Europe: analysis of a historical data set. *Hydrobiologia* 503: 21-28, <http://dx.doi.org/10.1023/B:HYDR.0000008483.63314.3c>
- Herborg L-M, Rushton SP, Clare AS, Bentley MG (2005) The invasion of the Chinese mitten crab (*Eriocheir sinensis*) in the United Kingdom and its comparison to Continental Europe. *Biological Invasions* 7: 959-968, <http://dx.doi.org/10.1007/s10530-004-2999-y>
- Holthuis LB (1953) On the dates of publication of W. de Haan's volume on the Crustacea of P.F. von Siebold's "Fauna Japonica". *The Journal of the Society for the Bibliography of Natural History* 3: 36-47, http://dx.doi.org/10.3366/jsbnh.1953.3.Part_1.36
- Holthuis LB, Sakai T (1970) Ph. F. von Siebold and Fauna Japonica. A history of early Japanese Zoology. Tokyo, Academic Press of Japan. I-xviii + part I, 1-132, (in English) + part II, 207-323, (in Japanese) + pls. I-XXXII
- Hong M-S, Ho H-P, Yu H-P (1993) Freshwater mitten crab of Taiwan. In China Fishery Monthly, China Fisheries Association, Taipei, No. 479: 9-20, pls. A and F
- Horwath JL (1989) Final Rule on importation of injurious wildlife: mitten crabs. *Federal Register, Rules and Regulations* 54: 22286-22289
- Hwang JJ, Mizue K (1985) Freshwater crabs of Taiwan. Bulletin of the Faculty of Fishery. *Nagasaki University Japan* 57: 1-22, pls. 1-2
- Hymanson Z, Wang J, Sasaki T (1999) Lessons from the home of the Chinese mitten crab. *IEP Newsletter* 12: 25-32
- Ingle RW, Andrews MJ (1976) Chinese crab reappears in Britain. *Nature*, London 263: 638, <http://dx.doi.org/10.1038/263638a0>
- Jensen GC, Armstrong DA (2004) The occurrence of the Japanese mitten crab, *Eriocheir japonica* (De Haan), on the west coast of North America. *California Fish and Game* 90: 94-99
- Jazdzewski K, Grabowski M (2011) Alien crustaceans along the southern and western Baltic Sea. In: Galil BS, Clark PF, Carlton JD (eds), In the wrong place - Alien marine crustaceans: distribution, biology and impacts, invading nature. Springer Series in Invasion Ecology 6, Dordrecht, Springer, pp 323-344
- Jin G, Li Z, Xie P (2001) The growth patterns of juvenile and precocious Chinese mitten crabs, *Eriocheir sinensis* (Decapoda, Grapsidae), stocked in the freshwater lakes of China. *Crustaceana* 74: 261-273, <http://dx.doi.org/10.1163/156854001505505>
- Kim HS (1973) Anomura and Brachyura. In: Illustrated encyclopedia of fauna and flora of Korea, 14. Samwha Publishing Co., Seoul, Korea, 694 pp
- Komai T, Yamasaki I, Kobayashi S, Yamamoto T, Watanabe S (2006) *Eriocheir ogasawaraensis*, a new species of mitten crab (Crustacea: Decapoda: Brachyura: Varunidae) from Ogasawara Islands, Japan, with notes on the systematics of *Eriocheir* De Haan, 1835. *Zootaxa* 1168: 1-20
- Kumar S, Tamura K, Jakobsen IB, Nei M (2001) MEGA2: Molecular Evolutionary Genetics Analysis software. Arizona State University, Tempe, Arizona, USA
- de Lafontaine Y (2005) First Record of the Chinese mitten crab (*Eriocheir sinensis*) in the St. Lawrence River, Canada. *Journal of Great Lakes Research* 31: 367-370, [http://dx.doi.org/10.1016/S0380-1330\(05\)70267-7](http://dx.doi.org/10.1016/S0380-1330(05)70267-7)
- Lai W, Lu J (1992) Study on the decapod crustacean community in Chiangjiang River estuary. *Transactions of the Chinese Crustacean Society* 3: 23-29
- Li M, Zheng H (2000a) Study on the morphology of *Eriocheir hepuensis*. *Journal of Zhejiang Ocean University* 19: 1-5
- Li M, Zheng H (2000b) The reproductive biology of *Eriocheir hepuensis*. *Journal of Zhejiang Ocean University* 20: 1-5
- Li M, Zheng H (2001) The growth and ecological characteristics of *Eriocheir hepuensis*. *Journal of Zhejiang Ocean University* 21: 1-5
- Linnaeus C (1758) *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis Synonymis, Locis*. Edition 10. Holmiae. 1: iii + 1-824
- Lowe S, Browne M, Boudjelass S, De Poorter M (2004) 100 of the World's Worst Invasive Species Database. Published by The Invasive Species Specialist Group (ISSG), a specialist group of the Species Survival Commission (SSC) of the World Conservation Union (IUCN), 12 pp
- Milne Edwards H (1853) De la famille des ocypodides (Ocypodidae), suite. *Annales des Sciences Naturelles series 3* (Zoology) 20: 163-228, Pls. 6-11
- Milne Edwards H (1854) Notes sur quelques Crustacés nouve-aux ou peu connus conservés dans la collection du Musé-um d'Histoire Naturelle. *Archives du Muséum d'Histoire Naturelle*, Paris 7: 145-188, pls 9-16
- Minchin D, Gollasch S (2002) Vectors – how exotics get around. In: Leppäkoski E, Gollasch S, Olenin S (eds), Invasive aquatic species of Europe: Distribution, impacts and management. Kluwer, Dordrecht, pp 183-192
- Mizzan L (2005) *Rithropanopaeus harrisii* (Gould, 1841) (Crustacea: Decapoda: Panopeidae) and *Eriocheir sinensis* H. Milne Edwards 1854 (Crustacea: Decapoda: Grapsidae): two new exotic crabs in the Venetian Lagoon. *Bollettino del Museo Civico di Storia Naturale di Venezia* 56: 89-95

- Murina VV, Antonovs AG (2001) Chinese crab, *Eriocheir sinensis* is an invader into the basin of the Sea of Azov. *Ekologiya i Moria* 55: 37-39
- Nepszy SJ, Leach JH (1973) First records of the Chinese mitten crab, *Eriocheir sinensis* (Crustacea: Brachyura) from North America. *Journal of the Fisheries Research Board of Canada* 30: 1909-1010, <http://dx.doi.org/10.1139/f73-310>
- Ng NK, Dai AY, Guo J, Ng PKL (1998) The complete larval development of the southern Chinese mitten crab, *Eriocheir hepuensis* Dai, 1991 (Decapoda, Brachyura, Grapsidae) reared under laboratory conditions. *Crustaceana* 71: 493-517, <http://dx.doi.org/10.1163/156854098X00400>
- Ng NK, Guo J, Ng PKL (1999) On the generic affinities of *Eriocheir leptognathus* Rathbun, 1913, and *Eriocheir formosa* Chan, Hung and Yu, 1995, with description of a new genus (Crustacea: Grapsidae: Varuninae). *Journal of Crustacean Biology* 19: 154-170, <http://dx.doi.org/10.2307/1549557>
- Ng PKL (1998) Crabs. In: FAO Species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 2. Cephalopods, crustaceans, holothurians and sharks. In: Carpenter KE and Niem VH (eds), Food and Agriculture Organisation, Rome, pp 1045-1155
- Ng PKL, Guinot D, Davie PJF (2008) Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. *Raffles Bulletin of Zoology*, Supplement 17: 1-286
- Panning A (1933) Die Chinesische Wollhandkrabbe in Deutschland. *Zoologischer Anzeiger*, 104(Supplement): 1-58
- Panning A (1938) Systematisches über *Eriocheir sinensis* H. Milne-Edwards. *Mitteilungen aus dem Hamburgischen zoologischen Museum und Institut* 47: 105-111
- Panning A (1939) The Chinese Mitten Crab. Report of the Board of Regents of the Smithsonian Institution (Washington) [1938] 3508, pp 361-375
- Paunovic M, Cacic P, Hegedis A, Kolarevic J, Lenhardt M (2004) A report of *Eriocheir sinensis* (H. Milne Edwards, 1854) [Crustacea: Brachyura: Grapsidae] from the Serbian part of the Danube River. *Hydrobiologia* 529: 275-277, <http://dx.doi.org/10.1007/s10750-004-5493-8>
- Peng W (1986) Preliminary study on the problem of variation of *Eriocheir sinensis* in Zhujiang river valley. *Information of Fisheries Science and Technology* 2: 19-22
- Peters N, Panning A (1933) Die Chinesische Wollhandkrabbe (*Eriocheir sinensis* H. Milne Edwards) in Deutschland. *Zoologischer Anzeiger* 104: 1-180
- Peters N (1938) Ausbreitung und Verbreitung der chinesischen Wollhandkrabbe (*Eriocheir sinensis* H.M.-Edw) in Europa im Jahre 1933 bis 1935. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 47: 1-31
- Petit G (1960) Le crabe chinois est parvenu en Méditerranée. *Vie et Milieu* 11: 133-136
- Petit G, Mizoule R (1974) En douze ans le 'crabe Chinois' n'a pu réussir son implantation dans les lagunes du Languedoc. *Vie et Milieu* 23: 181-186
- Rathbun MJ (1913) Description of new species of crabs of the families Grapsidae and Ocypodidae. *Proceedings of the United States National Museum* 46: 353-358, <http://dx.doi.org/10.5479/si.00963801.46-2030.353>
- Robbins RS, Sakari M, Nezami Baluchi S, Clark PF (2006) The occurrence of *Eriocheir sinensis* H. Milne Edwards, 1853 (Crustacea: Brachyura: Varunidae) from the Caspian Sea region, Iran. *Aquatic Invasions* 1: 32-34, Figs. 1-2
- Ruiz GM, Fegley L, Fofonoff P, Cheng Y, Lemaitre R (2006) First records of *Eriocheir sinensis* H. Milne Edwards, 1853 (Crustacea: Brachyura: Varunidae) for Chesapeake Bay and the mid-Atlantic coast of North America. *Aquatic Invasions* 1: 137-142, <http://dx.doi.org/10.3391/ai.2006.1.3.7>
- Sakai T (1983) Description of new genera and species of Japanese crabs, together with systematically and biogeographically interesting species (I). *Researches on Crustacea* 12: 1-23
- Schellenberg A (1928) Krebstiere oder Crustacea. II. Decapoda, Zehnfüßer. Die Tierwelt Deutschlands und der angrenzenden Meeressteile. Jena, Verlag Gustav Fischer, 10: iv + 146
- Schmidt RE, Daniels RA, Swift EL, Shadis IB (2009) Inferences on the biology of juvenile Chinese mitten crab (*Eriocheir sinensis*) from exuviae in a Hudson River tributary, New York, USA. *Aquatic Invasions* 4: 613-617, <http://dx.doi.org/10.3391/ai.2009.4.4.7>
- Shen CJ, Dai AY (1964) Illustration of animals in China, Crustacea 2. Beijing, People's Republic of China, 172 pp
- Shakirova FM, Panov VE, Clark PF (2007) New records of the Chinese mitten crab, *Eriocheir sinensis* H. Milne Edwards, 1853, from the Volga River, Russia. *Aquatic Invasions* 2: 169-173, <http://dx.doi.org/10.3391/ai.2007.2.3.3>
- Sherborn CD, Jentink JA (1895) On the dates of the Parts of Siebold's 'Fauna Japonica' and Giebel's 'Allgemeine Zoologie' (first edition). *Proceedings of the Zoological Society of London* 1895: 149-150
- Slyenko YV, Korneva LG, Rivier IK, Shcherbina KH, Papchenkov VG, Orlova MI, Theriault TW (2002) Caspian-Volga-Baltic invasion corridor. In: Leppakoski E, Olenin S, Gollasch S (eds), Invasive Aquatic Species of Europe. Distribution, Impacts and Management, Kluwer Academic Publishers, Dordrecht, pp 339-411
- Sui L, Zhang F, Wang X, Bossier P, Sorgeloos P, Hänfling B (2009) Genetic diversity and population structure of the Chinese mitten crab, *Eriocheir sinensis*, in its native range. *Marine Biology* 156: 1573-1585, <http://dx.doi.org/10.1007/s00227-009-1193-2>
- Sun H, Zhou K, Yang X (2003) Phylogenetic relationships of the mitten crabs inferred from mitochondrial 16S rDNA partial sequences (Crustacea, Decapoda). *Acta Zoologica Sinica* 49: 592-599
- Sun H, Wang G, Zhan D, Zhou K (2005) Mitochondrial DNA sequence variation in two geographical subspecies of the mitten crab, *Eriocheir japonica*. *Acta Zoologica Sinica* 51: 862-866
- Swofford DL (2002) PAUP * version 4.0b5 - Phylogenetic Analysis Using Parsimony (* and other methods). Sinauer Associates, Sunderland, Massachusetts, USA
- Tang B, Zhou K, Song D (2004) A neotype of *Eriocheir recta* (Crustacea, Decapoda). *Acta Zootaxa Sinica* 29: 255-259
- Tang B, Zhou K, Song S, Yang S, Dai A (2003) Molecular systematics of the Asian mitten crabs, genus *Eriocheir* (Crustacea: Brachyura). *Molecular and Phylogenetic Evolution* 29: 309-316, [http://dx.doi.org/10.1016/S1055-7903\(03\)00112-X](http://dx.doi.org/10.1016/S1055-7903(03)00112-X)
- Vigneux E, Keith P, Noël P (eds) (1993) Atlas préliminaire des Crustacés Décapodes d'eau douce de France. Coll. Patrimoines Naturels. 14, Paris: Secrétariat de la Faune et de la Flore, Laboratoire de Biologie des Invertébrés Marins et Malacologie Muséum National d'Histoire Naturelle, Conseil Supérieur de la Pêche, Ministère de l'Environnement, pp i-vi, 1-56, <http://dx.doi.org/10.1016/j.ympcv.2009.02.007>
- Xu JW, Chan TY, Tsang LM, Chu KH (2009) Phylogeography of the mitten crab *Eriocheir sensu stricto* in East Asia: Pleistocene isolation, population expansion and secondary contact. *Molecular Phylogenetics and Evolution* 52: 45-56
- Wall C, Limbert M (1983) A Yorkshire record of the Chinese Mitten Crab. *Naturalist* 108: 147
- Wang HZ, Wang HJ, Liang XM, Cui FB (2006) Stocking models of Chinese mitten crabs (*Eriocheir japonica sinensis*) in Yangtze lakes. *Aquaculture* 255: 456-465, <http://dx.doi.org/10.1016/j.aquaculture.2006.01.005>
- Yu HP, Ho PH (1986) Notes on the freshwater crabs of the genus *Eriocheir* (Crustacea, Decapoda, Grapsidae) from Taiwan. *Annals of Taiwan Museum* 29: 111-116
- Zaitsev Y, Öztürk B (eds) (2001) Exotic species in the Aegean, Marmara, Black, Azov and Caspian Seas. Turkish Marine Research Foundation, Istanbul, Turkey, 265 pp

Zhao N, Du N, Bao X, Zhang L (1988) Artificial Breeding, Propagation and Culture of Chinese Mitten Crab. Anhui Science and Technology Press, Hefei, People's Republic of China, pp 134-136

Zibrowius H (1991) On going modification of the Mediterranean marine fauna and flora by the establishment of exotic species. *Bulletin du Muséum d'Histoire Naturelle de Marseille* 51: 83-107

Appendix 1. Other *Eriocheir* COI sequences downloaded from GenBank (<99% similar to Iraqi specimen).

| Species in original publication | GenBank accession number | Species in original publication | GenBank accession number |
|---------------------------------|--------------------------|---------------------------------|--------------------------|
| <i>E. formosa</i> | AF105249 | <i>E. sinensis</i> | AF105247 |
| | AF317326 | | AF105248 |
| | AF516698 | | AF279269 |
| | F105250 | | AF317333 |
| <i>E. japonica</i> | FJ750332 | | AF317335 |
| | AF317330 | | AF435113 |
| | AF317331 | | AF435114 |
| | AF516700 | | AF435115 |
| | AY640095 | | AF435116 |
| | AY640096 | | AF435117 |
| | AY640097 | | AF435118 |
| | AY640098 | | AF435119 |
| | AY640099 | | AF516702 |
| | AY640100 | | AY640082 |
| | AY640101 | | AY640083 |
| | AY640102 | | AY640084 |
| | FJ455505 | | AY640085 |
| | FJ750312 | | AY640086 |
| | FJ750313 | | AY640087 |
| | FJ750314 | | DQ438943 |
| | FJ750315 | | DQ882062 |
| FJ750316 | | FJ750306 | |
| FJ750317 | | FJ750307 | |
| FJ750318 | | FJ750308 | |
| FJ750319 | | FJ750309 | |
| <i>E. japonica</i> (Okinawa) | AY640092 | | FJ750310 |
| | AY640093 | | FJ750311 |
| | AY640094 | | HM640253 |
| | FJ750329 | | HM640254 |
| <i>E. leptognathus</i> | AF516701 | | HM640255 |
| <i>E. ogasawaraensis</i> | FJ750330 | | HM640256 |
| | FJ750331 | | HM640257 |
| <i>E. recta</i> | AF317332 | | HQ534046 |
| | | | HQ534047 |
| | | | HQ534048 |
| | | | HQ534049 |