

(±)-1-hydroxy-6,6-dimethyl-1H,6H-naphtho[1,2-c]furan-3,9-trione

Author

Henderson, LC, Loughlin, WA, Jenkins, ID, Healy, PC

Published

2006

Journal Title

Acta crystallographica. Section E, Structure reports online

DOI

[10.1107/S1600536806009172](https://doi.org/10.1107/S1600536806009172)

Rights statement

© The Author(s) 2006. For information about this journal please refer to the journal's website. All articles published in Acta Crystallographica Section E are open access and distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. See <http://creativecommons.org/licenses/by/2.0/uk/legalcode>

Downloaded from

<http://hdl.handle.net/10072/14339>

Link to published version

<http://journals.iucr.org/e/journalhomepage.html>

Griffith Research Online

<https://research-repository.griffith.edu.au>


Luke C. Henderson,^a Wendy A. Loughlin,^b Ian D. Jenkins^a and Peter C. Healy^{b*}^aNatural Product Discovery and the Eskitis Institute, Griffith University, Nathan, Brisbane 4111, Australia, and ^bSchool of Science and the Eskitis Institute, Griffith University, Nathan, Brisbane 4111, Australia

Correspondence e-mail: p.healy@griffith.edu.au

Key indicators

Single-crystal X-ray study
 $T = 295\text{ K}$
Mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$
Disorder in main residue
 R factor = 0.060
 wR factor = 0.179
Data-to-parameter ratio = 12.9For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.**(±)-1-Hydroxy-6,6-dimethyl-1*H*,6*H*-naphtho-[1,2-*c*]furan-3,9-trione**The title compound, $\text{C}_{14}\text{H}_{12}\text{O}_4$, crystallizes as discrete molecular species which form hydroxy-to-ketone hydrogen-bonded dimers disposed about crystallographic centres of symmetry.

Comment

The 1-hydroxyphthalide unit in the title compound, (II), is present in several natural products which display biological activity. For example, madurahydroxylactone (Jutten *et al.*, 2002) and its derivatives exhibit activity as esterone sulfatase inhibitors. 1-Hydroxyphthalides also have been used as precursors for the synthesis of inhibitors of platelet aggregation (Sugimoto *et al.*, 1984) and GABA_B receptor antagonists (Donati *et al.*, 1989). 1-Hydroxyphthalides with a γ -ketone, as in compound (II), occur in natural product metabolites of *aspergillus duricaulis* (Achenbach *et al.*, 1985) and the basidiomycete *Hyphoderma radula* (Henkel *et al.*, 1997). We report here the structural elucidation of (II), prepared by regioselective reduction of anhydride (I) previously obtained from 4,4-dimethylcyclohexane-1,3-dione (Henderson *et al.*, 2006).Compound (II) crystallizes in the space group $P2_1/c$ as discrete molecular species and is isomorphous with 1-hydroxy-6,6-dimethyl-7,8-dihydronaphtho[1,2-*c*]furan-3,9(1*H*,6*H*)-dione (hyphodermin B) (Henderson *et al.*, 2006). All three rings and carbonyl atom O3 are coplanar. In the structure of hyphodermin B, the cyclohexyl ring is disordered with C7 modelled as two C atoms with 50% occupancy above and below the plane. The methyl groups on C6 lie above and below this plane with the Fourier synthesis showing an eclipsed conformation for the H atoms on these two groups. Carbonyl atom O4 is twisted slightly out of the plane of the molecule, the pseudo-torsion angle $\text{O4}-\text{C9}\cdots\text{C9b}-\text{C1}$ being $8.7(3)^\circ$. The geometry of the 1-hydroxyphthalimide compounds (Valente *et al.*, 1998; Khoo & Hazell, 1999; Paulus *et al.*, 1994). In these two structures and in (II), the molecules form $R_2^2(14)$ (Bernstein *et al.*, 1995) O—H \cdots O hydroxy-to-ketone hydrogen-bonded dimers about a crystallographic centre of symmetry (Table 2 and Fig. 1). This hydroxy-to-Received 14 February 2006
Accepted 10 March 2006




Figure 1

View of the major component of the molecule of (II), shown in its hydrogen-bonded dimer. The symmetry code of the primed atoms (') is $2 - x, 2 - y, 1 - z$. Displacement ellipsoids for non-H atoms are drawn at the 30% probability level and H atoms are shown as circles of arbitrary radii.

ketone dimerization mode is rare, with only six examples previously reported (Rath *et al.*, 2005, and references therein).

Experimental

Lithium tri-*tert*-butoxyaluminumhydride (165 mg, 0.64 mmol) was added to tetrahydrofuran (THF, 10 ml) at room temperature and stirred vigorously. The solution was then cooled to 273 K, anhydride (I) (Henderson *et al.*, 2006) (155 mg, 0.64 mmol) was added and the solution stirred for 4 h at 273 K. Ammonium chloride (5 ml) and HCl (1 M, 15 ml) were added and the aqueous phase extracted with dichloromethane (DCM, 3×25 ml). The combined organic phases were washed with brine (2×30 ml), dried (MgSO_4) and the solvent removed *in vacuo* to give a brown solid (143 mg). Purification by silica-gel chromatography (ethyl acetate-hexane, 1:1) gave compound (II) (38 mg, 24%). Slow evaporation of a chloroform solution gave pale-orange crystals (m.p. 452–453 K); λ_{max} (KBr/ cm^{-1}) 3421 (*m, br*), 2970 (*w, br*), 1767 (*s*), 1655 (*s*), 1607 (*m*); δ_{H} (200 MHz, CDCl_3) 1.57 (6H, *s*, $2 \times \text{CH}_3$), 6.48 (1H, *d*, J 10.2, H7), 7.04 (1H, *s*, H1), 7.08 (1H, *d*, J 10.2, H8), 7.84 (1H, *d*, J 8, H5), 8.08 (1H, *d*, J 8.4, H4); δ_{C} (100 MHz, CDCl_3) 185.7 (C9), 167.9 (C3), 159.2 (C7), 157.0 (C5a), 148.3 (C9b), 132.0 (C9a), 130.0 (C4), 129.1 (C5), 126.7 (C3a), 126.5 (C8), 97.5 (C1), 39.0 (C6), 29.9, 29.4 ($2 \times \text{CH}_3$); (ESMS, $-ve$) 243, (M^- , 100%), (ESMS, $+ve$) 245 (MH^+ , 30%), 267 (MNa^+ , 100%), 251 (MLi^+ , 100%); HRMS calculated $\text{C}_{14}\text{H}_{11}\text{O}_4$ 243.0657, found 243.06564.

Crystal data

$\text{C}_{14}\text{H}_{12}\text{O}_4$
 $M_r = 244.24$
 Monoclinic, $P2_1/c$
 $a = 10.4825$ (13) Å
 $b = 7.0735$ (17) Å
 $c = 16.081$ (3) Å
 $\beta = 90.556$ (12)°
 $V = 1192.3$ (4) Å³
 $Z = 4$

$D_x = 1.361$ Mg m^{-3}
 Mo $K\alpha$ radiation
 Cell parameters from 25 reflections
 $\theta = 12.7$ – 17.4 °
 $\mu = 0.10$ mm^{-1}
 $T = 295$ K
 Plate, pale orange
 $0.45 \times 0.40 \times 0.15$ mm

Data collection

Rigaku AFC-7R diffractometer
 ω - 2θ scans
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\text{min}} = 0.956$, $T_{\text{max}} = 0.985$
 2395 measured reflections
 2100 independent reflections
 1441 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.018$
 $\theta_{\text{max}} = 25.0$ °
 $h = -12 \rightarrow 12$
 $k = -8 \rightarrow 0$
 $l = -9 \rightarrow 19$
 3 standard reflections
 every 150 reflections
 intensity decay: 0.3%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.179$
 $S = 1.05$
 2100 reflections
 163 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0794P)^2 + 0.8739P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.60$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

O1—C1	1.468 (4)	O4—C9	1.231 (4)
O1—C3	1.359 (4)	C1—C9B	1.508 (4)
O2A—C1	1.343 (4)	C3—C3A	1.474 (4)
O2B—C1	1.3404	C3A—C9B	1.362 (4)
O3—C3	1.190 (4)	C7—C8	1.311 (5)
C1—O1—C3	111.1 (2)	O1—C3—O3	122.1 (3)
O1—C1—C9B	103.3 (3)	O3—C3—C3A	130.5 (3)
O2A—C1—C9B	112.3 (3)	C3—C3A—C9B	109.5 (3)
O2B—C1—C9B	110.98	C6—C7—C8	125.9 (3)
O1—C1—O2B	111.20	C7—C8—C9	121.8 (3)
O1—C1—O2A	107.8 (3)	O4—C9—C8	122.0 (3)
O1—C3—C3A	107.4 (3)	O4—C9—C9A	120.4 (3)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O2A—H2A \cdots O4 ⁱ	0.95	1.81	2.709 (4)	156

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

Significant residual electron density in the vicinity of the H atom bonded to C1 suggested the presence of minor enantiomeric disorder of the hydroxy group in the crystal structure. This was modelled with occupancy factors of 0.9 for the major component and 0.1 for the minor component. The carbon-bound H atoms were constrained as riding atoms, with $C-H = 0.95$ Å. $U_{\text{iso}}(\text{H})$ values were set at $1.2U_{\text{eq}}$ of the parent atom. The hydroxy H atom of the major component was located in a difference Fourier synthesis and constrained with $O-H = 0.95$ Å. The O and H atoms of the minor group were constrained with $C-O = 1.34$ Å and $O-H = 0.95$ Å.

Data collection: *MSC/AFC7 Diffractometer Control Software* (Molecular Structure Corporation, 1999); cell refinement: *MSC/AFC7 Diffractometer Control Software*; data reduction: *TEXSAN* for Windows (Molecular Structure Corporation, 2001); program(s) used to solve structure: *TEXSAN* for Windows; program(s) used to refine structure: *TEXSAN* for Windows and *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *TEXSAN* for Windows and *PLATON* (Spek, 2003).

We acknowledge financial support of this work by Griffith University and the Eskitis Institute for Cell and Molecular Therapies.

References

- Achenbach, H., Muehlenfeld, A. & Brillinger, G. U. (1985). *Liebigs Ann. Chem.* pp. 1596–1628.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Donati, C., Prager, R. H. & Weber, B. (1989). *Aust. J. Chem.* **42**, 787–795.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Henderson, L. C., Loughlin, W. A. & Jenkins, I. D., Healy, P. C. & Campitelli, M. R. (2006). *J. Org. Chem.* **71**, 2384–2388.
- Henkel, T., Mueller, H., Schmidt, D. & Wollweber, H. (1997). Ger. Offen. DE 19611366 A1 19970925 CAN 127:260060 AN 1997:651311 CAPLUS.
- Jutten, P., Schumann, W., Hartl, A., Heinisch, L., Grafe, U., Werner, W. & Ulbricht, H. Bioorg. (2002). *Med. Chem. Lett.* **12**, 1339–1342.
- Khoo, E. K. & Hazell, A. (1999). *Acta Cryst.* **C55**, 2070–2073.
- Molecular Structure Corporation (1999). *MSC/AF7 Diffractometer Control Software for Windows*. Version 1.02. MSC, 9009 New Trails Drive, The Woodlands, TX 77381, USA.
- Molecular Structure Corporation (2001). *TEXSAN for Windows*. Version 1.06. MSC, 9009 New Trails Drive, The Woodlands, TX 77381-5209, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Paulus, E. F., Domberger, K., Werner, W. & Fenske, D. (1994). *Acta Cryst.* **C50**, 2064–2067.
- Rath, N. P., Haq, W. & Balendiran, G. K. (2005). *Acta Cryst.* **C61**, o81–o84.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Sugimoto, A., Tanaka, H., Eguchi, Y., Ito, S., Takashima, Y. & Ishikawa, M. (1984). *J. Med. Chem.* **27**, 1300–1305.
- Valente, E. J., Fuller, J. F. & Ball, J. D. (1998). *Acta Cryst.* **B54**, 162–173.