

Effects of biochar application on soil greenhouse gas fluxes: A meta-analysis

Author

He, Yanghui, Zhou, Xuhui, Jiang, Liling, Li, Ming, Du, Zhenggang, Zhou, Guiyao, Shao, Junjong, Wang, Xihua, Xu, Zhihong, Bai, Shahla Hosseini, Wallace, Helen, Xu, Chengyuan

Published

2017

Journal Title

Global Change Biology Bioenergy

Version

Version of Record (VoR)

DOI

[10.1111/gcbb.12376](https://doi.org/10.1111/gcbb.12376)

Rights statement

© 2016 The Authors. Global Change Biology Bioenergy Published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Downloaded from

<http://hdl.handle.net/10072/123907>

Griffith Research Online

<https://research-repository.griffith.edu.au>

Effects of biochar application on soil greenhouse gas fluxes: a meta-analysis

YANGHUI HE¹, XUHUI ZHOU^{2,3}, LILING JIANG¹, MING LI¹, ZHENGANG DU², GUIYAO ZHOU², JUNJIONG SHAO^{2,3}, XIHUA WANG², ZHIHONG XU⁴, SHAHLA HOSSEINI BAI⁵, HELEN WALLACE⁵ and CHENGYUAN XU⁶

¹Coastal Ecosystems Research Station of Yangtze River Estuary, Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, The Institute of Biodiversity Science, School of Life Sciences, Fudan University, 2005 Songhu Road, Shanghai 200433, China, ²Tiantong National Field Observation Station for Forest Ecosystem, School of Ecological and Environmental Sciences, East China Normal University, Shanghai 200241, China, ³Center for Global Change and Ecological Forecasting, East China Normal University, Shanghai 200062, China, ⁴Environmental Futures Research Institute, School of Natural Sciences, Griffith University, Nathan, Brisbane, Qld 4111, Australia, ⁵Faculty of Science, Health, Education and Engineering, University of the Sunshine Coast, Maroochydore, DC, Qld 4558, Australia, ⁶School of Medical and Applied Sciences, Central Queensland University, Bundaberg, Qld 4670, Australia

Abstract

Biochar application to soils may increase carbon (C) sequestration due to the inputs of recalcitrant organic C. However, the effects of biochar application on the soil greenhouse gas (GHG) fluxes appear variable among many case studies; therefore, the efficacy of biochar as a carbon sequestration agent for climate change mitigation remains uncertain. We performed a meta-analysis of 91 published papers with 552 paired comparisons to obtain a central tendency of three main GHG fluxes (i.e., CO₂, CH₄, and N₂O) in response to biochar application. Our results showed that biochar application significantly increased soil CO₂ fluxes by 22.14%, but decreased N₂O fluxes by 30.92% and did not affect CH₄ fluxes. As a consequence, biochar application may significantly contribute to an increased global warming potential (GWP) of total soil GHG fluxes due to the large stimulation of CO₂ fluxes. However, soil CO₂ fluxes were suppressed when biochar was added to fertilized soils, indicating that biochar application is unlikely to stimulate CO₂ fluxes in the agriculture sector, in which N fertilizer inputs are common. Responses of soil GHG fluxes mainly varied with biochar feedstock source and soil texture and the pyrolysis temperature of biochar. Soil and biochar pH, biochar applied rate, and latitude also influence soil GHG fluxes, but to a more limited extent. Our findings provide a scientific basis for developing more rational strategies toward widespread adoption of biochar as a soil amendment for climate change mitigation.

Keywords: biochar, carbon dioxide, global warming potential, methane, nitrous oxide, soil greenhouse gas

Received 16 March 2016; accepted 22 April 2016

Introduction

The global average surface temperature has increased by 0.85 °C over the period 1880–2012 based on multiple independently produced datasets, and current projections suggest that the temperature is likely to increase by another 0.3–4.8 °C by the end of this century (IPCC, 2013). Global warming is mostly attributable to the increasing atmospheric concentrations of greenhouse gases (GHGs) due to human activities. The three main GHGs (i.e., CO₂, CH₄, and N₂O) in combination contribute to more than 90% of anthropogenic climate warming (Hansen *et al.*, 2000; IPCC, 2013).

Greenhouse gas mitigation strategies include reducing and avoiding emissions as well as enhancing the

removal of GHGs from the atmosphere (Smith *et al.*, 2008). Soil carbon (C) sequestration through biochar amendment has been proposed as an effective countermeasure for the rising concentration of atmospheric GHGs (Lal, 1999; Pan *et al.*, 2004; Smith *et al.*, 2008). Biochar is a carbon-rich, charcoal-like product produced by burning biomass in the absence of oxygen (Lehmann, 2007b; Laird *et al.*, 2009); it contains a high proportion of recalcitrant organic C and is stable for hundreds to thousands of years after it is applied to soil (Schmidt *et al.*, 2002). Biochar application to soils has the potential to mitigate global warming via soil C sequestration, and provide other benefits, such as improving soil fertility, retaining soil moisture, and increasing crop yields (Marris, 2006; Lehmann, 2007a; Laird, 2008; Woolf *et al.*, 2010; Mukherjee *et al.*, 2014; Reverchon *et al.*, 2014; Bai *et al.*, 2015a,b; Xu *et al.*, 2015a,b; Darby *et al.*, 2016).

Correspondence: Xuhui Zhou, tel. +86 21 54341275, fax +86 21 54341275, e-mail: xhzhou@des.ecnu.edu.cn

However, the precise effects of biochar application on soil GHG emissions remain controversial and appear very variable among many case studies (Cayuela *et al.*, 2014; Lorenz & Lal, 2014). Soil CO₂, CH₄, and N₂O fluxes increased significantly in some studies (Yanai *et al.*, 2007; Van Zwieten *et al.*, 2010; Jones *et al.*, 2011; Wang *et al.*, 2012), but substantially decreased or remained unchanged in others (Rogovska *et al.*, 2011; Feng *et al.*, 2012; Zheng *et al.*, 2012; Case *et al.*, 2014; Quin *et al.*, 2015). For example, a field trial in paddy soils amended with biochar produced from wheat straw induced a 12% increase in CO₂ emissions, but a 41.8% decrease in N₂O emissions (Zhang *et al.*, 2012b). Another field experiment in pasture showed no significant effects of biochar amendment on soil CO₂ and N₂O emissions in a pasture ecosystem (Scheer *et al.*, 2011). Thus, the efficacy of biochar for climate change mitigation is largely uncertain due to these variable effects on soil GHG emissions.

There are many hypotheses to explain why biochar may increase or decrease soil GHG fluxes. For example, increases in soil CO₂ emissions induced by biochar might be due to the labile C input and positive priming effects of biochar as well as increased belowground net primary productivity (BNPP) (Zimmerman *et al.*, 2011; Zhang *et al.*, 2012a), while the suppression of soil CO₂ emissions may be due to reduced enzymatic activity and the precipitation of CO₂ onto the biochar surface (Case *et al.*, 2014). Elevated CH₄ emissions could be attributed to the inhibitory effect of chemicals in the biochar on soil methanotrophs (Spokas, 2010). Reduced CH₄ emissions might be associated with decreased ratios of methanogenic archaea to methanotrophic proteobacteria, as the increase in oxygen supply due to biochar application supports a group of aerobic methanotrophs (Feng *et al.*, 2012).

There are also contradictory reports with respect to N₂O emissions. For example, increases in N₂O emissions may be ascribed to biochar-induced increases in soil water content, which favors denitrification, or the release of biochar embodied-N (Lorenz & Lal, 2014). In contrast, mechanisms that explain decreased N₂O emissions include (1) improved soil aeration, (2) increased soil pH, (3) enhanced N immobilization, and (4) a toxic effect induced by biochar organic compounds (polycyclic aromatic hydrocarbons) on nitrifier and denitrifier communities (Clough *et al.*, 2010, 2013; Taghizadeh-Toosi *et al.*, 2011; Hale *et al.*, 2012).

The contradictory reports of changes in size and even direction of soil GHG emissions when biochar is applied, and the diversity of mechanisms proposed, suggest that biochar effects may depend on many factors, including soil properties, experimental methods, artificial cultivation management, biochar application

rate, and biochar physicochemical properties (Hilscher & Knicker, 2011; Lorenz & Lal, 2014). These factors may determine to what extent biochar alters soil C and N transformation processes and consequently soil GHG emissions. However, how these factors contribute to the variable responses of soil GHG emissions to biochar application across the globe still remains unclear. If these factors are not adequately addressed, the effects of biochar application on mitigating global warming cannot be fully understood.

Recently, three meta-analyses on the effects of biochar application on soil GHG fluxes have been conducted. Two of them (i.e., Cayuela *et al.*, 2014, 2015) only emphasized the central tendency of soil N₂O fluxes under biochar addition, and the other by Liu *et al.* (2016) examined the response of CO₂ fluxes, soil organic C (SOC), and soil microbial biomass C (MBC) to biochar amendment. However, there is limited information on the simultaneous effects of biochar amendments on soil GHG fluxes and their global warming potential (GWP). It is necessary to compile all available data to synthesize results from individual studies to reveal the patterns of biochar-induced changes in soil GHG fluxes and to identify the major drivers for responses of GHG fluxes to biochar addition.

In this study, we compiled data from individual experimental studies that quantified the effect of soil biochar application on GHG fluxes across various ecosystems and then quantitatively evaluated the responses of soil CO₂, CH₄, and N₂O fluxes to biochar application under different environmental and experimental conditions using meta-analysis techniques. Our objectives were to (1) quantify the effect size of biochar amendment on soil GHG fluxes across studies; (2) examine whether environmental conditions, experimental methods, and biochar characteristics would influence the responses of soil GHG fluxes to biochar application; and (3) evaluate the response of GWP of soil GHGs to biochar application.

Materials and methods

Data sources

Publications were searched using Web of Science (1900–2015) with the following search terms: (biochar or black carbon or charcoal) and [soil greenhouse gases (GHGs) or CO₂ or CH₄ or N₂O or global warming potential (GWP)]. The selection criteria were as follows: (i) Experiments had at least one pair of data (control and treatment) and measured soil CO₂, CH₄, or N₂O fluxes; (ii) the method of biochar application was clearly described, including experimental duration, amount of biochar application, physico-chemical characteristics of biochar, and soil properties such as pH and C/N ratio; (iii) the means, standard deviations/errors, and sample sizes of variables in the

control and treatment groups could be extracted directly from tables, graphs, or contexts. In total, 91 research papers on biochar application were selected from more than 2000 published papers. The geographic distribution of the selected studies over the world is presented in Fig. 1. The studies contained multiple biochar application levels (Case *et al.*, 2012; Stewart *et al.*, 2013), biochar types (Spokas & Reicosky, 2009; Ameloot *et al.*, 2013), soil types (Wang *et al.*, 2011; Gomez *et al.*, 2014), or N fertilization levels (Barbosa De Sousa *et al.*, 2014; Sun *et al.*, 2014), which were treated as multiple independent studies.

Four categories of data were extracted from the literature of biochar application experiments: (1) soil GHG fluxes, including CO₂, CH₄, and N₂O fluxes; (2) soil properties, including pH, total C, total N, and C/N ratio; (3) biochar properties, including biochar feedstock types, pyrolysis temperature, rate of biochar applied, pH, total C, total N, and C/N ratio; and (4) other auxiliary variables, including latitude, longitude, experiment types (field, pot, and incubation), experimental duration, and N fertilization (whether or not). The variables listed in categories (2), (3), and (4) were used as explanatory factors (either categorical or continuous) of the variation in GHG fluxes in response to biochar application.

Analysis

We followed the methods used by Hedges *et al.* (1999) and Luo *et al.* (2006) to evaluate the responses of soil CO₂, CH₄, and N₂O fluxes to biochar application. A response ratio (RR, natural log of the ratio of the mean value of a variable in biochar treatment plots to that in control) was used to calculate effect sizes as below:

$$RR = \ln \frac{X_t}{X_c} = \ln(X_t) - \ln(X_c) \quad (1)$$

where X_t and X_c are means in the treatment and control groups, respectively. The variance (v) of each individual RR is estimated as:

$$v = \frac{S_t^2}{n_t X_t^2} + \frac{S_c^2}{n_c X_c^2} \quad (2)$$

where n_t and n_c are the sample sizes of the variable in treatment and control groups, respectively; S_t and S_c are the standard deviations for the treatment and control groups.

The mean response ratio (RR₊₊) was calculated from RR of individual pairwise comparisons between treatment and control as below,

$$RR_{++} = \frac{\sum_{i=1}^m \sum_{j=1}^k W_{ij} RR_{ij}}{\sum_{i=1}^m \sum_{j=1}^k W_{ij}} \quad (3)$$

where m is the number of groups and k is the number of comparisons in the i th group. The reciprocal of its variance (v) was considered as the weight (W) of each RR.

We used a bootstrapping method to obtain the 2.5% and 97.5% percentiles as the lower and upper limits of our 95% bootstrap confidence interval (CI) based on 5000 iterations (Adams *et al.*, 1997; Zhou *et al.*, 2014). When the 95% CI of RR₊₊ for soil GHG emissions overlapped with zero, biochar application had no significant impact on the variable. Otherwise, the biochar-induced response was considered as significance (Luo *et al.*, 2006). The percentage change of variables was calculated on the basis of $[\exp(RR_{++}) - 1]100\%$. The frequency distribution of the individual response ratio (RR) was tested by a normal test and fitted by a Gaussian function using Eqn (5) in SIGMAPLOT software (Systat Software Inc., San Jose, CA, USA).

$$y = \alpha \exp \left[-\frac{(x - \mu)^2}{2\sigma^2} \right] \quad (4)$$




Fig. 1 Global distribution of 137 study sites selected in this meta-analysis. Letters C, M, and N represent the sites with CO₂, CH₄, and N₂O measurements, respectively.

where x is RR of a variable; y is the frequency (i.e., the number of RR values); α is a coefficient showing the expected number of RR values at $x = \mu$; and μ and α are the mean and variance of the frequency distributions of RR, respectively.

In addition, global warming potential (GWP) was calculated when three soil GHG (i.e., CO₂, CH₄, and N₂O) fluxes were extracted simultaneously from one study (IPCC, 2007). It should be noted that the units of soil CO₂, CH₄, and N₂O fluxes were unified before the calculation of the GWP. The GWP (t CO₂ equivalent ha⁻¹) was then determined as follows:

$$\text{CO}_2 \times 1 + \text{CH}_4 \times 25 + \text{N}_2\text{O} \times 298 \quad (5)$$

The between-group heterogeneity (Q_b) across all data for a given response variable was calculated to further analyze the biochar effect among different subgrouping categories. A random-effect model was used to explore the soil and biochar properties and other auxiliary variables that may explain the response of soil GHG fluxes to biochar application. We also conducted metaregression analysis to examine the relationships between RR (GHGs) and continuous forcing factors. The correlations of RR (GHGs) among different variables were examined by correlation analysis applied in R (R Core Team, 2015).

The publication bias was tested by funnel plot method and assessed using Kendall's Tau (Moller & Jennions, 2001). If the mean effect had significant difference from zero (i.e., indicating the existence of publication bias), Rosenthal's fail-safe number was calculated (MetaWin 2.1; Rosenberg *et al.*, 1997) to estimate

whether our conclusion is likely to be affected by the non-published studies (Rosenberg, 2005).

Results

Effects of biochar application on soil greenhouse gas (GHG) fluxes

The individual response ratios (RRs) of soil GHG fluxes (i.e., CO₂, CH₄, and N₂O) all displayed normal/Gaussian distributions (Fig. S1). On average, biochar application significantly increased soil CO₂ fluxes by 22.14% with a mean weighted RR₊₊ of 0.20 [CI = (0.12, 0.31)], but decreased soil N₂O fluxes by 30.92% with a RR₊₊ of 0.37 [CI = (-0.48, -0.28)]. Soil CH₄ fluxes were not significantly affected by biochar application [RR₊₊ = -0.03, CI = (-0.35, 0.23)] (Fig. 2, Table S2). Publication bias for this analysis was not suggested by Rosenthal's method (Table S3).

The response of soil CO₂ flux to biochar application depended significantly on biochar properties, experimental method, nitrogen (N) fertilization, and latitude. Soil texture and biochar pH were the two most critical parameters affecting the response of soil CH₄ flux to biochar addition. Biochar-induced changes in soil N₂O fluxes were significantly associated with soil and




Fig. 2 The effect of biochar application on soil CO₂ (a), CH₄ (b), and N₂O (c) emissions differed with experimental method [including field studies (F), laboratory incubation (I), and pot experiments (P)] in unfertilized soils and N-fertilized soils, shown as weighted response ratio (RR₊₊). Mean effect and 95% CIs are shown. If the CI did not overlap with zero, the response was considered significant (*). Numerals indicate the number of observations. 'Overall' indicates the integrated biochar effect across N fertilization as compared with controls.

Table 1 Between-group variability (Q_b) among observations (n) suggesting their potential as predictive variables influencing soil greenhouse gas (GHG) emissions responses to biochar application

Variables	CO ₂		CH ₄		N ₂ O	
	n	Q_b	n	Q_b	n	Q_b
All studies	402	–	121	–	371	–
Role of N fertilization	402	13.43***	121	7.70**	371	0.37
Experimental method	402	19.52***	121	9.33**	371	2.34
Feedstock source	402	4.28	121	10.60**	371	19.37***
Soil texture	277	9.95*	86	115.98***	256	14.34**
Pyrolysis temperature (°C)	385	37.27***	110	6.85**	354	1.94
Biochar pH	327	25.08***	103	14.22***	317	3.05
Soil pH	390	0.55	117	1.62	351	10.19**
Applied rate [Lg (t ha ⁻¹)]	400	15.65***	120	4.53*	371	39.05***
Latitude (°)	401	50.44***	121	0.00	371	2.50
Soil C/N ratio	212	0.25	58	0.02	183	2.35
Duration (day)	402	0.02	121	0.51	371	1.62
Biochar C/N ratio	387	0.06	120	0.64	363	0.53

A variable with larger Q_b is a better predictor than a variable with smaller Q_b .

Statistical significance of Q_b : * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

biochar properties, that is, biochar feedstock type and applied rate, soil texture, and pH (Table 1).

Combined effect of biochar with N fertilization on soil GHG fluxes

The combination of biochar with N fertilizer application significantly decreased soil CO₂ and increased CH₄ fluxes, whereas it did not change soil N₂O fluxes (Figs 2 and S2). In unfertilized soils, biochar application significantly increased soil CO₂ fluxes by 43.3% with a RR_{++} of 0.36 [CI = (0.24, 0.50)], but decreased soil CO₂ fluxes by 8.6% in N-fertilized soils [$RR_{++} = 0.09$, CI = (-0.17, -0.02)] (Table S2). Meanwhile, biochar application significantly increased soil CH₄ fluxes by 11.6% with a RR_{++} of 0.11 in N-fertilized soils [CI = (0.01, 0.25)], but had no significant effect in unfertilized soils [CI = (-0.62, 0.27)]. Biochar application significantly reduced soil N₂O fluxes by 33.0% and 28.8% in both fertilized and unfertilized soils with RR_{++} of 0.4 [CI = (-0.53, -0.28)] and 0.34 [CI = (-0.51, -0.18)], respectively (Fig. 2).

Effects of biochar applying methods on GHG fluxes

Experimental methods (i.e., field studies, laboratory incubations, and pot experiments) had a significant effect on the response of soil CO₂ and CH₄ fluxes to biochar application, while it was not pronounced for N₂O fluxes (Table 1, Fig. S3). On average, biochar application significantly increased soil CO₂ fluxes by 30.34% in laboratory incubations, but had no changes under field studies and pot experiments. Biochar application significantly increased soil CH₄ fluxes by 25.4% in field studies, but did not change in laboratory incubations and pot experiments. In addition, experimental duration showed no significant effect on responses of soil GHG fluxes to biochar application (Fig. S4).

Interestingly, the effect of fertilization on GHG fluxes in biochar-amended soil appears closely related to experiment methodology. Only laboratory incubations showed a significant increase in CO₂ fluxes to biochar application in unfertilized soils compared to those in field and pot experiments, while there were no responses in fertilized soils. For CH₄ fluxes, only field studies showed significant positive responses to biochar application in fertilized soils, and other treatments did not exhibit any significant effects (Fig. 2).

Effects of soil and biochar properties on soil GHG emissions

The response of soil GHG fluxes to biochar application differed for biochar feedstock source (i.e., wood, herb, and biowaste, Table 1, Fig. 3a–c). Among all biochar feedstock sources, wood source had the smallest positive effect for CO₂ fluxes and negative effect for N₂O fluxes. Meanwhile, biowaste source induced the largest positive effect and negative effect for CO₂ and N₂O fluxes, respectively. The effects of biochar application on soil CH₄ fluxes were not significant among different feedstock sources.

The response of soil GHG fluxes to biochar application also varied with soil texture (Table 1, Fig. 3d–f). For CO₂ fluxes, positive effects of biochar application occurred in soils with coarse and medium texture, while no significant effects were found in fine texture. CH₄ fluxes showed a significant negative response to biochar amendment only in coarse soils. N₂O fluxes significantly decreased by biochar application in all soil types, but the smallest negative response occurred in medium soils.

Response ratios of soil GHG fluxes across all the studies were significantly correlated with biochar pyrolysis temperature (Tem), biochar pH (BpH), soil pH, and biochar application rate (App), and latitude (Lat) (Table 1, Fig. 4). The response of soil CO₂ and CH₄ fluxes to




Fig. 3 The effect of biochar application on soil CO₂ (a and d), CH₄ (b and e), and N₂O (c and f) emissions depended on biochar feedstock source and soil texture, shown as weighted response ratio (RR₊₊). Mean effect and 95% CIs are shown. If the CI did not overlap with zero, the response was considered significant (*). Numerals indicate the number of observations.

biochar amendment slightly decreased with pyrolysis temperature and biochar pH ($P < 0.001$), but increased with application rate and latitude of the study for soil CO₂ fluxes ($P < 0.001$). In addition, the responses of soil N₂O fluxes to biochar application revealed negative trends with soil pH ($P = 0.001$) and application rate ($P < 0.001$). Although these correlations were statistically significant, their contributions in explaining the variation in GHG flux responses were low ($0.04 < R^2 < 0.11$, Fig. 4).

Effects of biochar application on global warming potential (GWP)

With those data measured simultaneously for soil CO₂, CH₄, and N₂O fluxes, biochar application positively affected GWP [RR₊₊ = 0.44, CI = (0.22, 0.69)]. Meanwhile, biochar application significantly increased GWP by a mean response ratio of 0.69 [CI = (0.39, 0.99)] in unfertilized soils compared to a minor negative effect in N-fertilized soils [RR₊₊ = -0.08, CI = (-0.15, -0.03), Fig. 5a–c]. Interestingly, laboratory incubations showed significant positive responses of GWP to biochar application, while field and pot experiments exhibited no effects (Fig. 5d). The different responses between

laboratory incubations, field and pot experiments for all data were the same as those in unfertilized soils (Fig. 5d₁), while, in fertilized soils, there were no significant effects of biochar application on GWP (Fig. 5d₂). This pattern generally matched the effect of biochar on soil CO₂ fluxes.

Discussion

Responses of CO₂, CH₄, and N₂O fluxes to biochar application

On average, our meta-analysis showed that biochar application significantly increased soil CO₂ fluxes by 22.14%. Among individual studies, biochar application affected soil CO₂ fluxes with diverse magnitudes and even directions (Scheer *et al.*, 2011; Augustenborg *et al.*, 2012; Zhang *et al.*, 2012a). The stimulating effects of biochar application on soil CO₂ fluxes were usually ascribed to higher labile C mineralization and/or inorganic C release from biochar (Fig. 6; e.g., Jones *et al.*, 2011; Smith *et al.*, 2010; Zimmerman *et al.*, 2011). Furthermore, as suggested by Liu *et al.* (2016), biochar application enhanced soil organic C (SOC) by 40% and soil microbial biomass C (MBC) content by 18%. This




Fig. 4 Effects of biochar pyrolysis temperature, biochar pH, soil pH, applied rate, and latitude on response ratios of soil CO₂ emissions (a–e), CH₄ emissions (f–j), and N₂O emissions (k–o) to biochar application.

indicates that the stimulation of soil CO₂ fluxes might be associated with the higher SOC status and the more active soil microbial activities (Fig. 6).

Soil CO₂ fluxes declined with biochar pyrolysis temperature. Low pyrolysis temperature results in more microbial available C and nutrients in biochar than a high pyrolysis temperature, which promotes high soil microbial activities to decompose soil organic matter (SOM) and release more CO₂ from soil (Chan *et al.*, 2008; Novak *et al.*, 2010; Hale *et al.*, 2012). This results in the negative relationship between RR (CO₂) and biochar pyrolysis temperature and a positive relationship between RR (CO₂) and application rate (Fig. 4a, d). Meanwhile, high-temperature biochars may contain higher relative concentrations of toxic compounds (i.e., polycyclic aromatic hydrocarbons) (Nakajima *et al.*, 2007), which can affect soil microbial biomass and activity. In addition, the RR (CO₂) exhibited a negative correlation with biochar pH probably because biochar with pH < 7 had a relatively high input of labile C fractions and triggered a higher priming effect on soil C mineralization (Crombie *et al.*, 2015). Our results indicated that CO₂ fluxes did vary over time after biochar application. However, mechanisms involved in soil CO₂ stimulation after biochar application may differ in short term compared to long term. In short term, soil CO₂ stimulation may have been originated from the breakdown of organic C and the release of inorganic C contained in

the biochar (Jones *et al.*, 2011). In the long term, biochar can promote the rapid loss of humus and belowground C (Wardle *et al.*, 2008). Meanwhile, increased belowground NPP induced by biochar amendment may also cause the stimulation of CO₂ emissions during the long-time experiments (Major *et al.*, 2010).

In addition, biochar-induced changes in soil CO₂ fluxes significantly increased with latitude, which may be related to increase in soil temperature after biochar application (Bozzi *et al.*, 2015). The increasing temperature may induce the larger stimulation on soil microbes and thereby CO₂ fluxes, in the high-latitude soils, where microbial activities and soil respiration are strongly limited by temperature (Mikan *et al.*, 2002).

Biochar application had no significant effect on soil CH₄ fluxes in our meta-analysis, although individual studies showed diverse effects. In experimental studies, multiple factors (e.g., soil aeration and porosity, methanogens, and methanotrophs) have been proposed to explain the different effects of biochar application on soil CH₄ fluxes (Lehmann & Rondon, 2006; Karhu *et al.*, 2011), but the underlying mechanisms are still poorly understood (Lorenz & Lal, 2014). Soil CH₄ fluxes are largely determined by methanogens and methanotrophs at a microbial scale (Bodelier & Laanbroek, 2004). Therefore, decreased soil CH₄ fluxes under biochar application might be due to the higher ratios of methanogenic to methanotrophic bacteria observed in some studies




Fig. 5 Frequency distributions of response ratio (RR) of global warming potential (GWP, panel a) to biochar application, GWP in unfertilized soils (b) and N-fertilized soils (c). The sample size (n), weighted response ratio (RR_{w}), and 95% CIs are shown. The effect of biochar application on GWP differed with experimental method (d), and GWP differed with experimental method in unfertilized soils (d_1) and N-fertilized soils (d_2). Mean effect and 95% CIs are shown. If the CI did not overlap with zero, the response was considered significant (*). Numerals indicate the number of observations.

(Fig. 6; Feng *et al.*, 2012), and others suggested that improved soil aeration and CH_4 oxidation after biochar application suppressed soil CH_4 fluxes (Fig. 6; Karhu *et al.*, 2011). In contrast, the increased soil CH_4 fluxes under biochar application could be attributed to biochar compounds that inhibit the activity of methanotrophs (Spokas, 2013).

Biochar application decreased CH_4 fluxes in coarse soils, whereas it increased CH_4 fluxes in fine soils. Biochar application to the coarse soils is likely to improve soil aeration, thus making the soils more favorable for the aerobic methanotrophs communities and increases CH_4 oxidation (Van Zwieten *et al.*, 2009). However, in the fine-textured soils, the porous structure of biochar may be filled with a clay and fine silt fraction, which could offset the aeration effect. A weak stimulation of CH_4 fluxes induced by biochar amendment may be due to enhancing soil methanogenic archaea (Feng *et al.*, 2012). In addition, the biochar-induced effects on soil CH_4 fluxes decreased with biochar pH, probably

resulting from altered soil microbial community structure, especially the ratio of soil methanogenic to methanotrophic abundance (Anders *et al.*, 2013).

Our meta-analysis showed that biochar application decreased soil N_2O fluxes by 30.92%, consistent with another meta-analysis reported by Cayuela *et al.* (2014). This response was probably driven by the changes in the activity of the nitrifiers and denitrifiers that produce N_2O . Biochar application enhances soil aeration (absorbing/holding an excess of soil moisture) and reduces N leaching as a result of NH_4^+ and NO_3^- adsorption by biochar (Fig. 6; Bai *et al.*, 2015a; Reverchon *et al.*, 2014; Rogovska *et al.*, 2011; Steiner *et al.*, 2008; Yanai *et al.*, 2007). The enhanced soil aeration and reduced compaction may inhibit denitrification due to more oxygen being present, and the diminished N leaching may decrease the inorganic N pool available for soil nitrifiers and denitrifiers (Fig. 6). Moreover, biochar amendment stimulates the *nosZ* transcription (i.e., denitrifying bacteria gene markers), which suggests




Fig. 6 Potential mechanisms of soil greenhouse gas (GHG) fluxes in response to biochar amendment. The red line and blue line represent the positive and negative regulations, respectively.

that biochar mitigates N_2O fluxes by further reducing it to N_2 (Xu *et al.*, 2014). In addition, biochar facilitates the transfer of electrons to soil denitrifying microorganisms, which promotes the reduction of N_2O to N_2 (Fig. 6; Cayuela *et al.*, 2013).

Furthermore, our study found that biochar-induced decreases in N_2O fluxes were enhanced with increasing biochar application rate. Larger amounts of microbial available and active nutrients due to high biochar application rates may promote the complete denitrification to N_2 (Lorenz & Lal, 2014), which may largely contribute to the suppression of soil N_2O fluxes as well as high molar H : C_{org} ratio (Cayuela *et al.*, 2015).

Regulation of nitrogen (N) fertilization on biochar impacts

Our results showed that biochar application increased soil CO_2 fluxes by 43.33% in unfertilized soils, but decreased by 8.61% in N-fertilized soils, consistent with the meta-analysis of Liu *et al.* (2016). More available inorganic N source for soil microbes and/or plant roots could stimulate soil microbial C mineralization after N is added (Lu *et al.*, 2011; Zhou *et al.*, 2014), but the absorption of NH_4^+ and NO_3^- by biochar would decrease the soil inorganic N pool after N fertilizers were applied (Steiner *et al.*, 2008; Clough *et al.*, 2013). Therefore, immobilization of soil inorganic N induced by biochar application may be the main reason for the slight suppression of soil CO_2 fluxes in N-fertilized soils. In unfertilized soils, the significant stimulation of soil CO_2 fluxes was mainly explained by the relatively higher nutrient availability for soil

microbes and/or the priming effect on native soil C decomposition after biochar application (Wardle *et al.*, 2008; Smith *et al.*, 2010).

Biochar application increased soil CH_4 fluxes by 11.67% in N-fertilized soils, but had no significant effect on unfertilized soils. Soil CH_4 fluxes increased weakly under corn and strongly under rice cultivation with N fertilization, respectively, during the entire growing season (Zhang *et al.*, 2010, 2012b). Biochar input under N addition is likely to alleviate C limitation to microbes. Therefore, the activities of soil methanogenic archaea are enhanced and more CH_4 is produced. Alternatively, some studies showed that decrease in soil CH_4 fluxes could be partly explained by the facilitated CH_4 oxidation after biochar application (Karhu *et al.*, 2011; Yu *et al.*, 2012), and a more stimulatory effect of biochar on methanotrophic proteobacteria than on methanogenic archaea in unfertilized soils (Feng *et al.*, 2012).

The biochar-induced decrease in soil N_2O fluxes was not significantly different in unfertilized [28.82%, CI = (39.95%, 16.47%)] soils from those of N-fertilized soils [32.97%, CI = (41.14%, 24.42%)]. As N addition increased N_2O fluxes by 216% on average across the globe (Liu & Greaver, 2009), the quantity of soil N_2O fluxes mitigated by biochar application in N-fertilized soils is much larger than that in unfertilized soils. As mentioned above, this might be due to more soil NH_4^+ and NO_3^- absorbed by biochar after N fertilizer application, likely causing denitrification to decline (Russow *et al.*, 2008) and/or a facilitation of N_2O reduction to N_2 (Dalal *et al.*, 2003).

Biochar effects on soil GHG fluxes varying with experimental types

The effects of biochar application on soil CO₂ fluxes differed with experimental types. Our study found a significant positive response in unfertilized soils mainly in laboratory incubations, but not in field and pot experiments. The positive response of soil CO₂ fluxes in laboratory incubation is most likely due to the mineralization of the labile C fractions existed in biochar (Zimmerman *et al.*, 2011), as well as increased soil surface area due to pore structures which promotes microbial activity (Chia *et al.*, 2014). In field experiments, the nonsignificant difference in CO₂ fluxes between control and biochar treatments largely resulted from low application rates and/or high biochar labile C leaching due to rainfall (Kuz'yakov *et al.*, 2009; Spokas & Reicosky, 2009). In N-fertilized soils, there were no significant differences in biochar-induced changes in soil CO₂ fluxes among field studies, pot experiments, and laboratory incubations. The positive effects of biochar application on soil CO₂ fluxes as mentioned above may be offset by the absorption of soil inorganic N (NH₄⁺ and NO₃⁻) when biochar is applied (Steiner *et al.*, 2008; Wardle *et al.*, 2008; Smith *et al.*, 2010). Therefore, no changes were observed in soil CO₂ fluxes.

Across all studies, soil CH₄ fluxes showed a positive response to biochar application in field studies, but no significant changes in laboratory incubations and pot experiments. The positive effects in field studies mainly reported from the treatments with N fertilization. The increase in soil CH₄ fluxes under N addition probably resulted from the stimulation of soil microbial activities, especially the methanogenic archaea and methanotrophic bacteria (Bodelier & Laanbroek, 2004). As reported by Liu *et al.* (2016), biochar amendment significantly increased soil microbial biomass C (MBC) in the field experiments, whereas MBC decreased in controlled studies. This likely resulted from improving the availability of microbial habitats and the accessibility of microbial food resources in the field-based experiments compared to the controlled conditions especially under biochar amendment (Pietikainen *et al.*, 2000).

In contrast, the responses of soil N₂O fluxes to biochar application showed a consistent trend across all treatments (Fig. 4a–c). However, laboratory incubations showed greater N₂O flux decreases than field studies with respect to biochar application in unfertilized soils (Fig. 4b), likely due to the difference in mixing of biochar with soil in controlled and field studies. Biochar is mixed thoroughly with soils in most controlled studies, which enhances soil aeration, but in field studies biochar is applied to the soil surface (e.g., Scheer *et al.*,

2011; Wang *et al.*, 2012; Bamminger *et al.*, 2014; Case *et al.*, 2014).

Responses of GWP of soil GHGs to biochar application

Global warming potential (GWP) is a simplified index to estimate the potential future impacts of GHGs on the global climate system based on their radiative forcing and lifetimes (IPCC, 2013). Overall, biochar application significantly increased GWP by 46.22% [CI = (19.72%, 82.20%)]. The fluxes are governed by different mechanisms (Fig. 6), but largely resulting from the significant stimulation of soil CO₂ fluxes. The increased amount of soil CO₂ fluxes induced by biochar application was nearly a one thousand times the size of CH₄ or N₂O fluxes in most studies (e.g., Scheer *et al.*, 2011; Wang *et al.*, 2012; Zhang *et al.*, 2012a). In addition, biochar increased the GWP of soil GHGs in unfertilized soils, but decreased it in N-fertilized soils due to the suppression of soil CO₂ and N₂O fluxes under N addition.

Significant amounts of CO₂, CH₄, and N₂O were released to the atmosphere from agriculture, which accounted for nearly one-fifth of the annual increase in radiative forcing of climate change (Cole *et al.*, 1997). Soil GHG fluxes would increase substantially after N fertilizers were applied, especially in croplands (Hall & Matson, 1999; McSwiney & Robertson, 2005; Liu & Greaver, 2009; Zhou *et al.*, 2014). Agricultural GHG emissions from crop and livestock production were 5.3 Pg of carbon dioxide equivalents (CO₂ eq) in 2011 (FAO 2014). Tian *et al.* (2016) estimated that CH₄ and N₂O emissions in the agricultural ecosystems were 169 ± 26 and 4.9 ± 0.3 Tg N yr⁻¹, respectively. According to our estimates with a decrease of 7.69% for GWP under N fertilization, 0.41 Pg CO₂ eq yr⁻¹ could potentially be mitigated by biochar applied to agricultural soils in combination with N fertilizers. Moreover, biochar application would increase the average yield of 10% and nearly 14% in acidic soils (Jeffery *et al.*, 2011). Given that our study elicits that biochar application reduces CO₂ fluxes and GWP in N-fertilized soil, biochar therefore appears to be a good strategy to mitigate global warming in fertilized agro-ecosystems.

Implications for future experiments and land surface models

The compiled database in our meta-analysis was mainly obtained from laboratory incubations, and the results were different for the responses of soil GHG fluxes to biochar application compared to those from field studies (Scheer *et al.*, 2011; Spokas *et al.*, 2009; Fig 4). The lack of field-scale studies, especially those lasting at least two successive seasons (Lorenz & Lal, 2014), may

hamper our evaluation of soil GHG fluxes in response to biochar application in the longer term. In addition, most biochar application experiments had been conducted in North America, Europe, and China. There remains a dearth of field studies in other regions, including Africa, South-East Asia, and South America. Thus, long-term field experiments with biochar amendments are especially needed in these regions.

Nitrogen fertilization mediated the responses of soil GHG fluxes and their GWP to biochar application. Because N deposition increased from ~34 Tg N yr⁻¹ in 1860–100 Tg N yr⁻¹ in 1995 and is predicted to reach 200 Tg N yr⁻¹ in 2050 (Galloway *et al.*, 2008; IPCC, 2013), the interactive effects between biochar and N addition may dramatically influence soil microbial community structure and ecosystem functioning as well as soil GHG fluxes in the future (Liu *et al.*, 2016). To address this issue, biochar experiments with diverse types of N fertilization (e.g., fertilizer type and level) are needed to examine the potential nonlinear responses to biochar application.

In the nature, biochar is often produced by wildfire, and currently, industrially produced biochar application becomes more common, especially in agriculture. Our meta-analysis results from laboratory, pot, and field studies found significant effects of biochar application on soil GHG fluxes and their GWPs. These results may provide some insights into how the fire-generated biochar affects net climate forcing from soil GHG fluxes and offers recommendations for the development and improvement of land surface models. Tempo-spatial variability of soil GHG fluxes is mostly attributed to soil temperature, soil moisture, fire severity, aspect, and time since fire in wildfire models (Gathany & Burke, 2011). However, wildfire-produced and industrial biochar may play critical roles in shaping terrestrial ecosystem processes and affecting soil GHG fluxes. Thus, future land surface models may need to incorporate biochar-induced effects to natural ecosystem processes, especially soil GHG fluxes and their GWPs for better forecasting the feedback of terrestrial ecosystems to climate change. Additionally, the combined or interactive effects of N fertilization with biochar amendments can be incorporated into future land surface models to improve the predictions about N-mediated feedback of ecosystem C cycles to climate systems from soil GHG fluxes.

Acknowledgements

This research was financially supported by the National Natural Science Foundation of China (Grant No. 31370489), the Program for Professor of Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Learning, and 'Thousand Young Talents' Program in China. We would like to

acknowledge the work carried out by the researchers whose published data were used for this meta-analysis.

References

- Adams DC, Gurevitch J, Rosenberg MS (1997) Resampling tests for meta-analysis of ecological data. *Ecology*, **78**, 1277–1283.
- Ameloot N, De Neve S, Jegajeevagan K *et al.* (2013) Short-term CO₂ and N₂O emissions and microbial properties of biochar amended sandy loam soils. *Soil Biology and Biochemistry*, **57**, 401–410.
- Anders E, Watzinger A, Rempt F *et al.* (2013) Biochar affects the structure rather than the total biomass of microbial communities in temperate soils. *Agricultural and Food Science*, **22**, 404–423.
- Augustenborg CA, Hepp S, Kammann C, Hagan D, Schmidt O, Müller C (2012) Biochar and earthworm effects on soil nitrous oxide and carbon dioxide emissions. *Journal of Environment Quality*, **41**, 1203.
- Bai SH, Reverchon F, Xu CY *et al.* (2015a) Wood biochar increases nitrogen retention in field settings mainly through abiotic processes. *Soil Biology & Biochemistry*, **90**, 232–240.
- Bai SH, Xu CY, Xu Z *et al.* (2015b) Soil and foliar nutrient and nitrogen isotope composition ($\delta^{15}\text{N}$) at 5 years after poultry litter and green waste biochar amendment in a macadamia orchard. *Environmental Science and Pollution Research*, **22**, 3803–3809.
- Bai SH, Xu CY, Xu ZH *et al.* (2016) Wood base biochar alters inorganic N. *Acta Horticulturae*. doi: 10.17660/ActaHortic.2016.1109.24.
- Bamminger C, Marschner B, Juschke E (2014) An incubation study on the stability and biological effects of pyrogenic and hydrothermal biochar in two soils. *European Journal of Soil Science*, **65**, 72–82.
- Barbosa De Sousa M, Soares Santos RR, Gehring C (2014) Charcoal in Amazonian paddy soil-nutrient availability, rice growth and methane emissions. *Journal of Plant Nutrition and Soil Science*, **177**, 39–47.
- Bodelier PLE, Laanbroek HJ (2004) Nitrogen as a regulatory factor of methane oxidation in soils and sediments. *FEMS Microbiology Ecology*, **47**, 265–277.
- Bozzi E, Genesio L, Toscano P, Pieri M, Miglietta F (2015) Mimicking biochar-albedo feedback in complex Mediterranean agricultural landscapes. *Environmental Research Letters*, **10**, 084014.
- Case SDC, Mcnamara NP, Reay DS, Whitaker J (2012) The effect of biochar addition on N₂O and CO₂ emissions from a sandy loam soil – the role of soil aeration. *Soil Biology and Biochemistry*, **51**, 125–134.
- Case SDC, Mcnamara NP, Reay DS, Whitaker J (2014) Can biochar reduce soil greenhouse gas emissions from a *Miscanthus* bioenergy crop? *GCB Bioenergy*, **6**, 76–89.
- Cayuela ML, Sanchez-Monedero MA, Roig A, Hanley K, Enders A, Lehmann J (2013) Biochar and denitrification in soils: when, how much and why does biochar reduce N₂O emissions? *Scientific Reports*, **3**, 1732.
- Cayuela ML, Van Zwieten L, Singh BP, Jeffery S, Roig A, Sánchez-Monedero MA (2014) Biochar's role in mitigating soil nitrous oxide emissions: a review and meta-analysis. *Agriculture, Ecosystems & Environment*, **191**, 5–16.
- Cayuela ML, Jeffery S, Van Zwieten L (2015) The molar H: Corg ratio of biochar is a key factor in mitigating N₂O emissions from soil. *Agriculture, Ecosystems & Environment*, **202**, 135–138.
- Chan KY, Van Zwieten L, Meszaros I, Downie A, Joseph S (2008) Using poultry litter biochars as soil amendments. *Australian Journal of Soil Research*, **46**, 437–444.
- Chia CH, Singh BP, Joseph S, Graber ER, Munroe P (2014) Characterization of an enriched biochar. *Journal of Analytical and Applied Pyrolysis*, **108**, 26–34.
- Clough TJ, Bertram JE, Ray JL, Condon LM, O'callaghan M, Sherlock RR, Wells NS (2010) Unweathered wood biochar impact on nitrous oxide emissions from a bovine-urine-amended pasture soil. *Soil Science Society of America Journal*, **74**, 852.
- Clough T, Condon L, Kammann C, Müller C (2013) A review of biochar and soil nitrogen dynamics. *Agronomy*, **3**, 275–293.
- Cole CV, Duxbury J, Freney J *et al.* (1997) Global estimates of potential mitigation of greenhouse gas emissions by agriculture. *Nutrient Cycling in Agroecosystems*, **49**, 221–228.
- Crombie K, Masek O, Cross A, Sohi S (2015) Biochar – synergies and trade-offs between soil enhancing properties and C sequestration potential. *Global Change Biology Bioenergy*, **7**, 1161–1175.
- Dalal RC, Wang WJ, Robertson GP, Parton WJ (2003) Nitrous oxide emission from Australian agricultural lands and mitigation options: a review. *Australian Journal of Soil Research*, **41**, 165–195.
- Darby I, Xu CY, Wallace HM, Joseph S, Pace B, Bai SH (2016) Short-term dynamics of carbon and nitrogen using compost, compost-biochar mixture and organo-mineral biochar. *Environmental Science and Pollution Research*, **23**, 11267–11278.

- FAO (2014) *Agriculture, Forestry and Other Land Use Emissions by Sources and Removals by Sinks*. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Feng Y, Xu Y, Yu Y, Xie Z, Lin X (2012) Mechanisms of biochar decreasing methane emission from Chinese paddy soils. *Soil Biology and Biochemistry*, **46**, 80–88.
- Galloway JN, Townsend AR, Erisman JW *et al.* (2008) Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science*, **320**, 889–892.
- Gathany MA, Burke IC (2011) Post-fire soil fluxes of CO₂, CH₄ and N₂O along the Colorado Front Range. *International Journal of Wildland Fire*, **20**, 838–846.
- Gomez JD, Deneff K, Stewart CE, Zheng J, Cotrufo MF (2014) Biochar addition rate influences soil microbial abundance and activity in temperate soils. *European Journal of Soil Science*, **65**, 28–39.
- Hale SE, Lehmann J, Rutherford D *et al.* (2012) Quantifying the total and bioavailable polycyclic aromatic hydrocarbons and dioxins in biochars. *Environmental Science & Technology*, **46**, 2830–2838.
- Hall SJ, Matson PA (1999) Nitrogen oxide emissions after nitrogen additions in tropical forests. *Nature*, **400**, 152–155.
- Hansen J, Sato M, Ruedy R, Lacis A, Oinas V (2000) Global warming in the twenty-first century: an alternative scenario. *Proceedings of the National Academy of Sciences of the United States of America*, **97**, 9875–9880.
- Hedges LV, Gurevitch J, Curtis PS (1999) The meta-analysis of response ratios in experimental ecology. *Ecology*, **80**, 1150–1156.
- Hilscher A, Knicker H (2011) Carbon and nitrogen degradation on molecular scale of grass-derived pyrogenic organic material during 28 months of incubation in soil. *Soil Biology & Biochemistry*, **43**, 261–270.
- IPCC (2007) Technical Summary. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL), pp. 19–91. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- IPCC (2013) Summary for policymakers. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM), pp. 3–29. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Jeffery S, Verheijen FGA, Van Der Velde M, Bastos AC (2011) A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture Ecosystems & Environment*, **144**, 175–187.
- Jones DL, Murphy DV, Khalid M, Ahmad W, Edwards-Jones G, Deluca TH (2011) Short-term biochar-induced increase in soil CO₂ release is both biotically and abiotically mediated. *Soil Biology & Biochemistry*, **43**, 1723–1731.
- Karhu K, Mattila T, Bergström I, Regina K (2011) Biochar addition to agricultural soil increased CH₄ uptake and water holding capacity – results from a short-term pilot field study. *Agriculture, Ecosystems & Environment*, **140**, 309–313.
- Kuzyakov Y, Subbotina I, Chen HQ, Bogomolova I, Xu XL (2009) Black carbon decomposition and incorporation into soil microbial biomass estimated by C-14 labeling. *Soil Biology & Biochemistry*, **41**, 210–219.
- Laird DA (2008) The charcoal vision: a win-win-win scenario for simultaneously producing bioenergy, permanently sequestering carbon, while improving soil and water quality. *Agronomy Journal*, **100**, 178–181.
- Laird DA, Brown RC, Amonette JE, Lehmann J (2009) Review of the pyrolysis platform for coproducing bio-oil and biochar. *Biofuels Bioproducts & Biorefining-Biofpr*, **3**, 547–562.
- Lal R (1999) World soils and greenhouse effect. *IGBP Global Change Newsletter*, **37**, 4–5.
- Lehmann J (2007a) Bio-energy in the black. *Frontiers in Ecology and the Environment*, **5**, 381–387.
- Lehmann J (2007b) A handful of carbon. *Nature*, **447**, 143–144.
- Lehmann J, Rondon M (2006) Bio-char soil management on highly weathered soils in the humid tropics. *Biological Approaches to Sustainable Soil Systems*, **113**, 517–530.
- Liu L, Greaver TL (2009) A review of nitrogen enrichment effects on three biogenic GHGs: the CO₂ sink may be largely offset by stimulated N₂O and CH₄ emission. *Ecology Letters*, **12**, 1103–1117.
- Liu SW, Zhang YJ, Zong YJ *et al.* (2016) Response of soil carbon dioxide fluxes, soil organic carbon and microbial biomass carbon to biochar amendment: a meta-analysis. *Global Change Biology Bioenergy*, **8**, 392–406.
- Lorenz K, Lal R (2014) Biochar application to soil for climate change mitigation by soil organic carbon sequestration. *Journal of Plant Nutrition and Soil Science*, **177**, 651–670.
- Lu M, Zhou X, Luo Y, Yang Y, Fang C, Chen J, Li B (2011) Minor stimulation of soil carbon storage by nitrogen addition: a meta-analysis. *Agriculture Ecosystems & Environment*, **140**, 234–244.
- Luo YQ, Hui DF, Zhang DQ (2006) Elevated CO₂ stimulates net accumulations of carbon and nitrogen in land ecosystems: a meta-analysis. *Ecology*, **87**, 53–63.
- Major J, Lehmann J, Rondon M, Goodale C (2010) Fate of soil-applied black carbon: downward migration, leaching and soil respiration. *Global Change Biology*, **16**, 1366–1379.
- Marris E (2006) Putting the carbon back: Black is the new green. *Nature*, **442**, 624–626.
- McSwiney CP, Robertson GP (2005) Nonlinear response of N₂O flux to incremental fertilizer addition in a continuous maize (*Zea mays* L.) cropping system. *Global Change Biology*, **11**, 1712–1719.
- Mikan CJ, Schimel JP, Doyle AP (2002) Temperature controls of microbial respiration in arctic tundra soils above and below freezing. *Soil Biology & Biochemistry*, **34**, 1785–1795.
- Moller AP, Jennions MD (2001) Testing and adjusting for publication bias. *Trends in Ecology & Evolution*, **16**, 580–586.
- Mukherjee A, Lal R, Zimmerman AR (2014) Effects of biochar and other amendments on the physical properties and greenhouse gas emissions of an artificially degraded soil. *Science of the Total Environment*, **487**, 26–36.
- Nakajima D, Nagame S, Kuramochi H *et al.* (2007) Polycyclic aromatic hydrocarbon generation behavior in the process of carbonization of wood. *Bulletin of Environmental Contamination and Toxicology*, **79**, 221–225.
- Novak JM, Busscher WJ, Watts DW, Laird DA, Ahmedna MA, Niandou MAS (2010) Short-term CO₂ mineralization after additions of biochar and switchgrass to a Typic Kandiodult. *Geoderma*, **154**, 281–288.
- Pan GX, Li LQ, Wu LS, Zhang XH (2004) Storage and sequestration potential of top-soil organic carbon in China's paddy soils. *Global Change Biology*, **10**, 79–92.
- Pietikainen J, Kikkila O, Fritze H (2000) Charcoal as a habitat for microbes and its effect on the microbial community of the underlying humus. *Oikos*, **89**, 231–242.
- Quin P, Joseph S, Husson O *et al.* (2015) Lowering N₂O emissions from soils using eucalypt biochar: the importance of redox reactions. *Scientific Reports*, **5**, 16773.
- R Core Team (2015) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <https://www.R-project.org/> (accessed 6 November 2015).
- Reverchon F, Flicker RC, Yang H *et al.* (2014) Changes in delta N-15 in a soil-plant system under different biochar feedstocks and application rates. *Biology and Fertility of Soils*, **50**, 275–283.
- Rogovska N, Laird D, Cruse R, Fleming P, Parkin T, Meek D (2011) Impact of biochar on manure carbon stabilization and greenhouse gas emissions. *Soil Science Society of America Journal*, **75**, 871.
- Rosenberg MS, Adams DC, Gurevitch J (1997) *MetaWin: Statistical Software for Meta-Analysis with Resampling Tests*. Sinauer, Sunderland, MA.
- Rosenberg MS (2005) The file-drawer problem revisited: a general weighted method for calculating fail-safe numbers in meta-analysis. *Evolution*, **59**, 464–468.
- Russow R, Spott O, Stange CF (2008) Evaluation of nitrate and ammonium as sources of NO and N₂O emissions from black earth soils (Haplic Chernozem) based on ¹⁵N field experiments. *Soil Biology and Biochemistry*, **40**, 380–391.
- Scheer C, Grace PR, Rowlings DW, Kimber S, Van Zwieten L (2011) Effect of biochar amendment on the soil-atmosphere exchange of greenhouse gases from an intensive subtropical pasture in northern New South Wales, Australia. *Plant and Soil*, **345**, 47–58.
- Schmidt MWI, Skjemstad JO, Jager C (2002) Carbon isotope geochemistry and nanomorphology of soil black carbon: black chernozemic soils in central Europe originate from ancient biomass burning. *Global Biogeochemical Cycles*, **16**, 1123.
- Smith P, Martino D, Cai Z *et al.* (2008) Greenhouse gas mitigation in agriculture. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **363**, 789–813.
- Smith JL, Collins HP, Bailey VL (2010) The effect of young biochar on soil respiration. *Soil Biology & Biochemistry*, **42**, 2345–2347.
- Spokas KA (2010) Observed ethylene production from biochar additions. Available at: <http://www.biorenew.iastate.edu/fileadmin/www.biorenew.iastate.edu/biochar2010/Presentations/Spokas.pdf> (accessed 16 December 2012).
- Spokas KA (2013) Impact of biochar field aging on laboratory greenhouse gas production potentials. *GCB Bioenergy*, **5**, 165–176.
- Spokas KA, Reicosky DC (2009) Impacts of sixteen different biochars on soil greenhouse gas production. *Annals of Environmental Science*, **3**, 179–193.
- Spokas KA, Koskinen WC, Baker JM, Reicosky DC (2009) Impacts of woodchip biochar additions on greenhouse gas production and sorption/degradation of two herbicides in a Minnesota soil. *Chemosphere*, **77**, 574–581.
- Steiner C, Glaser B, Teixeira WG, Lehmann J, Blum WEH, Zech W (2008) Nitrogen retention and plant uptake on a highly weathered central Amazonian Ferralsol

- amended with compost and charcoal. *Journal of Plant Nutrition and Soil Science-Zeitschrift Fur Pflanzenernahrung Und Bodenkunde*, **171**, 893–899.
- Stewart CE, Zheng J, Botte J, Cotrufo MF (2013) Co-generated fast pyrolysis biochar mitigates green-house gas emissions and increases carbon sequestration in temperate soils. *GCB Bioenergy*, **5**, 153–164.
- Sun L, Li L, Chen Z, Wang J, Xiong Z (2014) Combined effects of nitrogen deposition and biochar application on emissions of N₂O, CO₂ and NH₃ from agricultural and forest soils. *Soil Science and Plant Nutrition*, **60**, 254–265.
- Taghizadeh-Toosi A, Clough TJ, Condon LM, Sherlock RR, Anderson CR, Craigie RA (2011) Biochar incorporation into pasture soil suppresses in situ nitrous oxide emissions from ruminant urine patches. *Journal of Environmental Quality*, **40**, 468–476.
- Tian H, Lu C, Ciais P *et al.* (2016) The terrestrial biosphere as a net source of greenhouse gases to the atmosphere. *Nature*, **531**, 225–228.
- Van Zwieten L, Singh B, Joseph S *et al.* (2009) *Biochar and Emissions of Non-CO₂ Greenhouse Gases from Soil*. Science and Technology. Earthscan, London.
- Van Zwieten L, Kimber S, Morris S, Downie A, Berger E, Rust J, Scheer C (2010) Influence of biochars on flux of N₂O and CO₂ from Ferrosol. *Australian Journal of Soil Research*, **48**, 555–568.
- Wang J, Zhang M, Xiong Z, Liu P, Pan G (2011) Effects of biochar addition on N₂O and CO₂ emissions from two paddy soils. *Biology and Fertility of Soils*, **47**, 887–896.
- Wang JY, Pan XJ, Liu YL, Zhang XL, Xiong ZQ (2012) Effects of biochar amendment in two soils on greenhouse gas emissions and crop production. *Plant and Soil*, **360**, 287–298.
- Wardle DA, Nilsson MC, Zackrisson O (2008) Fire-derived charcoal causes loss of forest humus. *Science*, **320**, 629–629.
- Woolf D, Amonette JE, Street-Perrott FA, Lehmann J, Joseph S (2010) Sustainable biochar to mitigate global climate change. *Nature Communications*, **1**, 118–124.
- Xu HJ, Wang XH, Li H, Yao HY, Su JQ, Zhu YG (2014) Biochar impacts soil microbial community composition and nitrogen cycling in an acidic soil planted with rape. *Environmental Science & Technology*, **48**, 9391–9399.
- Xu C-Y, Bai SH, Hao Y, Rachaputi RCN, Xu Z, Wallace HM (2015a) Peanut shell biochar improves soil properties and peanut kernel quality on a red Ferrosol. *Journal of Soils and Sediments*, **15**, 2220–2231.
- Xu CY, Bai SH, Hao Y, Rachaputi RCN, Wang H, Xu Z, Wallace H (2015b) Effect of biochar amendment on yield and photosynthesis of peanut on two types of soils. *Environmental Science and Pollution Research*, **22**, 6112–6125.
- Yanai Y, Toyota K, Okazaki M (2007) Effects of charcoal addition on N₂O emissions from soil resulting from rewetting air-dried soil in short-term laboratory experiments. *Soil Science and Plant Nutrition*, **53**, 181–188.
- Yu L, Tang J, Zhang R, Wu Q, Gong M (2012) Effects of biochar application on soil methane emission at different soil moisture levels. *Biology and Fertility of Soils*, **49**, 119–128.
- Zhang A, Cui L, Pan G *et al.* (2010) Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China. *Agriculture, Ecosystems & Environment*, **139**, 469–475.
- Zhang AF, Bian RJ, Pan GX *et al.* (2012a) Effects of biochar amendment on soil quality, crop yield and greenhouse gas emission in a Chinese rice paddy: a field study of 2 consecutive rice growing cycles. *Field Crops Research*, **127**, 153–160.
- Zhang AF, Liu YM, Pan GX, Hussain Q, Li LQ, Zheng JW, Zhang XH (2012b) Effect of biochar amendment on maize yield and greenhouse gas emissions from a soil organic carbon poor calcareous loamy soil from Central China Plain. *Plant and Soil*, **351**, 263–275.
- Zheng J, Stewart CE, Cotrufo MF (2012) Biochar and nitrogen fertilizer alters soil nitrogen dynamics and greenhouse gas fluxes from two temperate soils. *Journal of Environment Quality*, **41**, 1361.
- Zhou L, Zhou X, Zhang B, Lu M, Luo Y, Liu L, Li B (2014) Different responses of soil respiration and its components to nitrogen addition among biomes: a meta-analysis. *Global Change Biology*, **20**, 2332–2343.
- Zimmerman AR, Gao B, Ahn MY (2011) Positive and negative carbon mineralization priming effects among a variety of biochar-amended soils. *Soil Biology & Biochemistry*, **43**, 1169–1179.

Supporting Information

Additional Supporting Information may be found online in the supporting information tab for this article:

Figure S1. Frequency distributions of response ratios (RR) of soil CO₂, CH₄, and N₂O emissions to biochar application.

Figure S2. Frequency distributions of response ratios (RR) of soil CO₂, CH₄, and N₂O emissions to biochar application on unfertilized soils and N-fertilized soils.

Figure S3. The effect of biochar application on soil CO₂, CH₄, and N₂O emissions differed with experimental method.

Figure S4. Effects of experimental duration on response ratios of GHG emissions to biochar application.

Table S1. Response ratio (RR) and number of paired observations extracted from each of the papers.

Table S2. Percentage changes of soil greenhouse gas (GHG) emissions in response to biochar application.

Table S3. The Kendall's Tau for RR(CO₂), RR(CH₄), RR(N₂O) and RR(GWP) in different treatments.

Data S1. A list of 91 papers from which the data were extracted for this meta-analysis.