

The Potential of Apple Cider Vinegar in the Management of Type 2 Diabetes

Author

Morgan, Joanna, Mosawy, Sapha

Published

2016

Journal Title

International Journal of Diabetes Research

Version

Version of Record (VoR)

Rights statement

© The Author(s) 2016. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Downloaded from

<http://hdl.handle.net/10072/101108>

Link to published version

<http://article.sapub.org/10.5923.j.diabetes.20160506.02.html>

Griffith Research Online

<https://research-repository.griffith.edu.au>

The Potential of Apple Cider Vinegar in the Management of Type 2 Diabetes

Joanna Morgan^{1,2}, Sapha Mosawy^{1,2,*}

¹School of Medical Science, Griffith University, Gold Coast Campus, Queensland, Australia

²Menzies Health Institute Queensland, Griffith University, Gold Coast Campus, Queensland, Australia

Abstract Type 2 Diabetes represents a large burden on public health systems worldwide. The chronic metabolic condition is characterised by hyperglycaemia and insulin resistance and is frequently associated with obesity, hypertension and dyslipidaemia. There is a growing need for effective management techniques of these conditions that patients can utilise complementary to conventional therapy. Apple cider vinegar (ACV) has been the subject of growing interest in this field. The main component of ACV, acetic acid, has demonstrated effectiveness in reducing hyperglycaemia, correcting dyslipidaemia and assisting weight loss. The dominant polyphenol compound in ACV, chlorogenic acid may also be useful in managing the condition.

Keywords Apple cider, Diabetes, Vinegar, Hyperglycaemia, Dyslipidaemia

1. Introduction

In Australia 280 people develop type 2 diabetes each day which currently affects 1.7 million Australians. The burden on the public health system is estimated at \$14.6 billion [1]. Type 2 diabetes, accounting for 85% of all diabetes, is a complicated chronic metabolic condition characterised by insulin resistance and eventually insufficient insulin production resulting in abnormal glucose metabolism. The condition is generally associated with obesity, a sedentary lifestyle, hypertension and dyslipidaemia. Type 2 diabetes significantly increases the risk of cardiovascular disease. Management of the condition involves managing the risks of cardiovascular disease as well as managing blood glucose levels [2, 3].

Apple cider vinegar (ACV) may be able to play a role in the day-to-day management of type 2 diabetes as growing research has demonstrated that certain aspects of the beverage assist in controlling hyperglycaemia, as well as reducing cardiovascular disease risks through weight loss, lowering blood pressure and lowering blood lipids [4, 5].

Not all ACV, however, is made alike. There are several techniques utilised in commercial vinegar production, from slower more traditional methods to techniques that can produce ACV within a day. There are many other factors in production, from apple cultivar, yeast and bacterial cultures used, to whether the product was filtered and pasteurised

[4, 6, 7]. Few studies have examined the effect different production methods have on the final product and the presence and quantity of organic components. Commercial varieties also give limited information on package labelling regarding production methods. The present review focuses on ACV and its beneficial effects on type 2 diabetes.

2. Production of Apple Cider Vinegar

2.1. Two-Step Fermentation Process

ACV can be produced by a two-step fermentation process, and this process is characterised by the presence of acetic acid at a concentration equal to or above 4% [8]. Cider vinegars are typically 5-6% acetic acid [9]. The pH of vinegar will depend on acetic acid concentration and is typically between 2 – 3.5 [10].

Yeasts initially ferment the sugars or starch in raw materials to form ethanol, which is further fermented by acetic acid bacteria (AAB) to produce acetic acid. This can be accomplished with juices/mashes from apples, grapes, coconuts, rice, potato and others. If a starch is the initial raw material, it will first need to be hydrolysed into a sugar. Depending on the method used for the second fermentation, vinegar can be produced as quickly as within 24 hours or may be left for months to years to ferment [4]. Figure 1 shows the chemical equations for the 2-step fermentation process. The final product may be filtered and pasteurised prior to consumption. This process removes and destroys AAB, preventing formation of 'mother of vinegar'. Mother of vinegar develops when unpasteurised vinegar is allowed to remain in the product, forming an extracellular cellulose

* Corresponding author:

s.mosawy@griffith.edu.au (Sapha Mosawy)

Published online at <http://journal.sapub.org/diabetes>

Copyright © 2016 Scientific & Academic Publishing. All Rights Reserved

layer which can be seen as a layer on the surface of the liquid, or as a cloudy cobweb-like substance, making the fluid appear murky. It is not unique to ACV.

Production of ACV can occur spontaneously via the naturally occurring yeasts and bacteria on the surface of the fruit, allowing the beverage to be easily made in the home [11]. The product produced in the home will likely differ in microbiota, acetic acid content and other molecules given that the spontaneous process is not standardised. Filtering and pasteurisation may not be done and the 'mother' may be consumed or used to inoculate subsequent batches of vinegar.




Figure 1. Chemical equations for 2-step fermentation process

2.2. Other Vinegar Production Techniques

2.2.1. Orleans Process (Traditional)

Orleans process is an early traditional process, in which wine covered in a film (mother) of AAB, oxidises slowly in a barrel. The barrel has holes, allowing for air flow and wine is added beneath the mother. The mother causes the apparatus to become slimy and slows the rate of vinegar production and vinegar is removed through the bottom of the barrel [10].

2.2.2. Generator Process (Surface Culture/Quick Process)

The generator process is believed to date back to the 17th century. AAB are grown in a thick layer on a non-compacting material, such as beech wood shavings. A pump circulates the liquid, allowing a slow trickle over the bacterial culture while air is permitted to circulate through the apparatus. While generator fermentation is used commercially, it is considered to be slow and expensive [10, 12].

2.2.3. Submerged-Culture

In the submerged-culture generator, a mechanical system keeps the AAB submerged within the liquid in close contact with aeration. The Frings acetator is a popular submerged-culture generator. Submerged culture method was designed for efficient commercial use [10].

2.2.4. Maceration

Maceration is a process already utilised in wine-making in which the remaining pulp from extracted juice is left to soak in the juice for a period of time. The phenolic and flavour compounds within the skin and pulp are extracted via this process. It has been shown that maceration in combination

with the surface production method yielded the ACV with the highest phenolic content [6].

3. Production Methods, Apple Cultivar and End Product

The production method utilised may affect the final properties and composition of ACV. Different production techniques have been demonstrated to affect pH, acidity and phenolic content [6, 13]. Budak *et al.* [4] concluded that production method affected the ability of ACV to alter triglyceride levels in rats with some methods more effective than others. The total content of phenolic compounds in ACV and hence, production method, may also be relevant to the ability of the beverage to promote good health.

The variety of apple (cultivar) used may affect the phenolic content of the juice product [14] which, not surprisingly, will also carry over to the cider vinegar [7]. The level of ripeness that the apples achieved may also affect the final product. A study that examined the ripening stage of apples on phenolic compounds in apple cider (non-alcoholic) found that unripe apples yielded a product with a lower phenolic content compared to ripened apples. The apples used in production can also affect the microbial content of the end product, with organic apples found to produce a more heterogeneous product compared with conventional apples [15]. The variations in microbiota may in turn influence the organic components of ACV which may affect the health promoting properties.

4. Organic Components of ACV

Acetic acid is the most abundant compound. Organic acids from an analysis of a commercially produced ACV using high resolution H NMR spectroscopy are found in Table 1. ACV is well established that various types of phenolic compounds are found in cider apples, particularly the hydrocinnamic acid derivatives, oligomeric flavan-3-ols, dihydrochalcones, and flavonols [16]. The phenolic content of ACV will vary with cultivar and processing [17]. Phenolic content of ACV was determined to consist of gallic acid, catechin, epicatechin, chlorogenic acid, caffeic acid and *p*-coumaric acid. Chlorogenic acid is the dominant phenolic substance in ACV [6]. The total phenol content and chlorogenic acid content appear to vary significantly between different studies, possibly attributed to the different ACVs being used.

Dietary polyphenols are natural phytochemical compounds and include the phenolic acid chlorogenic acid, a hydroxycinnamic acid derivative. Studies demonstrate rapid absorption of the polyphenolic compounds from the intestine. Many healthful benefits are attributed to polyphenols, such as antioxidant, antiallergic, anti-inflammatory, anti-viral and anti-microbial, anti-proliferative, anti-mutagenic, anti-carcinogenic, free radical scavenging, and induction of

antioxidant enzymes [19, 20]. There is also some evidence of modulation of signalling pathways such as nuclear factor kappa-B (NF- κ B) and mitogen-activated protein kinases (MAPK) [16].

Table 1. Organic acids in ACV

Compound	Concentration (g/L)
Acetic Acid	50.9
Citric Acid	0.02
Formic Acid	0.28
Lactic Acid	0.38
Malic Acid	3.56
Succinic Acid	0.27
Fructose	6.83
Acetoin	0.21
2,3-Butanediol	0.37
Ethanol	1.03
Ethyl acetate	0.14

Values adopted from Caligiani et al (2007) [18]

5. Management of Hyperglycaemia Using ACV

Hemoglobin A1c (HbA1c) level measures the glycation of hemoglobin, accurately identifies the average plasma glucose concentration over the previous three months. A study which investigated the effect acetic acid had on HbA1c in type 2 diabetics, found that HbA1c values fell by 0.16% units over the course of the 12-week trial, compared with controls who did not ingest any vinegar, where HbA1c levels rose by 0.06% [21]. HbA1c in diabetic rats was also significantly lowered with ACV consumption [22].

Several mechanisms may explain the ways in which acetic acid lowers plasma glucose have been suggested. These include inhibition of disaccharidase activity [23-25] and/or decrease in the hydrolytic enzyme α -amylase [26], delayed gastric emptying [27, 28] and an enhanced glucose uptake and conversion to glycogen in the periphery [23, 29, 30].

Delayed gastric emptying was noted in healthy subjects who consumed white bread along with a white vinegar dressing which contained olive oil (18mmol acetic acid in 20g vinegar). White vinegar is an aqueous solution containing approximately 6% acetic acid. Gastric emptying rate was indirectly measured through consumption of paracetamol baked into white bread; the blood paracetamol level was lower in the vinegar group compared with the control group. During the postprandial phase subjects who had consumed vinegar had significantly lower blood glucose levels and the insulin response in these subjects was also noted to be significantly lower compared with the reference meal [28]. Paracetamol however, may be absorbed and metabolised at different rates. Other research has also found a link to acetic acid consumption with delayed gastric emptying [27].

Recent investigation found that ACV had a stronger ability to lower plasma glucose levels than acetic acid alone [26]. This study found that it was not until day 7 that ACV significantly reduced plasma glucose levels in diabetic mice. ACV had comparable antiglycemic effects to the positive control group treated with the anti-diabetic agent sulfonylurea Glibenclamide. It was found that ACV treated groups had a significant decrease in α -amylase. The ability of ACV to have a stronger effect than acetic acid alone suggests a role for other components of ACV in controlling hyperglycemia. Another study found that consuming two tablespoons of ACV prior to sleeping was found to reduce fasting glucose the following morning [25].

Furthermore, acetic acid was demonstrated to significantly decrease the activities of the disaccharidase sucrase, maltase, trehalase and lactase in Caco-2 cells, but did not affect the enzymes at transcriptional or translational levels. It was suggested that suppression of the disaccharidase may occur in the post-translational processes, such as trafficking of the enzymes to the cell membrane [24]. Consumption of 100mL ACV (5% acetic acid) in diabetic rats demonstrated a significant decrease in the activity of maltase, sucrase and lactase [23]. In addition, vinegar ingestion (10g) was found to have no effect on postprandial glycemia (PPG) when only monosaccharides were ingested while a meal of complex carbohydrates consumed with vinegar did result in decreased PPG, further indicating that a acetic acid may inhibit disaccharidase activity [31].

Glycogen uptake by the liver and skeletal muscle was found to be enhanced in mice fed a diet containing 2g acetic acid/kg, a concentration that corresponds to foods prepared with vinegar. Acetic acid ingestion may inhibit glycolysis through accumulation of glucose-6-phosphate and a corresponding increase in glycogen synthesis, which was seen in liver and skeletal muscle of rats supplemented with acetic acid [29, 32, 33], causing an anti-hyperglycemic effect. Modulation of GK, G6PD and PFK in the liver of rats consuming ACV has been associated with decreased plasma glucose levels [23].

The quantity of acetic acid needed to exert effects has been investigated and a significant dose-response relationship was found in a study that examined the effects of ingestion of 18, 23 or 28g of white vinegar (6% acetic acid; equivalent to 18, 23, 28 mmol acetic acid, respectively) [34]. Compared with the control, the highest concentration of acetic acid caused a significant decrease in plasma glucose and insulin response postprandially while the lower acetic acid concentrations did lower blood glucose and insulin response, it was not significant [34].

Chlorogenic acid has been demonstrated to have some antiglycemic effects that may be useful in the management of type 2 diabetes. 1mM of chlorogenic acid was found to significantly inhibit glucose-6-phosphatase (G-6-Pase) activity in rat hepatocytes. G-6-Pase promotes glucose production through catalyzing steps in both gluconeogenesis and glycogenolysis and inhibition of this step can decrease plasma glucose concentration [35]. Synthetic derivatives of

chlorogenic acid also have been shown to inhibit G-6-Pase [36]. Liver perfusion experiments, however, did not find a decrease in glucose production at various chlorogenic acid concentrations, perhaps due to insufficient uptake of chlorogenic acid by hepatocytes. However, 1mM chlorogenic acid was able to significantly reduce the plasma glucose peak during the oral glucose tolerance test in rats and this is thought to be due to reduced activity of Na⁺-dependant D-glucose transporters in brush-border membrane vesicles, as administration of chlorogenic acid intravenously was unable to achieve the same result [35, 37].

Insulin sensitivity was improved in human subjects with both insulin resistance or type 2 diabetes when 20g of ACV was consumed with a high-carbohydrate meal [38]. Animal studies also demonstrated results suggesting improved insulin sensitivity with chlorogenic acid infusion [39]. Improved insulin sensitivity results in increased glucose uptake and hence lowered plasma glucose levels. A diet supplemented with chlorogenic acid has also been shown to significantly lower insulin levels in mice [40].

6. Management of Hypertension and Obesity

Diabetes may affect the autonomic nervous system and endothelium which results in microvascular complication, which in turn impairs the autoregulation of blood flow. Diabetic subjects have been shown to have lower levels of the vasodilator nitric oxide and increased levels of the vasoconstrictor endothelin-1 which results in a state of vasoconstriction [41]. A consequence of elevated blood pressure is vascular damage which leads to cardiovascular disease.

Acetic acid combined with vinegar were found to significantly decrease blood pressure (21-30mmHg lower than the control) and renin activity compared with controls and subjects consuming only vinegar. A decrease in renin and the subsequent release in angiotension II may be the reason for lowered blood pressure. A decrease in aldosterone was also seen. Both rice vinegar and acetic acid were given at a concentration of 46.2g/L [42]. Rice vinegar is generally 4% acetic acid [43], therefore the acetic acid given alone would be more potent than the vinegar solution. It was suggested that acetic acid may cause an increase in calcium absorption, which in turn may cause an calcium influx into renin secretory cells, inhibiting renin secretion [42]. A combination of red wine vinegar and grape juice was also found to decrease activity of angiotensin converting enzyme (ACE) [44].

Type 2 diabetes is commonly associated with obesity and weight loss is considered to be an important component in the management of diabetes [2]. Acetic acid has been proposed to have a role in reducing food intake. This may occur as a result of the taste of acetic acid in vinegar and the nausea it may induce from ingestion [45]. A study found that supplementation of a meal with white vinegar increased the

subjective rating of satiety compared with a control group. Improved satiety may result in lowered food consumption and hence, weight loss [34]. A 12 week study found that ingestion of both 75g and 150g of acetic acid significantly reduced the bodyweight, body mass index (BMI), visceral fat and waist circumference in comparison with a control group [46].

Chlorogenic acid has been demonstrated to halt the cell cycle of mouse embryo 3T3-L1 preadipocytes and arrest the G1 phase, hence preventing proliferation. Preadipocytes were inhibited in this study in both time- and dose-dependant manner at a concentration of 100µM. Decreased preadipocyte differentiation is just one proposed method to reverse obesity [47]. Another recent study supplementing the diet of mice on a high fat diet with 0.02% (w/w) chlorogenic acid resulted in a significant 16% weight loss compared with the control group and increased adiponectin levels [40].

7. Management of Dyslipidaemia

Type 2 diabetes is frequently associated with dyslipidaemia. Part of management of the condition involves attempting to achieve normal blood levels of total cholesterol, low density lipoprotein (LDL), high density lipoprotein (HDL) and triglycerides. Dyslipidaemia is highly correlated with atherosclerosis.

7.1. Dyslipidaemia and ACV

Ingestion of ACV improved lipid profiles in both normal and diabetic rats, decreasing triglycerides, total cholesterol and LDL while increasing HDL. These effects became pronounced after 4 weeks of treatment [23]. Further animal studies found similar effects on plasma total cholesterol, triglycerides, HDL and LDL levels [22, 48]. Other research focused on healthy humans has found the same improvement in lipid profile with ingestion of 30ml of ACV (4% acetic acid) [49].

7.2. Dyslipidemia and Acetic Acid

Research studies examining the effect acetic acid has on blood lipids found that rats fed a diet supplemented with 1% (w/w) cholesterol combined with acetic acid had significantly lowered total cholesterol and triglyceride levels compared with controls. Acetic acid was found to lower liver ATP citrate lyase (ATP-CL) activity, liver 3-hydroxy-3-methylglutaryl-CoA content, all of which are involved in lipid synthesis. Liver mRNA levels of sterol regulatory element binding protein-1, ATP-CL and fatty acid synthase were also found to be decreased. Faecal bile content was found to be higher in the group fed acetic acid. Blood lipids in rats fed acetic acid were decreased by both the inhibition of lipogenesis in the liver and the increased increment of cholesterol in faecal bile acid [50].

7.3. Dyslipidemia and Chlorogenic Acid

Supplementation of mice on a high fat diet with

chlorogenic acid significantly lowered plasma triglyceride and total cholesterol concentrations compared with the control group on a high fat diet. Adipose tissue triglycerides were also found to be significantly lowered. Hepatic activity of HMG-CoA reductase was lowered and fatty acid β -oxidation levels increased with chlorogenic acid intake [40]. Studies using obese, hyperlipidemic and insulin resistant (fa/fa) Zucker rats which were infused with chlorogenic acid (5mg/Kg body weight/day) found significant decreases in fasting plasma cholesterol and triglycerides [39].

8. Conclusions

The ACV is a readily available product that is easily able to be incorporated into meals. Large body of research has demonstrated its beneficial properties as an entire product, as well as the abilities of the individual components acetic acid and chlorogenic acid. ACV may assist in controlling blood glucose and lipids, weight loss and hypertension and therefore may be helpful in the management of type 2 diabetes. ACV as a whole may be more effective than acetic acid alone, although there is little research directly comparing acetic acid and ACV. Consumption of the 'mother of vinegar' may also increase beneficial effects compared with ACV lacking this component. Production method of ACV has been shown to alter the components of ACV, which may in turn affect the beneficial qualities. Further investigation may be beneficial here to determine the extent of the effect of production method. Consumption of ACV may indeed be beneficial in the management of type 2 diabetes.

REFERENCES

- [1] Diabetes in Australia: Diabetes Australia; 2015 [cited 2016 August]. Available from: <https://www.diabetesaustralia.com.au/diabetes-in-australia>.
- [2] Type 2 diabetes in adults: management. UK: 2015 2 December 2015. Report No.
- [3] Type II diabetes in adults [Internet]. BMJ Best Practice. 2016 [cited August 2016]. Available from: <http://bestpractice.bmj.com/best-practice/welcome.html>.
- [4] Budak NH, Aykin E, Seydim AC, Greene AK, Guzel-Seydim ZB. Functional properties of vinegar. *J Food Sci*. 2014; 79(5): R757-64.
- [5] Samad A, Azlan A, Ismail A. Therapeutic effects of vinegar: a review. *Current Opinion in Food Science*. 2016; 8: 56-61.
- [6] Budak NH, Kumbul Doguc D, Savas CM, Seydim AC, Kok Tas T, Ciris MI, et al. Effects of apple cider vinegars produced with different techniques on blood lipids in high-cholesterol-fed rats. *J Agric Food Chem*. 2011; 59(12): 6638-44.
- [7] Heikefelt C. Chemical and sensory analyses of juice, cider and vinegar produced from different apple cultivars. 2011.
- [8] Food Standards Code; Standard 2.10.1. In: Zealand FSAN, editor. 1991.
- [9] The Vinegar Institute, FAQ 2016 [cited 2016]. Available from: <http://versatilevinegar.org/faqs/>.
- [10] Webb AD. Vinegar. *Kirk-Othmer Encyclopedia of Chemical Technology*: John Wiley & Sons, Inc.; 2000.
- [11] Aung MT, Lee P-R, Yu B, Liu S-Q. Cider fermentation with three *Williopsis saturnus* yeast strains and volatile changes. *Annals of Microbiology*. 2015; 65(2): 921-8.
- [12] Tan SC. Vinegar Fermentation. Louisiana USA: Louisiana State University; 2005.
- [13] Budak NH, Guzel-Seydim ZB. Antioxidant activity and phenolic content of wine vinegars produced by two different techniques. *J Sci Food Agric*. 2010; 90(12): 2021-6.
- [14] Zielinski AAF, Alberti A, Braga CM, da Silva KM, Canteri MHG, Mafra LI, et al. Effect of mash maceration and ripening stage of apples on phenolic compounds and antioxidant power of cloudy juices: A study using chemometrics. *LWT-Food Science and Technology*. 2014; 57(1): 223-9.
- [15] Stornik A, Skok B, Trcek J. Comparison of Cultivable Acetic Acid Bacterial Microbiota in Organic and Conventional Apple Cider Vinegar. *Food Technology and Biotechnology*. 2016; 54(1): 113.
- [16] Bahadoran Z, Mirmiran P, Azizi F. Dietary polyphenols as potential nutraceuticals in management of diabetes: a review. *Journal of diabetes & metabolic disorders*. 2013; 12(1): 1.
- [17] Mangas JJ, Rodriguez R, Suarez B, Picinelli A, Dapena E. Study of the phenolic profile of cider apple cultivars at maturity by multivariate techniques. *J Agric Food Chem*. 1999; 47(10): 4046-52.
- [18] Caligiani A, Acquotti D, Palla G, Bocchi V. Identification and quantification of the main organic components of vinegars by high resolution 1H NMR spectroscopy. *Anal Chim Acta*. 2007; 585(1): 110-9.
- [19] Mosawy S, Gaiz A, Karaksha A, Singh I. The Green Tea Extract Epigallocatechin Gallate Inhibits Human Platelet Function but not Plasma Coagulation. *International Journal of Prevention and Treatment*. 2016; 5(2): 17-21.
- [20] Mosawy S. Effect of the flavonol quercetin on human platelet function: A review. *Food and Public Health*. 2015; 5(1): 1-9.
- [21] Johnston CS, White AM, Kent SM. Preliminary evidence that regular vinegar ingestion favorably influences hemoglobin A1c values in individuals with type 2 diabetes mellitus. *Diabetes research and clinical practice*. 2009; 84(2): e15-e7.
- [22] Shishehbor F, Mansoori A, Sarkaki A, Jalali M, Latifi S. Apple cider vinegar attenuates lipid profile in normal and diabetic rats. *Pakistan journal of biological sciences: PJBS*. 2008; 11(23): 2634-8.
- [23] Halima BH, Sarra K, Houda BJ, Sonia G, Abdallah A. Antihyperglycemic, Antihyperlipidemic and Modulatory Effects of Apple Cider Vinegar on Digestive Enzymes in Experimental Diabetic Rats. *International Journal of Pharmacology*. 2016; 12(5).
- [24] Ogawa N, Satsu H, Watanabe H, Fukaya M, Tsukamoto Y,

- Miyamoto Y, et al. Acetic acid suppresses the increase in disaccharidase activity that occurs during culture of caco-2 cells. *The Journal of nutrition*. 2000; 130(3): 507-13.
- [25] White AM, Johnston CS. Vinegar ingestion at bedtime moderates waking glucose concentrations in adults with well-controlled type 2 diabetes. *Diabetes Care*. 2007; 30(11): 2814-5.
- [26] Iman M, Moallem SA, Barahoyee A. Effect of apple cider vinegar on blood glucose level in diabetic mice. *Pharmaceutical Sciences*. 2015; 20: 163.
- [27] Hlebowicz J, Darwiche G, Björgell O, Almér L-O. Effect of apple cider vinegar on delayed gastric emptying in patients with type 1 diabetes mellitus: a pilot study. *BMC gastroenterology*. 2007; 7(1): 1.
- [28] Liljeberg H, Björck I. Delayed gastric emptying rate may explain improved glycaemia in healthy subjects to a starchy meal with added vinegar. *European Journal of Clinical Nutrition*. 1998; 52(5): 368-71.
- [29] Fushimi T, Tayama K, Fukaya M, Kitakoshi K, Nakai N, Tsukamoto Y, et al. Acetic acid feeding enhances glycogen repletion in liver and skeletal muscle of rats. *The Journal of nutrition*. 2001; 131(7): 1973-7.
- [30] Salbe AD, Johnston CS, Buyukbese MA, Tsitouras PD, Harman SM. Vinegar lacks antiglycemic action on enteral carbohydrate absorption in human subjects. *Nutr Res*. 2009; 29(12): 846-9.
- [31] Johnston CS, Steplewska I, Long CA, Harris LN, Ryals RH. Examination of the antiglycemic properties of vinegar in healthy adults. *Ann Nutr Metab*. 2010; 56(1): 74-9.
- [32] Fushimi T, Sato Y. Effect of acetic acid feeding on the circadian changes in glycogen and metabolites of glucose and lipid in liver and skeletal muscle of rats. *British journal of nutrition*. 2005; 94(05): 714-9.
- [33] Fushimi T, Tayama K, Fukaya M, Kitakoshi K, Nakai N, Tsukamoto Y, et al. The efficacy of acetic acid for glycogen repletion in rat skeletal muscle after exercise. *International journal of sports medicine*. 2002; 23(03): 218-22.
- [34] Ostman E, Granfeldt Y, Persson L, Björck I. Vinegar supplementation lowers glucose and insulin responses and increases satiety after a bread meal in healthy subjects. *European Journal of Clinical Nutrition*. 2005; 59: 983.
- [35] Bassoli BK, Cassolla P, Borba-Murad GR, Constantin J, Salgueiro-Pagadigorria CL, Bazotte RB, et al. Chlorogenic acid reduces the plasma glucose peak in the oral glucose tolerance test: effects on hepatic glucose release and glycaemia. *Cell biochemistry and function*. 2008; 26(3): 320-8.
- [36] Hemmerle H, Burger H-J, Below P, Schubert G, Rippel R, Schindler PW, et al. Chlorogenic acid and synthetic chlorogenic acid derivatives: novel inhibitors of hepatic glucose-6-phosphate translocase. *Journal of medicinal chemistry*. 1997; 40(2): 137-45.
- [37] Welsch CA, Lachance PA, Wasserman BP. Dietary phenolic compounds: inhibition of Na⁺-dependent D-glucose uptake in rat intestinal brush border membrane vesicles. *The Journal of nutrition*. 1989; 119(11): 1698-704.
- [38] Johnston CS, Kim CM, Buller AJ. Vinegar improves insulin sensitivity to a high-carbohydrate meal in subjects with insulin resistance or type 2 diabetes. *Diabetes Care*. 2004; 27(1): 281-2.
- [39] de Sotillo DVR, Hadley M. Chlorogenic acid modifies plasma and liver concentrations of: cholesterol, triacylglycerol, and minerals in (fa/fa) Zucker rats. *The Journal of nutritional biochemistry*. 2002; 13(12): 717-26.
- [40] Cho AS, Jeon SM, Kim MJ, Yeo J, Seo KI, Choi MS, et al. Chlorogenic acid exhibits anti-obesity property and improves lipid metabolism in high-fat diet-induced-obese mice. *Food Chem Toxicol*. 2010; 48(3): 937-43.
- [41] Dokken BB. *The Pathophysiology of Cardiovascular Disease and Diabetes: Beyond Blood Pressure and Lipids*. *Diabetes Spectrum*. 2008; 21(3): 160-5.
- [42] Kondo S, Tayama K, Tsukamoto Y, Ikeda K, Yamori Y. Antihypertensive effects of acetic acid and vinegar on spontaneously hypertensive rats. *Bioscience, biotechnology, and biochemistry*. 2001; 65(12): 2690-4.
- [43] Ziedrich L. Rice vinegar for home canning 2012 [cited 2016 September]. Available from: <https://agardenerstable.com/2012/06/02/rice-vinegar-for-home-canning/>.
- [44] Honsho S, Sugiyama A, Takahara A, Satoh Y, Nakamura Y, Hashimoto K. A red wine vinegar beverage can inhibit the renin-angiotensin system: experimental evidence in vivo. *Biological and Pharmaceutical Bulletin*. 2005; 28(7): 1208-10.
- [45] Darzi J, Frost G, Montaser R, Yap J, Robertson M. Influence of the tolerability of vinegar as an oral source of short-chain fatty acids on appetite control and food intake. *International Journal of Obesity*. 2014; 38(5): 675-81.
- [46] Kondo T, Kishi M, Fushimi T, Ugajin S, Kaga T. Vinegar intake reduces body weight, body fat mass, and serum triglyceride levels in obese Japanese subjects. *Bioscience, biotechnology, and biochemistry*. 2009; 73(8): 1837-43.
- [47] Hsu C-L, Huang S-L, Yen G-C. Inhibitory effect of phenolic acids on the proliferation of 3T3-L1 preadipocytes in relation to their antioxidant activity. *Journal of agricultural and food chemistry*. 2006; 54(12): 4191-7.
- [48] Bárdos L, Bender B. Effect of apple cider vinegar on plasma lipids (model experiment in mice). *Potravinarstvo*. 2012; 6(1): 1-4.
- [49] Beheshti Z, Chan YH, Nia HS, Hajhosseini F, Nazari R, Shaabani M. Influence of apple cider vinegar on blood lipids. *Life Sci J*. 2012; 9(4): 2431-40.
- [50] Fushimi T, Suruga K, Oshima Y, Fukiharu M, Tsukamoto Y, Goda T. Dietary acetic acid reduces serum cholesterol and triacylglycerols in rats fed a cholesterol-rich diet. *British Journal of Nutrition*. 2006; 95(05): 916-24.