

## **The Functional Significance of Cognitive Empathy and Theory of Mind in Early and Chronic Schizophrenia**

### Author

Canty, Allana L, Cao, Yuan, Neumann, David, Shum, David HK

### Published

2021

### Journal Title

Psychiatry Research

### Version

Accepted Manuscript (AM)

### DOI

[10.1016/j.psychres.2021.113852](https://doi.org/10.1016/j.psychres.2021.113852)

### Rights statement

© 2021 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International Licence, which permits unrestricted, non-commercial use, distribution and reproduction in any medium, providing that the work is properly cited.

### Downloaded from

<http://hdl.handle.net/10072/402958>

### Griffith Research Online

<https://research-repository.griffith.edu.au>

## Journal Pre-proof

The Functional Significance of Cognitive Empathy and Theory of Mind in Early and Chronic Schizophrenia

Allana L. Canty , Yuan Cao , David Neumann , David H.K. Shum

PII: S0165-1781(21)00149-9  
DOI: <https://doi.org/10.1016/j.psychres.2021.113852>  
Reference: PSY 113852



To appear in: *Psychiatry Research*

Received date: 24 December 2020  
Accepted date: 28 February 2021

Please cite this article as: Allana L. Canty , Yuan Cao , David Neumann , David H.K. Shum , The Functional Significance of Cognitive Empathy and Theory of Mind in Early and Chronic Schizophrenia, *Psychiatry Research* (2021), doi: <https://doi.org/10.1016/j.psychres.2021.113852>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V.

## Highlights

- Self-reported cognitive empathy better in early than chronic schizophrenia
- Theory-of-mind (ToM) had predictive value for social functioning
- Cognitive empathy mediates relationship between ToM and social functioning

Journal Pre-proof

# The Functional Significance of Cognitive Empathy and Theory of Mind in Early and Chronic Schizophrenia

Author information removed for peer review

Allana L. Canty<sup>a</sup>, Yuan Cao<sup>b</sup>, David Neumann<sup>a</sup>, David H. K. Shum<sup>b\*</sup>

<sup>a</sup>School of Applied Psychology and Menzies Health Institute Queensland, Griffith University,  
Queensland, Australia

<sup>b</sup>Department of Rehabilitation Sciences, The Hong Kong Polytechnic University, Hung Hom,  
Kowloon, Hong Kong SAR, China

## Abstract

Theoretical models suggest that it is the interplay between social cognitive processes that result in adaptive social functioning in schizophrenia. This study explored the relative contributions of, and interplay between, cognitive empathy, affective theory of mind (ToM), neurocognition, and severity of clinical symptoms, in predicting the social functioning of individuals with schizophrenia. Clinical participants (early schizophrenia  $n = 26$ , chronic schizophrenia  $n = 32$ ) were administered an ecologically valid measure of ToM (viz., the Virtual Assessment of Mentalising Ability or VAMA) and the Empathy Quotient (EQ) as part of a larger neuropsychological and social functioning assessment battery. Results

indicated that individuals with early schizophrenia reported significantly better cognitive empathy than individuals with chronic schizophrenia. ToM was found to have added value in predicting both community functioning and functional capacity that was beyond that accounted for by cognitive empathy, clinical symptoms, and neurocognition for both clinical groups. Further, our results indicated that the capacity to demonstrate empathic understanding of another's situation (i.e., cognitive empathy) mediates the relationship between ToM and social functioning. Together, our findings highlight the intricate and compounding nature of social cognition constructs, and their effect on social functioning for individuals with schizophrenia.

**Key Words:** social cognition, ToM, early psychosis, virtual reality.

## 1. Introduction

Individuals with schizophrenia often experience severe interpersonal deficiencies that limit independent living, the initiation and maintenance of interpersonal relationships, and vocational functioning (Bellack et al., 2007; Couture et al., 2006; Green et al., 2008). This study extends upon previous research that confirms the individual contributions of theory of mind (ToM) and empathy to social functioning in schizophrenia (e.g., Michaels et al., 2014; Smith et al., 2014; Fett et al., 2011), by exploring the interplay of social cognitive processes in predicting adaptive functioning in early and chronic schizophrenia.

Social cognition is an umbrella term used to collectively describe advanced cognitive functions (including ToM and empathy) that underlie an individual's ability to process, store, and apply information about other people and social situations (Montag et al., 2011).

Empathy constitutes a set of distinct processes by which one person attends to the subjective experiences of another person (Bora et al., 2008; Zaki et al., 2012). Affective empathy relies on phylogenetically emotional contagion systems and refers to the vicarious experience of the emotional experiences of others (Decety, 2011; Michaels et al., 2014). In some models, affective empathy can be further broken down into empathic concern (having concerns or worries for another person) and personal distress (discomfort upon witnessing another's difficulty; Davis, 1983). Cognitive empathy, on the other hand, refers to a set of reflective processes that include understanding the emotional state of others, distinguishing another's feelings from one's own, and being able to integrate this information with social knowledge to adaptively guide interpersonal behaviour (Bernhardt & Singer, 2012; Decety, 2011; Shamay-Tsoory, 2011).

ToM refers to the ability to attribute mental states to others (Kalbe et al., 2007; Singer et al., 2009). Cognitive ToM has been defined as the ability to make inferences about another's thoughts/beliefs, while affective ToM is the inference of another's emotional state.

Given the theoretical similarities, affective ToM and cognitive empathy have been treated as interchangeable at times (Shamay-Tsoory et al., 2010). The main difference between the two constructs is that, the former only refers to an understanding of another's state, while the latter requires the active engagement of the person in the perspective of another person (see Table 1). ToM has been conceptualized as having first- and second-order components. First-order ToM refers to understanding the perspective of an immediate other, and second-order ToM refers to the ability to understand what a third person may be thinking that immediate others is thinking, feeling, or intending.

[Table 1 here]

### **1.1. Neurocognitive correlates of empathy**

Variation in the empathic response may reflect strength of mental flexibility. Decety and Jackson (2004) suggested that the capacity to adopt someone else's point of view requires effortful and controlled diversion of attentional resources away from default self-perspective and egocentricity and the ability to shift a course of thought or action according to the demands of a social situation. This theoretical reasoning has been supported by significant correlations between empathy and measures of cognitive flexibility (Grattan & Eslinger, 1989; Shamay-Tsoory et al., 2003). These results provide an early but promising indication that deficits in specific aspects of empathy (viz., cognitive empathy) may reflect a breakdown in neurocognitive processes. Research on possible relationships between ToM and cognitive abilities has shown mixed results. There are strong bodies of evidence confirming an interplay between social cognitive and cognitive processes (e.g., with executive functions, Greig et al., 2004; memory, Mehl et al., 2010), as well as the influence of cognitive abilities on social functioning in schizophrenia (Bowie et al., 2008; Green, 1996). As such, the current study sought to capture the influence of previously identified cognitive abilities (viz., mental flexibility, attention, verbal fluency, verbal response inhibition,

working memory, and estimated IQ) relative to ToM and empathy in predicting social functioning in individuals with schizophrenia.

## 1.2. Empathy and theory of mind in schizophrenia

The findings from research using samples of individuals with autism spectrum disorder (Mathersul et al., 2013), ventromedial prefrontal cortex (vmPFC) damage (Shamay-Tsoory et al., 2003) and schizophrenia (Shamay-Tsoory et al., 2007b) have provided preliminary support for the theoretical association between empathy and ToM. These results are also consonant with neuroimaging findings, which indicate that ToM and empathy are subserved by a shared neural network (Shamay-Tsoory, et al., 2007a). Several reviews and meta-analyses have confirmed that ToM is disrupted in early (e.g., Green et al., 2012) and chronic stages of schizophrenia (Bora et al., 2009; Fett et al., 2011; Sprong et al., 2007).

Results of previous research have consistently indicated that individuals with chronic schizophrenia report diminished cognitive empathy relative to healthy adults (Achim et al., 2011; Corbera et al., 2013; Haker et al., 2012; Lee et al., 2011; Michaels et al., 2014; Smith et al., 2012; Sparks et al., 2010). The finding that cognitive empathy is impacted in schizophrenia is echoed in studies using other assessment methods, including behavioural performance (Derntl et al., 2009; Derntl et al., 2012; Smith et al., 2014), fMRI (Lee et al., 2010; Smith et al., 2014; Horan et al., 2014a) and electrophysiological tasks (McCormick et al., 2012; Corbera et al., 2013; Horan et al., 2014b).

Research exploring affective empathy in schizophrenia has returned mixed results. Although some studies indicate that individuals with schizophrenia are self-reporting similar level of affective responsivity to that reported by healthy adults (Achim et al., 2011; Haker et al., 2012; Michaels et al., 2014; Singh et al., 2011), others have concluded that this clinical group reports diminished empathic concern and/or heightened personal distress (Lee et al., 2011; Shamay-Tsoory et al., 2007a, 2007b; Sparks et al., 2010).



Research has also started to explore the stability of empathic deficiencies across schizophrenia. Montag, Heinz, Kunz, and Gallinat (2007) found that ratings of empathic perspective taking, but not affective empathy, negatively correlated with duration of illness. Similarly, results of a meta-analysis revealed a significantly lower effect size for differences in cognitive empathy between individuals with first episode psychosis and healthy controls relative to those reported in studies comparing healthy controls with individuals with chronic schizophrenia (Achim et al., 2011). These results suggest that cognitive empathy may be less affected in the early stage of schizophrenia, but deteriorates with illness chronicity. Similar findings were reported for ToM; first-order cognitive ToM was found to be preserved in early psychosis (Canty et al., 2017b). Individuals with chronic schizophrenia were found to have poorer performance on all dimensions of a ToM task compared to those with early psychosis (Canty et al., 2017b).

### **1.3. Relationship between empathy, ToM and social functioning**

There is preliminary evidence to suggest that empathy (Michaels et al., 2014; Smith et al., 2014) and ToM (Fett et al., 2011; Pijnenborg et al., 2009; Zhu et al., 2007) account for significant incremental variance in social functioning after accounting for relevant neurocognitive and psychopathological variables. The integrative mediation model described by Ofir-Eyal and colleagues (2014) explains the decline in social functioning commonly observed in individuals with schizophrenia. According to this model, the interpersonal problems of persons with schizophrenia may emerge because their interpretation of the goals, intentions, and emotions of others preclude accurate empathic understanding and emotional displays congruent with the other persons' emotional experience. In turn, inaccurate empathising can reduce the fluidity, intimacy, and quality of social interactions (Harvey et al., 2013; Lee et al., 2011).

Only one study to date has attempted to explicitly test whether empathy mediates the relationship between social cognition (viz., emotion recognition and ToM) and social functioning. Utilising a sample of 30 individuals with schizophrenia or schizoaffective disorder, Sparks et al. (2010) found that impairment in the comprehension of sarcasm (viz., ToM) was associated with higher empathic personal distress (i.e., affective empathy), and lower recreational functioning. However, in this study, empathy could not be explored as a mediator of the association between social cognition and functional outcome due to a lack of common associations with functional outcome measures.

#### **1.4 Aims and hypotheses**

The primary aim of this study was to examine the added value of cognitive empathy above that of ToM, neurocognition, and severity of clinical symptoms in predicting social functioning in early and chronic schizophrenia. Based on the findings of Michaels et al. (2014) and Smith et al. (2012), it was anticipated that individuals with early schizophrenia will report their cognitive empathy as significantly higher than that of individuals with chronic schizophrenia. It was also expected that cognitive empathy would account for a unique portion of variance in social functioning for both clinical groups, after controlling for cognitive and affective ToM, neurocognition and clinical symptoms.

## **2. Method**

### **2.1. Participants**

All clinical participants had a DSM-5 diagnosis (APA, 2013) of schizophrenia and were recruited from either an early psychosis intervention or mental health rehabilitation program in Queensland, Australia. Diagnoses of schizophrenia were determined by medical records and consultation with the participant's treating psychiatrist. Clinical participants were stabilised on atypical antipsychotic medications for at least 1 month prior to participation. The early schizophrenia group included 15 inpatients and 11 outpatients (see Table 1). All

participants within this group had experienced their first psychotic episode within two years of participating in this study, with most individuals recovering from their first psychotic episode proximal to the time of participation (i.e., within 3 months of their first treatment by mental health services; Green et al., 2012; Saeedi et al., 2007). The chronic schizophrenia group included 15 inpatients and 18 outpatients. Chronic patients had experienced multiple psychotic episodes and were within 2 to 15 years of illness.

All clinical participants understood spoken English sufficiently to comprehend testing procedures and exhibited no physical or language impairment that could adversely affect task performance. Individuals with schizoaffective, schizophreniform, and bipolar disorder, IQ less than 70, or histories of neurological disorder/brain injury (e.g., epilepsy, brain tumour, previous head trauma requiring hospitalisation) were excluded. Sociodemographic, clinical variables, and medication dosage in terms of chlorpromazine equivalent (i.e., CPZ; Gardner et al., 2010) were gathered from medical records.

Consistent with earlier reports on the demographics of cohorts with schizophrenia (Ochoa et al., 2012), individuals with early schizophrenia were significantly younger than individuals with chronic schizophrenia ( $t(39.85) = 5.09, p < .001$ ). Individuals with early schizophrenia did not differ from individuals with chronic schizophrenia on estimated IQ ( $t(57) = -1.98, p = .053$ ). Symptom levels and CPZ equivalents are comparable to other studies using schizophrenia samples (e.g., Fretland et al., 2015).

All participants provided informed written consent to a protocol approved by the first author's institutional research ethics committee. Participants were individually administered all measures during a 2-hour assessment session. The EQ was administered after the ToM tasks and prior to the neurocognitive assessment battery.

## **2.2. Measures and procedure**

### **2.2.1. Symptom assessment**

Fretland et al.'s (2015) categorisation of the items of the Positive and Negative Syndrome Scale (PANSS) was used to assess the positive, negative and disorganised symptom severity for schizophrenia (Wallwork et al., 2012). Higher scores indicate higher symptom severity.

### **2.2.2. Empathy**

The Empathy Quotient (EQ; Baron-Cohen, 2003; Baron-Cohen & Wheelwright, 2004) was used to provide indices of cognitive empathy (items 25, 26, 44, 52, 54), emotional reactivity (items 4, 8, 12, 14, 35) and social skills (items 6, 27, 32, 50, 59; Muncer & Ling, 2006) using a forced-choice scale (viz., strongly agree to strongly disagree). Higher scores reflect better perceived empathic abilities. The EQ has been shown to be sensitive to variations in empathy abilities in the schizophrenia population (Bora et al., 2008; Didehbani et al., 2012), and the cognitive empathy subscale has a strong internal consistency ( $\alpha = .84$ , Muncer & Ling, 2006).

### **2.2.3. Theory of mind**

The Virtual Assessment of Mentalising Ability (VAMA) uses a virtual interface to simulate the demands of real-life social interactions (internal consistency ranged from 0.69 to 0.84; Canty et al., 2017a). Participants are required to navigate a virtual shopping centre and complete a list of errands (e.g., post a letter at the post office), during which they respond to multiple-choice questions about 10 social interactions that occur between the test-taker and his/her virtual 'friends'. Two neutral interactions are presented and are followed by questions that require non-mental state reasoning (e.g., "what size shirt did John want to try on?"). These serve as control questions to check accurate encoding of task content and ability to

undertake the VAMA. Performance on the VAMA is divided into indices of first- and second-order cognitive and affective ToM. For the purpose of this study, the first- and second-order ToM scores were pooled. ToM score in this study refers to the total pooled score from both cognitive and affective ToM indices. Higher scores indicate higher accuracy on this ToM task.

#### **2.2.4. Neurocognitive assessment**

Considering the effects of neuro-cognitive abilities on empathy (Decety & Jackson, 2004) and ToM (e.g., Greig et al., 2004), commonly used measures with adequate psychometric properties were selected to provide information about separate cognitive domains including attention (Digit Span subtest of the Wechsler Adult Intelligence Scale IV, Wechsler, 2008), mental flexibility (Trail Making Test: Part B; Reitan & Wolfson, 1985), verbal fluency (Controlled Oral Word Association Task; COWAT; Benton et al., 1978), verbal response inhibition (Hayling Test; Burgess & Shallice, 1997), working memory (Letter Number Sequencing subtest of the Wechsler Memory Scale III; LNS; Wechsler, 2003), and estimated IQ (i.e., as measured by the Vocabulary and Matrix Reasoning subtests from the Wechsler Abbreviated Scale of Intelligence II, Wechsler & Hsiao-pin, 2011).

#### **2.2.5. Social functioning assessment**

We used two measures of social functioning, a task-based measure of social skills, and a subjective self-report of community functioning. The Social Skills Performance Assessment (Patterson et al., 2001) is a behavioural measure of social skills used to provide an index of functional capacity (i.e., social skill; inter-rater reliability  $r = .89$ , test-retest reliability =  $.91$ ; McClure et al., 2007). Higher scores suggest better social skills. The Social Functioning Scale (Birchwood et al., 1990) is a 79-item questionnaire that requires subjective ratings within seven domains of community functioning: social engagement, interpersonal functioning, competence for independent living, performance of independent living tasks,

engagement in recreational hobbies, and prosocial behaviour. Total scores were used for measure of social functioning ( $\alpha = .80$ ; Birchwood et al., 1990). Higher scores indicate better self-reported community functioning.

### 2.3. Data Analyses

All raw data were screened according to the procedures outlined by Tabachnick and Fidell (2007). Given that IQ and neurocognitive abilities were significantly different between individuals with early and chronic schizophrenia and are dissociable from illness chronicity, a composite measure of neurocognition (including IQ) was used as a statistical control in all analyses. Group differences in empathic abilities, as assessed by the EQ, were explored using univariate ANCOVAs with the composite measure of neurocognition used as the covariate. Separate hierarchical multiple regressions were conducted for individuals with early and chronic schizophrenia to investigate the extent to which ToM and cognitive empathy account for incremental variance in functional capacity and community functioning, after controlling for neurocognitive abilities and severity of clinical symptoms. The composite measure of neurocognitive abilities was entered as Step 1. Positive, negative and disorganised symptoms were entered at Step 2. Total ToM performance on the VAMA (i.e., first- and second-order cognitive and affective ToM) and cognitive empathy on the EQ were entered at Steps 3 and 4, respectively. The mediated effect of affective ToM (i.e., composite of first- and second-order indices) on community functioning and functional capacity, via cognitive empathy, was tested using the Baron and Kenny's (1986) four-step approach, with neurocognition and clinical symptoms entered as covariates. In the absence of a sufficient sample size, tests of indirect effects (e.g., using the 95% bias-corrected bootstrap procedure; Preacher & Hayes, 2004) were considered inappropriate.

### 3. Results

#### 3.1. Demographic and clinical variables

The clinical and demographic characteristics of the individuals with early and chronic schizophrenia are presented in Table 2.

[Table 2 here]

#### 3.2. Self-Reported empathy in schizophrenia

Participants' mean ratings of cognitive empathy, emotional reactivity, and social skills, as measured by the EQ, are summarised in Figure 1. The ANCOVA results indicated that individuals with early schizophrenia rated their cognitive empathy ( $F(1, 50) = 10.83, p = .002, n_p^2 = .18$ ) and their emotional reactivity ( $F(1, 50) = 5.59, p < .05, n_p^2 = .10$ ) significantly better than did individuals with chronic schizophrenia. Significant differences were not observed between the two groups on self-reported ratings of social skills ( $F(1, 50) = .29, p = .59$ ).

[Figure 1 here]

#### 3.3. Factors associated with empathy in early and chronic schizophrenia

##### 3.3.1. Empathy and theory of mind

Significant moderate correlations were observed between cognitive empathy and second-order cognitive and total ToM for individuals with early schizophrenia, and with first-order cognitive and total ToM for individuals with chronic schizophrenia (Table 3).

Significant moderate correlations were also observed between social skills and second-order affective ToM for individuals with early schizophrenia (Table 3).

[Table 3 here]

### 3.3.2. Social cognitive correlates of social functioning]

Cognitive empathy scores were significantly and moderately associated with indices of functional capacity and community functioning in individuals with early and chronic schizophrenia (Table 3). Moderate significant correlations were observed between first- and second-order cognitive and affective ToM and functional capacity for both individuals with early and chronic schizophrenia (Table 4).

[Table 4 here]

### 3.4. Regression analyses

#### 3.4.1. Community functioning

When added to the model, cognitive empathy accounted for 9% incremental variance in community functioning for individuals with early schizophrenia ( $\Delta F(1, 16) = 78.73, p < .001$ ; Table 5), and 10% for individuals with chronic schizophrenia ( $\Delta F(1, 17) = 98.16, p < .05$ ). In the final models, both total ToM performance ( $\beta = .39, p < .001$ ) and cognitive empathy ( $\beta = .31, p < .001$ ) explained significant variance in community functioning for individuals with early and chronic schizophrenia (ToM  $\beta = .43, p < .001$ ; cognitive empathy  $\beta = .33, p < .001$ ).

#### 3.4.2. Functional capacity

When added to the model, cognitive empathy accounted for 5.9% incremental variance in functional capacity for individuals with early schizophrenia ( $\Delta F(1, 16) = 15.96, p = .001$ ; Table 6) and 7.9% for individuals with chronic schizophrenia ( $\Delta F(1, 17) = 18.62, p < .001$ ). In the final model, both total ToM performance ( $\beta = .52, p = .005$ ) and cognitive empathy ( $\beta = .48, p = .001$ ) explained significant variance in functional capacity for individuals with early and chronic schizophrenia (ToM  $\beta = .49, p < .001$ ; cognitive empathy  $\beta = .21, p < .01$ ).



[Table 5 and 6 here]

### 3.5. Mediation models

According to the statistical procedures described by Baron and Kenny (1986), mediation is present when the independent variable (affective ToM) significantly predicts the dependent variable (social functioning), as described in Path C, but this relationship is substantially minimised, or is no longer significant, when the proposed mediator (cognitive empathy) is added as an independent variable in the multiple regression model.

#### 3.5.1 Community functioning

##### 3.5.1.1. Early schizophrenia

As seen in Figure 2, results for Path A indicated that affective ToM was positively and significantly associated with cognitive empathy for individuals with early schizophrenia ( $\beta = .65, p < .001; R^2 = .74$ ). In the second model, which assessed Path B, a significant relationship was observed between cognitive empathy and community functioning ( $\beta = .63, p < .001; R^2 = .64$ ). Results for Path C (without the mediator) indicated a significant positive association between affective ToM and social functioning ( $\beta = .82, p < .001; R^2 = .67$ ). In the final model, the third analysis was repeated with the inclusion of the proposed mediator (cognitive empathy) in the model (Table 7) and was found to be a significant predictor of community functioning. The standardised beta-weight for affective ToM decreased from .82 ( $p < .001$ ) to .16 ( $p = .23$ ) in the fourth model, indicating that cognitive empathy mediated the relationship between affective ToM and community functioning for individuals with early schizophrenia.

[Figure 2]

[Table 7]

### 3.5.1.2. Chronic schizophrenia

The aforementioned steps were repeated for individuals with chronic schizophrenia (see Figure 3). Results for Path A indicated that affective ToM was positively and significantly associated with cognitive empathy ( $\beta = .78, p = .002; R^2 = .70$ ). Results for Path B indicated a highly significant relationship between cognitive empathy and community functioning ( $\beta = .58, p < .001; R^2 = .47$ ). Results for Path C (without the mediator) indicated a significant positive association between affective ToM and community functioning ( $\beta = .73, p < .01; R^2 = .63$ ). Cognitive empathy remained a significant predictor in the final model, but affective ToM no longer contributed significantly to the variance in community functioning. The standardised beta-weight for affective ToM decreased from .73 ( $p < .01$ ) to .11 ( $p = .23$ ) in the fourth model, indicating that cognitive empathy mediated the relationship between affective ToM and community functioning for individuals with chronic schizophrenia.

[Figure 3]

### 3.5.2. Functional capacity

#### 3.5.2.1. Early schizophrenia

As seen in Figure 4, results for Path A indicated that affective ToM was positively and significantly associated with cognitive empathy for individuals with early schizophrenia ( $\beta = .65, p < .001; R^2 = .74$ ). In the second model, which assessed Path B, a significant relationship was observed between cognitive empathy and functional capacity ( $\beta = .74, p < .001; R^2 = .76$ ). Results for Path C (without the mediator) indicated a significant positive association between affective ToM and functional capacity ( $\beta = .78, p < .001; R^2 = .79$ ). In the final model, the third analysis was repeated with the inclusion of the proposed mediator (cognitive empathy) in the model (Table 8) and was found to be a significant predictor of

functional capacity. The standardised beta-weight for affective ToM decreased from .78 ( $p < .001$ ) to .64 ( $p = .001$ ) but remained statistically significant in the fourth model. Although these results indicate that cognitive empathy may partially mediate the relationship between affective ToM and functional capacity, the inclusion of the mediator in the model did not have a large influence on the predictive value of affective ToM. Thus, the influence of cognitive empathy should be interpreted with caution.

[Figure 4]

[Table 8]

### 3.5.2.2. Chronic schizophrenia

The aforementioned steps were repeated for individuals with chronic schizophrenia (see Figure 5). Results for Path A indicated that affective ToM was positively and significantly associated with cognitive empathy ( $\beta = .78, p = .002; R^2 = .70$ ). Results for Path B indicated a highly significant relationship between cognitive empathy and functional capacity ( $\beta = .55, p < .001; R^2 = .68$ ). Results for Path C (without the mediator) indicated a significant positive association between affective ToM and functional capacity ( $\beta = .75, p = .002; R^2 = .71$ ). Cognitive empathy remained a significant predictor in the final model, but affective ToM no longer contributed significantly to the variance in functional capacity. The standardised beta-weight for affective ToM decreased from .75 ( $p = .002$ ) to .01 ( $p = .94$ ) in the fourth model, indicating that cognitive empathy mediated the relationship between affective ToM and functional capacity for individuals with chronic schizophrenia.

## 4. Discussion

This study examined the functional impact of cognitive empathy relative to that of ToM, neurocognition, and severity of clinical symptoms in early and chronic schizophrenia. Results indicated three key findings, which supported our hypotheses. First, individuals with early schizophrenia self-reported significantly better cognitive empathy than individuals with

chronic schizophrenia. Second, total ToM emerged as a stronger predictor of social functioning than cognitive empathy, neurocognition, and clinical symptoms. Third, there was evidence to suggest that affective ToM is a prerequisite of cognitive empathy, and that cognitive empathy mediates the association between affective ToM and social functioning.

#### **4.1. Self-reported empathy abilities in schizophrenia**

The findings that individuals with early schizophrenia report their cognitive empathy and emotional reactivity significantly higher than do individuals with chronic schizophrenia are consistent with the results reported by Achim et al. (2011) and Montag et al. (2007) and may indicate that cognitive empathy is less impacted in the early stage of illness.

Furthermore, these findings are consonant with evidence that individuals with chronic schizophrenia show diminished performance on behavioural and physiological measures of emotional reactivity, such as spontaneous mimicry of others' observable emotional expressions or behaviours (Sestito et al., 2015; Varcin et al., 2010).

#### **4.2. Empathy and social functioning**

Previous studies in the schizophrenia area suggest that some aspects of social cognition, particularly ToM, distinctly contribute to poor social functioning above and beyond the contributions of cognitive dysfunction and clinical symptoms (e.g., Roncone et al., 2002; Pinkham & Penn, 2006; Brüne et al., 2011). The current study extended this research by indicating that cognitive empathy, a key social cognitive process that has received relatively limited attention in schizophrenia, is a valuable consideration when explaining social functioning in early and chronic schizophrenia. Consonant with previous research (Michaels et al., 2014; Smith et al., 2012; Smith et al., 2014), cognitive empathy was found to account for incremental variance in social functioning (viz., community functioning and functional capacity) in early and chronic schizophrenia after controlling for neurocognition and clinical symptoms.

Although cognitive empathy was found to contribute significantly to social functioning in schizophrenia, ToM emerged as the strongest predictor of social functioning in the final models for both clinical groups. This result may reflect an advantage for ToM versus cognitive empathy in predicting social functioning. Previously it was proposed that ToM, or the ability to infer another's state, is a pre-requisite to other social cognitive abilities, including empathy (Kalbe et al., 2007; Singer et al., 2009). It may therefore be speculated that the ability to make such inferences about another is fundamental for social functioning.

#### **4.3. Mediation model**

As noted previously, the only previous study we are aware of, that attempted to assess the mediating influence of cognitive empathy on the relationship between social cognition and social functioning in schizophrenia found no evidence of common associations between these variables, and thus could not proceed with the mediation analysis (Sparks et al., 2010). According to Baron and Kenny's (1986) criteria, the present results indicated that cognitive empathy qualified as a mediator of the relationship between affective ToM and community functioning in early and chronic schizophrenia, and between affective ToM and social functioning for individuals with chronic schizophrenia. These results provide support for the adapted version of the integrative mediation model (Ofir-Eyal et al., 2014) and indicate that affective ToM is a prerequisite of cognitive empathy. That is, the capacity to understand and adapt one's affective behaviour to the subjective experience of those with whom we interact is secondary to being able to predict their thoughts, beliefs, and intentions. Although accurate mental state reasoning may promote appreciation and understanding of another's subjective experience, it is the latter social cognitive process that translates into positive functional outcomes. This idea is in line with results in the psychopathy literature, which suggests that mental state reasoning is dissociable from appreciating or having concerns for another person (van Dongen, 2020). Together, these results raise the hypothesis that affective ToM is a

'purer' ability, while cognitive empathy is at least partly driven by a need or desire to appreciate another's state. This motivational aspect may be a crucial link in translating ToM ability into positive social functioning.

#### **4.4. Implications and limitations**

Research that explores the relationships between social cognitive domains can develop clinical and theoretical understanding of the nature and functional impact of these abilities in psychotic illnesses. The present results indicate that cognitive empathy and ToM may have similar value as treatment targets to neurocognition, and possibly clinical symptoms, for functional recovery-oriented interventions. The consistency of findings across measures of community functioning (i.e., what a person actually does) and functional capacity (i.e., what a person can do) indicates that cognitive empathy may be a strong predictor of the extent to which an individual with schizophrenia can function adaptively and independently (Smith et al., 2012). Emerging evidence indicates that social cognitive deficits are amenable to psychosocial intervention in schizophrenia (Kurtz & Richardson, 2012). There is also growing interest in psychopharmacological approaches to enhancing social cognition (Carter et al., 2009; Goldman et al., 2011; Rosenfeld et al., 2011). The inclusion of cognitive empathy and ToM in these treatment development efforts may achieve greater improvements in social cognition and overall functioning in early and chronic schizophrenia.

Although our findings provide insight into the relationship between social cognitive domains and social functioning, they must be interpreted in the context of some limitations. First, the absence of a healthy control group within the design of the present study precludes discussion of whether the self-reported empathy of individuals with early and chronic schizophrenia is clinically impaired. However, because the focus of the study was on examining the predictive contributions of cognitive empathy and affective ToM on social

functioning of individuals with schizophrenia, we believe that the current patient data has the clinical and research implications as described above.

Second, it is important to note that the current findings are based on self-reported levels of empathy. Specifically, the EQ measures individuals' beliefs about their own empathic skills, and the ratings of people with deficits in awareness or knowledge of their cognitive abilities (meta-cognition) could be unreliable. Nevertheless, the EQ was identified as the most theoretical and psychometrically sound measure of cognitive empathy available at the time that this study was conceived and executed. Furthermore, studies using alternative methods, such as behavioural and functional neuroimaging tasks, suggest that individuals with schizophrenia show impairments in multiple processes that underlie empathy, including emotional contagion, emotion attribution, and empathic accuracy (Benedetti et al., 2009; Brüne et al., 2011; Derntl et al., 2009; Lee et al., 2011). It remains to be determined how the cognitive empathy subscale and other subscales of the EQ relate to these processes in schizophrenia. Considering that people with schizophrenia may have impaired insight and in turn under or over-represent their empathic abilities, it is important for future studies to measure their performance on behavioural tasks of empathy, in addition to traditional self-report measures such as the EQ.

Third, although significant associations were observed between cognitive empathy and indices of social functioning, the modest sample sizes may have limited the ability to detect meaningful relations between these indices and other processes assessed by the EQ. Notwithstanding, previous researchers have acknowledged the difficulties inherent in recruiting large samples of clinical participants, and proceeded to perform mediational analyses with less than optimal sample sizes ( $n = 41$ , Chung, Matthews, & Barch, 2011;  $n = 27$ , Ownsworth, Henderson, Chambers, & Shum, 2009).

Comprehensive and reliable assessment measures that capture the nature of ToM impairment have the potential to tailor treatment protocols for individuals with schizophrenia. The VAMA could be used to identify a person's strengths and weaknesses within ToM sub-processes (Canty et al., 2017b), as well as be used to assess clinical change. Expanding on earlier research (Bechi et al., 2013) which demonstrated that individuals with schizophrenia benefit from ToM training, future studies should examine how ToM assessment results and training translate into real-world outcomes (e.g., quality of relationships, employment, and community integration).

#### **4.5. Conclusion**

In conclusion, findings of our study provide further support for the functional significance of cognitive empathy in early and chronic schizophrenia. These findings also contribute to emerging evidence that cognitive empathy is less affected in the early stage of schizophrenia. There is evidence to suggest that affective ToM is a prerequisite of cognitive empathy, and that cognitive empathy mediates the relationship between affective ToM and social functioning. Efforts to target these processes in social cognitive training programs for schizophrenia may be a useful way to enhance the generalisability of these interventions to real-life settings. Further clarification of how cognitive and affective empathy are impacted, and continued efforts to develop theoretical, multidimensional measures of empathy, will contribute to a more sophisticated understanding of empathy processes in schizophrenia.

**Funding:** The first author was supported by an Australian Government's PhD scholarship. The second author is supported by a Postdoctoral Fellowships Scheme from The Hong Kong Polytechnic University (Work Program: YW4V, Project ID: P0031571).



### Author Statement

**Allana Cauty:** Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Software; Validation; Writing - original draft; Writing - review & editing.

**Yuan Cao:** Conceptualization; Investigation; Methodology; Validation; Writing - review & editing.

**David Neumann:** Conceptualization; Investigation; Methodology; Project administration; Software; Validation; Supervision; Writing - original draft; Writing - review & editing.

**David Shum:** Conceptualization; Investigation; Resources; Methodology; Project administration; Software; Validation; Supervision; Writing - original draft; Writing - review & editing.

### Conflict of interest

We have no conflict of interest to declare.

## References

- Achim, A. M., Ouellet, R., Roy, M.-A., & Jackson, P. L. (2011). Assessment of empathy in first-episode psychosis and meta-analytic comparison with previous studies in schizophrenia. *Psychiatry Research, 190*(1), 3-8. doi: 10.1016/j.psychres.2010.10.030
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders*. (5th ed.). Arlington, VA: Author. Saeedi, Addington, & Addington, 2007
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology, 51*(6), 1173-1182. doi: 10.1037//0022-3514.51.6.1173
- Baron-Cohen, S. (2003). *The essential difference: Men, women and the extreme male brain*. UK: Penguin Press Science.
- Baron-Cohen, S., & Wheelwright, S. (2004). The Empathy Quotient: An investigation of adults with Asperger syndrome or high functioning autism, and normal sex differences. *Journal of Autism and Developmental Disorders, 34*(2), 163-175. doi: 10.1023/B:JADD.0000022607.19833.00
- Bechi, M., Spangaro, M., Bosia, M., Zanoletti, A., Fresi, F., Buonocore, M., Cocchi, F., Guglielmino, C., Smeraldi, E., & Cavallaro, R. (2013). Theory of mind intervention for outpatients with schizophrenia. *Neuropsychological Rehabilitation, 23*, 383-400.
- Bellack, A. S., Green, M. F., Cook, J. A., Fenton, W., Harvey, P. D., Heaton, R. K., ...Wykes, T. (2007). Assessment of community functioning in people with schizophrenia and other severe mental illnesses: A white paper based on an NIMH-sponsored workshop. *Schizophrenia Bulletin, 33*(3), 805-822. doi: 10.1093/schbul/sbl035
- Benedetti, F., Bernasconi, A., Bosia, M., Cavallaro, R., Dallspezia, S., Falini, A., ...Smeraldi, E. (2009). Functional and structural brain correlates of theory of mind and empathy deficits in

- schizophrenia. *Schizophrenia Research*, 114(1), 154-160. doi: 10.1016/j.schres.2009.06.021
- Benton, A. L., Varney, N. R., & de Hamsher, K. (1978). Lateral differences in tactile directional perception. *Neuropsychological Research Journal*, 16(1), 109-114. doi:10.1016/0028-3932(78)90049-0
- Bernhardt, B. C., & Singer, T. (2012). The neural basis of empathy. *Annual Review of Neuroscience*, 35(1), 1-23. doi: 10.1146/annurev-neuro-062111-150536
- Birchwood, M., Smith, J., Cochrane, R., Wetton, S., & Copestake, S. (1990). The social functioning scale: The development and validation of a new scale. *British Journal of Psychiatry* 157, 853-859. doi. 10.1192/bjp.157.6.853
- Bora, E., Gökçen, S., & Veznedaroglu, B. (2008). Empathic abilities in people with schizophrenia. *Psychiatry Research*, 160(1), 23-29. doi: 10.1016/j.psychres.2007.05.017
- Bora, E., Yucel, M., & Pantelis, C. (2009). Theory of mind impairment in schizophrenia: Meta-analysis. *Schizophrenia Research*, 109(1), 1-9. doi: 10.1016/j.schres.2008.12.020
- Bowie, C., Leung, W., Reichenberg, A., McClure, M., Patterson, T., Heaton, R., & Harvey, P. D. (2008). Predicting schizophrenia patients' real-world behaviour with specific neuropsychological and functional capacity measures. *Biological Psychiatry*, 63, 505-511. doi: 10.1016/j.biopsych.2007.05.022
- Brüne, M., Schaub, D., Juckel, G., & Langdon, R. (2011). Social skills and behavioral problems in schizophrenia: The role of mental state attribution, neurocognition and clinical symptomatology. *Psychiatry Research*, 190(1), 9-17. doi: 10.1016/j.psychres.2010.03.015
- Burgess, P. W., & Shallice, T. (1997). *The Hayling and Brixton Tests*. Bury St Edmunds, UK: Thames Valley Test Company.
- Canty, A., Neumann, D., Fleming, J., & Shum, D. (2017a). Evaluation of a newly developed measure of theory of mind: The Virtual Assessment of Mentalising Ability.

*Neuropsychological Rehabilitation*, 27, 834-870. doi:

10.1080/09602011.2015.1052820

Canty, A. L., Neumann, D. L., & Shum, D. H. K. (2017b). Using virtual reality to assess theory of mind subprocesses and error types in early and chronic schizophrenia.

*Schizophrenia Research: Cognition*, 10, 15-19. doi: 10.1016/j.scog.2017.09.001

Carter, C. S., Barch, D. M., Gur, R., Gur, R., Pinkham, A., & Ochsner, K. (2009). CNTRICS final task selection: Social cognitive and affective neuroscience-based measures. *Schizophrenia Bulletin*, 35(1), 153-162. doi: 10.1093/schbul/sbn157

Chung, Y. S., Mathews, J. R., & Barch, D. M. (2011). The effect of context processing on different aspects of social cognition in schizophrenia. *Schizophrenia Bulletin*, 37(5), 1048-1056. doi: 10.1093/schbul/sbq012

Corbera, S., Wexler, B. E., Ikezawa, S., & Bell, M. D. (2013). Factor structure of social cognition in schizophrenia: Is empathy preserved? *Schizophrenia Research and Treatment*, 2013, 1-13. doi: 10.1155/2013/409205

Couture, S. M., Penn, D. L., & Roberts, D. L. (2006). The functional significance of social cognition in schizophrenia: A review. *Schizophrenia Bulletin*, 32, 44-63. doi: 10.1093/schbul/sb1029

Davis, M. H. (1983). Measuring individual differences in empathy: Evidence for a multidimensional approach. *Journal of Personality and Social Psychology*, 44(1), 113-126.

<https://doi.org/10.1037/0022-3514.44.1.113>

Decety, J. (2011). Dissecting the neural mechanisms mediating empathy. *Emotion Review*, 3(1), 92-108. doi: 10.1177/1754073910374662

Decety, J., & Jackson, P. L. (2004). The functional architecture of human empathy. *Behavioral and Cognitive Neuroscience Reviews*, 3(2), 71-100. doi: 10.1177/1534582304267187

Derntl, B., Finkelmeyer, A., Toygar, T. K., Hülsmann, A., Schneider, F., Falkenberg, D. I., & Habel, U. (2009). Generalized deficit in all core components of empathy in schizophrenia.

*Schizophrenia Research*, 108(1), 197-206. doi: 10.1016/j.schres.2008.11.009

Derntl, B., Finkelmeyer, A., Voss, B., Eickhoff, S. B., Kellermann, T., Schneider, F., & Habel, U.

(2012). Neural correlates of the core facets of empathy in schizophrenia. *Schizophrenia Research*, 136(1-3), 70-81. doi: 10.1016/j.schres.2011.12.018

Didehbani, N., Shad, M. U., Tamminga, C. A., Kandalaf, M. R., Allen, T. T., Chapman, S. B., &

Krawczyk, D. C. (2012). Insight and empathy in schizophrenia. *Schizophrenia Research*, 142(1-3), 246-247. doi: 10.1016/j.schres.2012.09.010

Fett, A.-K., Viechtbauer, W., Dominguez, M.-d., Penn, D. L., van Os, J., & Krabbendam, L. (2011).

The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: A meta-analysis. *Neuroscience and Biobehavioral Reviews*, 35(3), 573-588. doi: 10.1016/j.neubiorev.2010.07.001

Fretland, R. A., Andersson, S., Sundet, K., Andreassen, O. A., Melle, I., & Vaskinn, A. (2015).

Theory of mind in schizophrenia: Error types and associations with symptoms. *Schizophrenia Research*, 162(1), 42-46. doi: 10.1016/j.schres.2015.01.024

Gardner, D. M., Murphy, A. L., O'Donnell, H., Centorrino, F., & Baldessarini, R. J. (2010).

International consensus study of antipsychotic dosing. *The American Journal of Psychiatry*, 167, 686-693. doi: 10.1176/appi.ajp.2009.09060802

Goldman, M. B., Gomes, A. M., Carter, C. S., & Lee, R. (2011). Divergent effects of two different

doses of intranasal oxytocin on facial affect discrimination in schizophrenic patients with and without polydipsia. *Psychopharmacology*, 216(1), 101-110. doi: 10.1007/s00213-011-2193-8

Grattan, L. M., & Eslinger, P. J. (1989). Higher cognition and social behavior: Changes in cognitive

flexibility and empathy after cerebral lesions. *Neuropsychology*, 3(3), 175-185. doi: 10.1037/h0091764

Green, M. F. (1996). What are the functional consequences of neurocognitive deficits in

schizophrenia? *American Journal of Psychiatry*, 153(3), 321-330.

- Green, M. F., Bearden, C. E., Cannon, T. D., Fiske, A. P., Helleman, G. S., Horan, W. P., ... Nuechterlein, K. H. (2012). Social cognition in schizophrenia, Part 1: Performance across phase of illness. *Schizophrenia Bulletin*, *38*(4), 854-864. doi: 10.1093/schbul/sbq171
- Green, M. F., Penn, D. L., Bentall, R., Carpenter, W. T., Gaebel, W., Gur, R. C., ... Heinsen, R. (2008). Social cognition in schizophrenia: An NIMH workshop on definitions, assessment, and research opportunities. *Schizophrenia Bulletin*, *34*(6), 1211-1220. doi: 10.1093/schbul/sbm145
- Greig, T. C., Bryson, G. J., & Bell, M. D. (2004). Theory of mind performance in schizophrenia: Diagnostic, symptom, and neuropsychological correlates. *The Journal of Nervous and Mental Disease*, *192*(1), 12-18. doi: 10.1097/01.nmd.0000105995.67947.fc
- Haker, H., Schimansky, J., Jann, S., & Rössler, W. (2012). Self-reported empathic abilities in schizophrenia: A longitudinal perspective. *Psychiatry Research*, *200*(2-3), 1028-1031. doi: 10.1016/j.psychres.2012.04.004
- Harvey, P.-O., Zaki, J., Lee, J., Ochsner, K., & Green, M. F. (2013). Neural substrates of empathic accuracy in people with schizophrenia. *Schizophrenia Bulletin*, *39*(3), 617-628. doi: 10.1093/schbul/sbs042
- Horan, W. P., Pineda, J. A., Wynn, J. K., Iacoboni, M., & Green, M. F. (2014a). Some markers of mirroring appear intact in schizophrenia. *Cognitive, Affective, & Behavioral Neuroscience*, *14*(3), 1049-1060. doi: 10.3758/s13415-013-0245-8
- Horan, W. P., Wynn, J. K., Mathis, I., Miller, G. A., & Green, M. F. (2014b). Approach and withdrawal motivation in schizophrenia: An examination of frontal brain asymmetric activity: e110007. *PLoS One*, *9*(10). doi: 10.1371/journal.pone.0110007
- Kalbe, E., Grabenhorst, F., Brand, M., Kessler, J., Hilker, R., & Markowitsch, H. J. (2007). Elevated emotional reactivity in affective but not cognitive components of theory of mind: A psychophysiological study. *Journal of Neuropsychology*, *1*(Pt 1), 27.

- Kurtz, M. M., & Richardson, C. L. (2012). Social cognitive training for schizophrenia: A meta-analytic investigation of controlled research. *Schizophrenia Bulletin*, *38*(5), 1092-1104. doi: 10.1093/schbul/sbr036
- Lee, J., Zaki, J., Harvey, P. O., Ochsner, K., & Green, M. F. (2011). Schizophrenia patients are impaired in empathic accuracy. *Psychological Medicine*, *41*(11), 2297-2304. doi: 10.1017/S0033291711000614
- Lee, S. J., Lee, J.-M., Kang, D. H., Kim, C.-W., Gu, B. M., Park, J.-Y., ...Kwon, J. S. (2010). Multi-level comparison of empathy in schizophrenia: An fMRI study of a cartoon task. *Psychiatry Research: Neuroimaging*, *181*(2), 121-129. doi: 10.1016/j.psychresns.2009.08.003
- Mathersul, D., McDonald, S., & Rushby, J. A. (2013). Understanding advanced theory of mind and empathy in high-functioning adults with autism spectrum disorder. *Journal of Clinical and Experimental Neuropsychology*, *35*(6), 655.
- McClure, M. M., Bowie, C. R., Patterson, T. L., Heaton, R. K., Weaver, C., Anderson, H., & Harvey, P. D. (2007). Correlations of functional capacity and neuropsychological performance in older patients with schizophrenia: Evidence for specificity of relationships? *Schizophrenia Research*, *89*, 330-338.
- McCormick, L. M., Brumm, M. C., Beadle, J. N., Paradiso, S., Yamada, T., & Andreasen, N. (2012). Mirror neuron function, psychosis, and empathy in schizophrenia. *Psychiatry Research: Neuroimaging*, *201*(3), 233. doi: 10.1016/j.psychresns.2012.01.004
- measure of theory of mind: The Virtual Assessment of Mentalising Ability.
- Mehl, S., Rief, W., Mink, K., Lüllmann, E., & Lincoln, T. M. (2010). Social performance is more closely associated with theory of mind and autobiographical memory than with psychopathological symptoms in clinically stable patients with schizophrenia-spectrum disorders. *Psychiatry Research*, *178*(2), 276-283. doi: 10.1016/j.psychres.2009.10.004
- Michaels, T. M., Horan, W. P., Ginger, E. J., Martinovich, Z., Pinkham, A. E., & Smith, M. J.

- (2014). Cognitive empathy contributes to poor social functioning in schizophrenia: Evidence from a new self-report measure of cognitive and affective empathy. *Psychiatry Research*, 220(3), 803-810. doi: 10.1016/j.psychres.2014.08.054
- Montag, C., Dziobek, I., Richter, I. S., Neuhaus, K., Lehmann, A., Sylla, R., ... Gallinat, J. (2011). Different aspects of theory of mind in paranoid schizophrenia: Evidence from a video-based assessment. *Psychiatry Research*, 186(2), 203-209. doi: 10.1016/j.psychres.2010.09.006
- Montag, C., Heinz, A., Kunz, D., & Gallinat, J. (2007). Self-reported empathic abilities in schizophrenia. *Schizophrenia Research*, 92(1), 85-89. doi: 10.1016/j.schres.2007.01.024
- Muncer, S. J., & Ling, J. (2006). Psychometric analysis of the empathy quotient (EQ) scale. *Personality and Individual Differences*, 40(6), 1111-1119. doi: 10.1016/j.paid.2005.09.020
- Ochoa, S., Usall, J., Cobo, J., Labad, X., & Kulkarni, J. (2012). Gender differences in schizophrenia and first-episode psychosis: A comprehensive literature review. *Schizophrenia Research and Treatment*, 916198. doi:10.1155/2012/916198
- Ofir-Eyal, S., Hasson-Ohayon, I., & Kravetz, S. (2014). Affective and cognitive empathy and social quality of life in schizophrenia: A comparison between a parallel process model and an integrative meditation model. *Psychiatry Research*, 220(1-2), 51-57. doi: 10.1016/j.psychres.2014.06.049
- Owensworth, T., Henderson, L. F., Chambers, S. K., & Shum, D. (2009). Functional impairments and caregiver depressive symptoms in the context of brain tumour and other cancers: A mediating effect of strain. *Brain Impairment*, 10, 149-161. doi: 10.1375/brim.10.2.149
- Patterson, T. L., Goldman, S., McKibbin, C. L., Hughs, T., & Jeste, D. V. (2001). UCSD Performance-Based Skills Assessment: Development of a new measure of everyday functioning for severely mentally ill adults. *Schizophrenia Bulletin*, 27, 235-245. doi: 10.1093/oxfordjournals.schbul.a006870
- Pijnenborg, G. H. M., Withaar, F. K., Evans, J. J., Van Den Bosch, R. J., Timmerman, M. E., & Brouwer, W. H. (2009). The predictive value of measures of social cognition for community



- functioning in schizophrenia: Implications for neuropsychological assessment. *Journal of the International Neuropsychological Society*, 15(2), 239-247. doi:  
10.1017/S1355617709090341
- Pinkham, A. E., & Penn, D. L. (2006). Neurocognitive and social cognitive predictors of interpersonal skill in schizophrenia. *Psychiatry Research*, 143(2), 167-178. doi:  
10.1016/j.psychres.2005.09.005
- Preacher, K. J., & Hayes, A. F. (2004). SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods, Instruments, & Computers*, 36(4), 717-731. doi: 10.3758/BF03206553
- Reitan, R., & Wolfson, D. (1985). *The Halstead-Reitan Neuropsychological Test Battery: Theory and Clinical Interpretation*. Tuscon: Neuropsychology Press.
- Ronccone, R., Falloon, I. R. H., Mazza, M., De Risio, A., Pollice, R., Necozone, S., Morosini, P., & Casacchia, M. (2002). Is theory of mind in schizophrenia more strongly associated with clinical and social functioning than with neurocognitive deficits. *Psychopathology*, 35(6), 368-368. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12457019>
- Rosenfeld, A. J., Lieberman, J. A., & Jarskog, L. F. (2011). Oxytocin, dopamine, and the amygdala: A neurofunctional model of social cognitive deficits in schizophrenia. *Schizophrenia Bulletin*, 37(5), 1077-1087. doi: 10.1093/schbul/sbq015
- Sestito, M., Raballo, A., Umiltà, M. A., Amore, M., Maggini, C., & Gallese, V. (2015). Anomalous echo: Exploring abnormal experience correlates of emotional motor resonance in schizophrenia spectrum. *Psychiatry Research*, 229(1-2), 559-564. doi:  
10.1016/j.psychres.2015.05.038
- Shamay-Tsoory, S. G. (2011). The neural bases for empathy. *The Neuroscientist*, 17(1), 18-24. doi:  
0.1177/1073858410379268
- Shamay-Tsoory, S. G., Aharon-Peretz, J., & Levkovitz, Y. (2007a). The neuroanatomical basis of

affective mentalizing in schizophrenia: Comparison of patients with schizophrenia and patients with localized prefrontal lesions. *Schizophrenia Research*, 90(1), 274-283. doi: 10.1016/j.schres.2006.09.020

Shamay-Tsoory, S. G., Harari, H., Aharon-Peretz, J., & Levkovitz, Y. (2010). The role of the orbitofrontal cortex in affective theory of mind deficits in criminal offenders with psychopathic tendencies. *Cortex*, 46, 668-677.

Shamay-Tsoory, S. G., Shur, S., Barcai-Goodman, L., Medlovich, S., Harari, H., & Levkovitz, Y. (2007b). Dissociation of cognitive from affective components of theory of mind in schizophrenia. *Psychiatry Research*, 149(1), 11-23. doi: 10.1016/j.psychres.2005.10.018

Shamay-Tsoory, S. G., Tomer, R., Berger, B. D., & Aharon-Peretz, J. (2003). Characterization of empathy deficits following prefrontal brain damage: The role of the right ventromedial prefrontal cortex. *Journal of Cognitive Neuroscience*, 15(3), 324-337. doi: 10.1162/089892903321593063

Singer, T., Critchley, H. D., & Preuschoff, K. (2009). A common role of insula in feelings, empathy and uncertainty. *Trends in Cognitive Sciences*, 13(8), 334-340. doi: 10.1016/j.tics.2009.05.001

Singh, F., Pineda, J., & Cadenhead, K. S. (2011). Association of impaired EEG mu wave suppression, negative symptoms and social functioning in biological motion processing in first episode of psychosis. *Schizophrenia Research*, 130(1), 182-186. doi: 10.1016/j.schres.2011.04.004

Smith, M. J., Horan, W. P., Cobia, D. J., Karpouzian, T. M., Fox, J. M., Reilly, J. L., & Breiter, H. C. (2014). Performance-based empathy mediates the influence of working memory on social competence in schizophrenia. *Schizophrenia Bulletin*, 40(4), 824-834. doi: 10.1093/schbul/sbt084

Smith, M. J., Horan, W. P., Karpouzian, T. M., Abram, S. V., Cobia, D. J., & Csernansky, J. G.

- (2012). Self-reported empathy deficits are uniquely associated with poor functioning in schizophrenia. *Schizophrenia Research*, *137*(1-3), 196-202. doi: 10.1016/j.schres.2012.01.012
- Sparks, A., McDonald, S., Lino, B., O'Donnell, M., & Green, M. J. (2010). Social cognition, empathy and functional outcome in schizophrenia. *Schizophrenia Research*, *122*(1), 172-178.
- Sprong, M., Schothorst, P., Vos, E., Hox, J., & Van Engeland, H. (2007). Theory of mind in schizophrenia: Meta-analysis. *The British Journal of Psychiatry*, *191*(1), 5-13. doi: 10.1192/bjp.bp.107.035899
- Tabachnick, B. & Fidell, L. (2007). *Using Multivariate Statistics* (5th Ed.). Boston: Pearson.
- Van Dongen, J. D. M. (2020). The empathic brain of psychopaths: From social science to neuroscience in empathy. *Frontiers in Psychology*, *11*, 695.
- Varcin, K. J., Bailey, P. E., & Henry, J. D. (2010). Empathic deficits in schizophrenia: The potential role of rapid facial mimicry. *Journal of the International Neuropsychological Society*, *16*(4), 621-629. doi: 10.1017/S1355617710000329
- Wallwork, R. S., Fortgang, R., Hashimoto, R., Weinberger, D. R., & Dickinson, D. (2012). Searching for a consensus five-factor model of the Positive and Negative Syndrome Scale for schizophrenia. *Schizophrenia Research*, *137*(1), 246-250. doi:10.1016/j.schres.2012.01.031
- Wechsler, D. (2003). *Wechsler Memory Scale III*. San Antonio: The Psychological Corporation.
- Wechsler, D. (2008). *Wechsler Adult Intelligence Scale—Fourth Edition (WAIS—IV)*. San Antonio, TX: NCS Pearson.
- Wechsler, D., & Hsiao-pin, C. (2011). *WASI-II: Wechsler Abbreviated Scale of Intelligence*. San Antonio, TX: NCS Pearson.
- Zaki, J., Ochsner, K. N., & Ochsner, K. (2012). The neuroscience of empathy: Progress, pitfalls and

promise. *Nature Neuroscience*, 15(5), 675. doi: 10.1038/nn.3085

Zhu, C., Lee, T., Li, X., Jin, S., Wang, Y., & Wang, K. (2007). Impairments of social cues recognition and social functioning in Chinese people with schizophrenia. *Psychiatry and Clinical Neurosciences*, 61, 149-158. doi. 10.1111/j.1440-1819.2007.01630.x

Journal Pre-proof

Table 1. Key Difference Between Cognitive Empathy and Affective ToM

	Understanding another's state	Emotionally engaged in another's perspective
Cognitive empathy	✓	✓
Affective theory of mind	✓	X

Table 2. Descriptive Statistics for Demographic and Clinical Characteristics of Individuals with Early and Chronic Schizophrenia

Variable	Early schizophrenia M (SD)	Chronic schizophrenia M (SD)
Male:female	13:13	22:11
Age, years	23.19 (2.84)	31.64 (8.97)
Education, years	12.19 (2.02)	11.30 (1.70)
Estimated IQ (WASI)	99.04 (12.30)	92.79 (11.88)
Duration of illness, years	0.62 (1.21)	10.98 (6.16)
Age of onset, years	22.27 (3.11)	20.91 (5.63)
Number of psychotic episodes	1.62 (1.33)	9.13 (8.56)
PANSS		
Positive symptoms factor	8.00 (3.48)	11.70 (4.48)
Negative symptoms factor	11.76 (4.54)	13.63 (5.29)
Disorganisation symptoms factor	6.16 (2.82)	8.77 (2.88)
Chlorpromazine equivalent (daily)	466.58 (373.49)	627.14 (399.61)

Table 3

*Partial Correlations between the Virtual Assessment of Mentalising Ability and Empathy Quotient Subscales for Individuals with Early and Chronic Schizophrenia*

	Cognitive empathy	Emotional reactivity	Social skills
<i>Early schizophrenia</i>			
First-order cognitive ToM	.39	-.01	.46*
First-order affective ToM	.46*	-.04	.14
Second-order cognitive ToM	.52**	-.01	-.10
Second-order affective ToM	.30	-.21	-.53**
Total ToM	.65***	-.01	.08
<i>Chronic schizophrenia</i>			
First-order cognitive ToM	.65***	.09	.11
First-order affective ToM	.45*	.05	-.12
Second-order cognitive ToM	.44*	.46*	.04
Second-order affective ToM	.48*	.25	.18
Total ToM	.65***	.15	.16

*Note.* ToM = Theory of mind. Control variable = Neurocognition.

\*  $p < .05$ .

\*\*  $p < .01$ .

\*\*\*  $p < .001$ .

Table 4

*Partial Correlations between Empathy Quotient Subscales and Theory of Mind Subscales with Indices of Social Functioning for Individuals with Early and Chronic Schizophrenia*

	SFS	SSPA
<i>Early schizophrenia</i>		
Cognitive empathy	.66***	.61***
Emotional reactivity	.09	.02
Social skills	.19	.01
First-order cognitive	.55**	.56**
First-order affective	.48**	.52**
Second-order cognitive	.47**	.57**
Second-order affective	.30	.50**
Total ToM	.52**	.66***
<i>Chronic schizophrenia</i>		
Cognitive empathy	.67***	.68***
Emotional reactivity	.27	.21
Social skills	.36	.36

First-order cognitive	.48**	.62***
First-order affective	.42*	.52**
Second-order cognitive	.57***	.47**
Second-order affective	.18	.48**
Total ToM	.56***	.68***

---

*Note.* Control variable = Neurocognition. SFS = Social Functioning Scale. SSPA = Social Skills Performance Assessment.

\*\*  $p < .01$ .

\*\*\*  $p < .001$ .



Table 5

*Hierarchical Regression Analyses for Incremental Prediction of Community Functioning in Individuals with Early and Chronic Schizophrenia*

Variable	Step 1			Step 2			Step 3			Step 4		
	<i>B</i>	<i>SE</i>	$\beta$	<i>B</i>	<i>SE</i>	$\beta$	<i>B</i>	<i>SE</i>	$\beta$	<i>B</i>	<i>SE</i>	$\beta$
	<i>B</i>			<i>B</i>								
<i>Early</i>												
<i>Schizophrenia</i>												
<i>a</i>												
	3.9	2.2	.36	.80	2.4	.07	2.9	2.17	.26	.69	.87	.06
Neurocognition	9	9			8		3					
Positive symptoms				.05	2.8	.07	2.9	2.33	.32	.25	.95	.03
Negative symptoms					5		0					
Disorganised symptoms				1.7	1.7	.23	.11	1.46	.01	.67	.58	.09
Total ToM <sup>a</sup>				8.5	4.6	.52	.81	3.61	.48*	.81	1.58	.05
				8	8							
							6.1	1.68	.63**	4.1	1.83	.39
							3		*	3		*



$R^2$	.28	.44	.63	.73
$\Delta R^2$	.28	.16	.19	.10
$\Delta F$	8.5	1.7	19.2	98.1
	7	7	7	6

---

*Note.* <sup>a</sup> First- and second-order cognitive and affective ToM as measured by the Virtual

Assessment of Mentalising Ability.

\*  $p < .05$

\*\*  $p < .01$ .

\*\*\*  $p < .001$ .

Table 6

*Hierarchical Regression Analyses for Incremental Prediction of Functional Capacity in Individuals with Early and Chronic Schizophrenia*

Variable	Step 1			Step 2			Step 3			Step 4		
	<i>B</i>	<i>SE</i>	$\beta$	<i>B</i>	<i>SE</i>	$\beta$	<i>B</i>	<i>SE</i>	$\beta$	<i>B</i>	<i>SE</i>	$\beta$
	<i>B</i>			<i>B</i>								
<i>Early</i>												
<i>Schizophrenia</i>												
<i>a</i>												
	1.58	.5	.53*	.96	.59	.3	.12	.41	.04	.2	.31	.07
Neurocognition		5	*			2				0		
Positive symptoms				1.08	.68	.45	.25	.44	.11	.63	.33	.27
Negative symptoms				.13	.42	.07	.40	.27	.20	.2	.20	.15
Disorganised symptoms				.29	1.1	.0	.13	.67	.03	.9	.56	.21
Total ToM <sup>a</sup>							1.23	1.77	.54**	.9	.30	.52**
							3		*	8		

Cognitive empathy										.9	.27	.48**
										6		
$R^2$	.2		.53		.68						.74	
	8											
$\Delta R^2$	.2		.25		.15						.059	
	8											
$\Delta F$	8.		3.1		31.7						15.9	
	29		5		0						6	

---

*Chronic Schizophrenia*

	1.5	.46	.47*	1.2	.45	.4	.37	.34	.14	.0	.10	.02
Neurocogniti on	8		*	8		8					6	
Positive symptoms				.48	.43	.2	.03	.29	.02	.0	.10	.02
						8					2	
Negative symptoms				.35	.35	.1	.11	.24	.06	.1	.07	.07
						9					3	
Disorganised symptoms				.15	.63	.0	.27	.42	.09	.0	.13	.00
						5					1	
Total ToM <sup>a</sup>							1.4	.27	.61**	.8	.13	.49**
							0		*	7		*

Cognitive empathy				.7 4	.12	.21**
$R^2$	.35	.52	.71		.79	
$\Delta R^2$	.35	.18	.18		.079	
$\Delta F$	11.6	2.3	26.1		18.6	
	5	2	1		2	

---

*Note.* <sup>a</sup> First- and second-order cognitive and affective ToM as measured by the Virtual Assessment of Mentalising Ability.

Table 7

*Mediating Effect of Cognitive Empathy on the Relationship between Affective Theory of Mind and Community Functioning for Individuals with Early and Chronic Schizophrenia*

<i>Variable</i>	<i>B</i>	<i>B SE</i>	$\beta$	<i>t</i>	<i>p</i>
<i>Early Schizophrenia</i>					
Intercept	58.76	25.49			
Affective ToM	1.97	1.56	.16	1.26	.23
Cognitive empathy	8.59	.85	.74	10.12	< .001
<i>Chronic Schizophrenia</i>					
Intercept	62.87				
Affective ToM	1.31	1.05	.11	1.24	.23
Cognitive empathy	8.28	.58	.78	14.36	< .001

*Note.* Affective ToM = Composite of first- and second-order VAMA subscales. Dependent variable = Total scores on the Social Functioning Scale. Control variables = neurocognition and clinical symptoms.


Table 8

*Mediating Effect of Cognitive Empathy on the Relationship between Affective Theory of Mind and Functional Capacity for Individuals with Early and Chronic Schizophrenia*


<i>Variable</i>	<i>B</i>	<i>B SE</i>	$\beta$	<i>t</i>	<i>p</i>
<i>Early Schizophrenia</i>					
Intercept	3.48	8.46			
Affective ToM	2.05	.52	.64	3.94	.001
Cognitive empathy	.78	.28	.35	2.78	.01
<i>Chronic Schizophrenia</i>					
Intercept	16.84	3.25			
Affective ToM	.02	.18	.01	.08	.94
Cognitive empathy	1.64	.10	.75	16.66	< .001

*Note.* Affective ToM = Composite of first- and second-order VAMA subscales. Dependent variable = Total scores on the Social Skills Performance Assessment. Control variables = neurocognition and clinical symptoms.






*Figure 1.* Mean scores on the Empathy Quotient subscales provided by individuals with early and chronic schizophrenia. *Error bars represent  $\pm 1$  SE.*



*Figure 2.* Path model depicting influence of cognitive empathy on the relationship between affective theory of mind and community functioning for individuals with early schizophrenia. Path coefficients represent standardised beta-weights. The coefficient below the path from affective ToM to social functioning represents the direct effect without the mediator in the model, and the coefficient above the path represents the direct effect when the mediator is included in the model.

\*\*  $p < .01$ .


\*\*\*  $p < .001$ .



*Figure 3.* Path model depicting influence of cognitive empathy on the relationship between affective theory of mind and community functioning for individuals with chronic schizophrenia. Path coefficients represent standardised beta-weights. The coefficient below the path from affective ToM to social functioning represents the direct effect without the mediator in the model, and the coefficient above the path represents the direct effect when the mediator is included in the model.

\*\*  $p < .01$ .


\*\*\*  $p < .001$ .



*Figure 4.* Path model depicting influence of cognitive empathy on the relationship between affective theory of mind and functional capacity for individuals with early schizophrenia. Path coefficients represent standardised beta-weights. The coefficient below the path from affective ToM to social functioning represents the direct effect without the mediator in the model, and the coefficient above the path represents the direct effect when the mediator is included in the model.

\*\*  $p < .01$ .

\*\*\*  $p < .001$ .



*Figure 5.* Path model depicting influence of cognitive empathy on the relationship between affective theory of mind and functional capacity for individuals with chronic schizophrenia. Path coefficients represent standardised beta-weights. The coefficient below the path from affective ToM to social functioning represents the direct effect without the mediator in the model, and the coefficient above the path represents the direct effect when the mediator is included in the model.

\*\*  $p < .01$ .

\*\*\*  $p < .001$ .