

Resistance training in addition to aerobic activity is associated with lower likelihood of depression and comorbid depression and anxiety symptoms: A cross sectional analysis of Australian women

Author

Oftedal, Stina, Smith, Jordan, Vandelanotte, Corneel, Burton, Nicola W, Duncan, Mitch J

Published

2019

Journal Title

Preventive Medicine

Version

Accepted Manuscript (AM)

DOI

[10.1016/j.ypmed.2019.105773](https://doi.org/10.1016/j.ypmed.2019.105773)

Rights statement

© 2019 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International Licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, providing that the work is properly cited.

Downloaded from

<http://hdl.handle.net/10072/386694>

Griffith Research Online

<https://research-repository.griffith.edu.au>

Accepted Manuscript

Resistance training in addition to aerobic activity is associated with lower likelihood of depression and comorbid depression and anxiety symptoms: A cross sectional analysis of Australian women

Stina Oftedal, Jordan Smith, Corneel Vandelanotte, Nicola W. Burton, Mitch J. Duncan



PII: S0091-7435(19)30249-X
DOI: <https://doi.org/10.1016/j.ypmed.2019.105773>
Article Number: 105773
Reference: YPMED 105773
To appear in: *Preventive Medicine*
Received date: 6 March 2019
Revised date: 11 June 2019
Accepted date: 14 July 2019

Please cite this article as: S. Oftedal, J. Smith, C. Vandelanotte, et al., Resistance training in addition to aerobic activity is associated with lower likelihood of depression and comorbid depression and anxiety symptoms: A cross sectional analysis of Australian women, *Preventive Medicine*, <https://doi.org/10.1016/j.ypmed.2019.105773>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Resistance training in addition to aerobic activity is associated with lower likelihood of depression and comorbid depression and anxiety symptoms: A cross sectional analysis of Australian women

Authors: Stina Oftedal^{1,2}, Jordan Smith^{2,3}, Corneel Vandelanotte⁴, Nicola W Burton⁵, Mitch J Duncan^{1,2}

¹ School of Medicine & Public Health; Faculty of Health and Medicine, The University of Newcastle, University Drive, Callaghan NSW 2308, Australia

² Priority Research Centre for Physical Activity and Nutrition, The University of Newcastle, University Drive, Callaghan NSW 2308, Australia

³ School of Education, Faculty of Education and Arts, University of Newcastle, Callaghan, New South Wales, Australia

⁴ Physical Activity Research Group, Appleton Institute, School of Health, Medical and Applied Sciences, Central Queensland University, Rockhampton Queensland 4702, Australia

⁵ School of Applied Psychology, Griffith University, Brisbane 4122, Queensland, Australia

Corresponding author: Stina Oftedal, Advanced Technology Centre. Level 3. The University of Newcastle, University Drive, Callaghan NSW 2308, Australia, e-mail: stina.oftedal@newcastle.edu.au.

Key words: muscle-strengthening activity, physical activity, common mental health disorders

Word count:

- Abstract: 250 words
- Manuscript: 3500 words

Abstract

The mental health benefits of resistance training (RT) alone or beyond those provided by aerobic physical activity (PA) are unclear. This study aimed to determine the association between meeting recommendations for aerobic PA and/or RT, and symptoms of depression and/or anxiety. Participants were Australian female members of the 10,000 Steps project (n=5180, 50.0±11.5 years). Symptoms of depression and anxiety were determined using the Depression Anxiety Stress Score. Participants were grouped as ‘depression only’, ‘anxiety only’, ‘co-occurring depression and anxiety’ or ‘neither depression nor anxiety’ based on relevant subscale score (cut-points: depression≥14 points, anxiety≥10 points). The International Physical Activity Questionnaire-Long Form questionnaire was used to determine PA with an additional item to specify RT frequency. Participants were classified as adhering to ‘aerobic PA only’ (≥150 min PA/week), ‘RT only’ (RT≥2 days/week), ‘aerobic PA+RT’ (≥150 min PA/week+RT≥2 days/week), or ‘neither aerobic PA nor RT’ (<150 min PA/week+RT<2 days/week). Adjusted relative risk ratios (RRR [95%CI]) were estimated using multinomial logistic regression models. Relative to the ‘neither PA nor RT’ (n=2215), the probabilities of ‘depression only’ (n=317) and ‘co-occurring depression and anxiety’ (n=417) were lower for the ‘aerobic PA only’(n=1590) (RRR=0.74 [0.56 – 0.97] and RRR=0.76 [0.59-0.97] respectively), and ‘both PA+RT’ (n=974) groups (RRR=0.61 [0.43-0.86] and RRR=0.47 [0.33-0.67] respectively). There were no associations between adhering to one or both recommendations and ‘anxiety only’ (n=317), or between ‘RT only’ (n=401) and depression and/or anxiety. Prevention and treatment strategies including both aerobic PA and RT may provide additional benefits for depression with or without comorbid anxiety.

Keywords: common mental disorders, physical activity, muscle strengthening activity, anxiety, depression, aerobic activity

Introduction

Globally, approximately 322 million (4.4%) people had a depressive disorder in 2015, and 264 million (3.6%) had an anxiety disorder [1]. Depression is the leading global cause of disability, while anxiety is ranked sixth [1]. Approximately half of those diagnosed with one of these conditions have a comorbidity with the other [1, 2]. An estimated 12-32% of the population live with symptoms of varying severity, and women are about twice as likely to experience depression or anxiety in their lifetime as men [1, 3, 4]. Even mild symptoms of depression and anxiety can have debilitating effects on wellbeing, relationships, career and productivity [5-9].

The World Health Organization (WHO) recommends 18-64 year-olds participate in muscle-strengthening activities (i.e. resistance training: RT) on 2 or more days and either ≥ 150 minutes of moderate-intensity physical activity (PA), ≥ 75 minutes of vigorous-intensity PA, or an equivalent combination per week [10]. Yet relative to the proportion of adults who report meeting aerobic PA guidelines far fewer adults report meeting the RT recommendation. In the US, 23.0% of adults met the RT recommendation, 46.1% met the aerobic PA recommendation and 19.4% met both [12]. Similar patterns have been found in Australia [13], the UK [14] and Scotland [11]. The prevalence of insufficient PA is concerning given the contribution of physical inactivity to the global burden of disease [15], and the even higher rates of insufficient RT are particularly concerning given RT may confer unique health benefits [16-22].

Physical activity can have a role in the management of depression and anxiety and is protective against cardiovascular disease and metabolic disorders, which are common comorbidities [23, 24]. While RT can have benefits for both depression and anxiety [25, 26], the majority of prior work has focused on aerobic PA or aerobic PA and RT combined without exploring the differential associations between the modes [27-30]. The few

intervention studies comparing 'RT only' with 'aerobic PA only' have indicated equal effectiveness of the two modes in terms of reducing symptoms of depression or anxiety [25, 26]. A recent observational study found the prevalence of depressive symptoms was lower in those who met both versus one of the recommendations for aerobic PA and RT [31]. A combination of aerobic PA and RT may confer unique benefits over and above that of aerobic PA or RT alone, but few studies have explored this. Additionally, few studies have explored the effect of RT and aerobic PA on co-occurring symptoms of depression and anxiety [32]. This is a major gap, as co-morbidity is common and is associated with unique symptom and neurobiological characteristics, and more severe and protracted symptoms than those with either disorder alone [2]. Consequently, the relationship between PA and co-occurring symptoms of depression and anxiety may differ from that of depression or anxiety alone [2]. The aim of this study was to determine the association between meeting recommendations for aerobic PA or RT alone and combined, and symptoms of depression and anxiety, alone or co-occurring.

Methods

Study design and participants

This was a cross-sectional sample from the member database of the 10,000 Steps project (www.10000steps.org.au), a web-based PA promotion initiative which commenced in Rockhampton, Australia, in 2001 [33]. The Central Queensland University Human Research Ethics Committee provided approval for the current study (H15/09-210). All participants provided informed consent to participate via the online portal. As of November-December 2016, the project had approximately 330,000 members with 42,090 email addresses verified as valid at the time of the email-invitation to participate in the current study [34]. The overall response rate based completed surveys (n=6982) and the total number of verified email addresses was 16.5%. The majority of those who completed the survey were women

(n=5180, 74.2%). Given the higher response rate from women, and the higher prevalence of depression and anxiety in women [1], this study focuses on women only. Data analysis took place in November 2018.

Study measures

Depression and anxiety symptoms: Symptoms of depression and anxiety were assessed using relevant DASS-21 (Depression Anxiety Stress Score) questionnaire [35]. The depression and anxiety sub-scales have shown criterion validity (depression: Area Under the Curve [AUC]=0.77-0.91, anxiety: AUC=0.60-0.83) in comparison to DSM-IV diagnosis, and adequate construct validity [36]. Each domain comprises seven statements assessing symptom frequency last week (e.g. 'I felt I was close to panic', 'I felt I wasn't worth much as a person') and are rated on a 4-point scale (0=never, 1=sometimes, 2=often, 3=almost always). The score for each subscale is summed and multiplied by 2, making the maximum score for each domain 42. Scores were then categorized according to symptom severity (normal-extremely severe) [35]. These categories are not diagnostic but describe the continuum of severity observed in the population, i.e. 'mild' indicates symptoms above population average, not 'mild disorder'[35]. Participants were categorized into four mutually exclusive groups: 'depression only' or 'anxiety only' if responses indicated at least 'moderate' severity (depression \geq 14, anxiety \geq 10), and less than 'moderate' severity on the other subscale. The 'co-occurring depression and anxiety' group scored at least 'moderate' severity on both subscales. Those categorized as 'neither depression nor anxiety' scored below 'moderate' severity on both subscales.

Aerobic physical activity and resistance training Aerobic PA during the previous week was measured using the International Physical Activity Questionnaire Long Form (IPAQ-LF). It assesses frequency and duration of participation in each of walking, moderate, and vigorous intensity PA as part of work, transportation, housework, gardening, sport or leisure during the

last week [37]. Total MET-minutes were calculated according to standard IPAQ scoring protocols, using MET values of 3.3 for walking, 4.0 for moderate intensity activity and 8.0 for vigorous intensity PA [37]. Standard scoring procedures were used to subsequently classify participants' PA as 'low', 'moderate' or 'high' [37]. The IPAQ-LF measures "total activity" across multiple domains resulting in higher reported activity relative to only assessing leisure time activity [38]. Consequently the 'high' category (≥ 3 sessions of vigorous activity per week and a total of ≥ 1500 MET-minutes per week OR ≥ 7 sessions of any activity intensity and a total of ≥ 3000 MET-minutes per week) is considered to represent a total activity level comparable to that of meeting the PA recommendation of ≥ 150 minutes of aerobic PA per week [38, 39]. Participants were asked an additional item "How many days in the last 7 days have you participated in resistance/weight training" [40] and a frequency of ≥ 2 days per week was considered meeting the WHO recommendations [10]. Study participants were then categorized into the following PA groups: meeting 'aerobic PA only' (≥ 150 minutes aerobic PA/week), 'RT only' (RT ≥ 2 days/week), 'aerobic PA+RT' (≥ 150 minutes aerobic PA/week+RT ≥ 2 days/week), or 'neither aerobic PA nor RT' (< 150 minutes aerobic PA/week+RT < 2 days/week).

Socio-demographic and behavioral measures Household income was reported in seven categories and collapsed into categories of: '>AUD\$70,000 per annum'; ' \leq AUD\$70,000 per annum'; or 'unsure/ prefer not to state'. Employment status was categorized as 'employed' or 'not employed'. Cigarette-smoking status was assessed using a single item and categorized as 'current smoker' or 'non-smoker' [41, 42]. Number of alcoholic drinks typically consumed was reported, and categorized as: 'non-drinkers' (no alcohol in the last year); 'low risk' (1-2 standard drinks on a typical occasion); or 'high risk' (≥ 3 standard drinks on a typical occasion), in line with the Australian guidelines [43]. The Workforce Sitting Questionnaire [44] asked participants how long they spent sitting for each of transport, at work, watching

TV, using a computer or other devices and leisure on each of work and non-work days, and number of work days per week. Average daily sitting-time, calculated by summing sitting-time for work and non-work days and averaging over 7 days, was categorized as low (<8hrs/day) or high (≥ 8 hours/ day) [45]. Sleep dimensions were measured using three items (sleep duration, quality and latency) from the Pittsburgh Sleep Quality Index (PSQI) [46]. The PSQI has strong reliability and validity, and moderate structural validity in a variety of samples [47]. Sleep duration was classified as 'meeting recommendations' (7 to ≤ 9 hours if 18 to <65 years old, 7 to ≤ 8 hours if ≥ 65 years old), or 'not meeting recommendations' (<7 hours or >9 hours if 18 to <65 years old, >8 hours if ≥ 65 years old). Sleep quality was collapsed into categories of 'very good'; 'fairly good'; or 'fairly-to-very bad' [45]. Sleep latency was classified as normal (≤ 30 minutes) or long (>30 minutes), indicating a sleep disorder [48].

Participants self-reported diagnosis of 17 chronic diseases: high blood pressure, diabetes type 1 or 2, coronary heart disease, chronic obstructive pulmonary disease, cerebrovascular disease, cancer, asthma, high cholesterol, arthritis, insomnia, restless legs syndrome, sleep apnea, irritable bowel syndrome, kidney disease, arthritis, and mental illness other than depression or anxiety. Participants were categorized as ' ≥ 1 chronic conditions' or 'None'. Participants reported if they had ever received a diagnosis of depression and/or anxiety and were dichotomized as 'prior diagnosis of depression and/or anxiety' or 'none'. Self-reported height (cm) and weight (kg) were used to calculate body mass index (BMI) and subsequently categorized as >18.5, 18.5-24.9, 25.0-29.9 and ≥ 30.0 kg/m².

Statistical analysis

Descriptive statistics included means (SD) for continuous variables and count (%) for categorical variables, Cronbach's alpha calculated for DASS subscales. T-tests and Chi-square tests were used to assess differences in descriptive variables between the sample and

those excluded from analysis due to incomplete data. One-way ANOVA analysis was used to compare MET-minutes reported for each activity intensity, activity domain and RT days between the four activity groups. Three multinomial logistic regression models were used, and Model A represents the unadjusted association between PA groups and mental health symptom groups while Model B presents the association between PA groups and mental health symptom groups, adjusted for socio-demographic variables, sleep health measures, and sitting-time. All covariates were chosen *a priori* based on previous literature [49-53]. Model C included the same variables as Model B, plus chronic disease diagnosis, BMI category and diagnosis of depression and/or anxiety as covariates. Chronic disease and BMI can be conceptualized as being on the causal path between PA and mental health symptoms, and previous diagnosis of depression and/or anxiety is likely associated with current symptoms of depression and/or anxiety [54-57]. Controlling for these variables is therefore potentially over-adjusting the model, as it cancels out the effect of PA on mental health symptoms mediated through the influence of PA on chronic disease, BMI or previous depression and anxiety [58]. However, Model C was included as a sensitivity analysis. A separate sensitivity analysis using an alternative threshold (≥ 600 MET minutes) for meeting aerobic PA recommendations was also conducted. All analyses were conducted using Stata Version 14 (College Station, TX: StataCorp) in November 2018.

Results

The 5180 women included in the study were primarily middle-aged (50.0 ± 11.5 years) and employed (85.5%) and had at least one chronic disease diagnosis (62.9%). These are higher than average age (38 years), employment rate (56%) and chronic disease diagnosis (52%) in the Australian female population [59, 60]. The mean DASS-21 depression score was 6.0 (7.3) points and mean DASS-21 score was 4.2 (5.3) points, with 79.9% ($n=4129$) classified as 'neither depression nor anxiety', 6.1% ($n=317$) classified as 'depression only', 6.1% (317)

classified as ‘anxiety only’ and 8.1% (n=417) classified as ‘both depression and anxiety’.

Participant characteristics by mental health symptoms are described in **Table 1**. Women who were excluded because of incomplete data (n=3086, 37.3%) were younger, less likely to disclose their income or report an income of >AUD\$70,000 per annum, and more likely to be smokers and have a high-risk alcohol consumption or be non-drinkers than those who were included (**Supplemental Table 1**). They were also more likely to have a chronic disease diagnosis, a BMI \geq 30 kg/m² and meet ‘aerobic PA only’, and less likely to report a diagnosis depression and/or anxiety and meet ‘neither aerobic PA nor RT’. There were no differences by employment status; sitting-time; or sleep quality, duration or latency.

Aerobic physical activity and resistance training

Of the study sample (n=5180), 42.8% (n=4129) met ‘neither aerobic PA nor RT’, 30.7% (n=1590) met the ‘aerobic PA only’, 7.7% (n=401) met the ‘RT only’, and 18.8% (n=974) met ‘aerobic PA+RT recommendation’. **Figure 1** shows the distribution of PA groups by mental health symptom groups. A comparison of total MET-minutes and the specific intensities and activity domains that characterized PA groups are described in **Table 2**. Notably, the ‘aerobic PA+RT recommendation’ group reported higher total MET-minutes, higher leisure domain MET-minutes for all intensity levels, and higher total vigorous MET-minutes than the ‘aerobic PA only’ group. The ‘aerobic PA only’ group had higher total moderate MET-minutes than the ‘aerobic PA+RT recommendation’ group from higher work, transport, house and yard chores MET-minutes. The ‘aerobic PA+RT’ group had more days of RT than the ‘RT only’ group.




Figure 1: Percentage meeting physical activity recommendations for women ($n=5180$) who responded to the Australian 10,000 Steps Project 2016 survey by symptoms of neither depression nor anxiety ($n=4129$), anxiety only ($n=317$), depression only ($n=317$) and both depression and anxiety ($n=417$).

Aerobic physical activity, resistance training and mental health symptoms

The probability of depression only and co-occurring depression and anxiety was lower for those meeting ‘aerobic PA only’ (ARRR= 0.74, 95%CI: 0.56-0.97 and 0.76, 95%CI: 0.59-0.97 respectively) and ‘aerobic PA+RT’ (ARRR= 0.61, 95%CI 0.43-0.86 and 0.47, 95%CI: 0.33-0.67 respectively), relative to the ‘neither aerobic PA nor RT’ group (**Table 3**, Model B). The magnitude of association was greater for ‘aerobic PA+RT’ than for ‘aerobic PA only’. Adjusting for the presence of chronic disease, BMI and prior diagnosis of depression and anxiety attenuated the association between ‘aerobic PA only’ and depression (ARRR= 1.00, 95%CI 0.75-1.33) and co-occurring depression and anxiety (ARRR=0.87, 95%CI 0.67-1.13) (Table 3, Model C or full results including covariates in **Supplemental Table 2**).

Meeting ‘RT only’ was not associated with a change in the relative risk of depression and/or anxiety, and there were no associations between meeting activity recommendations and anxiety. In analysis with ≥ 600 MET minutes as the cut-off for meeting the aerobic PA

recommendation, the association between ‘aerobic PA only’ and depression was attenuated, while all other associations remained (**Supplemental Table 3**).

Discussion

In this study of adult women, we found an association between meeting ‘aerobic PA only’ or ‘aerobic PA+RT’ and depression and co-occurring depression and anxiety. There was evidence that meeting both recommendations had a stronger association with lowered risk of symptoms of depression than only meeting the aerobic PA recommendation. This suggests a synergistic relationship between aerobic PA and RT in terms of mental health benefits, and is consistent with a recent study [31]. It also aligns with research examining the relationship between aerobic PA, RT and physical health outcomes [16-22].

The stronger association observed for those meeting ‘aerobic PA+RT’ than for those meeting the ‘aerobic PA only’ may also simply reflect that more overall PA is better, which may also explain why the sensitivity analysis showed that ‘aerobic PA only’ was no longer associated with lower relative risk of depression when using a lower cut-off for meeting aerobic PA recommendations. Those meeting ‘aerobic PA+RT’ recommendations reported higher total MET-minutes than the ‘aerobic PA only’ group (6791 [4830] vs. 6331 [4075] MET-minutes respectively) in addition to meeting the RT recommendation (Table 2). However, other studies have not observed a dose-response relationship between aerobic PA and RT participation and depressive symptoms [31, 61]. Several other factors may also contribute to our finding. Those meeting ‘aerobic PA+RT’ also participated in substantially more vigorous activity than those meeting the ‘aerobic PA only’ (2292 [2165] vs. 1344 [1889] MET-minutes respectively). Vigorous PA may be more beneficial to mental health than low or moderate intensity PA through different physiological effects, but further research is needed [62, 63]. Previous studies have also shown that differences in mental health by activity domain. One study indicated that while leisure activity and transport activity had a positive association

with better mental health, work-related PA was associated with worse mental health [64]. Another study demonstrated that summing leisure and transport activity with house and garden activity attenuated the positive association between leisure and transport activity and mental well-being [65]. In the current study, those meeting ‘aerobic PA+RT’ also participated in more leisure activities, whereas those meeting the ‘aerobic PA only’ reported more work, household and garden activity (Table 2). Activity domain may therefore influence associations with depression and anxiety and warrants further investigation.

Both aerobic PA and RT have been shown to independently benefit mental health via anti-depressive and anxiolytic effects [27-31], however an association between ‘RT only’ and depression or anxiety was not identified in the current study. Interestingly, for women meeting ‘RT only’ has been shown to be associated with lower prevalence of depressive symptoms in those with the most severe symptoms of depression, not in those with mild or moderate symptoms [31]. The lack of an association between ‘RT only’ and depression in the current study could therefore be from combining moderate and severe symptoms. However, the proportion of the sample who were categorized as ‘RT only’ was only 7.7% (n=401), overall, limiting the statistical power available, as evidenced by wider confidence intervals around the point estimates for the ‘RT only’ group compared to the other groups (Table 3). Furthermore, the specific mechanisms that produce anti-depressive and anxiolytic effects are poorly understood, especially for RT, but may include both psychological, neurobiological and behavioral mechanisms [63]. The volume (load, length and intensity of sessions, duration of intervention) of RT necessary to elicit these effects is unclear [25, 26, 66].

We did not find an association between meeting one or both PA recommendations and relative risk of anxiety symptoms (Table 2). This is in contrast to other studies that have found that while the positive effects of PA are smaller for symptoms of anxiety than depression, they still exist [26, 27]. A possible explanation for this lack of association in the

current study may be differences in severity of anxiety symptoms (Table 1). On average, the anxiety only group had ‘moderate’ symptoms of anxiety (score: 10-14) whereas the co-occurring depression and anxiety group had ‘severe’ anxiety symptoms (score: 15-19). Evidence suggests the anti-depressive effect of aerobic PA and RT is greater in more severely depressed populations [25, 27], but whether or not there is a floor effect for the anxiolytic benefit has not been explored to the same extent. While a favorable effect of aerobic PA and RT on mental health has been shown in both non-clinical populations [26, 67] and participants with generalized anxiety disorder [32], further research is needed to explore the effect in populations with subclinical symptoms [26] and those with co-occurring depression and anxiety [2].

Strengths and limitations

The study included a relatively large sample of women where the proportion of participants meeting neither or both guidelines was similar to that reported in large Australian [13] and US [31] samples, and the mean DASS depression and anxiety subscale scores is similar to normative values [68]. Study strengths also include a focus anxiety and depression and their co-occurrence, and inclusion of a range of key covariates. The inclusion of sleep is a particular strength due to its strong association with mental health [69]. Adjusting for chronic disease, BMI and a prior diagnosis of depression and/or anxiety attenuated the association between meeting ‘aerobic PA only’ and depression and co-occurring depression and anxiety, though this could be due to over-adjusting. However adjustments did not appreciably alter the association between meeting ‘aerobic PA+RT’ and depression or co-occurring depression and anxiety.

The study included primarily middle-aged Australian women with an income above AUD\$70,000, and findings may not be representative of the wider female population, or generalizable to men. Furthermore, the IPAQ does not differentiate between aerobic PA and

RT, and some participants may have included RT activities in the IPAQ reporting and also when reporting RT. This may have underestimated the effect of aerobic activity. Participants were also part of a community-wide physical activity promotion program [33]. This may confound the prevalence of individuals meeting the aerobic PA guidelines, though the prevalence rate was similar to that previously reported in an Australian population [13] (52.6% vs 49.5% in the current study).

The referent group labelled 'neither depression nor anxiety' included participants with mild symptoms of depression and anxiety, and those with moderate and severe symptoms were combined as 'symptoms of depression and/or anxiety'. This grouping could potentially underestimate the association between mental health symptoms and PA [27, 31]. The DASS-21 assesses symptoms experienced in the last week, so they may be transient in nature as opposed to chronic, and not necessarily be associated with activity participation. Exploring associations between meeting PA recommendations and diagnosed depression and anxiety is an important next step.

Conclusion

Meeting recommendations for aerobic PA, or both aerobic PA and RT, was associated with a reduced risk of reporting symptoms of depression and co-occurring symptoms of depression and anxiety in women. The magnitude of association was stronger for those meeting both recommendations, indicating the potential added benefit of engaging in both types of PA. However, it remains unclear whether there is something unique about RT in particular, or whether additional PA of any kind would produce similar benefits. Prospective analyses of these cross-sectional relationships are needed. Clarifying the dose-response relationship between activity types, symptoms of depression and anxiety and their co-occurrence will also improve understanding of the application of PA to prevent and manage symptoms of depression and anxiety.

Word count:

Abstract: 250 words

Manuscript: 3500 words

Acknowledgements**Funding sources**

This work is in part supported by an NSW CVRN Collaborative Project Grant (101584).

Corneel Vandelanotte reports grants from National Heart Foundation of Australia (ID 100427) and National Health and Medical Research Council (APP1080186). Mitch J Duncan reports grants from National Heart Foundation of Australia (ID 100029), Diabetes Australia (Y17G-DUNM) and National Health and Medical Research Council (G1700197). The sponsors played no role in the collection, analysis or interpretation of the data; the writing of the report or the decision to submit the article for publication.

Conflicts of interest

The authors do not have any conflicts of interest to declare.

References

1. World Health Organisation, Depression and Other Common Mental Disorders: Global Health Estimates. . 2017. World Health Organisation: Geneva, Switzerland.
2. Kircanski, K., J. LeMoult, S. Ordaz, and I.H. Gotlib, Investigating the nature of co-occurring depression and anxiety: Comparing diagnostic and dimensional research approaches. *J Affect Disord.* 2017; 216:123-135. DOI: 10.1016/j.jad.2016.08.006
3. McLean, C.P., A. Asnaani, B.T. Litz, and S.G. Hofmann, Gender Differences in Anxiety Disorders: Prevalence, Course of Illness, Comorbidity and Burden of Illness. *J Psyc Res.*, 2011; 45(8):1027-1035. DOI: 10.1016/j.jpsychires.2011.03.006
4. Albert, P.R., Why is depression more prevalent in women? *J Psych Neurosc.* 2015; 40(4):219-221. DOI: 10.1503/jpn.150205
5. Haller, H., H. Cramer, R. Lauche, F. Gass, and G.J. Dobos, The prevalence and burden of subthreshold generalized anxiety disorder: a systematic review. *BMC Psychiatry.* 2014; 14:128-128. DOI: 10.1186/1471-244X-14-128
6. Shim, R.S., P. Baltrus, J. Ye, and G. Rust, Prevalence, treatment, and control of depressive symptoms in the United States: results from the National Health and Nutrition Examination Survey (NHANES), 2005-2008. *J Am Board Fam Med.* 2011; 24(1):33-8. DOI: 10.3122/jabfm.2011.01.100121
7. Karsten, J., C.A. Hartman, J.H. Smit, et al., Psychiatric history and subthreshold symptoms as predictors of the occurrence of depressive or anxiety disorder within 2 years. *Br J Psychiatry.* 2011; 198(3):206-12. DOI: 10.1192/bjp.bp.110.080572
8. Wittayanukorn, S., J. Qian, and R.A. Hansen, Prevalence of depressive symptoms and predictors of treatment among U.S. adults from 2005 to 2010. *Gen Hosp Psychiatry.* 2014; 36(3):330-336. DOI: <https://doi.org/10.1016/j.genhosppsy.2013.12.009>

9. Cuijpers, P., N. Vogelzangs, J. Twisk, A. Kleiboer, J. Li, and B.W. Penninx, Differential mortality rates in major and subthreshold depression: meta-analysis of studies that measured both. *Br J Psychiatry*. 2013; 202(1):22-7. DOI: 10.1192/bjp.bp.112.112169
10. World Health Organisation, *Global Recommendations on Physical Activity for Health*. 2010. World Health Organisation Geneva, Switzerland.
11. Strain, T., C. Fitzsimons, P. Kelly, and N. Mutrie, The forgotten guidelines: cross-sectional analysis of participation in muscle strengthening and balance & co-ordination activities by adults and older adults in Scotland. *BMC Public Health*. 2016; 16(1):1108. DOI: 10.1186/s12889-016-3774-6
12. Schoenborn, C.A., P.F. Adams, and J.A. Peregoy, Health behaviors of adults: United States, 2008-2010. *Vital Health Stat* 10. 2013; (257):1-184. DOI:
13. Bennie, J.A., Z. Pedisic, J.G.Z. van Uffelen, et al., The descriptive epidemiology of total physical activity, muscle-strengthening exercises and sedentary behaviour among Australian adults – results from the National Nutrition and Physical Activity Survey. *BMC Public Health*. 2016; 16(1):73. DOI: 10.1186/s12889-016-2736-3
14. UK Health and Social Care Information Centre, *Health Survey for England 2011: Summary of key findings*, UK Health and Social Care Information Centre, Editor. 2011. Leeds: UK.
15. Lee, I.M., E.J. Shiroma, F. Lobelo, P. Puska, S.N. Blair, and P.T. Katzmarzyk, Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012; 380(9838):219-29. DOI: 10.1016/s0140-6736(12)61031-9

16. Grontved, A., A. Pan, R.A. Mekary, et al., Muscle-strengthening and conditioning activities and risk of type 2 diabetes: a prospective study in two cohorts of US women. *PLoS Med.* 2014; 11(1):e1001587. DOI: 10.1371/journal.pmed.1001587
17. Grontved, A., E.B. Rimm, W.C. Willett, L.B. Andersen, and F.B. Hu, A prospective study of weight training and risk of type 2 diabetes mellitus in men. *Arch Intern Med.* 2012; 172(17):1306-12. DOI: 10.1001/archinternmed.2012.3138
18. Hardee, J.P., R.R. Porter, X. Sui, et al., The Role of Resistance Exercise on All-cause Mortality in Cancer Survivors. *Mayo Clin Proc.* 2014; 89(8):1108-1115. DOI: 10.1016/j.mayocp.2014.03.018
19. Stamatakis, E., I.M. Lee, J. Bennie, et al., Does Strength-Promoting Exercise Confer Unique Health Benefits? A Pooled Analysis of Data on 11 Population Cohorts With All-Cause, Cancer, and Cardiovascular Mortality Endpoints. *Am J Epidemiol.* 2018; 187(5):1102-1112. DOI: 10.1093/aje/kwx345
20. Bakker, E.A., D.C. Lee, X. Sui, et al., Association of Resistance Exercise, Independent of and Combined With Aerobic Exercise, With the Incidence of Metabolic Syndrome. *Mayo Clin Proc.* 2017; 92(8):1214-1222. DOI: 10.1016/j.mayocp.2017.02.018
21. Bakker, E.A., D.C. Lee, X. Sui, et al., Association of Resistance Exercise With the Incidence of Hypercholesterolemia in Men. *Mayo Clin Proc.* 2018; 93(4):419-428. DOI: 10.1016/j.mayocp.2017.11.024
22. Colcombe, S. and A.F. Kramer, Fitness Effects on the Cognitive Function of Older Adults: A Meta-Analytic Study. *Psychol Sci.* 2003; 14(2):125-130. DOI: 10.1111/1467-9280.t01-1-01430

23. Celano, C.M., D.J. Daunis, H.N. Lokko, K.A. Campbell, and J.C. Huffman, Anxiety Disorders and Cardiovascular Disease. *Current psychiatry reports*. 2016; 18(11):101-101. DOI: 10.1007/s11920-016-0739-5
24. Dhar, A.K. and D.A. Barton, Depression and the Link with Cardiovascular Disease. *Frontiers in Psychiatry*. 2016; 7:33. DOI: 10.3389/fpsyt.2016.00033
25. Gordon, B.R., C.P. McDowell, M. Hallgren, J.D. Meyer, M. Lyons, and M.P. Herring, Association of Efficacy of Resistance Exercise Training With Depressive Symptoms: Meta-analysis and Meta-regression Analysis of Randomized Clinical Trials. *JAMA Psychiatry*. 2018; 75(6):566-576. DOI: 10.1001/jamapsychiatry.2018.0572
26. Gordon, B.R., C.P. McDowell, M. Lyons, and M.P. Herring, The Effects of Resistance Exercise Training on Anxiety: A Meta-Analysis and Meta-Regression Analysis of Randomized Controlled Trials. *Sports Med*. 2017; 47(12):2521-2532. DOI: 10.1007/s40279-017-0769-0
27. Rebar, A.L., R. Stanton, D. Geard, C. Short, M.J. Duncan, and C. Vandelanotte, A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychol Rev*. 2015; 9(3):366-78. DOI: 10.1080/17437199.2015.1022901
28. Schuch, F.B., D. Vancampfort, J. Firth, et al., Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. *Am J Psychiatry*. 2018; 175(7):631-648. DOI: 10.1176/appi.ajp.2018.17111194
29. Mammen, G. and G. Faulkner, Physical Activity and the Prevention of Depression: A Systematic Review of Prospective Studies. *Am J of Prev Med*. 2013; 45(5):649-657. DOI: <https://doi.org/10.1016/j.amepre.2013.08.001>

30. Stubbs, B., D. Vancampfort, S. Rosenbaum, et al., An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. *Psychiatry Res.* 2017; 249:102-108. DOI: 10.1016/j.psychres.2016.12.020
31. Bennie, J.A., M.J. Teychenne, K. De Cocker, and S.J.H. Biddle, Associations between aerobic and muscle-strengthening exercise with depressive symptom severity among 17,839 U.S. adults. *Preventive Medicine.* 2019. DOI: <https://doi.org/10.1016/j.ypmed.2019.02.022>
32. Herring, M.P., M.L. Jacob, C. Suveg, and P.J. O'Connor, Effects of short-term exercise training on signs and symptoms of generalized anxiety disorder. *Ment Health Phys Act.* 2011; 4(2):71-77. DOI: <https://doi.org/10.1016/j.mhpa.2011.07.002>
33. Brown, W.J., K. Mummery, E. Eakin, and G. Schofield, 10,000 Steps Rockhampton: Evaluation of a Whole Community Approach to Improving Population Levels of Physical Activity. *J Phys Act Health.* 2006; 3(1):1-14. DOI: 10.1123/jpah.3.1.1
34. Haapasalo, V., H. de Vries, C. Vandelanotte, R.R. Rosenkranz, and M.J. Duncan, Cross-sectional associations between multiple lifestyle behaviours and excellent well-being in Australian adults. *Prev Med.* 2018; 116:119-125. DOI: 10.1016/j.ypmed.2018.09.003
35. Parkitny, L. and J. McAuley, The Depression Anxiety Stress Scale (DASS). *J Physiother.* 2010; 56(3):204. DOI: [https://doi.org/10.1016/S1836-9553\(10\)70030-8](https://doi.org/10.1016/S1836-9553(10)70030-8)
36. Lee, J., E.-H. Lee, and S.H. Moon, Systematic review of the measurement properties of the Depression Anxiety Stress Scales-21 by applying updated COSMIN methodology. *Quality of Life Research.* 2019. DOI: 10.1007/s11136-019-02177-x
37. Bauman, A., F. Bull, T. Chey, et al., The International Prevalence Study on Physical Activity: results from 20 countries. *Int J Behav Nutr Phys Act.* 2009; 6(1):21. DOI: 10.1186/1479-5868-6-21

38. Sjöström, M., P. Oja, M. Hagströmer, B.J. Smith, and A. Bauman, Health-enhancing physical activity across European Union countries: the Eurobarometer study. *Journal of Public Health*. 2006; 14(5):291-300. DOI: 10.1007/s10389-006-0031-y
39. Bauman, A., B.E. Ainsworth, F. Bull, et al., Progress and pitfalls in the use of the International Physical Activity Questionnaire (IPAQ) for adult physical activity surveillance. *J Phys Act Health*. 2009; 6 Suppl 1:S5-8. DOI:
40. Plotnikoff, R.C., K.S. Courneya, L. Trinh, N. Karunamuni, and R.J. Sigal, Aerobic physical activity and resistance training: an application of the theory of planned behavior among adults with type 2 diabetes in a random, national sample of Canadians. *Int J Behav Nutr Phys Act* 2008; 5:61-61. DOI: 10.1186/1479-5868-5-61
41. Stringhini, S., A. Dugravot, M. Shipley, et al., Health Behaviours, Socioeconomic Status, and Mortality: Further Analyses of the British Whitehall II and the French GAZEL Prospective Cohorts. *PLOS Medicine*. 2011; 8(2):e1000419. DOI: 10.1371/journal.pmed.1000419
42. Gordon, S., C. Vandelanotte, A.T. Rayward, B. Murawski, and M.J. Duncan, Sociodemographic and behavioral correlates of insufficient sleep in Australian adults. *Sleep Health*. 2018; 5(1):12-17. DOI: <https://doi.org/10.1016/j.sleh.2018.06.002>
43. National Health and Medical Research Council, Australian Guidelines to reduce health risk from drinking alcohol, N.H.a.M.R. Council, Editor. 2009. National Health and Medical Research Council: Canberra, Australia. 179.
44. Chau, J.Y., H.P. van der Ploeg, S. Dunn, J. Kurko, and A.E. Bauman, A tool for measuring workers' sitting time by domain: the Workforce Sitting Questionnaire. *Br J Sports Med*. 2011; 45(15):1216-22. DOI: 10.1136/bjsports-2011-090214
45. Duncan, M.J., C.E. Kline, C. Vandelanotte, C. Sargent, N.L. Rogers, and L. Di Milia, Cross-sectional associations between multiple lifestyle behaviors and health-related

- quality of life in the 10,000 Steps cohort. *PLoS One*. 2014; 9(4):e94184. DOI: 10.1371/journal.pone.0094184
46. Buysse, D.J., C.F. Reynolds, 3rd, T.H. Monk, S.R. Berman, and D.J. Kupfer, The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res.*, 1989; 28(2):193-213. DOI:
47. Mollayeva, T., P. Thurairajah, K. Burton, S. Mollayeva, C.M. Shapiro, and A. Colantonio, The Pittsburgh sleep quality index as a screening tool for sleep dysfunction in clinical and non-clinical samples: A systematic review and meta-analysis. *Sleep Med Rev.*, 2016; 25:52-73. DOI: 10.1016/j.smr.2015.01.009
48. Ohayon, M., E.M. Wickwire, M. Hirshkowitz, et al., National Sleep Foundation's sleep quality recommendations: first report. *Sleep Health*. 2017; 3(1):6-19. DOI: <https://doi.org/10.1016/j.sleh.2016.11.006>
49. Zhai, L., Y. Zhang, and D. Zhang, Sedentary behaviour and the risk of depression: a meta-analysis. *Br J Sports Med*. 2015; 49(11):705. DOI:
50. LeBlanc, M., C. Merette, J. Savard, H. Ivers, L. Baillargeon, and C.M. Morin, Incidence and risk factors of insomnia in a population-based sample. *Sleep*. 2009; 32(8):1027-37. DOI:
51. Verger, P., C. Lions, and B. Ventelou, Is depression associated with health risk-related behaviour clusters in adults? *Eur J Public Health*. 2009; 19(6):618-24. DOI: 10.1093/eurpub/ckp057
52. Allen, M.S., E.E. Walter, and C. Swann, Sedentary behaviour and risk of anxiety: A systematic review and meta-analysis. *J Affect Disord* 2019; 242:5-13. DOI: <https://doi.org/10.1016/j.jad.2018.08.081>

53. Kline, C.E., The bidirectional relationship between exercise and sleep: Implications for exercise adherence and sleep improvement. *Am J Lifestyle Med.* 2014; 8(6):375-379. DOI: 10.1177/1559827614544437
54. Kyu, H.H., V.F. Bachman, L.T. Alexander, et al., Physical activity and risk of breast cancer, colon cancer, diabetes, ischemic heart disease, and ischemic stroke events: systematic review and dose-response meta-analysis for the Global Burden of Disease Study 2013. *BMJ.* 2016; 354. DOI: 10.1136/bmj.i3857
55. Clarke, D.M. and K.C. Currie, Depression, anxiety and their relationship with chronic diseases: a review of the epidemiology, risk and treatment evidence. *Med J Aust.* 2009; 190(7 Suppl):S54-60. DOI:
56. Kelly, S.J., M. Daniel, E. Dal Grande, and A. Taylor, Mental ill-health across the continuum of body mass index. *BMC public health.* 2011; 11:765-765. DOI: 10.1186/1471-2458-11-765
57. van der Zee, M.D., D. van der Mee, M. Bartels, and E.J.C. de Geus, Tracking of voluntary exercise behaviour over the lifespan. *Int J Behav Nutr Phys Act.* 2019; 16(1):17. DOI: 10.1186/s12966-019-0779-4
58. Suttorp, M.M., B. Siegerink, K.J. Jager, C. Zoccali, and F.W. Dekker, Graphical presentation of confounding in directed acyclic graphs. *Nephrol Dial Transplant.* 2015; 30(9):1418-23. DOI: 10.1093/ndt/gfu325
59. Australian Bureau of Statistics, Census of Population and Housing 2016, A.B.o. Statistics, Editor. 2017: Canberra, Australia.
60. Australian Bureau of Statistics, National Health Survey: first results, 2014-15, Australian Bureau of Statistics, Editor. 2015. Australian Bureau of Statistics: Canberra, Australia. 56.

61. McDowell, C.P., R.K. Dishman, M. Hallgren, C. MacDonncha, and M.P. Herring, Associations of physical activity and depression: Results from the Irish Longitudinal Study on Ageing. *Experimental Gerontology*. 2018; 112:68-75. DOI: <https://doi.org/10.1016/j.exger.2018.09.004>
62. Teychenne, M., K. Ball, and J. Salmon, Physical activity and likelihood of depression in adults: A review. *Prev Med*. 2008; 46(5):397-411. DOI: <https://doi.org/10.1016/j.ypmed.2008.01.009>
63. Lubans, D., J. Richards, C. Hillman, et al., Physical Activity for Cognitive and Mental Health in Youth: A Systematic Review of Mechanisms. *Pediatrics*. 2016; 138(3):e20161642. DOI: 10.1542/peds.2016-1642
64. White, R.L., M.J. Babic, P.D. Parker, D.R. Lubans, T. Astell-Burt, and C. Lonsdale, Domain-Specific Physical Activity and Mental Health: A Meta-analysis. *Am J Prev Med* 2017; 52(5):653-666. DOI: <https://doi.org/10.1016/j.amepre.2016.12.008>
65. Peeters, G., Y.R. van Gellecum, J.G. van Uffelen, N.W. Burton, and W.J. Brown, Contribution of house and garden work to the association between physical activity and well-being in young, mid-aged and older women. *Br J Sports Med*. 2014; 48(12):996-1001. DOI: 10.1136/bjsports-2012-091103
66. O'Connor, P.J., M.P. Herring, and A. Carvalho, Mental Health Benefits of Strength Training in Adults. *Am J Lifestyle Med* 2010; 4(5):377-396. DOI: 10.1177/1559827610368771
67. Wipfli, B.M., C.D. Rethorst, and D.M. Landers, The anxiolytic effects of exercise: a meta-analysis of randomized trials and dose-response analysis. *J Sport Exerc Psychol*. 2008; 30(4):392-410. DOI:

68. Henry, J.D. and J.R. Crawford, The short-form version of the Depression Anxiety Stress Scales (DASS-21): construct validity and normative data in a large non-clinical sample. *Br J Clin Psychol.* 2005; 44(Pt 2):227-39. DOI: 10.1348/014466505x29657
69. Franzen, P.L. and D.J. Buysse, Sleep disturbances and depression: risk relationships for subsequent depression and therapeutic implications. *Dialogues in clinical neuroscience.* 2008; 10(4):473-481. DOI:

ACCEPTED MANUSCRIPT

Table 1: Sample characteristics of women who responded to the Australian 10,000 Steps Project 2016 survey by symptoms of depression and/or anxiety (n=5180)

Mental health symptoms of \geq moderate severity ¹		Neither (n=4129)	Anxiety ¹ (n=317)	Depression ² (n=317)	Depression and anxiety (n=417)
		Mean \pm standard deviation			
Age	Years	50.6 \pm 11.3	47.0 \pm 11.9	50.0 \pm 11.5	46.9 \pm 13.1
Depression score	DASS (0 – 42)	3.3 \pm 3.4	7.4 \pm 3.6	18.0 \pm 5.5	22.1 \pm 8.0
Anxiety score	DASS (0 – 42)	2.3 \pm 2.3	12.9 \pm 3.7	4.4 \pm 2.6	16.4 \pm 6.2
Chronic disease diagnoses	Count	1.1 \pm 1.3	1.7 \pm 1.6	1.4 \pm 1.4	1.8 \pm 1.7
		Percentage (%)			
Income	AUD\$ \leq 70,000	824 (20.0)	77 (24.3)	75 (23.7)	140 (33.6)
	AUD\$>70,000k	2686 (65.0)	193 (60.9)	197 (62.2)	210 (50.4)
Employment	Unsure/ prefer not to say	619 (15.0)	47 (14.8)	45 (14.2)	67 (16.0)
	Unemployed	574 (13.9)	45 (14.2)	49 (15.5)	85 (20.4)
Cigarette Smoking	Employed	3555 (86.1)	272 (85.8)	268 (84.5)	332 (79.6)
	Current smoker	158 (3.8)	33 (10.4)	12 (3.8)	40 (9.6)
Alcohol consumption	Non-smoker	3971 (96.2)	284 (89.6)	305 (96.2)	377 (90.4)
	High risk	897 (21.7)	101 (31.9)	57 (18.0)	113 (27.1)
Sitting-time	Low risk	2677 (64.8)	166 (52.4)	196 (61.8)	240 (57.6)
	Non-drinker	555 (13.4)	50 (15.8)	64 (20.2)	64 (15.4)
Sleep quality	\geq 8 hours/day	2786 (67.5)	237 (74.8)	239 (75.4)	329 (78.9)
	<8 hours/day	1343 (32.5)	80 (25.2)	78 (24.6)	88 (21.1)
Sleep duration meets recommendation ⁴	Fairly bad/very bad	868 (21.0)	122 (38.5)	142 (44.8)	239 (57.3)
	Fairly good	2432 (58.9)	169 (53.3)	152 (48.0)	154 (36.9)
Sleep latency	Very good	829 (20.1)	26 (8.2)	23 (7.3)	24 (5.8)
	No	1706 (41.3)	206 (64.5)	177 (55.8)	275 (65.9)
Chronic disease diagnosis	Yes	2423 (58.7)	141 (44.5)	140 (44.2)	142 (34.1)
	Long (>30 minutes)	477 (11.6)	89 (28.1)	77 (24.3)	152 (36.5)
Ever diagnosed with depression and/or anxiety	Normal (\leq 30 min)	3652 (88.5)	228 (71.9)	240 (75.7)	265 (63.6)
	One or more	2486 (60.2)	239 (75.4)	210 (66.3)	322 (77.2)
Body mass index	None reported	1643 (39.8)	78 (24.6)	107 (33.8)	95 (22.8)
	Yes	785 (19.0)	160 (50.5)	160 (50.5)	265 (63.6)
Body mass index	No	3344 (81.0)	157 (49.5)	157 (49.5)	152 (36.5)
	<18.5 kg/m ²	59 (1.4)	5 (1.6)	7 (2.2)	11 (2.6)
	18.5-24.9 kg/m ²	1664 (40.3)	110 (34.7)	108 (34.1)	101 (24.2)
	25.0-29.9 kg/m ²	1252 (30.3)	72 (22.7)	88 (27.8)	116 (27.8)
	\geq 30 kg/m ²	1154 (28.0)	130 (41.0)	114 (36.0)	189 (45.3)

¹ Cut-off for 'moderate' anxiety symptoms \geq 10 points; ² Cut-off for 'moderate' depression symptoms \geq 14 points; ³ The internal consistency in this sample (Cronbach's alpha) was 0.91 for the DASS-21 depression subscale and 0.79 for the anxiety subscale; ⁴ Recommended sleep is 7-9 hours of sleep per night for <65-year olds and 7-8 hours of sleep per night for \geq 65-year olds.

Table 2: Intensity and domain-specific aerobic activity by group for women who responded to the Australian 10,000 Steps Project 2016 survey (n=5180)

	Activity recommendations met			
	Neither recommendation	Aerobic physical activity only	Resistance training only	Aerobic physical activity and resistance training
Sample size (%)	2215 (42.8)	1590 (30.7)	401 (7.7)	974 (18.8)
Total MET-minutes	1474±819	6331±4075	1899±695	6791±4830
	Walking activity MET-minutes			
Work	109±239	777±1110	81±187	589±993
Transport	155±250	376±600	159±223	352±558
Leisure	318±358	674±664	428±362	773±719
<i>Total walking MET</i>	<i>567±488</i>	<i>1726±1410</i>	<i>655±439</i>	<i>1627±1444</i>
	Moderate activity MET-minutes			
Work	50±150	528±996	44±145	414±910
Transport (cycling)	22±144	104±480	23±123	75±435
House/ yard	240±280	945±1025	228±231	736±939
House chores	224±266	766±824	217±237	627±757
Leisure	57±153	235±457	166±279	445±596
<i>Total moderate MET</i>	<i>725±609</i>	<i>3261±2620</i>	<i>784±531</i>	<i>2871±2768</i>
	Vigorous activity MET-minutes			
Work	61±190	825±1702	43±128	692±1554
House/yard	139±283	752±1154	113±221	636±1035
Leisure	132±293	626±479	423±479	1704±1459
<i>Total vigorous MET</i>	<i>183±332</i>	<i>1344±1889</i>	<i>460±481</i>	<i>2292±2165</i>
	Resistance training			
Days per week	1.1±0.3	1.2±0.4	3.8±1.1	4.1±1.3
	Sitting-time (%)			
≥8 hours/day	78.1	61.9	72.1	60.4

Values are mean±SD unless otherwise noted

Table 3: Multinomial regression analysis for associations between depression and/or anxiety and meeting physical activity recommendations for women who responded to the Australian 10,000 Steps Project 2016 survey (n=5180)

Recommendations met	n (%)	Model A ¹ RRR (95% CI)	Model B ² ARRR ¹ (95% CI)	Model C ³ ARRR ¹ (95% CI)
Neither depression (DASS<14) nor anxiety (DASS<10) (n=4129)				
Meets neither activity recommendation	1684 (40.8)			
Aerobic physical activity only	1282 (31.1)			
Resistance training only	327 (7.9)			
Aerobic physical activity and resistance training	836 (20.3)			
Anxiety (DASS≥10 points) (n=317)				
Meets neither activity recommendation	145 (45.7)	1	1	1
Aerobic physical activity only	97 (30.6)	0.88 (0.67 – 1.15)	0.91 (0.69 – 1.21)	1.00 (0.75 – 1.33)
Resistance training only	23 (7.3)	0.82 (0.52 – 1.29)	0.89 (0.56 – 1.41)	0.95 (0.59 – 1.52)
Aerobic physical activity and resistance training	52 (16.4)	0.72 (0.52 – 1.00)	0.79 (0.57 – 1.11)	0.89 (0.64 – 1.27)
Depression (DASS≥14) (n=317)				
Meets neither activity recommendation	165 (52.1)	1	1	1
Aerobic physical activity only	88 (27.8)	0.70 (0.54 – 0.92)	0.74 (0.56 – 0.97)	0.78 (0.59 – 1.04)
Resistance training only	21 (6.6)	0.67 (0.41 – 1.05)	0.69 (0.43 – 1.12)	0.71 (0.43 – 1.14)
Aerobic physical activity and resistance training	43 (13.6)	0.52 (0.37 – 0.74)	0.61 (0.43 – 0.86)	0.66 (0.46 – 0.95)
Depression (DASS≥14) & anxiety (DASS≥10) (n=417)				
Meets neither activity recommendation	221 (53.0)	1	1	1
Aerobic physical activity only	123 (29.5)	0.73 (0.58 – 0.92)	0.76 (0.59 – 0.97)	0.87 (0.67 – 1.13)
Resistance training only	30 (7.2)	0.70 (0.47 – 1.04)	0.78 (0.51 – 1.18)	0.83 (0.54 – 1.30)
Aerobic physical activity and resistance training	43 (10.3)	0.39 (0.28 – 0.55)	0.47 (0.33 – 0.67)	0.55 (0.39 – 0.80)

¹ Model A: unadjusted relative risk ratio; ² Model B: adjusted relative risk ratio: age, smoking, alcohol use, household income, education level, employment, sitting-time, sleep quality, sleep duration, sleep latency; ³ Model C: adjusted for Model B + chronic disease diagnosis.

Highlights

Lower depression risk when meeting aerobic activity recommendation

Lower comorbid depression/anxiety risk when meeting aerobic activity recommendation

Lower depression risk when meeting aerobic and resistance training recommendations

Lower comorbid depression/anxiety risk when meeting both activity recommendations

Magnitude of risk reduction greater when meeting both activity recommendations

ACCEPTED MANUSCRIPT