

Using camera traps to study hyenas: challenges, opportunities, and outlook

Author

Dheer, A, Samarasinghe, D, Dloniak, SM, Braczkowski, A

Published

2022

Journal Title

Mammalian Biology

Version

Version of Record (VoR)

DOI

[10.1007/s42991-021-00188-1](https://doi.org/10.1007/s42991-021-00188-1)

Rights statement

© The Author(s) 2022, corrected publication 2022. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Downloaded from

<http://hdl.handle.net/10072/428810>

Griffith Research Online

<https://research-repository.griffith.edu.au>



Using camera traps to study hyenas: challenges, opportunities, and outlook

Arjun Dheer¹ · Dinal Samarasinghe² · Stephanie M. Dloniak³ · Alexander Braczkowski⁴

Received: 29 November 2020 / Accepted: 4 October 2021 / Published online: 17 March 2022
© The Author(s) 2022, corrected publication 2022

Abstract

The use of remote camera traps has accelerated rapidly in the field of large carnivore science since the 1990s. Members of the *Hyaenidae* are important components of functional ecosystems in Africa and parts of the Middle East and South Asia, and make good candidates for study using camera traps. However, camera trap studies of hyenas remain rare in the literature when compared to species like tigers *Panthera tigris*, leopards *Panthera pardus*, and snow leopards *Panthera uncia*. In this paper, we examine the published use of camera traps for hyenas ($n = 34$ studies implemented between 2007 and 2020) and examine the logistical challenges of using camera traps, such as individual identification, limited sexual dimorphism, and complex social structures, for studies of hyena population biology, behavioral ecology, and conservation. We highlight what these challenges may mean for data analyses and interpretation. We also suggest potential benefits of further camera trap studies of this taxonomic family, including new insights into social behavior, range extensions, and robust density estimates.

Keywords Camera trapping · Carnivores · Hyenas · Spatially explicit capture–recapture

Introduction

Since Karanth's (1995) seminal paper using camera traps to estimate tiger densities with spatially explicit capture–recapture (SECR) models (White et al. 1982), the application of camera traps in wildlife research has expanded rapidly.

Handling editors: Scott Y.S. Chui and Leszek Karczmarski.

This article is a contribution to the special issue on “Individual Identification and Photographic Techniques in Mammalian Ecological and Behavioural Research – Part I: Methods and Concepts” — Editors: Leszek Karczmarski, Stephen C.Y. Chan, Daniel I. Rubenstein, Scott Y.S. Chui and Elissa Z. Cameron.

✉ Arjun Dheer
arjdheer@gmail.com

¹ Department of Evolutionary Ecology, Leibniz Institute for Zoo and Wildlife Research, 10315 Berlin, Germany

² Wildlife Research and Nature Conservation Foundation (WRNCF), 487/4 Old Road, Kottawa, Pannipitiya 10230, Sri Lanka

³ Department of Integrative Biology, Michigan State University, East Lansing, MI 48824, USA

⁴ School of Environmental Science and Engineering, Southern University of Science and Technology, Shenzhen, China

Camera traps have since been used to answer a myriad of ecological and applied conservation questions including density estimation, spatiotemporal activity patterns, and occupancy, and on organisms ranging from insects (Suckling et al. 2020) to African elephants *Loxodonta africana* (Moolman et al. 2019). As such, camera traps have transformed scientific understanding of mammalian distributions and assemblages (Rowcliffe and Carbone 2008). In particular, camera traps have found utility by scientists studying large carnivores, as these animals typically have large home ranges, are difficult to detect and observe, and often occur at naturally low densities (Balme et al. 2009).

Within the African large carnivore guild, the majority of publications that employ camera trapping skew towards African leopards *Panthera pardus* (e.g. du Preez et al. 2014; Braczkowski et al. 2016). Historically, research on hyenas has generally been dominated by questions in behavioral ecology, with camera trapping studies only recently gaining traction (Fig. 1). Long-term hyena research projects have mostly occurred in protected areas where the animals are readily observable and, therefore, conducive for behavioral studies (Smith et al. 2017). Among the hyenas, spotted hyenas *Crocuta crocuta* have been the main focus of published literature, with studies of aardwolves *Proteles cristatus*, brown hyenas *Hyaena brunnea*, and striped hyenas *Hyaena hyaena* being scarcer.




Fig. 1 A literature review of 34 studies (from 2007 to 2020) recovered on Google Scholar showing the hyena species, countries, and research topics encompassing the use of remote camera traps (see Supplementary Material 1 for a list of the studies). We limited our search to studies between 1900 and 2020 and to the first ten pages of results. Search terms were as follows: Spotted hyena: camera trap; *Crocuta crocuta*: camera trap; Striped hyena: camera trap; *Hyaena*

hyaena: camera trap; Brown hyena: camera trap; *Hyaena brunnea*: camera trap; Aardwolf: camera trap; *Proteles cristata*: camera trap. We removed all duplicates and classified studies into one of six classes. The “Other” category included only one study (Cusack et al. (2017)), which examined animal species identification using deep convolutional neural networks

For the conservation and management of these species, more studies are needed in areas where hyenas cannot be observed directly, making camera traps crucial in understudied ecosystems. Hyenas possess several traits that make them excellent candidates for camera trapping studies. However, these same traits can also pose challenges for data collection, analysis, and interpretation towards answering questions regarding their ecology and conservation.

We reviewed 34 recently published (between 2007 and 2020) camera trap studies of hyenas (see Supplementary Material). Here, we outline the positive and negative aspects of using camera traps to study members of the family *Hyaenidae* and provide a suite of key recommendations on how to optimize their use. Furthermore, we highlight the potential conservation and management implications of further camera trap studies of hyenas and how these may contribute to wildlife conservation and research in Africa and Asia.

Challenges and opportunities

Individual identification

All four hyena species have patterned pelage: they are either spotted (as in the spotted hyena) or striped (as in the aardwolf, brown hyena, and striped hyena). The natural markings on the left and right sides of the body are different for each animal, which makes individuals distinguishable and allows for capture–recapture analyses. Spotted hyenas are born with solid black natal coats; the coat brightens and spots gradually appear around the age of 1 month (Kruuk 1972). Though cubs are difficult to identify due to the fairly uniform natal coat, this can still be done based on differences in ear notches, abrasions on the fur, and other incidental markings (Frank et al. 1991). In

aardwolves, stripes are present along their flanks, fore-quarters, and legs, and spots may be present on the neck (Sliwa 1996; Spagnuolo et al. 2022). Brown hyenas are most readily identified by looking at the stripe patterns along their fore- and hindlegs (Spagnuolo et al. 2022).

Striped hyenas, like aardwolves, have stripes along their flanks and legs and do not have a natal coat (Rieger 1979). In all hyena species, ear notches and facial scars can also be used for individual identification (Thorn et al. 2009; Alam et al. 2015). However, such markings tend to change size and shape over time (Mills 1982) and should ideally only be used as supplementary features for individual identification. In general, the longer fur of aardwolves, brown hyenas, and striped hyenas can make them more difficult to individually identify than spotted hyenas via camera trap (authors' observation). It is, therefore, prudent to include an uncertainty level for identification in population models (ideally one that can be tested and quantified, as in Johansson et al. 2020).

Associating left and right flank images to the same hyena from camera traps is challenging because images are passively collected without control over which individual is photographed, in contrast to active photo identification surveys where individuals are photographed from both sides. The spotted hyena's propensity to aggregate in groups, perform scent-marking behaviors, and closely investigate and manipulate or destroy unprotected camera traps with their jaws (authors' observation) poses serious practical challenges. In photographs captured during such social activities, it becomes extremely difficult to keep track of specific individuals and assign two flank patterns to the same hyena, especially away from places that the hyenas regularly visit and can be repeatedly captured at (e.g. communal dens or large carcasses). In addition, hyenas regularly rest in puddles (authors' observation), which can discolor and obscure their unique pelage patterns. Lastly, the spots on a spotted hyena's coat tend to fade with age (Frank 1986), which can pose difficulties for long-term studies unless the photographs are of high quality. Accordingly, there are potential context- and

species-specific inaccuracies that may influence accurate individual identification and, therefore, population estimates.

Because hyenas are patterned asymmetrically, it is possible to use both complete (double flank) and partial (single flank) identities. When using single flanks, identities can be assigned by identifying individuals separately based on the available flank profiles (right and left) and later selecting the flank profile with the highest number of individuals identified for the analysis (Kalle et al. 2011). In long-term studies using a single flank approach, it is advisable to use a cut-off date (e.g. 1 year) after which there is a comparison of the number of left and right flank shots and the side with a larger sample size is henceforth chosen for all analyses for the project. This allows for the largest possible sample size up to a time limit and avoids the possibility that the flank with more photographs is discarded, which an a priori decision might induce. However, it has been suggested that single flank approaches lead to a loss of precision and an introduction of bias (Meredith 2017; Augustine et al. 2018; Petersen et al. 2019). It is possible to avoid these issues with the use of spatial partial identity models, which can probabilistically resolve the complete identity of single flank samples or photographs that are too blurry to identify using Bayesian techniques (Augustine et al. 2018; Davis et al. 2020).

The decision to use a single flank or double flank approach for hyenas can also depend on the propensity of the animals to aggregate in large groups. For studies in high density areas where hyenas generally aggregate in larger groups (e.g. for many spotted hyena populations), we recommend using only a single flank to identify individuals (Fig. 2). Using the double flank method in such populations may cause mismatches of left and right flank patterns, and, therefore, possibly cause inaccuracies in capture–recapture analyses. The less social hyena species forage alone more often than spotted hyenas (Mills 1982; Bothma and Walker 1999; Watts and Holekamp 2007) and using two flanks for each individual is more suitable. However, in study areas where the less social hyena species regularly forage in

Fig. 2 Camera trap images of the same individual spotted hyena taken at two separate camera trap stations in the Lake Mburo National Park, Uganda, in 2018 (author provided), showing identifiable spot patterns on the right flank



groups or dyads, it is also prudent to use a single flank (e.g. for striped hyenas: Harihar et al. 2010).

Sexual dimorphism

Of the four species in the family *Hyaenidae*, only the spotted hyena has masculinized external female genitalia (termed a pseudopenis) that is accompanied by a pseudoscrotum (Frank et al. 1990; Cunha et al. 2014). Along with minimal sexual dimorphism (Frank et al. 1990), this makes assigning sex to individual spotted hyenas difficult using camera traps (Davis et al. 2018). Aardwolves, brown hyenas, and striped hyenas also exhibit limited sexual dimorphism (Mills 1990; Van Jaarsveld et al. 1995; Wagner et al. 2008), and their long fur makes it difficult to view their external genitalia (Tichon et al. 2017; Hardouin et al. 2021). An inability to sex hyenas from camera trap images has ramifications for both density and occupancy studies. In spotted hyenas, adult females and adult immigrant males differ in their spatial patterning, with males ranging further from and within clan territories (Boydston et al. 2001). The difficulty of identifying sex influences the scale parameter (σ) in camera trap-based density estimation studies and also prevents an estimation of sex ratio (ψ_{sex}), which is an indicator of population health (Holekamp and Smale 1995; Braczkowski et al. 2020). Space use and ranging patterns either do not vary with sex in the other three hyena species, or the differences were deemed non-significant (Wagner et al. 2007; Brekelmans 2012; Edwards et al. 2020). Within our review of camera trapping studies, only one (Hardouin et al. 2021) derived density estimates based on sex, and only for aardwolves.

Despite the apparent lack of sexual dimorphism in hyenas, there are ways to accurately sex them under certain circumstances. In male spotted hyenas, an erect penis has a pointed glans (Fig. 3A). In females, when the pseudopenis is erect, the glans is rounded (Fig. 3B). Therefore, if a camera trap image has a hyena with an erect or semi-erect

phallus, one is able to assign a sex. Furthermore, the structure and shape of the pseudopenis changes post-parturition: the female spotted hyena develops a striking patch of pink scar tissue on the back of the phallus, which becomes saggy (Fig. 3C; Frank and Glickman 1994). Moreover, in all hyena species, post-parturition females tend to have pendulous teats due to lactation and suckling which can suffice to distinguish the sexes on camera trap images (Fig. 3D; Kent and Hill 2013; Davidson et al. 2019; Tichon et al. 2020). Regardless, it must be noted that researchers using camera traps are unlikely to be able to sex the majority of individual hyenas with certainty.

Finally, there are alternative approaches that can be used to estimate hyena densities when mark-recapture approaches are unsuitable. For example, camera-based methods that use distance sampling (Howe et al. 2017) and random encounter methods (Cusack et al. 2015) can produce density estimates for carnivores with less individually discernible features, such as lions. Occupancy studies can also be used as proxies for population status without the need for quantified density or abundance estimates, which makes individual or sex-specific identification unnecessary (Schuette et al. 2013). If sex is to be included as a variable in camera trap studies of hyenas, we urge researchers to use it only when sex can be determined unequivocally to avoid accruing errors in datasets and models (Johansson et al. 2020).

Social systems

Hyenas have unusual social systems when compared to other carnivores. Spotted hyenas live in clans that can range in size from fewer than 10 to more than 130 members. Clans are fission–fusion societies in which individuals can aggregate into large groups or be alone (Smith et al. 2008). Brown hyenas also live in clans, though these tend to be smaller than those of the spotted hyena, and adult males may be resident in a clan or nomadic (Knowles




Fig. 3 Camera trap images of four individual spotted hyenas taken at three separate camera trap stations at an undisclosed location in Kenya in 2016 (author provided), showing identifiable secondary sex

characteristics: **A** the male penis, **B** the young female pseudopenis, **C** the female pseudopenis post-parturition, and **D** visible teats on a female due to lactation and suckling

et al. 2009; Spagnuolo et al. 2022). Aardwolves live in socially (but not sexually) monogamous pairs (Richardson 1987). Striped hyenas are incipiently social and form small, stable, polyandrous groups (Wagner et al. 2008). However, they have a degree of flexibility in their sociality that has been suggested to depend on carrion availability (Califf et al. 2020; Tichon et al. 2020). In contrast to solitary carnivores such as the leopard, these unique and plastic social systems pose problems with estimating spatial scale parameters, non-independence of detections, and the application of the same methods across species ranges (Foster and Harmsen 2012). Densities of highly social species such as the spotted hyena may also be overestimated in camera trap studies, which is counterproductive for conservation and management efforts (Treves et al. 2010; Cusack et al. 2015). Capture probability can be modeled as a function of social parameters such as sex and then added to a complete density model later in the analysis (Foster and Harmsen 2012). Clan and/or territory sizes in a study area should also be included as covariates that may influence capture probability, as seen in other camera trap studies of ungulates (Massei et al. 2018; Palmer et al. 2018). Clan size can be estimated by assessing the gradual increase (and asymptotic stagnation) in the number of different hyenas detected in a territory (Stratford et al. 2020). Territory size can be estimate using minimum convex polygons or kernel densities (Nilsen et al. 2008).

As long as the aforementioned issues are addressed, concurrent density estimates of hyenas and other carnivores at a single study site can be obtained through camera trapping. Owing to their social complexity and abundance, the number of spotted hyena detections on camera traps is often higher than the number for sympatric carnivores (Rich et al. 2017a, b; Loveridge et al. 2020). This high detectability even in studies where hyenas were “bycatch” augurs well for biologists who may focus on other carnivores but want to provide densities for hyenas (O’Brien and Kinnaird 2011). In areas where hyenas occur at low or moderate densities, camera trap stations can still be spaced a standard distance apart (e.g. ~2 km) to provide valuable baseline status estimates in less optimal habitats or at the edge of species’ ranges (Hardouin et al. 2021). Indeed, the first camera trapping study to provide density estimates for aardwolves used a multi-species approach (Msuha 2009). In studies where density estimates are not possible to produce due to a lack of recaptures, point observations can provide evidence of the hyena presence in areas where they were previously not known to exist (Maurya et al. 2018). Single observations and the presence/absence studies along the edges of known hyena ranges can improve distribution records and provide valuable data on long-term trends on local extinctions and recolonization by hyenas (Spencer et al. 2020).

Outlook

We contend that the scope for camera trap studies of hyenas is large and that they facilitate updated, accurate range maps and a better understanding of their ecology in areas where hyenas cannot be directly observed (Fig. 1). IUCN Red List conservation status assessments are based on global range and population estimates, which in the case of hyenas, are impossible to accurately develop without camera traps (Gupta et al. 2009). Newer techniques that couple camera trapping with machine learning have become more popular (Tabak et al. 2019), but due to the challenges we raised above, they may be more difficult to use with hyenas. In recognition of this, other methods can be applied, e.g. citizen science efforts such as the Hyaena Distribution Mapping Project which involved trained experts (Killea 2020).

Hyenas are widespread in habitats that are harder to monitor than the open areas of the Mara-Serengeti ecosystem. For example, a camera trap survey in Gabon’s Haut-Ogooue province (Bohm 2015) provided baseline abundance estimates for spotted hyenas after several decades of presumed local extinction. Another camera trap survey detected a spotted hyena population and demonstrated the importance of Odzala-Kokoua National Park, Republic of Congo, in the species’ conservation in Central Africa (Henschel et al. 2014). In Namibia, a camera trap survey in a fenced system recorded the highest-density brown hyena population known to science (Edwards et al. 2019). Edwards’ study highlights the potential for camera traps to detect the effects of fences and closed systems on socio-spatial behavior in hyenas and other carnivores. Camera traps were also used to provide a thorough analysis of social networks, clan size, density, and other features of a spotted hyena population in Namibia which would have been impossible to monitor using direct observations (Stratford et al. 2020). In Nepal, camera traps were used to provide the first-ever record of an active striped hyena den in the country (Khanal et al. 2017). A camera trap study in Israel also suggested that striped hyenas, which were traditionally been described as strictly solitary, are in fact more social than previously thought (Tichon et al. 2020). Novel findings via camera trapping may lead to the documentation of range extensions and ostensibly allow for better conservation outcomes for this misunderstood and declining family. We encourage the use of camera traps for studying hyenas based on a sound understanding of their unique traits, which provide both challenges and opportunities for ongoing research on their distribution, abundance, and conservation.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s42991-021-00188-1>.

Funding Open Access funding enabled and organized by Projekt DEAL.

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alam MS, Khan JA, Pathak BJ (2015) Striped hyena (*Hyaena hyaena*) status and factors affecting its distribution in the Gir National Park and Sanctuary, India. *Folia Zool* 64:32–39. <https://doi.org/10.25225/fozo.v64.i1.a.2015>
- Augustine BC, Royle JA, Kelly MJ, Satter CB, Alonso RS, Boydston EE, Crooks KR (2018) Spatial capture–recapture with partial identity: an application to camera traps. *Ann Appl Stat* 12:67–95. <https://doi.org/10.1214/17-AOAS1091>
- Balme GA, Hunter LT, Slotow R (2009) Evaluating methods for counting cryptic carnivores. *J Wildl Manag* 73:433–441. <https://doi.org/10.2193/2007-368>
- Bohm T (2015) Population ecology, conservation status and genetics of the spotted hyena (*Crocuta crocuta*) in the Odzala-Kokoua National Park, Republic of Congo, including an assessment of the status of spotted hyenas in southeast Gabon. PhD thesis, Free University Berlin.
- Bothma JDP, Walker C (1999) The striped hyaena. In: Bothma JDP, Walker C (eds) *Larger carnivores of the African savannas*. Springer, Berlin, Heidelberg, pp 200–211
- Boydston EE, Morelli TL, Holekamp KE (2001) Sex differences in territorial behavior exhibited by the spotted hyena (*Hyaenidae*, *Crocuta crocuta*). *Ethology* 107:369–385. <https://doi.org/10.1046/j.1439-0310.2001.00672.x>
- Brackzkowski AR, Balme GA, Dickman A, Fattebert J, Johnson P, Dickerson T, Macdonald DW, Hunter L (2016) Scent lure effect on camera-trap based leopard density estimates. *PlosOne* 11:p. e0151033. <https://doi.org/10.1371/journal.pone.0151033>
- Brackzkowski AR, Gopalaswamy AM, Elliot NB, Possingham HP, Bezzina A, Maron M, Biggs D, Allan JR (2020) Restoring Africa's Lions: start with good counts. *Front Ecol Evol* 8:138. <https://doi.org/10.3389/fevo.2020.00138>
- Brekelmans W (2012) Sex differences in home range size, scent mark rates and hormones in the Aardwolf (*Proteles cristatus*). PhD thesis, University of Utrecht
- Califf KJ, Green DS, Wagner AP, Scribner KT, Beatty K, Wagner ME, Holekamp KE (2020) Genetic relatedness and space use in two populations of striped hyenas (*Hyaena hyaena*). *J Mammal* 101:361–372. <https://doi.org/10.1093/jmammal/gyz165>
- Cunha GR, Risbridger G, Wang H, Place NJ, Grumbach M, Cunha TJ, Weldele M, Conley AJ, Barcellos D, Agarwal S, Bhargava A (2014) Development of the external genitalia: perspectives from the spotted hyenas (*Crocuta crocuta*). *Differentiation* 87:4–22. <https://doi.org/10.1016/j.diff.2013.12.003>
- Cusack JJ, Swanson A, Coulson T, Packer C, Carbone C, Dickman AJ, Kosmala M, Lintott C, Rowcliffe JM (2015) Applying a random encounter model to estimate lion density from camera traps in Serengeti National Park, Tanzania. *J Wildl Manag* 79:1014–1021. <https://doi.org/10.1002/jwmg.902>
- Cusack JJ, Dickman AJ, Kalyahe M, Rowcliffe JM, Carbone C, Macdonald DW, Coulson T (2017) Revealing kleptoparasitic and predatory tendencies in an African mammal community using camera traps: a comparison of spatiotemporal approaches. *Oikos* 126:812–822. <https://doi.org/10.5061/dryad.br86d>
- Davidson Z, Dupuis-Desormeaux M, Dheer A, Pratt L, Preston E, Gilicho S, MacDonald S, Doncaster CP (2019) Borrowing from Peter to pay Paul: Managing threatened predators of endangered and declining prey species. *PeerJ* 7:e7916. <https://doi.org/10.7717/peerj.7916>
- Davis CL, Rich LN, Farris ZJ, Kelly MJ, Di Bitetti MS, Blanco YD, Albanesi S, Farhadinia MS, Gholikhani N, Hamel S, Harmsen BJ (2018) Ecological correlates of the spatial co-occurrence of sympatric mammalian carnivores worldwide. *Ecol Lett* 21:1401–1412. <https://doi.org/10.1111/ele.13124>
- Davis RS, Stone EL, Gentle LK, Mgoola WO, Uzal A, Yarnell RW (2021) Spatial partial identity model reveals low densities of leopard and spotted hyaena in a miombo woodland. *J Zool* 313:43–53. <https://doi.org/10.1111/jzo.12838>
- du Preez BD, Loveridge AJ, Macdonald DW (2014) To bait or not to bait: a comparison of camera-trapping methods for estimating leopard *Panthera pardus* density. *Biol Cons* 176:153–161. <https://doi.org/10.1016/j.biocon.2014.05.021>
- Edwards S, Noack J, Heyns L, Rodenwaldt D (2019) Evidence of a high density brown hyena population within an enclosed reserve: the role of fenced systems in conservation. *Mammal Res* 64:519–527. <https://doi.org/10.1007/s13364-019-00432-7>
- Edwards S, Noack J, Heyns L, Rodenwaldt D, Tarjan LM (2020) Socio-ecology of a high density brown hyena population within an enclosed reserve. *Mammal Res* 65:223–233. <https://doi.org/10.1007/s13364-020-00477-z>
- Foster RJ, Harmsen BJ (2012) A critique of density estimation from camera trap data. *J Wildl Manag* 76:224–236. <https://doi.org/10.1002/jwmg.275>
- Frank LG (1986) Social organization of the spotted hyena (*Crocuta crocuta*) I Demography. *Anim Behav* 34:1500–1509. [https://doi.org/10.1016/S0003-3472\(86\)80220-2](https://doi.org/10.1016/S0003-3472(86)80220-2)
- Frank LG, Glickman SE (1994) Giving birth through a penile clitoris: parturition and dystocia in the spotted hyena (*Crocuta crocuta*). *J Zool* 234:659–665. <https://doi.org/10.1111/j.1469-7998.1994.tb04871.x>
- Frank LG, Glickman SE, Powch I (1990) Sexual dimorphism in the spotted hyena (*Crocuta crocuta*). *J Zool* 221:308–313. <https://doi.org/10.1111/j.1469-7998.1990.tb04001.x>
- Frank LG, Glickman SE, Licht P (1991) Fatal sibling aggression, precocial development, and androgens in neonatal spotted hyenas. *Science* 252:702–704. <https://doi.org/10.1126/science.2024122>
- Gupta S, Mondal K, Sankar K, Qureshi Q (2009) Estimation of striped hyena *Hyaena hyaena* population using camera traps in Sariska Tiger Reserve, Rajasthan, India. *J Bombay Nat Hist Soc* 106:284
- Hardouin M, Searle CE, Strampelli P, Smit J, Dickman AJ, Lobora AL, Rowcliffe JM (2021) Density responses of lesser-studied carnivores to habitat and management strategies in southern Tanzania's Ruaha-Rungwa landscape. *PLoS ONE* 16:e0242293. <https://doi.org/10.1371/journal.pone.0242293>

- Harihar A, Ghosh M, Fernandes M, Pandav B, Goyal SP (2010) Use of photographic capture recapture sampling to estimate density of striped hyena (*Hyaena hyaena*): implications for conservation. *Mammalia* 74:83–87. <https://doi.org/10.1515/MAMM.2009.072>
- Henschel P, Malanda GA, Hunter L (2014) The status of savanna carnivores in the Odzala Kokoua National Park, northern Republic of Congo. *J Mammal* 95:882–892. <https://doi.org/10.1644/13-mamm-a-306>
- Holekamp KE, Smale L (1995) Rapid change in offspring sex ratios after clan fission in the spotted hyena. *Am Nat* 145:261–278. <https://doi.org/10.1086/285739>
- Howe EJ, Buckland ST, Després-Einspinner ML, Kühl HS (2017) Distance sampling with camera traps. *Methods Ecol Evol* 8:1558–1565. <https://doi.org/10.1111/2041-210X.12790>
- Johansson Ö, Samelius G, Wikberg E, Chapron G, Mishra C, Low M (2020) Identification errors in camera-trap studies result in systematic population overestimation. *Sci Rep* 10:1–10. <https://doi.org/10.1038/s41598-020-63367-z>
- Kalle R, Ramesh T, Qureshi Q, Sankar K (2011) Density of tiger and leopard in a tropical deciduous forest of Mudumalai Tiger Reserve, southern India, as estimated using photographic capture-recapture sampling. *Acta Theriol* 56:335–342. <https://doi.org/10.1007/s13364-011-0038-9>
- Karanth KU (1995) Estimating tiger *Panthera tigris* populations from camera-trap data using capture—recapture models. *Biol Cons* 71:333–338. [https://doi.org/10.1016/0006-3207\(94\)00057-W](https://doi.org/10.1016/0006-3207(94)00057-W)
- Kent VT, Hill RA (2013) The importance of farmland for the conservation of the brown hyaena *Parahyaena brunnea*. *Oryx* 47:431–440. <https://doi.org/10.1017/S0030605312001007>
- Khanal C, Baniya S, Acharye M (2017) First confirmed record of striped hyaena (*Hyaena hyaena*) den in Nepal. *J Biodivers Endanger Species* 5:2. <https://doi.org/10.4172/2332-2543.1000195>
- Killea A (2020) Brown Hyena (*Hyaena brunnea*) Distribution: Nuances in Modeling a Generalist Species. Master's thesis, Duke University
- Knowles JC, Coeverden V, de Groot PJ, Wiesel I, Boag PT (2009) Microsatellite variation in Namibian brown hyenas (*Hyaena brunnea*): population structure and mating system implications. *J Mammal* 90:1381–1391. <https://doi.org/10.1644/08-MAMM-A-298R1.1>
- Kruuk H (1972) The spotted hyena: a study of predation and social behavior. University of Chicago Press
- Loveridge AJ, Sousa LL, Seymour-Smith J, Hunt J, Coals P, O'Donnell H, Lindsey PA, Mandisodza-Chikerema R, Macdonald DW (2020) Evaluating the spatial intensity and demographic impacts of wire-snare bush-meat poaching on large carnivores. *Biol Cons* 244:108504. <https://doi.org/10.1016/j.biocon.2020.108504>
- Massei G, Coats J, Lambert MS, Pietravallo S, Gill R, Cowan D (2018) Camera traps and activity signs to estimate wild boar density and derive abundance indices. *Pest Manag Sci* 74:853–860. <https://doi.org/10.1002/ps.4763>
- Maurya V, Singh JP, Naseem K, Mehra S, Dhakate PM, Verma N, Ansari AG (2018) Photographic evidence of Striped Hyena *Hyaena hyaena* (Mammalia: Carnivora: Hyaenidae) in Ramnagar forest division Uttarakhand, India. *J Threat Taxa* 10:13017–13019. <https://doi.org/10.11609/jott.3555.10.15.13017-13019>
- Meredith M (2017) SECR with unpaired cameras. Available online at http://mmeredith.net/blog/2017/SECR_unpaired_cameras.htm. Accessed 21 Apr 2021
- Mills MGL (1982) Factors affecting group size and territory size of the brown hyaena, *Hyaena brunnea* in the southern Kalahari. *J Zool* 198:39–51. <https://doi.org/10.1111/j.1469-7998.1982.tb02059.x>
- Mills MGL (1990) Kalahari Hyaenas: comparative behavioral ecology of two species. Unwin Hyman, London
- Moolman L, de Morney MA, Ferreira SM, Ganswindt A, Poole JH, Kerley GI (2019) And Then There Was One: a camera trap survey of the declining population of African Elephants in Knysna, South Africa. *Afr J Wildl Res* 49:16–26. <https://doi.org/10.3957/056.049.0016>
- Msuh MJ (2009) Human impacts on carnivore biodiversity inside and outside protected areas in Tanzania. PhD thesis, UCL (University College London)
- Nilsen EB, Pedersen S, Linnell JD (2008) Can minimum convex polygon home ranges be used to draw biologically meaningful conclusions? *Ecol Res* 23:635–639. <https://doi.org/10.1007/s11284-007-0421-9>
- O'Brien TG, Kinnaird MF (2011) Density estimation of sympatric carnivores using spatially explicit capture–recapture methods and standard trapping grid. *Ecol Appl* 21:2908–2916. <https://doi.org/10.1890/10-2284.1>
- Palmer MS, Swanson A, Kosmala M, Arnold T, Packer C (2018) Evaluating relative abundance indices for terrestrial herbivores from large-scale camera trap surveys. *Afr J Ecol* 56:791–803. <https://doi.org/10.1111/aje.12566>
- Petersen WJ, Savini T, Steinmetz R, Ngoprasert D (2019) Estimating Leopard Cat *Prionailurus bengalensis* Kerr, 1792 (Carnivora: Felidae) density in a degraded tropical forest fragment in north-eastern Thailand. *J Threat Taxa* 11:13448–13458. <https://doi.org/10.11609/jott.4553.11.4.13448-13458>
- Rich LN, Davis CL, Farris ZJ, Miller DA, Tucker JM, Hamel S, Farhadinia MS, Steenweg R, Di Bitetti MS, Thapa K, Kane MD (2017a) Assessing global patterns in mammalian carnivore occupancy and richness by integrating local camera trap surveys. *Glob Ecol Biogeogr* 26:918–929. <https://doi.org/10.1111/geb.12600>
- Rich LN, Miller DAW, Robinson HS, McNutt JW, Kelly MJ (2017b) Carnivore distributions in Botswana are shaped by resource availability and intraguild species. *J Zool* 303:9098. <https://doi.org/10.1111/jzo.12470>
- Richardson PRK (1987) Aardwolf mating system: overt cuckoldry in an apparently monogamous mammal. *S Afr J Sci* 83:405. https://doi.org/10.10520/AJA00382353_5361
- Rieger I (1979) A review of the biology of striped hyenas, *Hyaena hyaena* (Linné, 1758). *Säuget Mitt* 27:81–95
- Rowcliffe JM, Carbone C (2008) Surveys using camera traps: are we looking to a brighter future? *Anim Conserv* 11:185–186. <https://doi.org/10.1111/j.1469-1795.2008.00180.x>
- Schuette P, Wagner AP, Wagner ME, Creel S (2013) Occupancy patterns and niche partitioning within a diverse carnivore community exposed to anthropogenic pressures. *Biol Conserv* 158:301–312. <https://doi.org/10.1016/j.biocon.2012.08.008>
- Sliwa A (1996) A functional analysis of scent marking and mating behaviour in the aardwolf, *Proteles cristatus*. PhD thesis, University of Pretoria
- Smith JE, Kolowski JM, Graham KE, Dawes SE, Holekamp KE (2008) Social and ecological determinants of fission–fusion dynamics in the spotted hyaena. *Anim Behav* 76:619–636. <https://doi.org/10.1016/j.anbehav.2008.05.001>
- Smith JE, Lehmann KD, Montgomery TM, Strauss ED, Holekamp KE (2017) Insights from long-term field studies of mammalian carnivores. *J Mammal* 98:631–641. <https://doi.org/10.1093/jmammal/gyw194>
- Spagnuolo OSB, Lemerle MA, Holekamp KE, Wiesel I (2022) The value of individual identification in studies of free-living hyenas and aardwolves. *Mamm Biol (Special Issue)* 102(3) (In Press)
- Spencer K, Sambrook M, Bremner-Harrison S, Cilliers D, Yarnell RW, Brummer R, Whitehouse-Tedd K (2020) Livestock guarding dogs enable human-carnivore coexistence: first evidence of equivalent carnivore occupancy on guarded and unguarded farms. *Biol Cons* 241:108256. <https://doi.org/10.1016/j.biocon.2019.108256>

- Stratford K, Stratford S, Périquet S (2020) Dyadic associations reveal clan size and social network structure in the fission–fusion society of spotted hyaenas. *Afr J Ecol* 58:182–192. <https://doi.org/10.1111/aje.12641>
- Suckling DM, Stanbury ME, Lennon O, Colhoun KM, Chinellato F, El Sayed AM (2020) Kairomone and camera trapping New Zealand flower Thrips *Thrips Obscuratus*. *Insects* 11:622. <https://doi.org/10.3390/insects11090622>
- Tabak MA, Norouzzadeh MS, Wolfson DW, Sweeney SJ, VerCauteren KC, Snow NP, Halseth JM, Di Salvo PA, Lewis JS, White MD, Miller RS (2019) Machine learning to classify animal species in camera trap images: applications in ecology. *Methods Ecol Evol* 10:585–590. <https://doi.org/10.1111/2041-210X.13120>
- Thorn M, Scott DM, Green M, Bateman PW, Cameron EZ (2009) Estimating brown hyaena occupancy using baited camera traps. *Afr J Wildl Res* 39:1–10. <https://doi.org/10.3957/056.039.0101>
- Tichon J, Rotem G, Ward P (2017) Estimating abundance of striped hyenas (*Hyaena hyaena*) in the Negev Desert of Israel using camera traps and closed capture–recapture models. *Eur J Wildl Res* 63:5. <https://doi.org/10.1007/s10344-016-1069-y>
- Tichon J, Gilchrist JS, Rotem G, Ward P, Spiegel O (2020) Social interactions in striped hyena inferred from camera trap data: is it more social than previously thought? *Curr Zool* 66:345–353. <https://doi.org/10.1093/cz/zoaa003>
- Treves A, Mwima P, Plumptre AJ, Isoke S (2010) Camera-trapping forest woodland wildlife of western Uganda reveals how gregariousness biases estimates of relative abundance and distribution. *Biol Conserv* 143:521–528. <https://doi.org/10.1016/j.biocon.2009.11.025>
- Van Jaarsveld AS, Richardson PRK, Anderson MD (1995) Post-natal growth and sustained lactational effort in the aardwolf: life-history implications. *Funct Ecol* 9:492–497. <https://doi.org/10.2307/2390014>
- Wagner AP, Creel S, Frank LG, Kalinowski ST (2007) Patterns of relatedness and parentage in an asocial, polyandrous striped hyena population. *Mol Ecol* 16:4356–4369. <https://doi.org/10.1111/j.1365-294X.2007.03470.x>
- Wagner AP, Frank LG, Creel S (2008) Spatial grouping in behaviorally solitary striped hyaenas, *Hyaena hyaena*. *Anim Behav* 75:1131–1142. <https://doi.org/10.1016/j.anbehav.2007.08.025>
- Watts HE, Holekamp KE (2007) Hyena societies. *Curr Biol* 17:657–660. <https://doi.org/10.1016/j.cub.2007.06.002>
- White GC, Anderson DR, Burnham KP, Otis DL (1982) Capture–recapture and removal methods for sampling closed populations. Los Alamos National Laboratory. <https://doi.org/10.2307/2347719>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.