

## **Cadmium adsorption behavior of some soils from northern of Iran**

### Author

Esfandbod, Maryam, Forghani, Akbar, Adhami, Ebrahim, Rashti, Mehran Rezaei

### Published

2010

### Conference Title

19th World Congress of Soil Science: Symposium: Soil and Human Culture

### Rights statement

© 2010 Australian Soil Science Society. The attached file is reproduced here in accordance with the copyright policy of the publisher. Please refer to the conference's website for access to the definitive, published version.

### Downloaded from

<http://hdl.handle.net/10072/49002>

### Link to published version

<https://www.iuss.org/>

### Griffith Research Online

<https://research-repository.griffith.edu.au>

# Cadmium adsorption behavior of some soils from northern of Iran

Maryam Esfandbod<sup>A</sup>, Akbar Forghani<sup>B</sup>, Ebrahim Adhami<sup>C</sup> and Mehran Rezaei Rashti<sup>D</sup>

<sup>A</sup> Soil Science Department of, Guilan University, Rasht, Iran, Email [maryam.esfandbod@gmail.com](mailto:maryam.esfandbod@gmail.com)

<sup>B</sup> Faculty of Soil Science, Guilan University, Rasht, Iran, Email

<sup>C</sup> Soil Science Department, College of Agriculture, Yasouj University, Yasouj, Iran, Email [eadhami@mail.yu.ac.ir](mailto:eadhami@mail.yu.ac.ir)

<sup>D</sup> Department of Soil Science, Guilan University, Rasht, Iran, Email [mehran\\_325@yahoo.com](mailto:mehran_325@yahoo.com)

## Abstract

Cadmium adsorption behavior (Cd) of soils is an important process which exerts a major influence on its uptake by plant roots. Cadmium (Cd) sorption characteristics of 20 surface soils (0-30 cm) with a wide range of soil physicochemical properties, was investigated. Two grams of each sample were equilibrated for 24h at laboratory temperature ( $25 \pm 2^\circ\text{C}$ ) with 40ml, 0.01M  $\text{CaCl}_2$  containing 5 to 500 mg  $\text{Cd L}^{-1}$ . Langmuir, Freundlich and Temkin adsorption isotherms were used to evaluate the trend of Cd adsorption. The data of Cd adsorption was well fitted to Langmuir and Freundlich adsorption isotherms. However, the fit to Langmuir adsorption model was closer than that of Freundlich. Multiple stepwise regression analysis indicated that, the adsorption capacity of the soils for Cd increased with an increase in pH, CEC and calcium carbonate equivalent (CCE).

## Key Words

Adsorption, Cadmium, Freundlich, Isotherm, Langmuir, Temkin

## Introduction

Cadmium (Cd) contamination imposes an adverse effect on environmental quality and constitutes a serious threat not only to plants and animals but also to human lives (Martin-Garin et al. 2002). Cadmium adsorption and desorption by soils is one of the factors that control its bioavailability and transfer in the environment (Singh and Nayyar 1993). Langmuir (Ramachandran and Souza 1999; Singh and Nayyar 1993), Freundlich (Ramachandran and Souza 1999), and Temkin (Maftoun et al. 2004) equations have been commonly used to assess the pattern of heavy metals adsorption by soils. The objectives of the present study were to (1) evaluate the pattern of Cd adsorption of some soils from Guilan province, and (2) to determine the effect of soil properties on Cd adsorption.

## Methods

Twenty surface soil samples (0-30 cm) varying widely in physicochemical properties (Table 1) from Guilan province in northern part of Iran, were used in the present study. 2g subsample of each soil was placed in centrifuge tubes and 40ml of 0.01M  $\text{CaCl}_2$  containing different concentrations of Cd (ranging from 5 to 500 mg  $\text{Cd L}^{-1}$ ) was added and shaken for 1h at room temperature and allowed to equilibrate at  $25 \pm 2^\circ\text{C}$  for 24h. The suspensions were then shaken for additional 30 min. Samples were centrifuged for 10 min and the supernatant was analysed for Cd by atomic absorption spectrophotometry. The amount of Cd adsorbed was calculated as the difference between initial and final Cd concentrations. The adsorption data were fitted to Freundlich, Langmuir, and Temkin adsorption isotherms. The stepwise multiple regression procedure was used to study the relationship between the Cd sorption coefficients and the physicochemical properties of the soils.

**Table 1. Range of physicochemical characteristics of the soils.**

	Clay (%)	pH	OM (%)	CCE (%)	CEC ( $\text{cmol}_c \text{ kg}^{-1}$ )
Min.	8.00	4.21	1.53	1.59	14.18
Max.	52.00	7.61	9.77	16.51	58.46
Ave.	31.92	6.58	4.99	7.37	37.20

## Equations

Freundlich  $\text{Log } X = \log k_F + n_F \log C$  (1)

Langmuir  $C / X = [1 / (K_L b_L)] + (1 / b_L) C$  (2)

In Langmuir equation,  $(X / C)$  is termed the distribution coefficient ( $K_d$ ) (Bolt and Bruggenwert 1976):

$$X / C = K b = K_d \quad (3)$$

Temkin  $X = k_{T1} + k_{T2} \ln C \quad (4)$

Where  $X$  is the amount of Cd sorbed per unit weight of soil ( $\text{mg kg}^{-1}$  soil),  $C$  is Cd concentration in equilibrium solution ( $\text{mg L}^{-1}$ ),  $b_L$  is Cd adsorption maxima ( $\text{mg kg}^{-1}$  soil),  $K_L$  is a coefficient related to Cd bonding energy ( $\text{L mg}^{-1}$ ),  $K_d$  is Cd distribution coefficient ( $\text{L kg}^{-1}$  soil),  $k_F$  is Cd adsorption capacity ( $\text{mg kg}^{-1}$  soil),  $n_F$  is rate of Cd adsorption,  $k_{T1}$  and  $k_{T2}$  are Cd adsorption coefficients.

## Results

Langmuir and Freundlich isotherms were well fitted to the adsorption data (Table 2).

**Table 2. Freundlich and Langmuir adsorption isotherm parameters and coefficients of determination ( $r^2$ ).**

	Freundlich Eq.			Langmuir Eq.			
	$r^2$	$k_F$	$n_F$	$r^2$	$K_L$	$b_L$	$K_d$
Min.	0.82 <sup>a</sup>	29.7	0.22	0.86 <sup>a</sup>	0.01	1105	17.10
Max.	0.98 <sup>a</sup>	1072.9	0.72	1.00 <sup>a</sup>	0.18	5000	702.10
Ave.	0.92 <sup>a</sup>	504.5	0.41	0.97 <sup>a</sup>	0.07	3506	236.60

<sup>a</sup> Significant at  $p \leq 0.001$

$k_F$  value ranged from 29.7 to 1072. The  $K_F$  value indicates the quantity of adsorption when the equilibrium concentration is  $1 \text{ mg L}^{-1}$  (Kariman and Cox, 1978). Multiple linear regression analysis between  $K_F$  and soil properties (Eq. (5) and Eq.(6)) indicated that 68 percent of  $K_F$  variation could be explained by cation exchange capacity (CEC) and pH. As it can be clearly seen from these equations, adsorbed Cd increased linearly with increase in pH and CEC. However these properties showed significant negative correlation with  $n_F$  coefficient, similar result has been reported by Ramachandran and Souza (1999).

$$k_F = -1658.47 + 328.97 \text{ pH} \quad r^2 = 0.60 \quad p \leq 0.001 \quad (5)$$

$$k_F = -2163.45 + 342.58 \text{ pH} + 11.17 \text{ CEC} \quad r^2 = 0.68 \quad p \leq 0.001 \quad (6)$$

Clay and OM surfaces are the main sorbent of Cd; however, the nonsignificant correlation found between  $k_F$  with clay and OM might be due to the fact that the adsorption of Cd by tested soils could be more affected by the kind of clay and OM than the amount of clay and OM.

$K_L$  coefficient was correlated with some soil properties. Multiple stepwise regression analysis showed that the simple regression equation could adequately describe such relations:

$$K_L = 0.001 + 0.009 \text{ CCE} \quad r^2 = 0.63 \quad p \leq 0.001 \quad (7)$$

$$K_L = -0.21 + 0.04 \text{ pH} \quad r^2 = 0.56 \quad p \leq 0.001 \quad (8)$$

$$K_L = -0.123 + 0.006 \text{ CCE} + 0.022 \text{ pH} \quad r^2 = 0.71 \quad p \leq 0.001 \quad (9)$$

These findings are similar to those reported by Maftoun et al. (2004) and Ramachandran and Souza (1999).

In the present study, based on soil pH, and initial Cd concentration, the Cavallaro and McBride solubility diagram (Cavallaro and McBride 1978) indicates the possibility of solid phase Octavite ( $\text{CdCO}_3$ ) precipitation in some soils. The calculated Langmuir  $b_L$ , ranged from 1105 to 5000 (Table 2). The relatively high  $b_L$  values in the present study revealed that the soil removed a large amount of Cd from solution.

Stepwise regression analysis between  $b_L$  and soil properties showed the following relationships:

$$b_L = 1133.05 + 63.80 \text{ CEC} \quad r^2 = 0.48 \quad p \leq 0.001 \quad (10)$$

$$b_L = -2482.02 + 68.79 \text{ CEC} + 521.62 \text{ pH} \quad r^2 = 0.75 \quad p \leq 0.001 \quad (11)$$

which, indicates that CEC and pH are the main soil properties controlling maximum Cd adsorption in the studied soils.

The distribution coefficient,  $K_d$  shows the ratio of the Cd concentration in the soil to the concentration of  $\text{Cd}^{+2}$  in the solution phase.  $K_d$  values varied greatly between soils, ranging from 17.10 to 702.10 ( $\text{L kg}^{-1}$  soil) and showed a highly significant correlation with CCE ( $r^2 = 0.63$ ,  $p \leq 0.001$ ) and pH ( $r^2 = 0.52$ ,  $p \leq 0.001$ ). These results clearly demonstrate that pH and, to a greater extent CCE, contribute to the variation of  $K_d$  in the soils studied (Eq. (12)). A portion of the removal of Cd from the solution phase may have been a result of precipitation of  $\text{CdCO}_3$ . Similar results have been reported by Yin et al. (2002) and Sauve et al. (2000).

$$K_d = -435.31 + 23.03 \text{ CCE} + 76.38 \text{ pH} \quad r^2 = 0.70 \quad p \leq 0.001 \quad (12)$$

## Conclusion

From results reported herein, it is concluded that the adsorption of Cd increased with increasing solution Cd

concentration and the soils varied in Cd sorption capacity. The results showed that CEC and pH were the most important soil properties affecting Cd adsorption in these soils. Alkaline soils favoured high adsorption of Cd and possibly precipitation as CdCO<sub>3</sub>, which may lead to less bioavailability of Cd; whereas, acid soils adsorbed less Cd, and more Cd was present in solution, which may result in high availability. Although, the Cd adsorption data showed a highly significant fit to Freundlich and Langmuir isotherms, Comparison of coefficients of determination of Langmuir and Freundlich equation demonstrated that Langmuir adsorption isotherm has a better fit than Freundlich.

## References

- Bolt GH, Bruggenwert MGM (Eds.) (1976) Soil chemistry. In 'Basic elements-Part A. (Elsevier Scientific Publishing Co: Amsterdam. the Netherlands)
- Cavallaro N, McBride MB (1978) Copper and cadmium adsorption characteristics of selected acid and calcareous soils. *Soil Sci. Soc. Am. J.* **42**, 550-556.
- Karimian N, Cox FR (1978) Adsorption and extractability of molybdenum in relation to some chemical properties of soil. *Soil Sci. Soc. Am. J.* **42**, 757-761.
- Maftoun M, Rassooli F, Ali-Nejad Z, Karimian N (2004) Cadmium sorption behavior in some highly calcareous soils of Iran. *Commun. Soil Sci. Plant Anal* **35**, 1271-1282.
- Martin-Garin A, Gaudet JP, Charlet L, Vitart X (2002) A dynamic study of the sorption and the transport processes of cadmium in calcareous sandy soils. *Waste Manage* **22**, 201-207.
- Ramachandran V, Souza TJD (1999) Adsorption of cadmium by Indian soils. *Water Air Soil Pollut* **111**, 225-234.
- Sauve S, Hendershot W, Allen HE (2000) Solid-solution partitioning of metals in contaminated soils: Dependence on pH, total metal burden and organic matter. *Environ. Sci. Technol* **34**, 1125-1131.
- Singh SP, Nayyar VK (1993) Adsorption of cadmium on alkaline and calcareous soils. *J. Indian Soc. Soil Sci* **41**, 270-273.
- Sukreeyapongse O, Holm PE, Strobel BW, Panichsakpatana S, Magid J, Hansen HCB (2002) pH-dependent release of cadmium, copper and lead from natural and sludge-amended soils. *J. Environ. Qual* **31**, 1901-1909.
- Yin Y, Impellitteri CA, You SJ, Allen HE (2002) The importance of organic matter distribution and extract soil: Solution ratio on the desorption of heavy metals from soils. *Sci. Total Environ* **287**, 107-119.