

Self-Assessed Competence of Experienced Expatriate Nurses in a Rural and Remote Setting

Author

Aqtash, Salah, Robb, Walter Frederick, Hunter, Lyndal Heather, Almuhtasib, Majdi, Hamad, Abeer, Brownie, Sharon Mary

Published

2017

Journal Title

SAGE Open Nursing

Version

Version of Record (VoR)

DOI

[10.1177/2377960817702382](https://doi.org/10.1177/2377960817702382)

Rights statement

© The Author(s) 2017. This article is distributed under the terms of the Creative Commons AttributionNonCommercial 3.0 License (<http://www.creativecommons.org/licenses/by-nc/3.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Downloaded from

<http://hdl.handle.net/10072/377968>

Griffith Research Online

<https://research-repository.griffith.edu.au>

Self-Assessed Competence of Experienced Expatriate Nurses in a Rural and Remote Setting

SAGE Open Nursing
Volume 3: 1–17
© The Author(s) 2017
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/2377960817702382
journals.sagepub.com/home/son



Salah Aqtash^{1,2}, Walter Frederick Robb², Lyndal Heather Hunter²,
Majdi Almuhtasib³, Abeer Hamad³, and Sharon Mary Brownie^{4,5,6}

Abstract

We aimed to measure the self-assessed level of competence among nurses working in the public hospitals of Al-Gharbia Region, a remote rural region of United Arab Emirates, and to explore the factors associated with the nurses' self-perceived competency. The Nurse Competency Scale, which measures the self-assessed level of competency of nurses, has been validated in a variety of clinical settings, in facilities of various sizes, and in small and large cohorts. However, its application among an expatriate nursing workforce working in small hospitals and health facilities in remote and rural areas has not been examined. We used the Nurse Competency Scale to survey the nursing workforce in Al-Gharbia's public hospitals in United Arab Emirates. All 435 practicing registered nurses with more than 3 months clinical experience in the network were invited to participate. Data were collected electronically and analyzed by international collaborators. Statistical analysis included analysis of variance, Kruskal–Wallis, multiple linear regression, χ^2 test of independence, and Cronbach's α . Totally, 189 responses were analyzed (43.4% response rate). Overall self-assessed levels of competence were uniformly "very good" across all competence categories. The overall score (84.3) was higher than those found in most other studies. Frequency of use was the most outstanding variable influencing self-assessed competence. Total years of experience were the next significant variable. Some items of the scale were not yet applicable to activities in the region, particularly those relating to supervision of students. The high scores achieved by expatriate nurses in the small hospitals of Al-Gharbia reflect well on the rigor of the recruitment process, ongoing cross-training and functional competency assessment. Policies and practices aimed at recruiting experienced expatriate nurses and providing opportunities to use competencies continue to be critical in providing a quality nursing service in a rural and remote region.

Keywords

professional competence, nursing services, continuing nurse education, self-assessment, United Arab Emirates

Date received: 10 August 2016; revised: 18 February 2017; accepted: 4 March 2017

The United Arab Emirates (UAE) has a high-quality health-care system, characterized by well-equipped, well-staffed, licensed hospitals; qualified and licensed health-care providers; and universal medical insurance coverage. The competency of nurses is taken seriously, with high stakes (including legal and immigration consequences) for malpractice. This study investigated self-assessed competence among an expatriate nursing workforce in Al-Gharbia, the geographically largest and least populated region in the Emirate of Abu Dhabi. In part, it represents an effort to find a relevant, useful, and practical assessment tool that may assist in maintaining high standards of nursing competency

¹College of Health Sciences, School of Nursing, Walden University, Minneapolis, MN, USA

²School of Nursing and Midwifery, Griffith University, Queensland, Australia

³Al-Gharbia Hospitals, Abu Dhabi Health Services Company (SEHA), Abu Dhabi, UAE

⁴School of Nursing and Midwifery, Aga Khan University, Nairobi, Kenya

⁵Office Pro-Vice Chancellor Health, Griffith University, Australia

⁶Oxford PRAXIS Forum, Oxford University, Oxford, UK

Corresponding Author:

Sharon Mary Brownie, Griffith University, University Drive, Gold Coast Campus, Gold Coast, Queensland 4222, Australia.

Email: s.brownie@griffith.edu.au



in Al-Gharbia. This study reports a survey conducted in 2015 in the region's public hospitals network, which used the Nurse Competence Scale (NCS), a popular, validated survey instrument (Meretoja, Isoaho, & Leino-Kilpi, 2004).

This study has relevance for the emerging field of self-assessed competence as a marker for professional nursing competence more generally. It is relevant to modern, well-resourced regional hospitals with in-service training and preceptor programs, rural and remote health-care systems where nurses provide services across multiple clinical areas, often with limited frequency, and the management of expatriate nurses.

Background

Al-Gharbia. The Al-Gharbia region, Abu Dhabi, sits at the border between the UAE and Saudi Arabia and is largely sand desert. It is part of the Arabian Peninsula's "empty quarter" (Figure 1). In 2014, Al-Gharbia had 8.9 persons per square kilometer and a total population of 313,000 people, of which 9.4% were UAE citizens. The remaining 90.6% were noncitizens, mainly employed in oil and gas production, regional development, and accompanying services. Of noncitizens, 84.4% were male and predominantly working age, with few aged over 60 years (Statistics Centre Abu Dhabi, 2015).

Al-Gharbia is medically well serviced. The region has seven rural townships with modern, licensed hospitals that are well-equipped and resourced. The public hospitals network provides primary, secondary, and selective tertiary care. It comprises six hospitals with an aggregate 358 bed capacity, three primary care clinics, and a family medicine center. The range and level of services has improved in recent years and more patients seek treatment locally. The highest risk patients are transferred to the city of Abu Dhabi.



Figure 1. Al-Gharbia region, Abu Dhabi Emirate, UAE.

The region has experienced rapid population expansion, and the public hospitals network has a proactive approach to resourcing. At the time of this survey, there were 585 budgeted nursing positions; 435 were filled and the remainder in process. Staffing is sufficient in the context of current occupancy rates and support staffing levels are good. Most patients are young, healthy, working-age expatriate men living away from family. This population does not have the same demands as a more gender- and age-distributed population. There are fewer female patients who are nursed on separate floors. Cultural values do not permit male nurses to work in female wards.

UAE National Licensure for Nurses

Sound professional competence is of vital interest to the UAE, and professional licensure is the gatekeeper. All practicing health professionals are licensed, registered, and hold medical indemnity cover. UAE entry standards for expatriate nursing staff include a minimum nursing education of 3 years and at least 2 years of experience. Nurses must pass a written examination in English before licensing, which is necessary to obtain a work permit. Nursing registration and licensure are under the jurisdiction of the Health Authority of Abu Dhabi, Dubai Health Authority, and the UAE Ministry of Health (Brownie, Hunter, Aqtash, & Day, 2015; UAE Ministry of Health, Health Authority Abu Dhabi, & Dubai Health Authority, 2014).

Nursing Competency in Al-Gharbia

Competency requirements for nurses are determined by the Nation, the Emirate, and the employer, with input from the nursing profession (UAE Ministry of Health et al., 2014). The UAE National Qualifications Framework (competency model) has established requirements for assessing the competence of student nurses graduating in the UAE (UAE National Qualifications Authority, 2012); however, Al-Gharbia had only one Emirati nursing graduate at the time of this study. The public hospitals network nursing management (all registered nurses [RNs] with BScN and MScN degrees) has a well-established competency validation program based on nurse self-assessment of mandatory competencies and in-depth discussion with nurse managers as needed. This validation is separate from annual staff performance reviews (a human resources function). Competency validation is performed on an individual basis. The competency matrix defines the nature and scope of mandatory, general and unit-specific competencies and dictates the frequency of revalidation of a nurse's competency. Mandatory competencies are validated by a preceptor or senior nurse during orientation

and repeated yearly. General competencies are validated during probation and may be validated again during the course of employment. Unit-specific competencies are assessed in real nursing situations during probation and revalidated every second year. Competency assessment also includes examination at licensing, post in-service education, and competency validation by a preceptor in real-life situations. This competency validation program is designed to meet the region's need for generalist nurses with specialist capability.

There are significant challenges in maintaining optimal nursing competency. Although Al-Gharbia has low population and high cultural diversity, rapid population growth means the public hospitals network undergoes constant expansion. The network depends on recruitment and retention of skilled, experienced expatriate nurses. In larger hospitals, nurses provide care based on their area of specialty, whereas the scope of practice is extended in smaller hospitals, and nurses are multi-credentialed. Some clinical environments have low patient volumes and some medical events occur with low frequency (e.g., pregnancy and postpartum care). In response to low patient volumes (e.g., few postpartum care and high-risk specialist services), nurses are required to have competencies validated in more than one specialty. Around three quarters of nurses are cross-trained, and most have medical/surgical experience. In this remote, rural health system, nurses must undertake a range of duties and exercise sound autonomous judgement. Distance and desert conditions require that health facilities provide combined services, and nurses in this region must be flexible, responsive, and multitasked.

Literature Review

The definition, measurement, and validation of nursing competence remain controversial (Garside & Nhemachena, 2013). Different methodologies have been used to define the concept of competency and describe the factors that constitute it. Researchers have employed expert interviews, questionnaires, focus groups, and expert panels to survey and gain consensus, often using a Delphi process (E. Hurme, 2009; F. Hurme, 2007; Meretoja, Isoaho et al., 2004). Some nations have also drafted policy, guidelines, and standards and instituted regulatory and credentialing processes to ensure competency (Brownie et al., 2015).

Measurement of a nebulous construct is difficult, and there is no consensus regarding the best way to measure competence. Providing hard evidence of competency is challenging for all layers of the nursing workforce and is costly in time, effort, and funding. Managers and researchers are often left to design their own tools and use multiple methods to confirm the results of their competency assessment. In seven focus groups with RNs and

midwives across three hospitals in Australia, Fereday and Muir-Cochrane (2006) showed that the performance review process provided nurse clinicians opportunity to self-assess their ability and comfortability in performing their work and the chance to benchmark their own assessment against continual feedback collected from their peers and unfamiliar challenges. This self-declaration of competence has become a requirement for continuing nursing registration in parts of Australia, as part of a multimethod approach of defining competence by a set of outcome standards (Fereday & Muir-Cochrane, 2006; McCready, 2007; Nurses Board of South Australia, 2000).

As there is no set way of interpretation, the validation process may take many forms, including direct and indirect observation. Simulation drills are also used (Inman, 2016). Supervisors or preceptors (Butler et al., 2011) are deployed to observe skills and attitudes, question the knowledge base, coach, mentor, and ensure compliance. Checklists are used to aid these processes.

In the nursing profession, the frequency of use (FOU) of a competency and years of experience are positively correlated with self-assessed competency (Hengstberger-Sims et al., 2008; Meretoja, Isoaho, et al., 2004; O'Leary, 2012; Salonen, Kaunonen, Meretoja, & Tarkka, 2007). Additionally, self-assessment of competence is gaining traction as an important indicator of competence. The NCS (Meretoja, Isoaho, et al., 2004) is a prominent scale among various validated competence self-assessment tools, including the Six Dimensional Scale of Nursing Performance (Schwirian, 1978), European Healthcare Training and Accreditation Network (ETHAN) Questionnaire (Cowan, Wilson-Barnett, Norman, & Murrells, 2008), and the Competency Inventory for RNs (Liu, Yin, Ma, Lo, & Zeng, 2009). The NCS was developed, tested, and validated in a Finnish University Hospital (Meretoja, Isoaho, et al., 2004). The intended use was nurses and managers assessing nurse competence using self-assessment. The NCS has good internal consistency and good content validity; however, "concurrent and predictive validity testing with large samples was needed in the future" (Meretoja, Isoaho, et al., 2004, p. 131).

The NCS has been used in numerous studies by the Meretoja group in both large and small cohorts (Meretoja & Koponen, 2012; Numminen, Leino-Kilpi, Isoaho, & Meretoja, 2015). Underlying indicators of competent nursing practice have been validated (Meretoja, Eriksson, & Leino-Kilpi, 2002). The NCS has been tested in intensive care, high dependency, and emergency units (Salonen et al., 2007); different hospital environments (Meretoja, Leino-Kilpi, & Kaira, 2004); and with graduating student nurses (Kajander-Unkuri et al., 2014; Wangenstein, Johansson, Björkström, & Nordström, 2012). It has been translated from Finnish into English (Meretoja, Numminen, Isoaho, & Leino-Kilpi, 2015), Italian (Dellai, Mortari, & Meretoja,

2009), German (Müller, 2013), Norwegian (Wangensteen et al., 2012; Wangensteen, Johansson, & Nordström, 2015), and Lithuanian (Istomina et al., 2011) and tested in Australia (Hengstberger-Sims et al., 2008), Iran (Bahreini, Shahamat, Hayatdavoudi, & Mirzaei, 2011), Ireland (McCarthy, Cornally, O'Mahoney, White, & Weathers, 2013), and Switzerland (Müller, 2013). Supplementary questions have been added (McCarthy et al., 2013), and it has inspired the development of new tools (Donilon, 2013).

The construct validity of the NCS has received criticism, with previous studies finding redundancy and that they could not confirm the seven category (across 73 competency items) model (Müller, 2013; Wangensteen et al., 2015). In a reanalysis of 2007 dissertation data, Wangensteen (2010) could not confirm the original NCS seven category model. Müller used confirmatory factor analysis and Rasch modeling with data generated in a 2007 study, and found a "reduced set" of six categories and 54 items that "removes redundancy among items, is free from item bias and constitutes six uni-dimensional scales" (Müller, 2013, p. 1410). Missing values are also a recurring feature with the NCS (Numminen et al., 2015; Wangensteen, 2010; Wangensteen et al., 2015), and the reasons for this phenomenon have not been addressed. Treatment for missing values has been rejection of responses with values under 50% or substitution with the mean.

The NCS has mainly been tested in large teaching hospitals in metropolitan settings, where nurses operate within distinct clinical environments and have predictable nursing responsibilities. Increased perceived competency has been associated with increased FOU of a competency item (Meretoja, Isoaho, et al., 2004; Meretoja & Leino-Kilpi, 2003; Numminen, Meretoja, Isoaho, & Leino-Kilpi, 2013). Nurses usually perceive their competency as "good" (Meretoja, Isoaho et al., 2004), and managers generally perceive their nurses to have higher competency than do nurses themselves (Numminen et al., 2015).

There is limited research into rural/remote nursing competencies, including measurement or assessment. The available literature suggests rural nurses are often required to be skilled and knowledgeable generalists, capable of providing patient-centric nursing regardless of clinical specialty (Hunsberger, Baumann, Blythe, & Crea, 2009; Montour, Baumann, Blythe, & Hunsberger, 2009). Lovelace (2015) also noted that rural nurses have a greater scope of practice than their urban counterparts and are generalists who "must have at least some specialty knowledge in multiple areas of nursing." Rural nurses must also know when to assess their patients, stabilize them, and transfer them safely to an advanced level of care for more specialized treatment (Lovelace, 2015, p 1). In remote areas, this can involve longer distances, limited ambulances, or other transport

and poor roads. Lovelace suggested that "independence, flexibility and a strong sense of adventure" (p. 1) are necessary characteristics for rural nurses (Lovelace, 2015). Resourcefulness and innovativeness are important, as nurses must deal with absence of economies of scale and sometimes poor access to resources (including Internet and library access). Rural nurses usually live and work in the same small communities, meaning they are visible and lack anonymity. In rural communities, a nurse's competence can even be judged on their children's behavior (Bushy, 2002). Personal characteristics, behaviors, and attitudes become more important in this context and understanding the community and cultural competencies are critical in rural and remote environments (Morgan & Reel, 2003).

In her doctoral dissertation, Hurme (2007) used a Delphi panel of rural nursing experts to identify rural nursing competencies most associated with critical access hospitals. She highlighted similarities and differences between urban and rural nursing competencies and identified competencies specific to the rural environment in a Louisiana population. These included the importance of possessing knowledge of and ability to assess patients across the life span and being certified in both pediatric advanced life support and advanced cardiac life support. However, there are no examples of the NCS having been tested in rural and remote settings.

Study Purpose

The primary purpose of the present study was to measure the self-assessed level of competence (SALC) among nurses working in the Al-Gharbia public hospitals network and explore factors associated with nurses' self-perceived competency. The secondary purpose was to use a popular, validated tool that permitted the results to be compared with those from other countries to help identify areas for future professional development.

Research Questions

The research questions for this study were (a) What is the overall level of self-assessed competence? (b) How do nurses rate themselves in specifically defined areas of competence? (c) What differences in self-assessed competency occur between clinical areas and facilities? (d) What variables are associated with nurses' perception of their competence?

Method

Design

A cross-sectional study design was used to allow the public hospitals network to determine the current level

and nature of competence of the region's nursing workforce. A literature review identified several scales that could have been used for the survey (Cowan et al., 2008; Gardner, Smyth, Renison, Cann, & Vicary, 2012; Lakanmaa et al., 2014; Liu et al., 2009). Permission to use the Clinical Competence Questionnaire was initially obtained (Liou & Cheng, 2014). However, the present research team selected the English version of the NCS for two main reasons. First, the NCS has been used in many studies worldwide, including several European countries and Iran in the Middle East, and in a number of languages. In those studies, the NCS appeared to be stable and had good Cronbach's α values compared with other scales. Secondly, the NCS was the best fit for this study, as it focuses on (a) self-perceived competency and (b) FOU of the competencies. These two areas were the focus of competency concern for the nursing leadership in Al-Gharbia. As previously mentioned, nurses in smaller facilities (critical access facilities) often do not use some of their competencies for periods of time. This may negatively affect nurses' perception of being competent as well as their ability to actually perform certain skills safely. The NCS was regarded as the best tool to capture these two areas.

Use of the NCS allows comparison of the present results with those from other countries, benchmarking Al-Gharbia's expatriate nursing workforce competency in an international context. The NCS has not previously been validated in a small hospital context, in rural or remote conditions, in an expatriate or multireadentialed workforce, and in the Arab world or across an entire nursing workforce. Although this context makes Al-Gharbia unique, individual factors are experienced internationally. The development of the NCS was guided by Benner's stages of clinical competence framework. This suggests that to acquire and develop clinical skills and competence, nurses should pass through five levels of proficiency: novice, advanced beginner, competent, proficient, and expert (Benner, 1984).

Sample

This study used a purposive sample of 435 RNs, which was the public hospitals network entire nursing workforce. This permitted maximum population validity within the network's nursing workforce. Inclusion criteria were RNs who had worked in the network for a minimum of 3 months and who provided direct nursing care. Newly hired nurses (those who had worked in the network for less than 3 months), practical nurses (UAE Ministry of Health et al., 2014), and those working entirely in administrative positions were excluded. This is consistent with approved NCS methodology.

Ethical Considerations

This study was approved by the Institutional Research Ethics Committee of the public hospitals network. Copyright permission to implement the NCS was obtained from the publishers, and NCS methodology was followed as required (Meretoja, Isoaho et al., 2004). Informed consent was obtained from all participating nurses. At the beginning of the questionnaire, respondents placed a checkmark next to a statement indicating their agreement to join the study and complete the survey. Participation was voluntary and no incentives were offered. Respondents were guaranteed anonymity. The survey was administered online via Google Forms and did not include identifying material.

Survey Instrument

Permission to use the NCS was contingent on using the complete scale. No modifications to the scale (Meretoja, Isoaho et al., 2004) were made. The NCS measures self-perceived competency in 73 nursing items grouped into seven competency categories (areas of competence). The NCS uses a Visual Analogue Scale (VAS) with a range from 0 to 100. Respondents rate their level of competence by placing a mark on a line with 0 at one end and 100 at the other. The VAS is suited to a paper environment and is not able to be used in Google Forms. Instead, respondents were asked to rate their competency out of 100. A 4-point Likert-type scale is used to measure the FOU of each NCS competency item.

Competence measure. NCS competency items are grouped into seven categories: helping role (seven items), teaching-coaching (16 items), diagnostic functions (seven items), managing situations (eight items), therapeutic interventions (10 items), ensuring quality (six items), and work role (19 items; Meretoja, Isoaho et al., 2004). For each item, respondents were asked to place a number from 0 to 100 in the designated box, with 0 being very low and 100 very high. This measure reflects the SALC. "Item SALC" represents the SALC for each item. "Category SALC" means the average of Item SALCs within that category, and "Overall SALC" refers to the average of the seven Category SALCs. These SALC measures can be calculated for a respondent, for sub-groups of respondents or for all respondents. The term "SALC" is used throughout this article to introduce rigor to terminology and avoid confusion with actual competence. For analysis, the NCS methodology breaks the SALC measure into four overlapping integer groups: 0 to 25 is low, 25 to 50 quite good, 50 to 75 good, and 75 to 100 very good. In the present sample, this was modified slightly for clarity, with 0 to 24, 25 to 49, 50 to 74, and 75 to 100 as the integer ranges for the groups.

Frequency of use. A 4-point Likert-type scale (0 = *not applicable in my work*, 1 = *used very seldom*, 2 = *used occasionally*, 3 = *used very often in my work*) was used to capture the FOU of each item (Meretoja, Isoaho et al., 2004). Respondents were asked to indicate “the frequency with which this item is actually used in clinical practice.”

Data Collection

An explanatory e-mail was sent to all RNs in the public hospitals network inviting participation in the survey. The NCS was coded in Google Forms and made available for completion online between May 26 and July 16, 2015. Nurses were then e-mailed a link to the website where the survey could be completed. Two reminder e-mails were sent.

Statistical Analysis

Statistical analyses were performed with the R package (R x64 3.2.2). Microsoft Excel was used for summary statistics and graphics.

Categorical variables. Demographic variables included gender, age-group, highest level of education, marital status, ethnicity/nationality, total years of experience, years of experience in the public hospitals network, current position, facility, and clinical area. These were collected as categorical variables. Intervals between points on the FOU scale are not equal and therefore cannot be interpreted as numerical values. These data were captured, stored as text, and analyzed as categorical variables.

Missing values. Respondents with 50% or more missing values were excluded, consistent with NCS methodology. If a respondent was missing SALC for every item in a category, their Category SALC was recorded as “not available.” Otherwise, their Category SALC was the average of the available Item SALC. Overall SALC was calculated as the average of the available Category SALC. If a variable was used in an analysis and a respondent had a missing value for that variable, that respondent was excluded from the analysis. For the multiple linear models for each item, only respondents with Item SALC, Item FOU, and all demographics were included.

Merged responses. As some response categories had small counts, similar demographic groups were merged for analysis. For “age-group,” nurses aged ≤ 25 years and 26–35 years were merged. For “marital status,” separated and divorced nurses were merged. For “clinical area,” merging occurred between critical care and operating room, medical/surgical and managerial positions

(managerial positions were nurses who worked in senior charge nurse capacity and had a medical/surgical background), outpatients and emergency, and pediatrics and obstetrics. The four nonhospital facilities were merged. Decisions regarding merging clinical area response categories were made by the lead researcher, a nurse, and proper consideration of Al-Gharbia conditions.

Aggregated data. When an analysis found small numbers of responses in cells, data were aggregated to increase the strength of statistical analysis, where the test required the expected values of each cell to be ≥ 5 . This occurred in the pediatrics/obstetrics clinical area and in the first-two FOU levels.

Statistical tests. Statistical tests used to assess the significance of relationships included: analysis of variance (ANOVA) to test the association of individual demographics with SALC; the Kruskal–Wallis test, a non-parametric test for differences in median where the numbers of responses were small and not normally distributed; the χ^2 test of independence where two or more categorical variables were included; Cronbach’s α to test internal consistency; and Tukey’s multiple comparisons of means for post hoc analysis of differences.

Because FOU only occurs at the item level, testing the association of FOU with SALC required a multiple linear model for each of the 73 items, with FOU and demographic variables as independent variables and item SALC as the dependent variable. The significance of models was tested with regression ANOVA, and an independent variable was deemed to be significant if at least 1 of its levels was $p < .01$.

Results

In total, 227 responses were received from all network facilities and clinical areas (response rate 52.2%). After excluding responses with 50% or more missing values, 189 responses were analyzed, representing 43.4% of the nurse workforce.

Respondent Characteristics

Participating RNs were predominantly female (73.5%), married (76.7%), and aged under 45 years (78.3%; Table 1). The workforce was well qualified, with the majority holding a bachelor’s degree in nursing (69%) or higher (5.3%). Just 16.9% held recognized diploma qualifications, including nurses from countries with qualifications that do not easily equate (e.g., Tunisia and Lebanon) but who had achieved professional licensing and registration in Abu Dhabi. RNs had substantial nursing experience; only 7.9% had less than 5 years of

Table 1. Respondent Characteristics.

	<i>n</i>	Percentage ^a
Total	189	100.0
Gender		
Female	139	73.5
Male	36	19.0
Missing	14	7.4
Age-group		
35 Years or less	78	41.3
36–45 Years	70	37.0
Over 45 years	26	13.8
Missing	15	7.9
Highest level of education		
Diploma	32	16.9
Bachelor's degree	131	69.3
Postgraduate degree	10	5.3
Missing	16	8.5
Marital status		
Divorced/separated	4	2.1
Married	145	76.7
Single	25	13.2
Missing	15	7.9
Ethnicity/nationality		
Filipino	65	34.4
Indian	60	31.7
Middle Eastern (Jordan, Syrian, Lebanon, Palestine, Iraq . . .)	43	22.8
North African (Egypt, Tunisia, Morocco . . .)	7	3.7
Missing	14	7.4
Total years of work experience		
5 Years or less	15	7.9
6–10 Years	52	27.5
11–15 Years	38	20.1
16–20 Years	40	21.2
21–25 Years	15	7.9
More than 25 years	13	6.9
Missing	16	8.5
Years of experience in the hospitals network		
5 Years or less	67	35.4
6–10 Years	66	34.9
11–15 Years	23	12.2
16–20 Years	11	5.8
More than 20 years	7	3.7
Missing	15	7.9
Current position		
Charge nurse	33	17.5
Managerial position	15	7.9
Staff nurse	124	65.6
Missing	17	9.0

(continued)

Table 1. Continued.

	<i>n</i>	Percentage ^a
Facility		
Hospital 1	19	10.1
Hospital 2	28	14.8
Hospital 3	10	5.3
Hospital 4	67	35.4
Hospital 5	13	6.9
Hospital 6	22	11.6
Primary care clinics/home care	17	9.0
Missing	13	6.9
Current clinical area		
Critical care/operating room	42	22.2
Medical/surgical	55	29.1
Outpatient/emergency department	64	33.9
Pediatrics/obstetrics	14	7.4
Missing	14	7.4

^aPercentages may not add to 100% due to rounding.

experience and 6.9% had more than 25 years of experience. The main nationality groups were Filipino (34.4%), Indian (31.7%), and Middle Eastern (22.8%); few RNs came from North Africa (3.7%). The RN workforce was well settled with 64.6% having worked in Al-Gharbia for more than 5 years. Most RNs worked in a hospital environment and 9% worked in nonhospital facilities. All RNs had direct patient contact (65.6% staff nurses and 17.5% charge nurses), and all RNs in managerial positions (7.9%) had medical/surgical experience. There were few RNs in pediatrics/obstetrics (7.4%), reflecting the region's demographic characteristics. Nurses worked mainly in outpatient or emergency care (33.9%), medical or surgical units (29.1%), or critical care or the operating room (22.2%).

Overall SALC

The overall SALC for the study sample was very good at 84.3.

SALC in Defined Areas

Categories. All RNs rated themselves as very good (the highest competence level) in every category. They assessed themselves as most competent in management situations and least competent in ensuring quality and teaching-coaching situations (Figure 2). At the item level, mean SALC scores were mostly in the very good range and occasionally were in the good range (Table 2).

Reliability analysis within categories. Cronbach's α for the seven categories ranged from .88 for "ensuring quality"




Figure 2. Self-assessed level of competence by category.

to 0.94 for “work role” (Table 3). A maximum α value of .90 has been recommended (Streiner, 2003). Higher values indicate that there may be redundant variables and suggest the questionnaire can be shortened (Tavakol & Dennick, 2011). This will be considered in more detail if the questionnaire is reused.

Low SALC items. Items where individual RNs recorded SALC less than 50 (excluding nonresponses), were “coordinating student nurse mentoring in the unit” (19.6%), “supporting student nurses in attaining goals” (15.5%), and “taking student nurse’s level of skill acquisition into account in mentoring” (14.1%). These items also had the greatest nonresponse.

Differences in Competency Ratings in Nurses Working in Different Clinical Areas and Facilities

The four clinical areas did not show significant differences in overall SALC (ANOVA; $p = .177$). Although SALC means for all network facilities were in the broad classification of very good ($=75+$), there were significant differences (ANOVA; $p = .0001$), as shown in Figure 3. Post hoc analysis of these differences using Tukey’s multiple comparisons of means with a 95% family-wise confidence level showed there were no significant differences between pairs of hospitals, but that RNs from primary care clinics had significantly lower scores than four of the six hospitals. Nonhospital-based facilities had the lowest overall SALC (75.7), with ensuring quality (71.6), “work role” (72.5), and “therapeutic interventions” (73.0) having the lowest category SALCs.

Item FOU

For the majority of RNs, “not applicable” items were related to student nurses. These items were

“coordinating student nurse mentoring in the unit” (50.0%), “supporting student nurses in attaining goals” (49.7%), and “Taking student nurse’s level of skill acquisition into account in mentoring” (44.4%). The next item most frequently identified as not applicable was “mentoring novices and advanced beginners” (15.8%). Both “orientation for new nurses” and “guiding staff members” were not applicable for over 6.5% of RNs. All other items were not applicable for fewer than 6.5% of RNs (Table 2).

Items with the highest FOU were “documentation of patient care” (87.0%), “equipment maintenance” (86.4%), “individualized patient education” (84.5%), “planning patient care” (82.3%), “ability to recognize life threatening situations” (79.4%), “modifying care plans to suit needs” (78.1%), and “finding optimal timing for patient education” (78.1%).

Differences in FOU by Clinical Area

Fourteen items had significantly different FOU across clinical areas (χ^2 ; $p < .01$). Nurses in outpatient, emergency, critical care, and operating room environments reported the highest FOU for most competency items. Pediatrics/obstetrics had the highest FOU for “coaching others in duties within my responsibility area” and “professional identity serves as resource in nursing.” Medical/surgical RNs reported the lowest FOU for all significant items, except “coordinating multidisciplinary team’s nursing activities,” where pediatric/obstetric nurses had a lower FOU.

Variables That Influenced Nurses’ Perception of Their Competence

To determine the association between FOU and SALC in the context of demographic variables, regression models were produced for each item. All models were significant (ANOVA, highest p value was .0058). FOU was a significant variable for 62 of the 73 items. For 34 of these items, it was the only significant variable. Total years of experience was significant for 20 items, facility for seven items, ethnicity for six items, education for five items, years in the network for two items, and age-group for one item (Table 2). Gender, marital status, current position, and clinical category were not significant in any model. No variables were significant for seven items (Table 2).

For four items, FOU was not significant, but other variables were: “years of experience” was significant for “evaluating critically own philosophy in nursing”; years of experience and “years in the network” were significant for “developing documentation of patient care,” with over 15 years in the network having a negative association; years of experience was significant for

Table 2. SALC, FOU, and Significant Variables Associated With SALC.

Item	SALC			FOU		Linear model variables								
	n	Mean	SD	Rated < 50 (%)	n	“Not applicable” (%)	“Very often” (%)	F	A	Ed	Et	Yr	Yr N	Fa
Overall	189	84.3	11.2											
Helping role	189	84.5	10.7											
Planning patient care according to individual needs	185	87.8	10.5	0.0	186	0.0	82.3	*	*			*		
Supporting patients' coping strategies	187	85.0	13.6	1.6	185	0.5	69.2	*						
Evaluating critically own philosophy in nursing	184	82.3	14.6	1.6	184	0.5	50.5					*		
Modifying the care plan according to individual needs	185	89.0	10.3	0.0	183	0.0	78.1	*			*	*		
Utilizing nursing research findings in relationships with patients	181	77.0	18.7	3.9	182	1.6	32.4	*						
Developing the treatment culture of my unit	183	83.0	14.5	1.6	184	0.5	52.7	*			*	*		
Decision-making guided by ethical values	183	87.4	11.6	0.5	183	0.5	71.0	*						
Teaching-coaching	189	83.3	12.1											
Mapping out patient education needs carefully	183	85.8	15.4	2.2	182	1.1	74.2							
Finding optimal timing for patient education	185	85.6	13.8	1.6	183	0.0	78.1	*						
Mastering the content of patient education	187	85.9	12.5	1.6	184	0.5	73.9							
Providing individualized patient education	181	88.5	10.5	0.6	181	0.0	84.5	*						
Coordinating patient education	186	84.9	14.4	2.2	184	0.5	64.7	*						
Able to recognize family members' needs for guidance	187	84.5	13.6	1.6	186	0.0	53.2	*				*		*
Acting autonomously in guiding family members	182	82.3	15.3	3.8	181	1.1	47.0	*						
Taking student nurse's level of skill acquisition into account in mentoring	156	70.1	30.9	14.1	178	44.4	25.8	*				*		
Supporting student nurses in attaining goals	155	70.4	32.7	15.5	179	49.7	29.1	*						
Evaluating patient education outcome together with patient	187	84.0	15.9	2.7	184	0.5	59.8	*				*		

(continued)

Table 2. Continued.

Item	SALC			FOU			Linear model variables							
	n	Mean	SD	Rated < 50 (%)	n	"Not applicable" (%)	"Very often" (%)	F	A	Ed	Et	Yr	Yr N	Fa
Evaluating patient education outcomes with family	185	83.6	14.5	1.6	184	1.1	50.0	*	*	*				
Evaluating patient education outcome with care team	186	83.5	16.2	4.3	182	1.6	54.4	*						*
Taking active steps to maintain and improve my professional skills	186	87.8	13.6	1.6	183	0.5	76.5	*		*				
Developing patient education in my unit	186	85.4	14.5	1.6	184	0.5	69.6							
Developing orientation programs for new nurses in my unit	182	81.7	22.0	5.5	184	9.2	53.8	*						*
Coaching others in duties within my responsibility area	182	84.3	19.1	3.8	185	3.8	56.8	*						
Diagnostic functions	189	85.4	11.9											
Analyzing patients' well-being from many perspectives	184	84.8	13.4	1.6	184	1.1	65.2	*	*	*				
Able to identify patients' need for emotional support	187	86.2	14.1	1.1	188	1.6	69.1	*						*
Able to identify family members' need for emotional support	187	84.5	13.3	0.5	184	1.1	55.4	*						
Arranging expert help for patient when needed	187	86.2	14.8	1.6	186	2.7	65.1	*				*		
Coaching other staff members in patient observation skills	181	83.6	16.2	2.8	183	2.2	55.7	*						
Coaching other staff members in use of diagnostic equipment	183	83.9	16.8	2.7	182	3.3	56.6	*		*				
Developing documentation of patient care	185	88.8	11.6	0.5	184	0.5	87.0					*	*	
Managing situations	189	86.9	11.4											
Able to recognize situations posing a threat to life early	185	89.2	11.0	0.5	184	0.5	79.3							
Prioritizing my activities flexibly according to changing situations	183	89.1	11.0	0.5	183	1.1	75.4							
Acting appropriately in life-threatening situations	185	90.5	10.2	0.0	180	1.1	71.7							
Arranging debriefing sessions for the care team when needed	181	82.4	17.3	2.8	183	6.0	41.5	*						

(continued)

Table 2. Continued.

Item	SALC			FOU			Linear model variables							
	n	Mean	SD	Rated < 50 (%)	n	"Not applicable" (%)	"Very often" (%)	F	A	Ed	Et	Yr	Yr N	Fa
Coaching other team members in mastering rapidly changing situations	184	80.9	18.8	4.3	184	4.3	41.8	*						
Planning care consistently with resources available	187	85.5	15.0	2.7	185	0.5	64.3	*						
Keeping nursing care equipment in good condition	186	91.0	12.1	0.5	184	1.1	86.4	*						*
Promoting flexible team cooperation in rapidly changing situations	185	88.1	12.6	1.1	182	0.0	72.5	*						
Therapeutic interventions	189	84.0	12.3											
Planning own activities flexibly according to clinical situation	188	87.6	11.9	0.5	186	0.0	72.6	*						
Making decisions concerning patient care taking the particular situation into account	186	86.7	13.1	0.5	182	0.0	67.6	*						
Coordinating multidisciplinary team's nursing activities	187	85.5	14.0	0.5	183	3.3	62.8	*						
Coaching the care team in performance of nursing interventions	183	84.1	15.2	1.6	182	3.3	51.6	*				*		
Updating written guidelines for care	181	83.9	15.7	2.8	184	4.3	53.3	*						
Providing consultation for the care team	177	81.2	19.4	4.5	181	4.4	49.7	*						
Utilizing research findings in nursing interventions	181	78.5	18.8	3.3	181	3.3	30.9	*				*		
Evaluating systematically patient care outcomes	183	85.8	14.2	1.6	179	1.1	73.7	*						
Incorporating relevant knowledge to provide optimal care	181	87.1	12.1	0.6	179	0.6	76.0	*				*		
Contributing to further development of multidisciplinary clinical paths	180	80.4	17.7	3.3	174	2.9	36.8	*		*		*		*
Ensuring quality	183	82.0	13.5											
Committed to my organization's care philosophy	179	86.5	15.2	2.2	178	1.7	70.8	*				*		
Able to identify areas in patient care needing further development and research	181	81.6	15.5	1.7	180	1.7	39.4	*						

(continued)

Table 2. Continued.

Item	SALC				FOU		Linear model variables							
	n	Mean	SD	Rated < 50 (%)	n	“Not applicable” (%)	“Very often” (%)	F	A	Ed	Et	Yr	Yr N	Fa
Evaluating critically my unit's care philosophy	181	83.5	14.1	1.1	179	2.2	49.2	*			*	*		
Evaluating systematically patients' satisfaction with care	180	88.4	11.6	0.6	177	1.1	75.7							
Utilizing research findings in further development of patient care	179	79.9	18.7	3.4	179	2.2	38.5	*		*		*		
Making proposals concerning further development and research	181	73.9	21.2	7.2	177	4.5	20.9	*				*		
Work role	180	84.1	12.5											
Able to recognize colleagues' need for support and help	178	88.5	11.4	0.6	175	1.1	76.6							*
Aware of the limits of my own resources	178	89.0	11.9	0.6	175	0.0	73.1	*				*		
Professional identity serves as resource in nursing	178	85.7	14.1	1.7	179	1.7	60.9	*			*			
Acting responsibly in terms of limited financial resources	176	81.8	20.4	4.5	175	6.3	50.9	*						
Familiar with my organization's policy concerning division of labor and coordination of duties	176	85.8	14.6	1.1	174	0.6	71.3	*						
Coordinating student nurse mentoring in the unit	153	65.7	34.5	19.6	176	50.0	22.2	*					*	
Mentoring novices and advanced beginners	163	77.3	24.2	6.7	171	15.8	32.2	*						
Providing expertise for the care team	171	83.0	17.3	2.9	174	4.6	53.4	*						
Acting autonomously	172	84.4	17.3	2.9	170	2.4	64.1	*					*	
Guiding staff members to duties corresponding to their skill levels	173	84.0	18.3	3.5	175	8.0	58.3	*						
Incorporating new knowledge to optimize patient care	175	85.9	13.9	1.1	174	1.1	62.6	*						
Ensuring smooth flow of care in the unit by delegating tasks	173	86.2	13.9	1.2	171	5.3	66.7	*				*		
Taking care of myself in terms of not depleting my mental and physical resources	175	84.8	16.1	2.3	176	0.0	63.6	*						
Utilizing information technology in my work	175	88.1	12.7	0.6	169	0.0	74.6	*						

(continued)

Table 2. Continued.

Item	SALC			FOU		Linear model variables								
	n	Mean	SD	Rated < 50 (%)	n	"Not applicable" (%)	"Very often" (%)	F	A	Ed	Et	Yr	Yr N	Fa
Coordinating patients' overall care	176	88.0	13.7	1.1	174	1.1	77.6	*						
Orchestrating the whole situation when needed	169	83.7	16.8	2.4	172	3.5	51.7	*						
Giving feedback to colleagues in a constructive way	173	86.5	14.0	1.2	170	0.0	60.6	*						
Developing patient care in multidisciplinary teams	175	84.8	16.3	2.9	175	0.6	62.3	*				*		
Developing work environment	175	86.3	14.6	2.3	176	0.0	66.5	*						

Note. SALC = self-assessed level of competence; FOU = frequency of use; n/a = not applicable; SD = standard deviation; F = FOU; A = age; Ed = education; Et = ethnicity; Yr = total years of experience; Yr N = years in the network; Fa = facility.

*Percentage of respondents who rated themselves < 50 on the Visual Analogue Scale.

*At least one level of the variable was significant at $p < .01$.

Table 3. Reliability of Categories in the Scale.

Category	Cronbach's α
Helping role	.89
Teaching-coaching	.92
Diagnostic functions	.92
Managing situations	.92
Therapeutic interventions	.93
Ensuring quality	.88
Work role	.94




Figure 3. Boxplot of overall self-assessed level of competence score by facility.

“incorporating relevant knowledge to provide optimal care”; and facility was significant for “able to recognize colleagues’ need for support and help,” with hospital RNs scoring higher than those in clinics.

Discussion

Overall SALC

The Overall SALC in Al-Gharbia was 84.3 or very good (Table 2). This was high compared with almost all other studies of a reasonable sample size that used the NCS. It was most similar to an Iranian study, in which RNs with BScN qualification working in a range of areas achieved 87.03 in “Type 1 University Hospital” and 71.07 in “Type 2 University Hospital” (terms not defined; Bahreini et al., 2011).

In Finland University Hospital(s), results were 56.1 for RNs with limited work experience (Salonen et al., 2007), 63.7 for RNs in medical/surgical environments (Meretoja, Isoaho et al., 2004; Meretoja, Leino-Kilpi, et al., 2004), 63.9 for 81 staff nurses working in a variety of environments (Meretoja & Leino-Kilpi, 2003), and 66.0 for RNs with broad range of experience in four clinical environments (medical, surgical, pediatric/obstetric, and psychiatric; Meretoja et al., 2015). Previous studies showed Overall SALC was 59.0 for those aged 20–29 years, 65.0 for those aged 30–39 years, 70.1 for

those aged 40 years and over (Meretoja et al., 2015), and 66.7 for graduating student nurses in their final clinical placement (Kajander-Unkuri et al., 2014).

Understanding self-assessed competency in nurses in Al-Gharbia. The high overall SALC in this study population may be understood in the context of the nursing workforce in Al-Gharbia. All expatriate nurses undergo rigorous testing as part of the due diligence for their appointment. It is Abu Dhabi's practice to screen all nurses for qualifications, experience, character, and fluency in English before licensing, registration, immigration, and appointment. These are experienced nurses and Meretoja et al. (2015) found self-assessed competence increased with age, with those aged 40 years and over having higher ratings than those aged 20–29 years by a difference of 11.1 percentage points. Their rating for younger staff (59.0 for 20- to 29-year olds) was consistent with that of RNs with limited work experience (56.1) found by Salonen et al. (2007).

In Abu Dhabi, nurses are interviewed before employment and deployment. There are strong motivators for nurses to maintain and develop competencies. Residence in the UAE is dependent upon employment, and work permits are renewed at the end of a 3-year period. Expatriates may not become citizens and must leave the UAE after employment ceases. Most expatriates send remittances home to support families, meaning their sense of dedication to work is heightened by a desire to remain in the UAE and earn money to send home. The Al-Gharbia nursing workforce is stable, with most nurses staying more than 5 years.

A culture of competence and professionalism exists among nurses in Al-Gharbia, who access continuing practice development opportunities as offered. Nursing management provides extensive orientation, training, and testing to ensure continued competence. Management also screens nurses entering the region with clinical scenarios and has an established program of continuing education and performance appraisals. Nurses in Al-Gharbia are accustomed to assessment post in-service training.

In Al-Gharbia, as with many rural or remote settings, nurses can see that their presence makes a difference to health care. Nurses invariably have ready access to supplies and the latest equipment necessary to do their job. Hospitals and clinics are modern, clean, and well supported by cleaning and administration staff. With a younger, generally healthy population, nurses may also have less exposure to conditions that can create "burn out."

Self-Assessed Competence in Defined Areas

In this study, there were no significant differences by clinical area, meaning the overall SALC (84.3) can be

compared with that in studies involving a range of clinical areas (63.7; Meretoja et al., 2015) and medical/surgical areas (66.0; Meretoja, Isoaho et al., 2004). If permission to use the scale had allowed flexibility for modification, redundant items could have been removed. Student-related competencies are not required in Al-Gharbia, where all expatriate nurses are educated and experienced. Therefore, it is unsurprising that participating nurses were least confident in items relating to students (SALC: 65.7–70.4; Table 2). The poorest Item SALC was for coordinating student nurse mentoring in the unit, with 50% of RNs indicating it was not applicable. However, it is interesting that around a quarter of responses indicated the FOU for student-related items was "very often." These items might have been misinterpreted to mean new staff. RNs with more regional experience were involved in orienting and coaching new staff on probation and other staff in tasks such as the "use of diagnostic equipment" and "mastering rapidly changing situations," where SALC was reported as very good.

In addition, items relating to family were expected to have low FOU (except in female wards). However, around half of the RNs indicated FOU was very often for these items. Nurses also rated their overall SALC in these items as very good which might relate to prior experience. These results indicate there may be issues with the practical application of the NCS and may also point to reliability and validity issues. These concerns are discussed in the limitations section and will be elaborated in a future article. RNs in this study sample indicated comparative weakness (still very good) in "utilizing research findings in relationships with patients" (77.0), "Utilizing research findings in nursing interventions" (78.5), "utilizing research findings in further development of patient care" (79.9), and "making proposals concerning further research and development" (72.9). These were also lower frequency items, with 31–39% of respondents rating FOU as very often.

Limitations

Data collection. The decision to collect age, overall years of experience, and years of experience in a categorical format reduced the ability to be definitive about the strength of the association of these variables with SALC.

Face validity. The NCS has been tested for face validity as a measure of self-assessed competence and stands as a subjective measure. This study did not assume that it represented other measures of competence, such as objective tests or supervisor assessment.

Construct validity. The construct validity of the NCS has received criticism for redundancy and nonconfirmation

of the seven category model (Müller, 2013; Wangensteen et al., 2015). In this study, the high Cronbach's α for the categories also suggested a level of inbuilt redundancy. In addition, several NCS items were not relevant for the present study population, specifically those related to students and research. Items about student supervision and management are found in the NCS "teaching-coaching" and "work role" categories. Items relating to research are in the "helping role," "therapeutic interventions," and "ensuring quality" categories.

Bias. There remains a possibility of a compliance bias if participating nurses were trying to please nursing management; however, response rates were similar to other studies that used the NCS. If Al-Gharbia nurses felt under duress to respond, a much higher response rate and lower missing values would be expected.

Missing values. As for other NCS studies, missing values were a feature of survey returns (Kajander-Unkuri et al., 2014; Meretoja, Isoaho et al., 2004; Meretoja & Leino-Kilpi, 2003; Meretoja, Leino-Kilpi, et al., 2004; Numminen et al., 2013; Salonen et al., 2007). This adds complexity to the analyses and reduced instrument practicality and user friendliness.

Practicality of the NCS

There is no clear published methodology for online implementation of the NCS (Meretoja, Isoaho et al., 2004) and no definition, scope, or examples given for any item to aid nurses in understanding item statements. The lack of clear instructional methodology required to implement the NCS as a management tool suggests that the prime audience for NCS is researchers of nursing competency rather than nursing managers or nurses themselves.

Implications for Nursing Management

This study provided preliminary benchmarking material, affirmation for nursing management regarding the impact of training and competence screening policies and insight as to where to focus management attention. The study also highlighted that although FOU is clearly significant, it is possible for nurses operating in low-frequency event environments to achieve high self-perceived competence.

When a health system takes proactive responsibility for encouraging actual nursing competence (e.g., prescreening nurses, assessing and reassessing applied knowledge and skills, and targeting in-service training to address the known challenges of the population and catchment area), nurses' professional self-esteem is likely to be high. Given that Al-Gharbia remains reliant

on an expatriate nursing workforce, the message to management is that their investment in ensuring the competency of expatriate nurses is as vital as investment in modern equipment and infrastructure, and this investment is paying off. Such positive management behavior may contribute to nurse migration to this remote rural region and encourage retention and workforce stability.

Conclusions

The NCS was implemented to measure the SALC among RNs working in the public hospitals network in Al-Gharbia, UAE. This population produced a higher overall SALC score (84.3) than found in most previous studies. Public hospitals network nurses rate their professional competence as very good, and FOU is the variable most associated with SALC; results consistent with a stable nursing labor force. Al-Gharbia's remote, rural, and desert environment appears to encourage resilience and adaptability.

The public hospitals network management encourages continual revalidation of the knowledge, skills, and competencies of its nurses. In this population, NCS items relating to students might have been open to interpretation and the associated FOU improbable. Even so, the reported SALC might be valid. As UAE-trained graduates and students are introduced into Al-Gharbia's nursing workforce, such items will become more relevant. The present exploration of factors associated with nurses' self-perceived competency found that the FOU of a competency item and "years of experience" had the most influence on self-assessment of competency.

This experience of using the NCS and the corresponding analyses provides insights for future managerial guidance. If implementing the NCS again, the present authors would consider scoping or explaining each item, removing irrelevant items, and developing strategies to address and minimize missing data. Collecting numerical data rather than categorical data for variables such as age, years of experience, and FOU would also enhance the analyses.

While use of the NCS as a management tool remains to be fully tested, the NCS provides guidance for directing future effort. Nursing management can investigate the adequacy of documentation, application of research into practice, and heightened use of existing data to support the sustainability of sound nursing outcomes. The alignment of the NCS with the existing performance-oriented competency matrix can also be examined.

Author Contributions

All the authors have contributed equally to this work.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

- Bahreini, M., Shahamat, S., Hayatdavoudi, P., & Mirzaei, M. (2011). Comparison of the clinical competence of nurses working in two university hospitals in Iran. *Nursing & Health Sciences, 13*(3), 282–288. doi:10.1111/j.1442-2018.2011.00611.x
- Benner, P. (1984). *From novice to expert: Excellence and power in clinical nursing*. Menlo Park, CA: Addison-Wesley.
- Brownie, S., Hunter, L., Aqdash, S., & Day, G. (2015). Establishing policy foundations and regulatory systems to enhance nursing practice in the United Arab Emirates. *Policy, Politics and Nursing Practice, 16*(1–2), 38–50. doi:10.1177/1527154415583396
- Bushy, A. (2002). International perspectives on rural nursing: Australia, Canada, USA. *Australian Journal of Rural Health, 10*, 104–111.
- Butler, M., Cassidy, I., Quillinan, B., Fahy, A., Bradshaw, C., Touhy, D., . . . Tierney, C. (2011). Competency assessment methods—Tool and processes: A survey of nurse preceptors in Ireland. *Nurse Education in Practice, 11*, 298–303. doi:10.1016/j.nepr.2011.01.006
- Cowan, D., Wilson-Barnett, D. J., Norman, I., & Murrells, T. (2008). Measuring nursing competence: Development of a self-assessment tool for general nurses across Europe. *International Journal of Nursing Studies, 45*(6), 902–913. doi:10.1016/j.ijnurstu.2007.03.004
- Dellai, M., Mortari, L., & Meretoja, R. (2009). Self-assessment of nursing competencies—Validation of the Finnish NCS instrument with Italian nurses. *Scandinavian Journal of Caring Sciences, 23*(4), 783–791. doi:10.1111/j.1471-6712.2008.00665.x
- Donilon, D. (2013). *Nurses perceptions of their competence in managing patient situations in acute care* (Open Access Dissertations, Paper 13). University of Rhode Island, USA. doi:10.1111/jocn.12057
- Fereday, J., & Muir-Cochrane, E. (2006). The role of performance feedback in the self-assessment of competence: A research study with nursing clinicians. *Collegian, 13*(1), 10–15. doi:10.1016/S1322-7696(08)60511-9
- Gardner, A., Smyth, W., Renison, B., Cann, T., & Vicary, M. (2012). Supporting rural and remote area nurses to utilise and conduct research: An intervention study. *Collegian, 19*, 97–105.
- Garside, J., & Nhemachena, J. (2013). A concept analysis of competence and its transition in nursing. *Nurse Education Today, 33*(5), 541–545. doi:10.1016/j.nedt.2011.12.007
- Hengstberger-Sims, C., Cowin, L., Eagar, S., Gregory, L., Andrew, S., & Rolley, J. (2008). Relating new graduate nurse competence to frequency of use. *Collegian, 15*, 69–76. doi:10.1016/j.colegn.2008.02.003
- Hunsberger, M., Baumann, A., Blythe, J., & Crea, M. (2009). Sustaining the rural workforce: Nursing perspectives on worklife challenges. *Journal of Rural Health, 25*(1), 17–25. doi:10.1111/j.1748-0361.2009.00194.x
- Hurme, F. (2007). *Competencies for rural nursing practice: A dissertation*. Louisiana State University and Agricultural and Mechanical College.
- Hurme, E. (2009). Competencies for nursing practice in a rural critical access hospital. *Online Journal of Rural Nursing and Health Care, 9*(2), 67–81.
- Inman, F. (2016). Using simulation drills to improve competency and team skill for rural perinatal nurses. *Journal of Obstetric, Gynecological and Neonatal Nursing, 45*, s9–s10.
- Istomina, N., Suominen, T., Razbadauskas, A., Martinkėnas, A., Meretoja, R., & Leino-Kilpi, H. (2011). Competence of nurses and factors associated with it. *Medicina (Kaunas), 47*(4), 230–237.
- Kajander-Unkuri, S., Meretoja, R., Katajisto, J., Saarikoski, M., Salminen, L., Suhonen, R., . . . Leino-Kilpi, H. (2014). Self-assessed level of competence of graduating nursing students and factors related to it. *Nurse Education Today, 34*(5), 795–801. doi:10.1016/j.nedt.2013.08.009
- Lakanmaa, R.-L., Suominen, T., Perttilä, J., Ritmala-Castrén, M., Vahlberg, T., & Leino-Kilpi, H. (2014). Basic competence in intensive and critical care nursing: Development and psychometric testing of a competence scale. *Journal of Clinical Nursing, 23*(5/6), 799–810. doi:10.1111/jocn.12057
- Liou, S.-R., & Cheng, C.-Y. (2014). Developing and validating the clinical competence questionnaire: A self-assessment instrument for upcoming baccalaureate nursing graduates. *Journal of Nursing Education and Practice, 4*(2), 56.
- Liu, M., Yin, L., Ma, E., Lo, S., & Zeng, L. (2009). Competency inventory for registered nurses in Macao: Instrument validation. *Journal of Advanced Nursing, 65*(4), 893–900. doi:10.1111/j.1365-2648.2008.04936.x
- Lovelace, D. (2015). Rural nursing: Nurses enjoy greater scope of practice when serving less populated communities. *Advance for Nurses*. Retrieved from <http://nursing.advanceweb.com/Features/Articles/Rural-Nursing.aspx#AnchorComment>
- McCarthy, G., Cornally, N., O'Mahoney, C., White, G., & Weathers, E. (2013). Emergency nurses: Procedures performed and competence in practice. *International Emergency Nursing, 21*(1), 50–57. doi:10.1016/j.ienj.2012.01.003
- McCready, T. (2007). Portfolios and the assessment of competence in nursing: A literature review. *International Journal of Nursing Studies, 44*(1), 143–151. doi:10.1016/j.ijnurstu.2006.01.013
- Meretoja, R., Eriksson, E., & Leino-Kilpi, H. (2002). Indicators for competent nursing practice. *Journal of Nursing Management, 10*(2), 95–102. doi:10.1046/j.0966-0429.2001.00299.x
- Meretoja, R., Isoaho, H., & Leino-Kilpi, H. (2004). Nurse competence scale: Development and psychometric testing. *Journal of Advanced Nursing, 47*(2), 124–133. doi:10.1111/j.1365-2648.2004.03071.x

- Meretoja, R., & Koponen, L. (2012). A systematic model to compare nurses' optimal and actual competencies in the clinical setting. *Journal of Advanced Nursing*, 68(2), 414–422. doi:10.1111/j.1365-2648.2011.05754.x
- Meretoja, R., & Leino-Kilpi, H. (2003). Comparison of competence assessments made by nurse managers and practising nurses. *Journal of Nursing Management*, 11(6), 404–409. doi:10.1046/j.1365-2834.2003.00413.x
- Meretoja, R., Leino-Kilpi, H., & Kaira, A. (2004). Comparison of nurse competence in different hospital work environments. *Journal of Nursing Management*, 12(5), 329–336. doi:10.1111/j.1365-2834.2004.00422.x
- Meretoja, R., Numminen, O., Isoaho, H., & Leino-Kilpi, H. (2015). Nurse competence between three generational nurse cohorts: A cross-sectional study. *International Journal of Nursing Practice*, 21, 350–358. doi:10.1111/ijn.12297
- Montour, A., Baumann, A., Blythe, J., & Hunsberger, M. (2009). The changing nature of nursing work in rural and small community hospitals. *Rural and Remote Health*, 9(1), 1089–1089.
- Morgan, L., & Reel, S. (2003). Developing cultural competence in rural nursing. *Online Journal of Rural Nursing and Health Care*, 3(1 Spring), 28–37.
- Müller, M. (2013). Nursing competence: Psychometric evaluation using Rasch modelling. *Journal of Advanced Nursing*, 69(6), 1410–1417. doi:10.1111/jan.12009
- Numminen, O., Leino-Kilpi, H., Isoaho, H., & Meretoja, R. (2015). Congruence between nurse managers' and nurses' competence assessments: A correlation study. *Journal of Nursing Education and Practice*, 5(1), 142–150. doi:10.5430/jnep.v5n1p142
- Numminen, O., Meretoja, R., Isoaho, H., & Leino-Kilpi, H. (2013). Professional competence of practising nurses. *Journal of Clinical Nursing*, 22(9/10), 1411–1423. doi:10.1111/j.1365-2702.2012.04334.x
- Nurses Board of South Australia. (2000). *Your practice, continuing competence and self-regulation*. Adelaide, Australia: Author.
- O'Leary, J. (2012). Comparison of self-assessed competence and experience among critical care nurses. *Journal of Nursing Management*, 20, 607–614.
- Salonen, A., Kaunonen, M., Meretoja, R., & Tarkka, M. (2007). Competence profiles of recently registered nurses working in intensive and emergency settings. *Journal of Nursing Management*, 15(8), 792–800. doi:10.1111/j.1365-2934.2007.00768.x
- Schwirian, P. (1978). Evaluating the performance of nurses: A multidimensional approach. *Nursing Research*, 27, 347–351.
- Statistics Centre Abu Dhabi. (2015). *Statistical yearbook of Abu Dhabi 2015*. Retrieved from <http://www.scad.ae/en/Pages/ThemesReleases.aspx?ThemeID=4>
- Streiner, D. (2003). Starting at the beginning: An introduction to coefficient alpha and internal consistency. *Journal of Personality Assessment*, 80, 99–103. doi:10.1207/S15327752JPA8001_18
- Tavakol, M., & Dennick, R. (2011). Making sense of Cronbach's alpha. *International Journal of Medical Education*, 2, 53–55. doi:10.5116/ijme.4dfb.8dfd
- United Arab Emirates Ministry of Health, Health Authority Abu Dhabi, & Dubai Health Authority. (2014). *Professional Qualification Requirements Chapter 3: Nurses and Midwives*.
- United Arab Emirates Ministry of Health, Health Authority Abu Dhabi, & Dubai Health Authority. (2014). *Healthcare professionals qualification requirements*. Retrieved from [http://www.moh.gov.ae/en/Services/Documents/Evaluation/\(PQR\)%202014-1.pdf](http://www.moh.gov.ae/en/Services/Documents/Evaluation/(PQR)%202014-1.pdf) (accessed 10 August 2016)
- Wangensteen, S. (2010). *Newly graduated nurses' perception of competence, critical thinking and research utilization*. Karlstad, Sweden: Karlstad University.
- Wangensteen, S., Johansson, I., Björkström, M., & Nordström, G. (2012). Newly graduated nurses' perception of competence and possible predictors: A cross-sectional survey. *Journal of Professional Nursing*, 28(3), 170–181. doi:10.1016/j.profnurs.2011.11.014
- Wangensteen, S., Johansson, I., & Nordström, G. (2015). Nurse competence scale—Psychometric testing in a Norwegian context. *Nurse Education in Practice*, 15(1), 22–29. doi:10.1016/j.nepr.2014.11.007.